

UNIVERSITY OF TORONTO



3 1761 00979550 1

"The greatest and most important Literary Enterprise of the day."—CRITIC

CABINET EDITION OF THE
ENCYCLOPÆDIA METROPOLITAN

Thoroughly Revised and abundantly Illustrated,

Publishing in Volumes, each containing a Complete Treatise, Crown Octavo, Cloth.

Volumes now Ready.

Vol. I., price 2s., cloth lettered,
INTRODUCTORY DISSERTATION ON THE SCIENCE OF METHOD.
SAMUEL TAYLOR COLERIDGE. With a Synopsis. Sixth Edition.

Vol. II., price 5s., cloth lettered,
UNIVERSAL GRAMMAR, or the Pure Science of Language. By Sir J
STODDART, Knt., LL.D. Third Edition, Revised.
"The soundest and most philosophical treatise of the kind in the English language."—*North British Rev*

Vol. III., price 3s., cloth lettered,
LOGIC. By the Most Rev. RICHARD WHATELY, D.D., Archbishop of Dub
The Original Edition, with Synopsis and Index by the Editor.

Vol. IV., price 3s. 6d., cloth lettered,
RHETORIC. By the Most Rev. RICHARD WHATELY, D.D., Archbishop
Dublin. The Original Edition, with Synopsis and Index by the Editor.

Vol. V., price 6s., cloth lettered,
HISTORY OF THE CHRISTIAN CHURCH in the First Century. By
Right Rev. SAMUEL HINDS, D.D., Bishop of Norwich, and the Rev. J. H. NEWMAN, I
Third Edition, Revised.
"Erudite research is here combined with independent thought."—*Scottish Guardian*.

Vol. VI., price 4s., cloth lettered,
POLITICAL ECONOMY. By NASSAU WILLIAM SENIOR, Esq., late Profes
of Political Economy in the University of Oxford. Third Edition.

Vol. VII., price 2s. 6d., cloth lettered,
HISTORY OF THE JEWS, from the time of Alexander the Great to the l
struction of Jerusalem by Titus, A.M. 3595, B.C. 409 to A.D. 70. By the Venerable Archdea
HALE, A.M., Master of the Charter-house. Fourth Edition.

Vol. VIII., price 6s., cloth lettered,
SACRED HISTORY AND BIOGRAPHY, from the Antediluvian Period to
Time of the Prophet Malachi, A.M. 1 to A.M. 3607, B.C. 397. Edited by the Rev. F. A. C
D.D., LL.D.
"Pleasing and good, and worthy of general attention."—*United Presbyterian Magazine*.

Vol. IX., price 7s. 6d., cloth lettered,
HISTORY OF GREEK LITERATURE. By the Hon. Sir T. N. TALFOUR
D.C.L., the Right Rev. C. J. BLOMFIELD, D.D., Bishop of London; R. WHITCOMBE, Esq., M
Trinity College, Cambridge; E. POCOCKE, Esq.; the Rev. J. B. OTTLEY, M.A.; and the R
H. THOMPSON, M.A., St. John's College, Cambridge.
"Full of information, expressed in the choicest language."—*Athenæum*.

Vol. X., price 5s., cloth lettered,
MORAL AND METAPHYSICAL PHILOSOPHY. SYSTEMS OF PHILOS
PHY ANTERIOR TO THE TIME OF CHRIST. By the Rev. F. D. MAURICE, M.A. Th
Edition. Re-written.
"Complete and satisfactory."—*Literary Gazette*.

Vol. XI., price 5s., cloth lettered,
INTRODUCTION TO UNIVERSAL HISTORY—ON THE USES OF HISTORY
AS A STUDY, AND ON THE SEPARATION OF THE EARLY FACTS OF HISTORY FR
Sir JOHN STODDART, LL.D. Re-written.
"We cannot speak too highly in its praise."—*Literary Gazette*.

THOMAS
Medical
Alders

John Murray
1886
Cabinet Edition of the Encyclopædia Metropolitana—continued.

Vol. XII., price 8s. 6d., cloth lettered,
AN ANTIQUITIES. By WILLIAM RAMSAY, M.A., Professor of Humanity
the University of Glasgow. With Map, and One Hundred and Thirty Illustrations.
Third Edition.
prises all the results of modern improved scholarship within a moderate compass."—*Athenæum*.

Vol. XIII., price 10s. 6d., cloth lettered,
NY. By JOHN HUTTON BALFOUR, M.D., F.L.S., F.R.S.E., Professor of
tany in the University of Edinburgh. Third Edition, revised and enlarged by JOSEPH WIL-
MS, M.D., Lecturer on Botany at the original School of Medicine, Dublin. Numerous
gravings.
aps the most masterly digest of the science which has yet appeared."—*Witness*.

Vol. XIV., price 3s. 6d., cloth lettered,
TRO-METALLURGY: Containing an Account of the most Improved
thods of Depositing Copper, Silver, Gold, and other Metals. Third Edition, with
merous Illustrations. By JAMES NAPIER, F.C.S.
this book at hand, we don't think it possible to fail in obtaining satisfactory results."—*Art Journal*.

Vol. XV., price 9s., cloth lettered,
DRY OF GREECE from the Earliest Times to the Close of the Pello-
nesian War. By E. POCOCKE, Esq.; Sir T. N. TALFOURD, D.C.L.; Rev. J. B. OTTLEY, A.M.;
the late J. T. RUTT, Esq. With upwards of One Hundred Engravings, from the most
hentic sources.
ether a gem."—*Glasgow Citizen*.

Vol. XVI., price 6s., cloth lettered,
OGRAPHY. By ROBERT HUNT, F.R.S., Professor of Physical Science in
Metropolitan School of Science, London. Fifth Edition, Numerous Engravings.
prove of infinite service to those engaged in the pursuit of this entertaining science."—*Art Journal*.

Vol. XVII., price 3s., cloth lettered,
INARY ART: A Practical Treatise on the Diseases of the Horse. By
C. SPOONER, Esq., with 50 Engravings.

Vol. XVIII., price 8s., cloth lettered,
ORIENTAL HISTORY, comprising the History of the Egyptians, As-
sians, Phœnicians, Persians, &c. By the Rev. Professor EADIE, D.D., LL.D. With
merous Illustrations.
respects a thoroughly good book."—*Dr. Kitto*.

Vol. XIX., price 8s. 6d., cloth lettered,
KEY OF THE ROMAN REPUBLIC. By the Rev. THOS. ARNOLD, D.D.,
Head Master, Rugby School; Sir THOS. N. TALFOURD, D.C.L.; the Rev. Professor JEREMIE,
others. With Numerous Illustrations.

Vol. XX., price 7s. 6d., cloth lettered,
AL ANTIQUITIES AND GEOGRAPHY, including the Language, Geo-
hy, and Early History of Palestine. Edited by the Rev. F. A. COX, D.D., LL.D. With
s and Numerous Illustrations.
ost readable book on the subject in our language."—*Baptist Magazine*.

Vol. XXI., price 12s. 6d., cloth lettered,
LLURGY. A Practical Treatise on the Chemistry of the Metals. By
ARTHUR PHILLIPS, Esq., F.C.S. Second Edition, revised. Illustrated by nearly 200
avings.
qua non to every metallurgist."—*Mining Journal*.

Vol. XXII., price 4s., cloth lettered,
RY OF THE CHRISTIAN CHURCH in the Second and Third Centuries.
ANES AMHAUX JEREMIE, D.D., Regius Professor of Divinity, Cambridge.
a admirable text-book."—*Evangelical Magazine*.

Vol. XXIII., price 8s., cloth lettered,
RY OF GREECE AND MACEDONIA, from the Age of Xenophon to
bsorption of Greece in the Roman Empire, including the History of Alexander the Great
his Successors. By Dr. LYALL, Dean of Canterbury, Rev. G. C. RENOUEAU, B.D., &c. &c.
Numerous Illustrations.
assertations are full, luminous, and charmingly written."—*Eclectic Review*.

Vol. XXIV., price 10s. 6d., cloth lettered,
RY OF ROMAN LITERATURE. By the Rev. THOS. ARNOLD, D.D.,
Head Master, Rugby School; Rev. HENRY THOMPSON, M.A.; Rev. Dr. NEWMAN; Rev. J.
TLEY, M.A.; and Rev. J. M. NEALE, Warden of Sackville College.

ENCYCLOPÆDIA METROPOLITANA :

OR

System of Universal Knowledge ;

ON A METHODICAL PLAN

PROJECTED BY SAMUEL TAYLOR COLERIDGE.

SECOND EDITION, REVISED.

B O T A N Y.

THIRD EDITION, REVISED.

Price 12s. 6d., uniform with this Volume,

A MANUAL OF GEOLOGY;

PRACTICAL AND THEORETICAL.

BY JOHN PHILLIPS, M.A., F.R.S., F.G.S.,

DEPUTY READER IN GEOLOGY IN THE UNIVERSITY OF OXFORD.

WITH NUMEROUS ENGRAVINGS.

207
3

Mackenzie's
A *June 1858.*

MANUAL OF BOTANY:

BEING AN

INTRODUCTION TO THE STUDY

OF THE

STRUCTURE, PHYSIOLOGY, AND CLASSIFICATION OF PLANTS.

BY

JOHN HUTTON BALFOUR, M.D., F.L.S., F.R.S.E.,

PROFESSOR OF BOTANY AND MEDICINE IN THE UNIVERSITY OF EDINBURGH,
AND REGIUS KEEPER OF THE BOTANIC GARDENS.

Third Edition, Revised and Enlarged,

BY JOSEPH WILLIAMS, M.D.,

LECTURER ON BOTANY, ORIGINAL SCHOOL OF MEDICINE, DUBLIN.

Illustrated by Numerous Engravings.

70061
5-16-58

LONDON AND GLASGOW:

RICHARD GRIFFIN AND COMPANY,

PUBLISHERS TO THE UNIVERSITY OF GLASGOW.

1855.

GLASGOW:
PRINTED BY BELL AND BAIN.

PREFACE TO THE THIRD EDITION.

THE object of the Editor of this—the third edition of the “MANUAL OF BOTANY,” has been to render it deserving of the same degree of public approval which has so signally marked its two predecessors. With this view many alterations and amendments, which the advanced state of Botanical Science rendered requisite, have been introduced throughout its pages, to which such additions have also been made, as it is confidently anticipated will constitute the present “Manual” the most complete volume on the subject of which it treats.

The great success of the “MANUAL OF BOTANY,” notwithstanding the existence of many rival works, has induced the Publishers to extend the number of the present edition, and to offer it to the Botanical World at a considerable reduction of price.

March, 1855.

PREFACE TO THE FIRST EDITION.

IN the compilation of this Manual of Botany, the object has been to give a comprehensive, and, at the same time, condensed view of all departments of the science. Attention is directed, first, to the Elementary structure of plants, and the functions of the simplest tissues, and then to the Compound organs, and the functions which they perform. In the consideration of these subjects, the works of Jussieu and Henfrey have served as a model. The application of Physiology to Agriculture, both as regards the cultivation of plants and their diseases, is brought under notice; the works of Liebig, Mülder, and Johnston having been consulted. In the important subject of Classification, much aid has been derived from the standard work of Lindley. The system adopted is that of De Candolle, but in the arrangement and definition of the natural orders, Walker Arnott has been chiefly followed. Many important hints have been derived from Henslow's excellent Syllabus, as well as from the systematic work of Endlicher. In detailing the properties of plants, care has been taken to notice all those which are important in a medical and economical point of view—Christison, Royle, Burnett, and Lindley, supplying valuable data. In the chapter on the Geographical distribution of plants, a very general view is given of the

principal facts brought forward by Meyen, Schouw, Humboldt, Berg-
haus, Watson, and Forbes; and in Fossil Botany, the labours of
Brongniart, Ansted, and Hooker have been made available. The
Publishers placed at the Author's disposal, the wood-cuts of Jussieu's
Cours Elementaire, and some from Beudant's Geology, and, in addi-
tion to these, there are others taken from Raspail, St. Hilaire,
Schleiden, Amici, and Maout. By combining together in the Manual,
information which the student has to acquire by the consultation of
several volumes, it is hoped that the work may be a useful text-book.

EDINBURGH, *March*, 1849.

INTRODUCTORY REMARKS.

It has too often been supposed that the principal object of Botany is to give names to the vegetable productions of the globe, and to arrange them in such a way that these names may be easily found out. This is a most erroneous view of the science, and one which was perhaps fostered by some of the advocates of the Linnæan system. The number of species collected by a botanist is not considered now-a-days as a measure of his acquirements, and names and classifications are only the mechanism by means of which the true principles of the science are elicited. The views in regard to a natural system proposed by Ray and Jussieu did much to emancipate botany from the trammels of artificial methods, and to place it in its proper rank as a science. Their labours have been ably carried out by De Candolle, Brown, Endlicher, Lindley, Hooker, Arnott, and many others. The relative importance of the different organs of plants, their structure, development, and metamorphoses, are now studied upon philosophical principles. The researches of Gaudichand, Mirbel, and others, as to the structure and formation of wood; the observations of Schleiden and Mohl on cell development; the investigations of Brown, Schleiden, Fritzsche, Amici, Meyen, Griffith, and others, into the functions of the pollen, the development of the ovule, and the formation of the embryo; the experiments of Schultz, Decaisne, and Thuret, on the movements observed in the cells, vessels, and spores of plants, and various other physiological inquiries, have promoted much our knowledge of the alliances and affinities of plants. Thus the labours of vegetable anatomists and physiologists all tend to give correct views of the relation which plants bear to each other, and of the great plan on which they were formed by the All-wise Creator.

The Botanist, in accomplishing the ends he has in view, takes an enlarged and comprehensive view of the vegetation with which the earth is clothed. He considers the varied aspects under which plants appear in the different quarters of the globe, from the Lichen on the Alpine summits, or on the Coral reef, to

the majestic Palms, the Bananas and Baobabs of tropical climes—from the minute aquatics of our northern pools to the gigantic Victoria of the South American waters—from the parasitic fungus, only visible by the aid of the microscope, to the enormous parasite discovered by Raffles in the Indian Archipelago.

It is interesting to trace the relation which all these plants bear to each other, and the mode in which they are adapted to different climates and situations. The lichens are propagated by spores (seeds) so minute as to appear like thin dust, and so easily carried by the wind that we can scarcely conceive any place which they cannot reach. They are the first occupants of the sterile rock and the coral-formed island—being fitted to derive the greater part of their nourishment from the atmosphere and the moisture suspended in it. By degrees they act on the rocks to which they are attached, and cause their disintegration. By their decay a portion of vegetable mould is formed, and in progress of time a sufficient quantity of soil is produced to serve for the germination of the seeds of higher plants. In this way the coral island is, in the course of years, covered with a forest of coco-nut trees. Thus it is that the most despised weeds lay the foundation for the denizens of the wood; and thus, in the progress of time, the sterile rock presents all the varieties of meadow, thicket, and forest.

The Creator has distributed his floral gifts over every part of the globe, from the poles to the equator. Every climate has its peculiar vegetation, and the surface of the earth may be divided into regions characterized by certain predominating tribes of plants. The same thing takes place on the lofty mountains of warm climates, which may be said to present an epitome of the horizontal distribution of plants. Again, if we descend into the bowels of the earth, we find there traces of vegetation—a vegetation, however, which flourished at distant epochs of the earth's history, and the traces of which are seen in the coal, and in the fossil plants which are met with in different strata. By the labours of Brongniart especially, these fossil remains have been rendered available for the purposes of science. Many points have been determined relative to their structure, as well as in regard to the climate and soil in which they grew, and much aid has been afforded to the Geologist in his investigations.

The bearings which Botany has on Zoology are seen when we consider the lowest tribe of plants, such as Diatomaceæ. These bear a striking resemblance to the lowest animals, and have been figured as such by Ehrenberg and others. The observations of Thwaites on Conjugation have confirmed the view of the vegetable nature of many of these bodies. There appear, however, to be many productions which occupy a sort of intermediate territory between the animal and vegetable kingdom, and for the time being the Botanist and Zoologist must consent to joint occupancy.

The application of botanical science to Agriculture and Horticulture has of late

attracted much attention, and the chemistry of plants has been carefully examined by Liebig, Mülder, and Johnston. The consideration of the phenomena connected with germination, and the nutrition of plants, has led to important conclusions as to sowing, draining, ploughing, the rotation of crops, and the use of manures.

The relation which Botany bears to Medicine has often been misunderstood. The medical student is apt to suppose that all he is to acquire by his botanical pursuits, is a knowledge of the names and orders of medicinal plants. The object of the connection between scientific and mere professional studies is here lost sight of. It ought ever to be borne in mind by the medical man, that the use of the collateral sciences, as they are termed, is not only to give him a great amount of general information, which will be of value to him in his after career, but to train his mind to that kind of research which is essential to the student of medicine, and to impart to it a tone and a vigour which will be of the highest moment in all his future investigations. What can be more necessary for a medical man, than the power of making accurate observations, and of forming correct distinctions and diagnoses? These are the qualities which are brought into constant exercise in the prosecution of the botanical investigations, to which the student ought to turn his attention, as preliminary to the study of practical medicine. In the prosecution of his physiological researches, it is of the highest importance that the medical man should be conversant with the phenomena exhibited by plants. For no one can be reckoned a scientific physiologist who does not embrace within the range of his inquiries all classes of animated beings; and the more extended his views, the more certain and comprehensive will be his generalizations.

To those who prosecute science for amusement, Botany presents many points of interest and attraction. Though relating to living and organized beings, the prosecution of it calls for no painful experiments nor forbidding dissections. It adds pleasure to every walk, and affords an endless source of gratification, which can be rendered available alike in the closet and in the field. The prosecution of it combines healthful and spirit-stirring recreation with scientific study; and its votaries are united by associations of no ordinary kind. He who has visited the Scottish Highlands with a botanical party, knows well the feelings of delight connected with such a ramble—feelings by no means of an evanescent nature, but lasting during life, and at once recalled by the sight of the specimens which were collected. These apparently insignificant remnants of vegetation recall many a tale of adventure, and are associated with the delightful recollection of many a friend. It is not indeed a matter of surprise, that those who have lived and walked for weeks together in a Highland ramble, who have met in sunshine and in tempest, who have climbed together the misty summits, and have slept in the miserable shieling—should have such scenes indelibly impressed on their

memory. There is, moreover, something peculiarly attractive in the collecting of *alpine* plants. Their comparative rarity, the localities in which they grow, and frequently their beautiful hues, conspire in shedding around them a halo of interest far exceeding that connected with lowland productions. The alpine *Veronica* displaying its lovely blue corolla on the verge of dissolving snows; the Forget-me-not of the mountain summit, whose tints far excel those of its namesake of the brooks; the *Woodsia*, with its tufted fronds, adorning the clefts of the rocks; the snowy *Gentian* concealing its eye of blue in the ledges of the steep crags; the alpine *Astragalus* enlivening the turf with its purple clusters; the *Lychnis* choosing the stony and dry knoll for the evolution of its pink petals; the *Sonchus*, raising its stately stalk and azure heads in spots which try the enthusiasm of the adventurous collector; the pale-flowered *Oxytropis* confining itself to a single British cliff; the *Azulea* forming a carpet of the richest crimson; the *Saxifrages*, with their white, yellow, and pink blossoms, clothing the sides of the streams; the *Saussurea*, and *Erigeron* crowning the rocks with their purple and pink capitula; the pendent *Cinquefoil* blending its yellow flowers with the white of the alpine *Cerastiums* and the bright blue of the stony *Veronica*; the stemless *Silene* giving a pink and velvety covering to the decomposing granite; the yellow *Hieracia*, whose varied transition forms have been such a fertile cause of dispute among botanists; the slender and delicate grasses, the chickweeds, the carices, and the rushes, which spring up on the moist alpine summits; the graceful ferns, the tiny mosses, with their urn-like thecæ, the crustaceous dry lichens, with their spore-bearing apothecia; all these add such a charm to highland botany, as to throw a comparative shade over the vegetation of the plains.

Many are the important lessons which may be drawn from the study of plants, when prosecuted in the true spirit of Wisdom. The volume of Creation is then made the handmaid of the volume of Inspiration, and the more that each is studied, the more shall we find occasion to observe the harmony that subsists between them. It is only Science, falsely so called, which is in any way opposed to Scripture. Never, in a single instance, remarks Gaussen, do we find the Bible in opposition to the just ideas which Science has given us regarding the form of our globe, its magnitude, its geology, and the productions which cover the surface. "The invisible things of God from the creation of the world are clearly seen, being understood by the things that are made, even his eternal power and Godhead." The more minutely we examine the phenomena of the material world, and the more fully we compare the facts of science with Revealed Truth, the more reason shall we have to exclaim, in adoring wonder, with the Psalmist of old, "O Lord! how manifold are thy works! in wisdom hast thou made them all; the earth is full of thy riches."

TABLE OF CONTENTS.

	PAGE
PREFACE.....	vii
INTRODUCTORY REMARKS.....	ix
PART I.—VEGETABLE ANATOMY, ORGANOGRAPHY, AND PHYSIOLOGY	1
CHAPTER I.—ELEMENTARY ORGANS, OR VEGETABLE TISSUES	1
SECTION I.—CELLULAR TISSUE.....	3
1. Form and Arrangement of Cells.....	3
2. Contents of Cells.....	8
3. Development and Functions of Cells	12
SECTION II.—VASCULAR TISSUE	15
1. Form and Arrangement of Vessels	15
2. Development and Functions of Vessels.....	21
Tabular Arrangement of Vegetable Tissues.....	23
CHAPTER II.—COMPOUND ORGANS FORMED BY THE TISSUES	24
SECTION I.—ORGANS OF NUTRITION OR VEGETATION.....	25
Structure, Arrangement, and Special Functions.....	25
General Integument	25
Stomata	27
Hairs	30
Stem or Ascending Axis	35
Forms of Stems.....	35
Internal Structure of Stems	39
Exogenous or Dicotyledonous Stem	39
Anomalies in its Structure.....	50
Endogenous or Monocotyledonous Stem.....	53
Acrogenous or Acotyledonous Stem.....	59
Formation of the different parts of Stems, and their special Functions	61
Root or Descending Axis	65
Structure of Roots	65
Forms of Roots	67
Functions of Roots	69
Leaves and their Appendages	71
Structure of Leaves.....	71
Venation of Leaves.....	75
Forms of Simple Leaves.....	77
Forms of Compound Leaves.....	81
Petiole or Leaf-stalk	84
Stipules	87
Anomalous Forms of Leaves and Petioles	89
Structure and Form of Leaves in the Great Divisions of the Vegetable Kingdom.....	90
Phyllotaxis, or the Arrangement of Leaves on the Axis.....	91
Leaf-buds	97
Vernation.....	99
Special Functions of Leaves.....	109

	PAGE
SECTION II.—GENERAL VIEW OF THE FUNCTIONS OF THE NUTRITIVE ORGANS.....	111
1. Food of Plants, and Sources whence they derive their Nourishment	111
Chemical Composition of Plants	111
Organic Constituents and their Sources	113
Inorganic Constituents and their Sources	116
Chemical Composition of Soils.....	122
Application of Manure	124
Various kinds of Manure	124
Epiphytic and Parasitic Plants	129
2. Absorption and Circulation of Fluids	130
3. Respiration of Plants.....	142
Effects of Certain Gases on Living Plants	146
4. Products and Secretions of Plants.....	149
SECTION III.—ORGANS OF REPRODUCTION	159
Structure, Arrangement, and Functions	159
1. Inflorescence or the arrangement of the flowers on the axis	160
Tabular View of Inflorescence or Anthotaxis	173
2. Bracts or Floral leaves	174
3. The Flower and its Appendages	176
External Floral Whorls, or the Floral Envelopes.....	180
Calyx	180
Corolla	185
Inner Floral Whorls, or the Essential Organs of Reproduction	195
Stamens.....	195
Pistil	217
Ovule	233
4. Functions of the Floral Envelopes	239
5. Functions of the Stamens and Pistil; Fertilization or Fecundation.....	245
Embryology.....	251
6. Fruit or the Pistil arrived at maturity	260
Fruits which are the produce of a single flower	271
Fruits which are the produce of several flowers united.....	277
Tabular arrangement of Fruits.....	279
7. Maturation of the Pericarp	280
Ripening of Fruits	281
Grafting	283
8. Seed or Fertilized Ovule arrived at Maturity	286
Embryo	295
9. Functions of the seed	303
Germination	304
Vitality of Seeds.....	307
Direction of Plumule and Radicle	311
Proliferous Plants	315
Duration of the Life of Plants	316
10. General Observations on the Organs of Plants, and on the mode in which they are arranged.....	318
Tabular View of the Organs	318
Symmetry of Organs.....	321
Teratology	323
SECTION IV.—SOME GENERAL PHENOMENA CONNECTED WITH VEGETATION.....	330
1. Vegetable Irritability	330
2. Temperature of Plants	335
3. Luminosity of Plants	336
4. Colours of Plants.....	337
5. Odours of Plants	342
6. Diseases of Plants	343
PART II.—SYSTEMATIC BOTANY, TAXONOMY, OR THE CLASSIFICATION OF PLANTS	349
CHAPTER I.—SYSTEMS OF CLASSIFICATION.....	349
Nomenclature and Symbols	353
Linnaean System	354
Natural System	356
System of Jussieu	359
System of De Candolle.....	359
System of Endlicher	360
System of Lindley	361

CHAPTER II.—ARRANGEMENT AND CHARACTERS OF THE CLASSES AND
NATURAL ORDERS 364

SECTION I.—PHANEROGAMOUS PLANTS 364

Class I.—Dicotyledones or Exogenæ 364

Subclass 1.—Thalamifloræ 365

1. Ranunculaceæ 365	21. Tremandraceæ 380	41. Malpighiaceæ 395
2. Dilleniaceæ 367	22. Tamaricaceæ 380	42. Aceraceæ 396
3. Magnoliaceæ 367	23. Frankeniaceæ 381	43. Sapindaceæ 396
4. Anonaceæ 368	24. Elatinaceæ 381	44. Rhizophoraceæ 397
5. Menispermaceæ 368	25. Caryophyllaceæ 381	45. Meliaceæ 397
6. Berberidaceæ 369	26. Vivianaceæ 383	46. Cedrelaceæ 398
7. Cabombaceæ 370	27. Malvaceæ 383	47. Ampelidææ 399
8. Nymphaeaceæ 370	28. Sterculiaceæ 386	48. Geraniaceæ 400
9. Nelumbiaceæ 371	29. Byttneriaceæ 387	49. Linaceæ 401
10. Sarraceniacææ 371	30. Tiliaceæ 387	50. Balsaminaceæ 401
11. Papaveraceæ 372	31. Dipterocarpaceæ 388	51. Oxalidaceæ 402
12. Fumariaceæ 372	32. Chlénaceæ 389	52. Tropæolaceæ 403
13. Cruciferae 373	33. Ternstroemiaceæ 389	53. Pittosporaceæ 403
14. Capparidaceæ 376	34. Olacaceæ 390	54. Brexiaceæ 403
15. Resedaceæ 376	35. Aurantiaceæ 391	55. Zygophyllaceæ 404
16. Flacourtiaceæ 377	36. Hypericaceæ 392	56. Rutaceæ 404
17. Cistaceæ 377	37. Guttiferæ 393	57. Xanthoxylaceæ 406
18. Violaceæ 378	38. Marcgraviaceæ 394	58. Simarubaceæ 406
19. Droséraceæ 379	39. Hippocrateææ 394	59. Ochnaceæ 407
20. Polygalaceæ 379	40. Erythroxylaceæ 395	60. Coriariaceæ 407

Subclass 2.—Calficifloræ 408

61. Stackhousiaceæ 408	74. Vochysiaceæ 425	87. Turneraceæ 435
62. Celastraceæ 408	75. Combretaceæ 425	88. Portulacaceæ 435
63. Staphyleaceæ 409	76. Melastomaceæ 425	89. Paronychiaceæ 436
64. Rhamnaceæ 409	77. Alangiaceæ 426	90. Crassulaceæ 436
65. Anacardiaceæ 410	78. Philadelphiceæ 426	91. Ficoideæ 437
66. Amyridaceæ 412	79. Myrtaceæ 427	92. Cactaceæ 438
67. Conaraceæ 413	80. Onagraceæ 429	93. Grossulariaceæ 439
68. Leguminosæ 413	81. Haloragaceæ 430	94. Saxifragaceæ 440
69. Moringaceæ 419	82. Loasaceæ 430	95. Bruniaceæ 441
70. Rosaceæ 420	83. Cucurbitaceæ 430	96. Hamamelidaceæ 441
71. Calycanthaceæ 423	84. Papayaceæ 433	97. Umbelliferae 441
72. Lythraceæ 423	85. Belvisiaceæ 433	98. Araliaceæ 445
73. Rhizophoraceæ 424	86. Passifloraceæ 434	99. Cornaceæ 445

Subclass 3.—Corollifloræ 446

100. Loranthaceæ 446	115. Epacridaceæ 465	130. Hydrophyllaceæ 478
101. Caprifoliaceæ 447	116. Columellaceæ 465	131. Convolvulaceæ 479
102. Rubiaceæ 448	117. Styracaceæ 466	132. Cordiaceæ 481
103. Valerianaceæ 452	118. Ebenaceæ 467	133. Boraginaceæ 481
104. Dipsacaceæ 452	119. Aquifoliaceæ 467	134. Solanaceæ 483
105. Calyceraceæ 453	120. Sapotaceæ 468	135. Orobanchaceæ 486
106. Compositæ 453	121. Myrsinaceæ 469	136. Scrophulariaceæ 487
107. Brunoniaceæ 459	122. Jasminaceæ 469	137. Labiatæ 489
108. Goodeniaceæ 460	123. Oleaceæ 470	138. Verbenaceæ 491
109. Stylidiaceæ 460	124. Asclepiadaceæ 471	139. Acanthaceæ 492
110. Campanulaceæ 461	125. Apocynaceæ 473	140. Lentibulariaceæ 492
111. Lobeliaceæ 462	126. Loganiaceæ 474	141. Primulaceæ 493
112. Gesneraceæ 463	127. Gentianaceæ 476	142. Plumbaginaceæ 494
113. Ericaceæ 463	128. Bignoniaceæ 477	143. Plantaginaceæ 495
114. Vacciniaceæ 464	129. Polemoniaceæ 478	

Subclass 4.—Monochlamydeæ 496

A.—Angiospermeæ 496

a.—Spermogeenæ. Having true seeds, with a Dicotyledonous Embryo.

144. Nyctaginaceæ 496	152. Proteaceæ 505	160. Santalaceæ 509
145. Amaranthaceæ 497	153. Elæagnaceæ 506	161. Aristolochiaceæ 509
146. Chenopodiaceæ 498	154. Penæaceæ 506	162. Nepenthaceæ 511
147. Phytolaccaceæ 499	155. Thymelæaceæ 507	163. Datisceææ 511
148. Polygonaceæ 499	156. Aquillariaceæ 507	164. Empetraceæ 512
149. Begoniaceæ 501	157. Chailletiaceæ 508	165. Euphorbiaceæ 512
150. Lauraceæ 502	158. Samydaceæ 508	166. Urticaceæ 515
151. Myristicaceæ 504	159. Homaliaceæ 509	167. Ceratophyllaceæ 520

	PAGE		PAGE		PAGE
168. Podostemaceæ	520	172. Lacistemaceæ	521	176. Amentaceæ	523
169. Stilaglinaceæ	520	173. Chloranthaceæ	522	177. Juglandaceæ	527
170. Monimiaceæ	521	174. Saururaceæ	522	178. Garryaceæ	527
171. Atherspermaceæ	521	175. Piperaceæ	522		
b.—Sporogenæ. Having spore-like seeds, with an Acotyledonous Embryo.					
179. Rhizanthæ					528
B.—Gymnospermæ					
180. Coniferæ	529	181. Cycadaceæ	532		
Class II.—Monocotyledones or Endogenæ					
Subclass 1.—Dictyogenæ					
182. Dioscoreaceæ	534	183. Similacæ	534	184. Trilliaceæ	535
Subclass 2.—Petaloidæ					
a.—Perianth adherent, Ovary inferior, Flowers usually hermaphrodite					
185. Hydrocharidaceæ	535	189. Musaceæ	541	193. Amaryllidaceæ	544
186. Orchidaceæ	536	190. Iridaceæ	542	194. Hypoxidaceæ	545
187. Zingiberaceæ	539	191. Burmanniaceæ	543	195. Bromeliaceæ	545
188. Marantaceæ	540	192. Ilæmodoraceæ	543		
b.—Perianth free, Ovary superior, Flowers usually hermaphrodite					
196. Liliaceæ	546	200. Xyridaceæ	549	203. Commelynacæ	553
197. Melanthaceæ	548	201. Juncaceæ	550	204. Allismaceæ	553
198. Gilliesiaceæ	549	202. Palmæ	550	205. Butomaceæ	554
199. Pontederiaceæ	549				
c.—Flowers incomplete, often unisexual, without a proper perianth, or with a few verticillate scales					
206. Pandanaceæ	554	208. Naladaceæ	556		
207. Araceæ	555	209. Restiaceæ	557		
Subclass 3.—Glumaceæ					
210. Cyperaceæ	557	211. Gramineæ	558		
SECTION II.—CRYPTOGAMOUS PLANTS					
Class III.—Acotyledones					
Subclass 1.—Acrogenæ or Cormogenæ					
212. Equisetaceæ	563	214. Marsileaceæ	566	216. Musci	567
213. Filices	564	215. Lycopodiaceæ	567	217. Hepaticæ	569
Subclass 2.—Thallogenæ or Cellulares					
218. Lichenes	570	219. Fungi	572	220. Algæ	576
PART III.—GEOGRAPHICAL BOTANY, OR THE DISTRIBUTION OF PLANTS OVER THE GLOBE					
I.—EPIRRHEOLOGY, OR THE INFLUENCE OF VARIOUS EXTERNAL AGENTS ON PLANTS					
1. Effects of Temperature					
2. Effects of Moisture					
3. Effects of Soil, Light, and other Agents					
II.—DISSEMINATION OF PLANTS					
1. Agents employed in their Dissemination					
2. General and Endemic Distribution of Plants					
3. Conjectures as to the mode in which the Earth was originally clothed with Plants					
4. Distribution of Plants considered Physiognomically and Statistically					
Physiognomy of Vegetation					
5. Phyto-geographical Division of the Globe					
Horizontal Range of Vegetation					
Schouw's Phyto-geographic Regions					
Meyen's Phyto-geographical Zones					
Vertical Range of Vegetation					
PART IV.—FOSSIL BOTANY					
Character and arrangement of Fossil Plants					
Fossiliferous Rocks					
Fossil Plants of different Strata					
Reign of Acrogens					
Fossil Plants of the Carboniferous system					
APPENDIX					
I.—ON THE USE OF THE MICROSCOPE IN BOTANICAL RESEARCHES					
II.—ON COLLECTING AND EXAMINING PLANTS, AND ON THE FORMATION OF A HERBARIUM					
GLOSSARY					

BOTANY.

PART I.

VEGETABLE ANATOMY, ORGANOGRAPHY, AND PHYSIOLOGY.


1. BOTANY is that branch of science which comprehends the knowledge of all that relates to the Vegetable Kingdom. It embraces a consideration of the external configuration of plants, their structure, the functions which they perform, the relations which they bear to each other, and the uses to which they are subservient. It has been divided into the following departments:—1. *Structural Botany*, or *Organography*, which has reference to the textures of which plants are composed, and to the forms of their various organs. 2. *Physiological Botany*, in which plants are considered in their living or active state, and while performing certain vital functions. 3. *Systematical Botany*, or *Taxonomy*, the arrangement and classification of plants. 4. *Geographical Botany*, or the distribution of plants over the globe. 5. *Fossil Botany*, or the nature of the plants found in a fossil state in the various geological formations. And, 6. *Microscopical Botany*, or that minute investigation of vegetable tissues by which we determine their atomic construction.

CHAPTER I.

ELEMENTARY ORGANS, OR VEGETABLE TISSUES.

2. In their earliest and simplest state, plants consist of minute vesicles, formed by an elastic transparent membrane, which is composed of a

substance called *Cellulose*. This substance is of general occurrence, and constitutes the basis of vegetable tissues. The chemical formula representing it is $C^{24} H^{21} O^{21}$, or $2 (C^{12} H^{10} O^{10}) + HO$.* It was long considered as an essential vegetable product, not to be found in animal structures; but Schmidt, Kölliker, and others, have recently detected it in the tissues of the ascidia, and other moluscous animals. It is allied to starch, to which it is convertible by the action of heat, the addition of sulphuric acid, or caustic alkali. It becomes yellow on the addition of iodine, and when acted upon by iodine and sulphuric acid assumes a blue colour; iodide of amidin being produced in consequence of the conversion of the cellulose into starch by the super-added acid. The membrane formed by it is permeable by fluids, and becomes altered in the progress of growth, so as to acquire various degrees of consistence. It must not be confounded with *ligneous* or *woody tissue*, which is but a modification of cellulose encrusting the walls of the original membranous cells, and conferring on them hardness and inflexibility. Lignine is not coloured by iodine, it yields on analysis a result indicating an excess of hydrogen above that required to form water with the oxygen, besides traces of nitrogen. Pure cellulose is a ternary compound of carbon and the elements of water, being closely allied in composition, if not actually isomeric with starch. In the advanced stages of growth, plants consist of two kinds of tissue, *Cellular* and *Vascular*, which, under various modifications, constitute their *Elementary organs*. These, by their union, form the *Compound organs*, by which the different functions of plants are carried on.

3. The elementary organs consist of *vesicles* and *tubes*, varying in form and size, and united in different ways. Vascular structures are ever confirmatory of the pre-existence of cell growth, though cellular development may proceed to infinite consummation without the formation of tubes. According as we ascend in the scale of vegetable life we shall find the diversity of the cellular and vascular structures increase, both as regards their relation to each other, and the purposes to which they are subservient. The vesicles are cavities surrounded by a membrane, their length not much exceeding

 1 their breadth (fig. 1); while the tubes are similar cavities more or less elongated (figs. 3, 4). The sphere must be regarded as the fundamental type of the vegetable cell, however it may become subsequently changed. While in a ratio to the changes occurring in cells, is their capability of forming tubes in a great measure determined.


* For the meaning of these and other chemical symbols, see Chap. II. Sect. I. Div. 2, on the Food of Plants.

Fig. 1.—Vesicles.


SECTION I.—CELLULAR TISSUE.

1.—FORM AND ARRANGEMENT OF CELLS.

4. **Cellular Tissue** is formed by the union of minute vesicles or bladders, called *cells*, *cellules*, or *utricles*. This tissue is often called Parenchyma ($\pi\alpha\rho\acute{\alpha}$, beside or between, and $\chi\epsilon\upsilon\mu\alpha$, any thing effused or spread out, tissue). The derivation of Parenchyma is, by some given $\pi\alpha\rho\acute{\alpha}$, through, and $\epsilon\gamma\chi\epsilon\omega$, I infuse. The terms *Parenchymatous*, *Areoler*, *Utricular*, and *Vesicular*, when applied to vegetable tissues, may be considered as synonymous. The individual cells of which it is composed, when allowed to develop themselves equally in all parts of their circumference, are usually of a more or less rounded form (figs. 5, 6, 7); but, when pressed upon during the progress of development, they become more elongated in one direction than in another (fig. 2), and often assume angular or polyhedral forms (fig. 8).



5. The following names have been applied by Morren, and other authors, to the tissue made up of variously-formed cells:—1. *Parenchyma*, a general name for cellular tissue, but often applied to that consisting of dodecahedral cells (figs. 8, 12, 13), which, when cut in one direction, exhibit an hexagonal form (figs. 14, 15), and hence the tissue is sometimes called *hexagonienchyma* ($\epsilon\kappa\alpha\gamma\omega\gamma\iota\omicron\varsigma$, six-angled); it is



seen in the pith of the Elder, and in young palm stems. 2. *Sphaerenchyma* ($\sigma\phi\alpha\iota\sigma\kappa\alpha$, a sphere), spheroidal cells (fig. 5). 3. *Merenchyma* ($\mu\eta\rho\upsilon\omega$, to revolve), ellipsoidal cells (fig. 6). 4. *Ovenchyma* ($\omega\omicron\nu$, anegg), oval cells. Round, elliptical, and oval cells, are common in herbaceous

Fig 2.—Fusiform or spindle-shaped cell. Figs. 3, 4.—Tubes or vessels.
Figs. 5, 6, 7, 8.—Cells, or utricles, separate and combined.
Figs. 9, 10, 11, 12, 13.—Figures representing the forms of cells.

plants. 5. *Conenchyma* (κῶνος, a cone), conical cells, as hairs. 6. *Columnar* cellular tissue, divided into *Cylindrenchyma* (κύλινδρος, a cylinder), cylindrical cells, as in Chara (fig. 17 a), and *Prismenchyma* (πρίσμα, a prism), prismatical cells, seen in the bark of some plants (fig. 10). When compressed, prismatical cells form the *muriform* (*murus*, a wall, like bricks of a building) tissue of the medullary rays of woody stems, and when much shortened they assume a tabular form, constituting *Pinenchyma* (πίναξ, a table), tabular cells (fig. 11), or square cells (fig. 9). 7. *Prosenchyma* (¶ 29), or *Atractenchyma* (ἄτρακτος, a spindle), fusiform or spindle-shaped cells, seen in bark and wood (fig. 2). 8. *Colpenchyma* (κόλπος, a sinus or fold), sinuous or waved cells, as in the cuticle of leaves. 9. *Cladenchyma* (κλάδος, a branch), branched cells, as in some hairs. 10. *Actinenchyma* (ἄκτιν, a ray), stellate or radiating cells, as in Juncus and Musa (fig. 16). 11. *Dactalenchyma* (δαίδαλος, entangled), entangled cells, as in some Fungi.



14



15



16


6. The size of cells varies not less than their figure in different plants, and in different parts of the same plant. They are frequently seen from $\frac{1}{300}$, $\frac{1}{300}$, to $\frac{1}{1000}$ of an inch in diameter. In cork, which is cellular, Hooke found more than a thousand in the length of an inch. The size of the cell varies considerably, according to the nature of the structure in which it is situated. In many succulent vegetables, and in the pith of some aquatic plants, large cells ranging from $\frac{1}{30}$ to $\frac{1}{30}$ of an inch in diameter occur. Mohl declares the general average of cells to vary from $\frac{1}{160}$ to $\frac{1}{800}$, while the cells in spores of Fungi have been computed at $\frac{1}{4000}$ of an inch in diameter.

7. Each cell consists originally of a separate membrane, but in the progress of growth the walls of contiguous cells may become united. When cells are united by their extremities (fig. 17), their partitions are occasionally absorbed so as to form continuous tubes. When cells are united in a rectilinear manner, those in contiguous rows are either directly opposite to each other, that is, are placed at the same height (fig. 18), or are alternate, from being placed at different heights (fig. 19); cells sometimes communicate with each other laterally (fig. 20 a a). Isolated cells, as spores of sea-weeds, occasionally

Figs. 14, 15.—Hexagonal cells.

Fig. 16.—Branching cells of *Vicia Faba*. l l, Intercellular lacunae.

have free filaments, or cilia (*cilium*, an eyelash) developed on their surface.



8. The simplest kinds of plants, as mushrooms and sea-weeds, are composed entirely of cellular tissue, and are called *Cellulares*. The pulpy and succulent parts of all plants contain much cellular tissue, and the object of horticultural operations is to increase the quantity of this tissue in ordinary fruits and vegetables. The pith of trees and plants during their early development is cellular; so also are cotton and rice-paper. The cell may be considered as the ultimate structural element of all organism. In the simplest form of generation, as in unicellular algæ, it is adequate to all the purposes of plant life. Proportionate to the vital activity of each cell, is its functional vegetable power, whether it be for the secretion of special products, or the organization of tissues. The life of the entire plant being dependent on the aggregate operation of its cells, may either find in the simple cell its representation, each cell being of the same physiological importance; or, in the different and unequal action of its cells, their varied physical and chemical natures, and the diversity of the purposes for which they are subservient, may the explanation of the compound nature of many vascular plants rest. Schacht has observed that the higher we ascend in the scale of vegetable life, the more unequal in their individual importance do the separate cells of the same plant become. In the lowest tribes of vegetables, multicellular plants are unicellular in vitality. As we advance, the organism becomes more complex, until, at length, the varied processes of absorption, assimilation, nutrition, and reproduction are dependent on the regular action of different cells, each having an unequal, though a vitally important influence on the perfection of the plant.

9. In general, no visible openings can be detected in cells, although fluids pass readily into and out of them. Harting and Mulder, however, state, that they have observed perforations in the cells of *Hoya carnosa*,

Figs. 17, 18, 19.—Cells united together by their extremities.

Fig. 20.—Elongated thickened cells, from the root of the Date palm. *a a*, Canals of communication.

Asclepias syriaca, *Cycas revoluta*, Virginian spiderwort, and Traveller's joy. In one cell (from a *Euphorbia*), having a transverse diameter of 0.03777 millimetres,* they counted 45 minute holes. In some mosses, also, openings have been found in the cells.

10. **Porous cells** are those in which the membrane appears thickened at certain parts, with thin rounded spots intervening, which, when viewed by transmitted light, appear like perforations or pores (figs. 21, 28).

The unequal deposit of the internal encrusting cellulose, and the addition of sclerogenous matter, are the chief causes of this condition. The pores of contiguous cells usually correspond as regards position, and sometimes the membrane becomes absorbed between them, so as to allow a direct communication by means of lateral canals, as is seen in the cells from the root of the Date (fig. 20 *a a*).

When porous cells are united end to end, so as to form tubes, they have been denominated *articulated Bothrenchyma* ($\beta\acute{o}\theta\rho\epsilon\omicron\varsigma$, a pit), on account of the pits or depressions in their thickened walls (fig. 22), which frequently imparts a bead-like appearance to the canals so formed.

11. **Fibrous or Spiral cells** are those in which there is a spiral elastic fibre coiled up in the inside of the membrane (fig. 23). When united they form *fibro-cellular tissue*, or *Inenchyma* ($\iota\nu\epsilon\varsigma$, fibres). These



21


22



23



24



25



26

cells generally consist of membrane and fibre combined, but the former appears to be sometimes absorbed wholly or partially during the progress of growth. The membrane, in some instances, is easily dissolved by water, and then the elastic close convolutions of the fibre spring out with considerable force, as in the outer covering of the seeds of *Collomia* and *Salvia*. Spiral cells abound in many of the Orchidaceous plants, and in the Cactus tribe. They are also found in the inner covering of anthers, in the spore-cases of many of the lower tribes of plants, and in the coats of the seed of *Acanthodium spicatum*. The spiral filaments sometimes exhibit peculiar movements when placed in water, which bear a close similitude to the ciliary action observable

* A millimetre is about 1.25th of an English inch.

Fig. 21.—Porous cell, from the Elder (*Sambucus nigra*).


Fig. 22.—Articulated Bothrenchyma, or Taphrenchyma, from Mistletoe, having a monilliform appearance. Figs. 23, 24, 25.—Spiral, annular, and reticulated cells, from Mistletoe (*Viscum album*).

Fig. 26.—Scalariform and dotted cells, from Elder (*Sambucus nigra*).

on the muscous membrane of animal tubes. In some instances they appear to be due to the hygroscopic properties of the cell, by which the filaments encircling its inner walls are expanded; in others, to be the result of an active process of nutrition, and the progress of cell growth, which are inexplicable by physical causes. The fibre varies from about $\frac{1}{1000}$ to $\frac{1}{100}$ of an inch in diameter; it is solid, and presents either a circular, an elliptic, or a quadrangular section. The coils of the fibre sometimes separate from each other, and become broken up and united in various ways, so as to appear in the form of rings, bars, or dots, thus giving rise to *annular* (fig. 24), *reticulated* (fig. 25), *scalariform* and *dotted* cells (fig. 26), which constitute the *spurious* or *imperfect Inenchyma* of authors.

12. In certain parts of plants cells are placed closely together, and compressed so as to touch each other by flat surfaces, filling up space completely, and leaving no intervals; they then form the *perfect Parenchyma* of Schleiden (figs. 8, 27). In lax tissues, however, the cells retain a rounded shape, and then touch each other at certain points only, leaving intervals of various sizes and shapes, and forming the *imperfect Parenchyma* of Schleiden (figs. 7, 28). These intervals, when of moderate size and continuous, are called *intercellular passages* or *canals*; when large, irregular, and circumscribed, *intercellular spaces*, or *Lacunæ* (fig. 16 l l).

13. A difference of opinion prevails as to the mode in which cells are united together. Some maintain that the cell-walls in the young state unite together directly, and become agglutinated, more or less,



according to their places of contact. Others, as Mohl and Henfrey, hold that there is an intercellular matter which acts as a sort of cement, or *Collenchyma* ($\kappaόλλα$, glutinous matter). In sea-weeds, the cells, of which the entire plant is composed, are placed at a distance from each other (fig. 29 a a), and the intervals are filled up by this intercellular substance (fig. 29 b), which thus forms a large part of

Fig. 27.—Cellular tissue, from pith of Elder.

Fig. 28.—Porous Merenchyma, from Houseleek (*Sempervivum tectorum*). a, Intercellular canal.

their bulk. In the higher classes of plants, when the cells touch each other, the layer of intercellular matter must be very thin, except in the intercellular canals or spaces. Mirbel looks upon it as the



remains of the mucilaginous fluid in which the cells were originally developed, and which has become thickened to a greater or less degree, as in the root of the Date (fig. 30), where *a a a* indicate the cells, and *b b b* the interposed substance.


2.—CONTENTS OF CELLS.

14. The external membrane of cells is composed of the unazotised substance called *Cellulose*, and in their interior a mucilaginous matter is contained, which undergoes changes in the progress of growth. This mucilaginous matter is the *Protoplasm* (πρωτοσ, first, and πλάσμα, formative matter) of Mohl, the *Cytoblastema* (κύτος, a cell, and βλάστημα, a germ) of some authors. It is at first homogeneous, but ultimately assumes a granular form. The appearance of granules may be regarded as the earliest evidence of formative power. It contains nitrogen in its composition, or is azotised, and it assumes a brownish colour when acted upon by iodine. It forms a mucilaginous layer on the inner surface of the cell-wall, and thus gives rise to the *internal utricle* of Harting and Mulder, the *primordeal utricle* of Mirbel. This inner membrane is visible in the young state of the cell, and under the action of tincture of iodine may be made to contract and separate from the outer cell-wall. It may also be rendered distinct by the action of strong hydrochloric acid, and by diluted sulphuric acid. When the process of lignification or thickening has advanced, this utricle disappears, in consequence of becoming incorporated with the cell-wall.

15. In certain cells the membrane continues throughout to be formed of a thin layer of cellulose, while in others it becomes thickened by the deposition of matter on its inner side. These secondary deposits, or incrustations, are sometimes of a gelatinous consistence; at other times they are hard. In the latter case, the incrusting matter is looked upon as a modification of cellulose, and has received the name

Fig. 29.—Cellular tissue of Sea-weed (*Himanthalia lorea*). *a a*, Cells. *b*, Intercellular matter.
Fig. 30.—Central portion of young root of Date. *a a a*, Thickened cells. *b b b*, Intercellular substance of Mirbel.

of *lignine* (*lignum*, wood), or *sclerogen* (σκληρός, hard, and γένναιον, to generate). On making sections of such cells, in a transverse (fig. 31) or longitudinal direction (fig. 32), the successive layers may be seen either continuous all round, or leaving parts of the membrane uncovered. Cells of this kind are well seen under the microscope in thin sections of the hard shell of the Coco-nut, or of *Attalea funifera*, and of the hard seed of the Ivory Palm. In all cell-deposits there is a tendency to a spiral arrangement. When the deposition is uniform over the whole surface, this arrangement may not be detected; but when interruptions take place, then the continued coil becomes evident. In spiral cells the fibre seems to be formed before the full development of the cell, the coils of the fibre being at first in contact, and afterwards separated, whereas the secondary thickening layers are deposited after the cell is fully formed. According to the observations of Barry, Agarth, and others, the filamentous origin of fibrous structures is not only recognizable in the earliest stage of cell growth, but the interweaving of these filaments constitute the cell walls.



16. Each cell is found to contain, at some period of its existence, a small body, called a *nucleus* (fig. 33 *n n n*), in which there are often one or two, rarely more, minute spots, called *nucleoli*. The nucleus is of a round or oval shape, granular and dark, or homogeneous and transparent, bearing some resemblance to a smaller internal cell. Nucleoli are not always present. They are either vesicles and granules contained in the nucleus, or minute cavities in its substance. The latter view is supported by Barry, who holds that a peculiar substance, called *hyaline* (ὑαλός, glass), is developed there, which, according to him, is the origin of the nucleus. The nucleus is situated at different parts of the cell. It is either free in its cavity, or connected with its wall by mucilaginous threads, or imbedded in the substance of the membrane. The addition of acetic acid often renders the nucleus distinct.




17. **Starchy matter** is found in cells, which constitute the tissue called, by Morren, *Perenchyma* (πρήσα, a sac.) Starch exists in the form of granules, which are minute cells, (perhaps nuclei, as Mùlder states,) in which nutritious matter is stored up. This matter may be deposited in such a way as to give the appearance of striæ surrounding a point or hilum, which is considered as an opening into the cell. The grains of starch are well seen in the cells of the potato (fig. 34).

Fig. 31.—Transverse section of cells from pulp of Pear.

Fig. 32.—Longitudinal section of the same.

Fig. 33.—Nucleated cells from the Beet.

In wheat (fig. 35), and in maize (fig. 36), the form of the granules, and the successive layers of deposit, are also seen. The grains in the



stem of *Nuphar luteum* show the centripetal formation, that is, the increase by layers deposited within each other. The addition of iodine causes the grains of starch to assume a blue colour, and marks the difference between them and the walls of the cell containing them. Schleiden affirms starch to be the most widely diffused substance in the vegetable kingdom; its presence may be regarded as in a measure indicating the age of the cell. With its formation in many cells, we have a limitation of vital activity, by which the organism is brought into a condition wherein the power of germination may be preserved for an extreme period.

18. **Crystals** are found in the interior of cells. They probably owe



their origin to the union between the acids produced or taken up by plants, as oxalic, phosphoric, malic and carbonic, and the alkaline matter, as lime and potash, absorbed from the soil and circulating in the sap. The crystals usually lie loose in the cells (figs. 37, 38); but, according to Payen, they are sometimes found in a distinct tissue, and suspended from the wall of a large cell (fig. 39)—filling what some have supposed to be the base of an undeveloped hair. The crystals are of different sizes and forms. Occasionally, a single large crystal nearly fills a cell, but in general there are numerous crystals united together. Sometimes the crystals radiate from a common point (figs. 40, 41), and form a conglomerate mass; at other times they lie parallel, and have the appearance of bundles of fine needles (figs. 37, 38). To the latter, the names of *Raphides* ($\rho\alpha\phi\iota\varsigma$, a needle), or *acicular* crystals (*acus*, a needle), was originally given. It has been said that these crystals exist also in the intercellular spaces; but this seems to depend on the mode in which the section of the plant is made, for when raphidian cells (fig. 42 *rrrrr*) are situated close to a lacuna, the crystals may easily be pushed into it accidentally by the knife. Raphides consist principally of phosphate and oxalate of lime. They abound in some plants, especially Cacti, and they are common in Squill, and in the


Fig. 34.—Cell of Potato, containing striated starch grains.

Fig. 35.—Grains of starch of Wheat.

Fig. 36.—Grains of starch of Malze.

Fig. 37.—Cellular tissue of *Aurum maculatum*. *c*, Cells containing chlorophyll. *rr*, Raphidian cells.

official Turkey Rhubarb, the latter of which owes its grittiness to their presence. One hundred grains of rhubarb root contain about 30 or



40 grains of oxalate of lime crystals. Acicular crystals may be easily seen by making a section of any Lilaceous plant, as the hyacinth, and spreading the thick mucilaginous matter of the cells on the field of the microscope. Radiating raphides are seen in the sepals of *Geranium robertianum* and *lucidum*; the crystals, consisting of oxalate of lime, fill the whole of the cells in the middle of the sepal, their size varying from $\frac{1}{2000}$ to $\frac{1}{1500}$ of an inch. Quekett found them in all the species of *Pelargonium* and *Morisonia* that he examined, and he thinks that they are as general as the beautiful markings in the cuticle of the petals of these plants. Clustered crystals have been detected in Malvaceous plants, and in the sepals of the strawberry; numerous acicular crystals have been observed in *Fuchsias*, and solitary cubical crystals in the superficial cells of the sepals of *Prunella vulgaris* and *Dianthus caryophyllus*.^r

In the outer covering of the seed of *Ulmus campestris*, the sinuous boundaries of the compressed cells are traced out completely by minute rectangular crystals adhering to each other. Unger detected oxalate of lime crystals in *Ficus bengalensis* and *Calathea zebrina*.

19. **Chlorophylle** (χλωρὸς, green, and φύλλον, a leaf), or the green colouring matter of plants, floats in the fluid of cells, accompanied by starch grains. It differs from starch in being confined to the superficial parenchyma, and in being principally associated with the phenomena of active vegetable life. It has a granular form (fig. 39 *u*, 42 *c*), is soluble in alcohol, appears to be analogous to wax in its composition, and is developed under the agency of light; subterranean




Fig. 38.—Cells of *Aurum maculatum*. Clusters of raphides in a large oval cell surrounded by smaller cells.

Fig. 39.—Cellular tissue, from leaf of *Ficus elastica*. *c*, A large cell. *r*, An agglomeration of crystals suspended in a sac by a tube, *t*. *u*, Utricles filled with grains of chlorophylle.

Fig. 40.—Cells of Beet with conglomerate radiating crystals, *a*. *b*, Separate crystals of different forms.


Fig. 41.—Conglomerate crystals of oxalate of lime from Rhubarb.

Fig. 42.—Cellular tissue of *Colocasia odora*. *c c*, Cells with grains of chlorophylle. *r r r r*, Raphidian cells projecting into alacuna or intercellular space.

vegetable structures which present a white appearance, and possess a bland taste, become altered in both respects, when freely exposed to the action of light and air. Its granules are usually separate, but sometimes they unite in masses (fig. 37 c). Other kinds of colouring matter are also produced during vegetation, and occur in the form of fluids or of granules in the interior of cells.

20. **Oils and Resinous matter** are found in the interior of cells, as well as in intercellular spaces. The cavities containing them are denominated *cysts*, *reservoirs of oil*, and *receptacles of secretions*. They are easily detected in the rind of the orange and lemon, in the myrtle tribe, and in *Hypericum*. When small portions of the fresh leaf of *Schinus molle* are thrown on water, the resinous matter, by its rapid escape, causes them to move by jerks, and the surface of the fluid is covered with the exudation. In the bark of the Fir tribe there are cavities with thick walls containing turpentine. In the fruit of *Umbelliferæ*, canals occur called *vittæ* (*vitta*, a head-band, from surrounding the fruit), containing oil.

21. **Air cells**, or cavities containing air, consist either of circumscribed spaces surrounded by cells (fig. 43), or of lacunæ formed by the rupture or disappearance of the septa between a number of contiguous cells, as in grasses, *Equisetum*, Umbelliferous plants, and pith of Walnut. They are often large in aquatic plants, and serve the purpose of floating them, as in *Pontederia*, *Trapa*, *Aldrovanda*, and sea-weeds.



3.—DEVELOPMENT AND FUNCTIONS OF CELLS.

22. The subject of *Cell-development*, or *Cytogenesis* ($\kappa\acute{\upsilon}\tau\omicron\varsigma$, a cell, and $\gamma\acute{\iota}\nu\epsilon\sigma\iota\varsigma$, origin), which has given rise to great diversity of opinion among physiologists, is still involved in much obscurity. By some it is affirmed that the first appearance of vegetable tissue is in the form of a mucilaginous fluid, which, gradually thickening, becomes hollowed into a number of small cavities constituting the future cells. Schleiden believes that the cell is formed from the nucleus, to which he gave the name of *Cytoblast* ($\kappa\acute{\upsilon}\tau\omicron\varsigma$, a cell, and $\beta\lambda\alpha\sigma\tau\omicron\varsigma$, a germ), or *cell-germ*, from its supposed generative function. This cytoblast, according to him, is the part first formed. It acts by attracting the mucilaginous matter in which it lies, and forming around itself a sort of gelatinous covering. There is thus produced round the nucleus a closed utricle, which increases in size by the assimilation of the fluid in which it is placed. The development usually takes place on one side, the new cell appearing in the form of a transparent vesicle rising from the surface, and leaving the nucleus attached to the other side of the

Fig. 43.—Air-cells in *Ranunculus aquatilis*.

utricule (fig. 33). The cytoblast is thus enclosed in the utricule, and may ultimately disappear by absorption, leaving a non-nucleated cell. The membrane surrounding the nucleus is converted gradually into cellulose, and thus the perfect cell is formed. According to Mohl, the nucleus is at first retained in the centre of the cell by means of mucous threads, and afterwards becomes fixed to the sides. Occasionally, the nucleus becomes imbedded in a duplication of the cell-wall. This process of cell-development, according to Ascherson, is similar to what takes place when oil is mingled with a mucilaginous or albuminous fluid, each minute molecule of oil becoming surrounded by a thin film of membrane. In this view the cell is originally of a more or less globular form, and all the varieties of shape afterwards seen are due to changes in the progress of growth.

Barry affirms that a minute pellucid globule (*hyaline*) is first seen in the formative matter. This absorbs and assimilates new matter, enlarges and becomes granular, thus forming the cytoblast of Schleiden, after it has prepared a nucleolus for itself. The outer part of the cytoblast rises in the form of a membrane to produce a cell; another portion of it is concerned in the formation of the contents of the cell; and what is left of the cytoblast in the cell-wall becomes the nucleus of the cell. This nucleus (not the cytoblast of Schleiden) remaining on the cell-wall, is not absorbed, but becomes the source whence cytoblasts are formed. Thus, according to Barry, the substance of the larger body is not deposited around the smaller, but the smaller is transformed into the larger; the nucleolus becomes the cytoblast, and the cytoblast becomes a nucleated cell.

As regards the development of cells from nuclei, the present state of our knowledge does not warrant us in stating more than that there is a protoplasm, or soft organizable matter, which is contained in cells, or in the spaces between them; that in this matter a nucleus is produced, either around previously existing nucleoli, or from the granules of the protoplasm; and that the nucleus has the power of developing new cells, which become nucleated, increase in size, and escape from the parent cell, by rupture or absorption. In the production of young cells, the nucleus of the parent cell sometimes divides into two, each part having the power of giving rise to a new cell. There is thus a constant multiplication of cells by an intra-cellular or endogenous (*ἐνδον*, within, and *γεννᾶειν*, to generate) process.

23. It is supposed by some that cells may arise without a nucleus, by the simple aggregation of granular matter, which becomes enveloped in a membrane, and thus forms a cell with granular contents. In such cells, a body similar to a nucleus may be afterwards formed, and may assume the function of the cytoblast of Schleiden, as far as the subsequent endogenous development of new cells is concerned. Some physiologists maintain that the cytoblast is never concerned in

cytogenesis, but only takes part in the various chemical and other changes which occur in the contents of the cell during its growth and nutrition. Mohl and Henfrey state that new cells are produced by the division of the primordial utricle (§ 14), which gradually folds inwards about the middle, forming an annular constriction, and ultimately a complete separation of the utricle into two parts. Each of these afterwards becomes covered by a permanent cell-wall. Henfrey has supported this view by observations made on the hairs of *Tradescantia* and of *Achimenes grandiflora*, in which he has traced the gradual formation of a septum.

24. Naegele maintains that new cells are produced by the division of the primordial utricle, or mucilaginous sac, as he calls it, and its contents into two or four portions, each of which encloses a free nucleus. From each of these portions, a cell, with its outer layer of cellulose, originates, while the parent cell becomes dissolved and disappears. The outer layer of the new cells is formed, according to him, round, and by the separate portion of the divided utricle. The mode of division he does not explain. This view does not appear to differ much from that adopted by Unger, who traces in *Algæ* the development of new cells, by a *fissiparous* (*fissus*, split, and *pario*, I produce), or *merismatic* (*μεριστικός*, division) separation of the old ones into four divisions, in the same way as occurs in pollen grains. In some of the most simple plants, multiplication takes place by a sort of sprouting of new cells from old ones, like buds from a stalk.

25. The various theories of cell-development may be therefore reduced to the following: 1. The Endogenous formation, within a parent cell; 2. the Exogenous, without, or on the outside of the cell; 3. Merismatic, or by division of cells; 4. Isolated, or the independent formation of cells in a protoplasm; and 5. Cell formation by *conjugation*, in which the cells, being similar, those uniting may be either free and isolated, as in most *Diatomaceæ*, and *Desmidiaceæ*, or they may be links of a many-celled filament, as in the *Zygnemaceæ*. Naegele asserts the conjugations of dissimilar cells in the *Vancheria*, which results in the formation of spores.*

26. The formation of cells from nuclei, and their fissiparous division, are by some attributed to different electrical currents excited by the chemical actions going on in the cell. Cells are produced with great rapidity, especially in the case of fungi. Lindley calculates that the cells of *Bovista gigantea* have been produced at the rate of more than sixty-six millions in a minute, and Ward has noticed a similar occur-

* For a full view of the subject of the development and growth of cells, the following works may be consulted:—Schleiden on *Phytogenesis*, and Mohl on the *Structure of the Vegetable Cell*, translated in *Taylor's Scientific Memoirs*, Vols. II. and IV.; Naegele on *Vegetable Cells*, *Ray Society's Reports*, 1846; Sharpey, *Anatomy*; M. Barry, *Physiology of Cells*, &c., in *Philosophical Transactions*, 1840; and on *Nucleus of Cells*, in *Jameson's New Philosophical Journal* for September, 1847; *Carpenter's Physiology*; Brown on the *Phenomenon of Rejuvenescence in Nature*, *Ray Society's Reports*, 1853.

rence in *Phallus impudicus*. In warm climates, at the commencement of the wet season, the production of cells in the higher classes of plants proceeds with astonishing rapidity.

27. The organized cells of plants appear to be the more immediate seats of the various changes which constitute the functions of nutrition and reproduction. In cellular plants they are the only form of elementary tissue produced throughout the whole of life. They absorb nourishment through their walls, elaborate secretions, and give rise to new individuals. In the newly-formed tissue of vascular plants, cells alone at first exist. Fluid matters are absorbed by them, and are transmitted from cell to cell by a process of transudation. The name of Endosmose (ἐνδον, inwards, μάα, μᾶ, I seek), and Exosmose (ἐξω, outwards), were given by Dutrochet to the process of transudation, which leads to the motions of fluids of different densities placed on opposite sides of animal and vegetable membranes. This process appears to be of universal occurrence in plants, being concerned in the movements of the sap, the opening of seed-vessels, and many other phenomena. The capsule of the Elaterium, for instance, opens with great force by a process of endosmose going on in the cells, and such is also the case with that of the Balsam. The power which cells possess of absorbing fluids is well seen in sea-weeds, which, after being dried, can easily be made to assume their natural appearance by immersion in fluids. It is also observable in the spores of the Equisetum, the teeth of Mosses, the seed-vessels of some Fig-mari-golds, the Rose of Jericho (*Anastatica*), and some Lycopodiums.

Various organic secretions, which are necessary for growth and nourishment, are formed by the internal membrane of cells. It is in cells that the azotised and unazotised matters are deposited, which are afterwards applied to the purposes of vegetable life. In them we meet with the proteine compounds, albumen, fibrine, and caseine, consisting of carbon, oxygen, hydrogen, and nitrogen, with proportions of sulphur and phosphorus; as well as starch, gum, sugar, oil, and colouring matters, in which no nitrogen occurs. Some of the organic matters found in plants have been artificially formed by chemical means, while others have only as yet been met with in the living organism. Spiral cells sometimes contain air.

SECTION II.—VASCULAR TISSUE.

1.—FORM AND ARRANGEMENT OF VESSELS.

28. **Vascular Tissue**, or *Angienchyma* (ἀγγος, a vessel), consists of tubes, whose length greatly exceeds their breadth. These may be formed of membrane only, or of membrane altered in various ways by deposits of fibre, or thickening matter in the interior.

29. **Woody Fibre, or Ligneous Tissue.** *Pleurenychyma* (πλευρά, a rib, from its firmness), (fig. 44,) consists of tubes, or, according to some, elongated cells, of a fusiform (*fusus*, a spindle) or spindle-like shape (fig. 3), having their walls thickened so as to give great firmness. This form of tissue does not exist in cellular plants. Some have called this tissue *Proenchyma* (πρός, close to, in reference to the close apposition of the tubes), a term, however, generally applied to



shortened fusiform cells only. Pleurenychymatous vessels lie close together, overlap each other, and, by their union, give strength and solidity to the plant. Their membrane becomes thickened by successive deposits of layers of cellulose and sclerogen, and in a transverse section the tubes present the appearance of concentric circles, occasionally with intervals, where the ligneous matter is deficient (fig. 45). The wood of trees is made up of fibres or tubes of this kind, and they are found in the inner bark, and in the veins of leaves. The woody fibres may be separated from the cellular parts of plants by maceration. In this way Flax and Hemp are procured, as well as the Bast used for mats. The strength of the woody fibres of different plants varies. Thus, New Zealand Flax, the produce of *Phormium tenax*, is superior in tenacity to Common Hemp; while the latter, in its turn, excels Common Flax, as well as Pita Flax, which is the produce of *Agave americana*. Linen is formed from woody tissue. Cotton, on the other hand, consists of elongated cells or hairs, the membrane of which becomes contracted in the process of drying, so as to appear twisted when viewed under the microscope. By this character mummy cloth was shown to be composed of linen. Woody fibres, in fabric, form muslin, lace, &c., some fine India muslins only are formed from woody fibre; other muslins are made of cotton; when reduced to small fragments, they constitute the pulp whence paper is made.

30. In its ordinary form, *Pleurenychyma* has no definite markings on its walls; but in some instances these present themselves in the form of simple discs (fig. 46), or of discs with smaller circles in the centre (fig. 47). The latter occurs in the wood of Firs, Pines, and Winter's bark, and has received the name of *glandular* or *punctated* woody tissue. These markings are formed by concave depressions on the outside of the walls of contiguous tubes, which are closely applied to each other, forming lenticular cavities between the vessels, like two watch-glasses in apposition, and when viewed by transmitted light they appear like discs (fig. 46). In the centre of the depression

Fig. 44.—Woody fibres (*Pleurenychyma*), from *Clematis vitalba*.


Fig. 45.—Transverse section of the same.

there is a canal, often funnel-shaped, and the part of the tube corresponding to it being thus thinner than the surrounding texture, gives the aspect of a smaller circle in the centre (fig. 47). When a thin section is made through two parallel lines of punctations, the slits or fissures are seen which give rise to the appearances mentioned (fig. 48). That these markings are cavities between the fibres was proved by Quekett in the case of fossil pine wood, where he separated lenticular masses of solid matter from the discs. There is sometimes observed a thickening layer, in the form of a spiral fibre, surrounding the discs more or less completely. The discs are usually arranged in single rows, but they occur also in double and triple rows, more particularly in *Araucaria* and *Altingia*.


31. **Fibro-Vascular Tissue**, or *Trachenchyma* (*trachea*, windpipe; *τεαχύς*, rough), is formed of membranous tubes tapering at each end, less firm than *Pleurenchyma*, and either having a fibre coiled up spirally in their interior, or having the membrane marked with rings, bars, or dots, arranged in a more or less spiral form.

32. **True Spiral Vessels**, (*spiroidea*, *tracheæ*), constituting the typical form, present themselves as elongated tubes clustered together, overlapping each other at their conical extremities, and having a spiral fibre or fibres surrounding the interior of the cylinder (fig. 49). Their outer membrane is thin, and consists of pure cellulose. At the point where they overlap, it is sometimes absorbed, so as to allow direct communication between the vessels. The fibre or spiral filament is generally single, forming *simple tracheæ* (fig. 50); but sometimes numerous fibres, varying from two to more than twenty, are united together, assuming the aspect of a broad ribband (fig. 51), and constituting *Pleio-tracheæ* (*πλειών*, more). The fibre is elastic, and can be unrolled. This can be seen by taking the leaf of a *Pelargonium*, and after making a superficial cut round the stalk, pulling the parts gently asunder, when the fibres will appear like the threads of a cobweb.


33. Spiral vessels were first noticed as early as 1661, by Henshaw

- 
- Fig. 46.—Woody fibres with circular spots where the membrane is thin (*Bignonia*).
 Fig. 47.—Punctated woody tissue, with a double circle or disc, from common Scotch fir.
 Fig. 48.—Longitudinal section of the same, showing the union between the fibres and the mode in which the circles are formed.
 Fig. 49.—Two spiral vessels united.
 Fig. 50.—Simple trachea.
 Fig. 51.—Spiral vessel with a ribband of united fibres (*Pleio-trachea*) from the Banana.

They occur principally in the higher classes of plants, and are well seen in annual shoots, as in *Asparagus*; in the stems of *Bananas* and *Plantains*, where the fibres may be pulled out in handfuls, and used as tinder, in many aquatics, as *Nelumbium* and *Nymphæa*, and in *Liliaceous* plants. In hard woody stems, they are principally found in the sheath



52



53

surrounding the pith, and they are traced from it into the leaves. They are rarely found in the wood, bark, or pith. Spiral vessels occasionally exhibit a branched appearance. This may arise from the union of separate vessels in an angular or jointed manner, as where a leaf or branch is given off (fig. 52 *a a*), or it may depend on a regular division of the fibres, as is seen in the *Mistletoe*, *Longleek*, and *Gourd* (fig. 53).

34. The fibre is on the inside of the membrane. Quekett has shown this in silicified spiral vessels, where the mark of the spiral was on the outside of the mineral matter filling the tube. The fibre usually turns from left to right, if we suppose the observer placed in the axis of the tube (fig. 54), or from right to left, if we suppose him looking at the vessel in its natural position. The fibre retains its direction throughout the length of the vessel. When examined under the microscope, there is often the appearance of the crossing of fibres (fig. 54), in consequence of the transparency of the membrane, and the observer seeing the fibre on each side of the vessel at the same time. In twining plants, the direction of the fibre does not always correspond with that of the stem. The coils of the spiral fibre may be close together (fig. 50), or be separated (fig. 55). Sometimes they become united together, and to the membrane of the tube, so that they cannot be unrolled. Such vessels are called *closed* trocheæ, or *closed* ducts.



54



55

35. **False or Spurious Tracheæ**, the *ducts* of some authors, are vessels in which the internal fibre does not form a complete spiral coil. The chief varieties are *annular*, *reticulated*, and *scalariform* vessels, or ducts. In annular vessels (*annulus*, a ring), the fibres form complete rings round the tubes (fig. 56). They resemble the tracheæ of animals more than spiral vessels do. The rings are by no means regular; they may be horizontal or inclined, simple, or forked (fig. 57), placed near to each other or separated by considerable intervals, the inter-


Fig 52.—Spiral vessels, united so as to have a branched appearance.

Fig. 53.—Branching fibre, from spiral vessels of Gourd (*Cucurbita Pepo*).

Fig. 54.—Spiral vessels. Coils seen on both sides.

Fig. 55.—Coils of fibre, much separated in trachea of Gourd.

mediate spaces being sometimes occupied by a fibre of an elongated spiral form, which is continuous with the rings or distinct from them (fig. 58). All these forms are easily recognized in the common Balsam. Occasionally, the ring becomes very much thickened in a direction perpendicular to the walls of the vessel, so as to leave only a small space in the centre, as in some of the Cactus tribe. When separate fibres cross each other, forming a kind of net-work on the walls of the tubes (fig. 59), the vessels become *reticulated* (*reticulum*, a net); and the name *dotted* is sometimes applied when the fibre is so broken up as to leave small isolated portions adhering to the membrane (fig. 60). In scalariform vessels (*scala*, a ladder), there are short horizontal lines or bars, composed of fibre, arranged along the sides of the tubes, at nearly equal distances, like the steps of a ladder, and presenting a striated appearance. In some cases, as in the Vine (fig.



61), they are composed of tubes united to each other by thin, broad, oblique extremities; at other times they taper like spiral vessels. They generally assume a prismatic form, the angles being unmarked by lines, as is seen in Ferns (fig. 62).

36. Porous Vessels.—Another kind of vessel common in plants is the *porous* vessel, so called from the appearance of pores on its surface.

Figs. 56, 57, 58.—Annular vessels from the stem of the Common Balsam.

Fig. 59.—Spiral vessel. Wide coil, and fibre dividing.


Fig. 60.—Vessel showing rings of fibre and dots.

Fig. 61.—Scalariform vessel from the Vine.

Fig. 62.—Prismatic scalariform vessel from Royal Fern (*Osmunda regalis*).

The tissue formed by porous vessels has received the name of *Vusiform tissue*, *Pitted tissue*, *Bothrenchyma*, or *Taphrenchyma* (βόθρος or τάφος, a pit). The vessels are of large size, and are easily observed in the Vine (fig. 63), Sugar Cane, Bamboo, Gourd (fig. 98 *ter*), and other plants, in which the sap circulates rapidly. They consist of cylinders more or less elongated, in which the thickening matter is so deposited as to leave part of the membrane uncovered, thus giving rise to the porous or pitted appearance. The uncovered portions of membrane are

sometimes absorbed in old vessels, and a direct communication is established between them. The pores have sometimes a boarded aspect, which, according to Schleiden, depends on air contained in the cavities between contiguous vessels. Porous vessels are usually united to each other by a broad and often oblique septum.



37. This kind of vessel occasionally presents a beaded appearance, as if formed by porous cells, with distinct constrictions at their point of union (figs. 64, 65). This articulated *Bothrenchyma* is by some considered as a form of cellular tissue (§ 10, fig. 22). To vessels exhibiting contractions of this kind, whether spiral or porous, the terms *moniliform* (*monile*, a necklace), or *vermiform* (*vermis*, a

worm), have been applied; and the tissue composed of spiral, annular, or porous moniliform vessels, has been denominated *phleboidal* (φλέψ, φλεβός, a vein).


38. **Laticiferous vessels** form the tissue called *Cinenchyma* (κινέω, I move, from certain movements of their contents, to be afterwards noticed). They are the Milk-vessels, and the Proper vessels of old authors; and of late years they have been particularly examined and described by Schultz. They consist of long, branched, homogeneous tubes, having a diameter of about $\frac{1}{1400}$ of an inch, which unite or anastomose freely (fig. 66), thus resembling the vessels of animals. At first the tubes are very slender and uniformly cylindrical (fig. 67 *a*), but afterwards they enlarge and present irregular distensions at different parts of their course (figs. 67 *b*, 68), so as to give rise to an articulated appearance. Their walls vary in thickness, and are not

Fig. 63.—Porous vessel (*Bothrenchyma*) from the Vine, showing its connection with woody fibres, and the broad septa or partitions of the vessel itself.

Fig. 64.—Porous vessel from Traveller's Joy (*Clematis vitalba*).

Fig. 65.—Moniliform porous vessels from the Common Balsam.

marked by any depressions or fibres. These vessels are met with in the inner bark, and they contain a granular fluid called *latex*, which is at first transparent, but often becomes of a white, yellow, or reddish colour. Endlicher and Unger state that they are formed by cells united in a linear series, their septa being obliterated; while Meyen and Schleiden maintain, that at a very early period the currents of latex may be seen in the intercellular canals, and that ultimately a



separate membrane is developed to form the vessels. The tissue can be easily examined in the India-rubber tree, in Dandelion, Lettuce, and Celandine, and in various species of *Ficus* and *Euphorbia*.

2.—DEVELOPMENT AND FUNCTIONS OF VESSELS.

39. The simple cell is the state in which vegetable tissue first makes its appearance. It is the primary form of all the textures subsequently produced in vascular plants. To the elongation of cells, and the deposition of thickening layers and fibres in their interior, the various vessels owe their origin. Thus when cells are developed as continuous branching tubes, which anastomose freely, *Cinenchyma* is formed; when they are elongated, as spindle-shaped tubes, and their walls are thickened and hardened by depositions of ligneous matter, they give rise to *Pleurenchyma*; and when elongated membranous tubes are strengthened by spiral fibres, the different kinds of *Fibro-vascular* tissue are produced. The spiral vessel may be considered as the type of the last-mentioned tissue, and all its varieties may be traced to changes

Fig. 66.—Laticiferous vessels (*Cinenchyma*) from *Euphorbia dulcis*.
Figs. 67, 68.—Vessels of Latex from Celandine (*Chelidonium majus*).

taking place in the development of the fibre. The coil may be broken in consequence of the fibre adhering to the membrane, and the latter increasing rapidly in growth; or the fibre may be deposited irregularly, in consequence of interruptions in growth. This view of the formation of vessels is confirmed by finding in the same tube a complete spiral fibre in one part, annular fibres, either complete or with their ends overlapping, at another, and bars or dots at a third portion. In the case of some vessels, their formation can be distinctly traced to cells placed end to end, the partitions between which have been absorbed. The moniliform or beaded appearance often presented by the different kinds of vessels, more especially the Porous, plainly indicates this mode of formation.

40. As in cells, so in vessels, the walls are composed of cellulose, and there are usually no visible perforations; the communication between them taking place by imbibition or endosmose. In some instances, when vessels are closely applied to each other, especially when they overlap, the membrane becomes absorbed, and direct communication takes place. This has been seen in spiral and porous vessels. The pits or depressions on the walls of vessels, and the thinning of the tissue at particular points, appear to serve the purpose of allowing the rapid transmission of fluids; and, according to some, they permit the passage of small cells from the interior, which become developed as tubes, and form branching vessels.

41. Pleurencyhma, in its early state, contains fluids, and conveys them from one part of the plant to another. In the progress of growth, the secondary deposits obliterate the vessels, as in the perfect or heart wood of ordinary trees. These deposits are often of a very hard nature, and assume particular colours in different kinds of trees. From the firmness of this tissue, it is well fitted to give solidity to the stems and to strengthen the leaves of plants. In Spiral vessels, the fibre adds to their elasticity, and keeps the tubes always pervious. The fibre, when once formed, does not increase much in thickness, and the secondary deposits do not obliterate the canal. Various opinions have prevailed regarding the contents of these vessels. The name *Tracheæ*, given by Grew and others, was partly from their structure, and partly from the idea that they contained air. The accurate experiments of Bischoff lead to the conclusion that the perfect spiral vessels convey air, which often contains a large amount of oxygen in its composition. Hales showed that air was evolved from the vessels of the Vine when cut, and Decandolle thought that part of the air in these vessels was derived from the pores of the leaves. Other authors look upon these vessels as conveying fluids, while a third set maintain that both air and fluids are present, the air being derived in part from decompositions going on in the interior of the plant. The other kinds of vascular tissue, and especially the porous vessels, are the means

by which the fluids taken up by the roots of plants are conveyed to the leaves, and to all parts of the plants. Laticiferous vessels contain, according to Schultz, the elaborated sap or latex on its return from the leaves to the bark. This latex is either transparent or opaque, colourless or coloured. These vessels, when examined with the microscope in the living plant, exhibit movements in their fluid contents of a peculiar kind, which will be considered under *Cyclosi*.

42. The cell has been already shown to be the type of all the tissues of plants, and to be the basis of all vegetable structure. It is of equal importance as regards function. In the lowest plants, as the *Protococcus nivalis*, or the Alga found in red snow, and the various species of *Palmella*, *Nostoc*, and *Hæmatococcus*, cells constitute the whole substance, and perform all the functions of life; they absorb and assimilate, thus performing the functions of nutrition and secretion, and they form new cells, thus reproducing individuals like themselves. When a more complex structure exists, as in the higher tribes of plants, certain cells are appropriated for absorption, others are concerned in assimilation, and others in forming and receiving secretions. When a certain degree of solidity appears to be required to support the stem, leaves, and flowers, ligneous matter is deposited, and woody fibre formed. When the transmission of fluids and air is carried on rapidly, the elastic fibres of the fibro-vascular tissue seem to keep the elongated cells and vessels pervious, and when the elaborated sap is conveyed continuously without interruption, anastomosing tubes occur in the form of laticiferous vessels.

TABULAR ARRANGEMENT OF VEGETABLE TISSUES.

- A.—Cellular Tissue (Parenchyma), composed of membrane, or of membrane and fibre, having the form of vesicles whose length does not greatly exceed their breadth.
1. Membranous Cellular Tissue; cells formed by membrane alone, of various thickness, but without markings on it.
 2. Porous Cellular Tissue; cells formed by membrane, which has been unequally thickened in such a way as to leave rounded depressions at regular intervals.
 3. Fibrous Cellular Tissue (Inenchyma); cells formed by membrane and fibre; occasionally formed by fibre alone.
 - a. Spiral Cells, with a complete spiral fibre inside.
 - b. Dotted Cells, with opaque spots, which are isolated portions of fibre.
- B.—Vascular or Tubular Tissue (Angienchyma), composed of cylindrical tubes, which are more or less continuous, and usually overlap each other, or are united by broad oblique extremities.
- I. Membranous Vascular Tissue; tubes formed by membrane alone, of various thickness, but without markings on it.
 1. Ligneous Tissue (Pleurenchyma), composed of fusiform tubes with thickened walls.
 2. Laticiferous Tissue (Cinenchyma), composed of tubes which anastomose, often present irregular dilatations, and convey a peculiar fluid, called Latex.

- II. Porous Vascular Tissue; tubes formed by membrane, which becomes thickened by spiral deposits, in such a way as to leave rounded depressions at regular intervals.
 1. Vasiform Tissue, or Porous vessels (Bothrenchyma or Taphrenchyma); large tubes, usually ending in broad extremities, with pits or circular markings on their walls. This tissue sometimes exhibits contractions at regular intervals, as if formed of porous cells laid end to end, and then is called Moniliform, or Beaded (Articulated Bothrenchyma).
 2. Punctated Tissue (Glandular Woody Tissue); fusiform woody tubes, with depressions and markings on their walls, presenting the appearance either of a single or double circular disc.
- III. Fibro-Vascular Tissue, composed of tubes in which the thickening matter is deposited in the form of spiral fibres, rings, bars, or dots.
 - a. Perfect Fibro-Vascular Tissue, composed of tubes, in which there is a complete spiral fibre.
 1. Spiral Vessels (Tracheæ, Trachenchyma), in which the spiral fibre is elastic, and may be unrolled.
 2. Closed Spiral vessels, or closed Tracheæ, in which the spiral fibre is brittle, or its coils so united to each other, and to the membrane, that they cannot be unrolled.
 - b. Imperfect Fibro-Vascular Tissue, composed of tubes marked by rings, lines, or dots, but without a complete fibre inside.
 1. Annular Vessels or Ducts, having fibres in the form of detached rings, which are occasionally united by portions of fibre.
 2. Reticulated Vessels, having fibres which cross each other, or are disposed so irregularly as to form a net-work.
 3. Scalariform Vessels, having their walls marked by isolated portions of fibre, in the form of ladder-like bars.
 4. Dotted Vessels, having their walls marked by isolated portions of fibre in the form of opaque dots or points.

Any of the vessels included under the Fibro-vascular tissue, may exhibit contractions at regular intervals, so as to become moniliform.

CHAPTER II.

COMPOUND ORGANS FORMED BY THE TISSUES.

43. Some plants consist of cells only, which continue throughout life to produce new cells, and to perform all the vital functions. The great mass of flowering plants, however, although originally cellular, produce organs composed of cells and vessels variously arranged, and covered by an epidermis. These compound Organs may be divided into *Nutritive*, or those concerned in the nourishment of the plant; and *Reproductive*, or those which are employed in the production of new individuals. The former consist of the stem, root, and leaves; the latter, of the flower and fruit.

SECTION I.—ORGANS OF NUTRITION OR VEGETATION.

1.—STRUCTURE, ARRANGEMENT, AND SPECIAL FUNCTIONS.

44. Under this head will be considered the tissues of which the various nutritive organs are composed, the mode in which the parts are arranged, and the particular function which each of the organs performs.

GENERAL INTEGUMENT.

45. **General Integument** is the name given to the external cellular covering of plants. It can be easily detached from young leaves and stems, usually in the form of a colourless transparent membrane. By prolonged maceration it has been shown to consist frequently of two layers; a superficial, called *Cuticle* or *Pellicle* (fig. 69 *pp*), and a deep layer, usually called the *Epidermis* (fig. 69 *ee*).

46. **The Superficial Pellicle** (*pellis*, skin) is a very thin continuous membrane, which is spread over all parts except the openings of the stomata; in some cases entering these openings, and lining the cavities beneath them. It is formed from the epidermal cells below it; Treviranus, Schleiden, and Payen, considering it as a secretion on the outside of the cells, while Mohl and Henfrey look upon it as composed of the altered primary walls of the cells. Mitscherlich regards it as a corky substance which preserves the humidity of the plant, in preventing the evaporation of moisture. This substance is by him considered as next to cellulose, the most important constituent of the cell-wall. In fig. 70, it is represented as detached from the leaf of the cabbage, forming a sheath over the hairs, *h h h h*, and leaving slits, *s s*, corresponding to the openings of the stomata. This pellicle appears to be similar to the intercellular substance surround-




Fig. 69.—General integument of a leaf of *Iris germanica*. *pp*, The Cuticular pellicle with slits, *f*, lying upon the proper epidermis, *ee*, formed of hexagonal cells, and furnished with stomata, *s s*.

Fig. 70.—Pellicle of Cabbage, detached by maceration, covering the hairs, *h h h h*, and having openings, *s s*, corresponding to the stomata.

ing cells, and to the *definite mucus* which is conspicuous in some seaweeds (fig. 29 *b*). It is possible that this matter, in place of being produced on the outside of cells, may be formed within them, and ultimately deposited externally by passing through their parietes. On the inner surface of the pellicle the impressions of the epidermal cells are sometimes observed (fig. 69 *p*). The pellicle is the only layer of integument which is present in aquatic plants, and in some of the lower tribes.

47. **The Epidermis** (*ἐπί*, upon, and *δέρμα*, skin), (fig. 69 *e e*), is extended over all the parts of plants exposed to the air, except the stigma. The internal cavities of seed-bearing organs are lined by a delicate membrane, termed *Epithelium*. On the extremities of newly-formed roots, the integument consists of loose cells, which are considered either as being the ordinary cellular tissue of the plant, or as being an imperfectly-formed epidermis, which has received the name of *Epiblema* (*ἐπί*, upon, and *βλήμα*, wound, as being the tissue which first covers wounds). This latter kind of tissue occupies the place of the epidermis, in the parts of plants which are always under water. On the aerial roots of Orchidaceous plants, there is an epidermal layer consisting of spiral cells (fig. 23), containing air.

48. The epidermis is usually formed by a layer or layers of compressed cells, which assume a more or less flattened tabular shape, and have their walls bounded by straight or by flexuous lines. Fig.



e

71


69 *e e*, represents an epidermis formed of regular hexagonal cells; fig. 72, one composed of irregular hexagons; while in fig. 71, the boundaries of the cells, *e*, are flexuous and wavy. The cells of the epidermis are so intimately united together, as to leave no intercellular spaces (fig. 74 *e e*).

49. The epidermis is sometimes thin and soft, at other times dense and hard. In the former case it may be easily detached from the


subjacent cells; in the latter, the cells become thickened by deposits, and sometimes the layers are so produced as to leave uncovered spots, which communicate with the interior of the cell by canals passing through the thickening layers, as in *Cycas*. In *Rochea falcata* (fig. 73), the epidermis, *e e*, consists of two layers of cells—the outer ones

Fig. 71.—Epidermis, from lower surface of the leaf of Madder (*Rubia tinctorum*). *e*, Cell of the Epidermis. *s*, Stoma.

large, the inner small. The cells of epidermis are usually filled with colourless fluid, but they sometimes contain resinous and other substances. Waxy matter is occasionally found in the epidermis, silica is



72



73

met with in the integument of grasses and Equisetums, and carbonate of lime in that of Chara. The colour of the epidermis generally depends on that of the subjacent parenchymatous cells, from which it can be separated as a colourless layer. The epidermal cells are usually larger than those of the tissue below them; but sometimes, for instance in *Ficus elastica*, they are smaller.

50. **Stomata** (στόμα, a mouth) are openings existing between some of the cells of the epidermis on parts exposed to the air. They consist usually of two semilunar cells surrounding an oval slit or orifice (figs. 69 *s s*, 71 *s*), which have been considered as resembling the lips and the orifice of the mouth. Stomata open or close according to the state of moisture or dryness in the atmosphere. This alteration results from the hygroscopic character of these marginal cells. By examining, under the microscope, thin strips of epidermis in a moist and dry state, it will be seen that in the former case the lips are distended, they assume a crescentic or arched form, and leave a marked opening between them; while in the latter, they approach each other, and close the orifice.

51. The cells surrounding the openings of stomata are sometimes more numerous, as in *Marchantia*. In *Ceratopteris thalictroides*, Allman observed stomata formed by three cells; two of which, in their open condition, are crescentic and concave inside, while the third surrounds them, except at a small space at the end of the long axis of the stoma, and has on this account been called *peristomatic* (περί, around). In *Equisetum*, the stomata, which are about $\frac{1}{200}$ of an inch in their greatest diameter, consist of four pieces; two of which are


Fig. 72.—Epidermal layer, from upper surface of a leaf of *Ranunculus aquatilis* when growing out of water. *ee*, Epidermal cells. *ssss*, Stomata.

Fig. 73.—Vertical section of lower epidermis of the leaf of *Rochea falcata*. *ee*, Double epidermal layer, with very large external cells, small internal ones, pierced by a stoma, *s*, which communicates with a lacuna, *l*. *p*, Parenchyma of the leaf.

arched and thick at their outer convex margin, becoming thin at their inner concave edge, where two other bodies occur, having numerous processes like the teeth of a comb, hence called *pectinate* (*pecten*, a comb). Occasionally the stomatic cells become united, so as to appear in the form of an uninterrupted rim; and at other times the stoma is a minute orifice in the walls of a cavity.

52. Stomata communicate with intercellular spaces (figs. 73 *s*, 74 *s*), the connection being sometimes kept up by means of a funnel-shaped prolongation of the cuticle, called, by Gasparrini, a *cistoma* (*κίστη*, a cyst or bag, and *στόμα*, a mouth). They are scattered over the surface of the epidermis in a variable manner. Sometimes they are placed at regular intervals corresponding to the union of the epidermal cells (fig. 69 *s*); at other times they are scattered without any apparent order (figs. 71, 72); and in other instances they are united in sets of two or three, or in clusters at particular points, as may be seen in *Begonia*, *Saxifraga* (fig. 75 *s s*), and *Proteacæ*.

53. Stomata occur on the green parts of plants, especially on the leaves and their appendages. They are not usually found in cellular plants, nor in plants always submerged, nor in pale parasites. This is




not, however, a universal rule, for stomata have been detected in *Marchantia* and some other *Cellulares*; also in the submerged leaves of *Eriocaulon setaceum*, according to Griffith, and in the pale parasite *Orobanche Eryngii*, according to Duchartre. They do not exist in roots, nor in plants kept long in darkness so as to be blanched or etiolated, and they are rare or imperfectly developed in succulent plants.

Fig. 74.—Vertical section of epidermis, from the lower surface of the leaf of Madder, showing the intimate union of the epidermal cells, *e e*, the loose subjacent parenchyma, *p*, with intercellular canals, *m*, and lacuna, *l*. *s*, Stoma.

Fig. 75.—Epidermis of leaf of *Saxifraga sarmentosa*, showing clusters of stomata, *s s*, surrounded by large epidermal cells, *e e*. The cells among which the stomata occur are very small.

54. Stomata vary in their form. The oval form is very common, and may be easily seen in Liliaceous plants; the spherical occurs in *Oncidium altissimum* and the Primrose, the quadrangular in *Yucca* and *Agave*. In the *Oleander*, in place of stomata, there are cavities in the epidermis protected by hairs (fig. 76 s).



55. The development of stomata has been traced by Mirbel and Mohl. In the *Hyacinthus orientalis*, they appear first between the epidermal cells in the form of quadrangular spaces containing granular matter, which gradually collects towards the centre, where a septum or partition is formed. This septum ultimately splits, leaving a slit or opening which constitutes the stoma. Mohl has traced this process throughout the same leaf in different stages of growth. In *Marchantia*, Mirbel found several tiers of cells forming the stoma, and he supposed that the opening was produced by the absorption of a central cell, leaving the others to form the rim or border.

56. The number of stomata varies in different parts of plants. They are most abundant on the under surface of leaves exposed to the air, and are often entirely wanting on the upper surface, more especially when it has a dense shining cuticle. In floating leaves the stomata, when present, are on the upper surface only. When plants usually under water are made to grow for some time in the air, their leaves exhibit stomata. When leaves grow vertically, the stomata are often equal in number on both sides. The number of stomata varies from a few hundreds to many thousands on a surface of one inch square. The following table exhibits the number of stomata in the leaves of a few plants:—

STOMATA IN ONE INCH SQUARE.

	Upper side.		Under side.
Mistletoe,	200	200
Tradescantia,	2,000	2,000
Rheum palmatum,	1,000	40,000
Crinum amabile,	20,000	20,000
Aloe,	25,000	20,000
Clove-pink,	38,500	38,500
Yucca,	40,000	40,000
Mezereon,	None.	4,000
Pæony,	None.	13,000


Fig. 76.—Vertical section of lower epidermis of the leaf of *Nerium Oleander*. *e*, Epidermis composed of several layers of cells. *p*, Parenchyma of the leaf. *s*, Cavity filled with hairs, which may represent a stoma.

	Upper side.	Under side.
Vine,.....	None.	13,600
Holly,	None.	63,600
Cherry-laurel,	None.	90,000
Lilac,	Few.	160,000

57. Appendages of the Epidermis, or Appendicular Organs.—

The epidermis frequently exhibits projections or papillæ on its surface, in consequence of some cells being enlarged in an outward direction (fig. 73 *e e*). When these assume an elongated or conical form they constitute *hairs* (*pili* or *villi*), as seen in (fig. 77 *h h h*).

Hairs, then, are composed of one or more transparent delicate cells proceeding from the epidermis, and covered with the cuticle (fig. 70). They are erect (fig. 78 *a*), or oblique, or they lie parallel to the surface, and are appressed. Sometimes they are formed of a single cell, which is simple and undivided (fig. 78 *a*), or forked (fig. 78 *b*), or branched (fig. 78 *c*); at other times they are composed of many cells either placed end to end, as in moniliform or neck-lace-like hairs (fig.



79), or united together laterally, and gradually forming a cone, as in compound hairs (fig. 80), or branched (fig. 81). When several hairs proceed from a common centre, they become stellate (*stella*, a star), or radiated (fig. 82). The latter arrangement occurs in the hairs of the Mallow tribe, and is well seen in those of *Deutzia scabra*. When stellate hairs are placed closely together, so as to form a sort of mem-

Fig. 77.—Young root of Madder, showing cellular processes, *h h h*, equivalent to hairs. *e*, The epidermal cells which are not elongated.

Fig. 78.—Hairs formed by single cells of the epidermis. *a*, Simple hair. *b*, Bifurcated hair of *Sisymbrium Sophia*. *c*, Branched hair of *Arabis alpina*.

Figs. 79—82.—Compound hairs formed by the union of several cells. *e*, Epidermis. 79. Moniliform hair, from *Lychinis chalcadonica*.

branous expansion (fig. 83), a *scale* or *scurf* is produced. To such expansions of the epidermis the name *lepis* (λεπίς, a scale), is applied, and the surface is said to be *lepidote*. These scales have sometimes a beau-



80




81



82

tiful silvery appearance, as in *Elæagnus*. Surrounding the base of the leaves of Ferns, a brown chaffy substance occurs, consisting of elongated cells, to which the name of *ramentaceous hairs*, or, *ramenta* (*ramentum*, a shaving), has been given. In Palms also a similar substance occurs, called *reticulum* (*reticulum*, a net), or *mattulla*, (*matta*, a mat). *Prickles* or *aculei*, as in the Rose (fig. 191 a), are hardened hairs connected with the epidermis, and differ from spines or thorns, which have a deeper origin. *Setæ* are bristles or stiff hairs, and the surfaces on which they occur are said to be *setose* or *setaceous*. Some hairs, as those of *Drosera*, or sundew, have one or more spiral fibres in their interior.




83

58. Various names have been given to the different forms of hairs: they are *clavate* or *club-shaped* (*clava*, a club), gradually expanding from the base to their apex; *capitate*, having a distinct rounded head; *rough* or *scabrous*, with slight projections on their surface; *hooked* or *uncinate* (*uncus*, a hook), with a hook at their apex pointing downwards and to one side; *barbed* or *glochidiate* (γλωχίς, a barb), with two or more hooks around the apex; *shield-like* or *peltate* (*pelta*, a buckler), when attached by their middle, and projecting horizontally

Fig. 80.—Partitioned, unbranched hair, from stem of *Bryonia alba* 81. Partitioned, branched hair, from flower of *Nicandra anomala*. 82. Stellate, a star-like hair, from leaf of *Althæa rosea*.

Fig. 83.—Scale or scaly hair, from leaf of *Hippophaë rhamnoides*.

on either side (fig. 84), as in many cruciferous plants; *ciliated* (*cilium*, an eye-lash), when surrounding the margin of leaves. On the pod of the Cowitch (*Mucuna pruriens*), hairs are produced with projections on their surface, which cause great irritation when applied to the skin. In Venus' Fly-trap (*Dionaea muscipula*), stiff hairs exist on the blades of the leaf (fig. 186 e), which, when touched, cause their closure.



59. Hairs occur on various parts of plants; as the stem, leaves, flowers, seed-vessels and seeds, and even in the interior of vessels. Cotton is the hair surrounding the seeds of *Gossypium herbaceum*. Hairs are developed occasionally to a great extent on plants exposed to elevated temperatures, as well as on those growing on lofty mountains. When occurring on the organs of reproduction, they seem to be connected with fertilization, as the hairs on the style of *Goldfussia* or *Ruellia*, and the retractile hairs of *Campanula*. Different parts of plants are transformed into hairs; as may be seen in the flowering stalks of *Rhus Cotinus*, and in the calyx of *Compositæ*.

60. Names are given to the surfaces of plants according to the presence or absence of hairs, as well as the nature of the hairs which cover them. The following are the more important terms:—*Glabrous*, smooth, having no hairs; *hairy* (*pilosus*), furnished with hairs; *pubescent*, covered with soft, short, downy hairs; *villous*, having long, weak, often oblique hairs; *sericeous*, covered with long, closely appressed hairs, having a silky lustre; *hispid* (*hispidus*, *hirtus*), covered with long stiff hairs not appressed; *hirsute*, having long tolerably distinct hairs; not stiff nor appressed; (*velvety*, *velutinus*), with a dense covering of short down, like velvet; *tomentose*, covered with crisp, rather rigid, entangled hairs like cotton, which form a sort of felt (*tomentum*); *woolly*, with long curled and matted hairs like wool; *bearded* or *stupose*, (*στύπη*, tow), when hairs occur in small tufts.


61. The hairs which are most frequently met with in plants are called *lymphatic*, from their not being connected with any peculiar secretion. Those, on the other hand, which have secreting cells at their base or apex, are denominated *glandular*, and are not to be distinguished from glands, under which therefore they will be considered. Lymphatic hairs occur on parts exposed to the air, and are wanting in blanched plants. On young roots, cellular projections of the cuticle are seen (fig. 77), which may be called *radical* hairs. Young leaves and buds are frequently thickly covered with protecting hairs. In this instance the hairs arise chiefly from the veins; and as the leaves increase in size, and the veins are separated, the hairs become scattered,

Fig. 84.—Peltate hair of *Malpighia*, *p p*, arising from epidermis, *e*. *g*, The gland which communicates with the hair.

and apparently less abundant. On the parts of the flower, coloured hairs occur which have been called *corolline*.

62. **Glands** are collections of cells forming secretions. The term has been vaguely applied to all excrescences occurring on the surfaces of plants. They are either *stalked* (*petiolate*, *stipitate*) or *not stalked* (*sessile*).


The former may be called glandular hairs, having the secreting cells at the apex. Stalked glands, or glandular hairs, are either composed of a single cell, with a dilatation at the apex (fig. 85 *a*), or of several cells united together, the upper one being the secreting organ (fig. 85 *b*). In place of a single terminating



85

secreting cell, there are occasionally two (fig. 85 *c*) or more (fig. 85 *d*).

Hairs sometimes serve as ducts through which the secretion of glands is discharged; these are glandular hairs, with the secreting cells at the base. Such hairs are seen in the nettle (fig. 86), in *Loasa* or *Chili* nettle, and in *Malpighia* (fig. 84), and are commonly called *stings*. In the nettle they are formed of a single conical cell, dilated at its base (fig. 86 *b*), and closed at first at the apex, by a small globular button placed obliquely (fig. 86 *s*). This button breaks off on the slightest touch, when the sharp extremity of the hair enters the skin, and pours into the wound the irritating fluid which has been pressed out from the elastic epidermal cells at the base. When a nettle is grasped with violence, the sting is fractured, and hence no injury is done to the skin. The globular apex of glandular hairs sometimes forms a viscid secretion, as in the Chinese primrose and sundew. The hairs of the latter plant, by this secretion, detain insects which happen to light on them.



86

Fig. 85.—Glandular hairs. *e*, Epidermis. *a*, Hair formed by a single cell from *Sisymbrium* chilense. *b*, Hairs formed of several cells terminated by a secreting cell, from flower-stalk of *Antirrhinum majus*. *c*, Hair composed of several cells, terminated by two secreting cells united laterally, from flower-stalk of *Lysimachia vulgaris*. *d*, Compound hair, terminated by several secreting cells united end to end, from *Geum urbanum*.

Fig. 86.—Conical hair of *Urtica dioica*, or common nettle, ending in a button or swelling *s*, with a dilatation or bulb at its base *b*, which is surrounded by epidermal cells *ue*. In the hair are currents of granular matter *ff*.

63. When glands are sessile, they consist of epidermal cells either surrounding a cavity, or enclosing small secreting cells. In fig. 87, is represented a gland taken from the flower-stalk of *Dictamnus albus*, cut vertically, to show the cavity surrounded by cells, and filled with a greenish oil; while in fig. 88, there is a gland with a short thick stalk, full of cells, taken from *Rosa centifolia*. These figures show the transition from sessile to stalked glands. Some of the superficial cells of the epidermis are sometimes slightly elevated above the rest, and contain peculiar fluids. In the Ice-plant, the appearance of small pieces of ice on the surface is produced by cells containing a clear fluid, which is said to have an alkaline reaction, while that of the tissue around the vesicles is acid; in the Chick-



pea, similar superficial cells contain a subacid fluid. Glandular depressions or pits occur, surrounded by secreting cells. At the base of the petals of the Crown-imperial, for instance, cavities are seen containing a honey-like fluid, secreted by what are called *nectariferous* glands. Cavities containing saccharine matter, surrounded by small thin-walled cells, are met with in the leaves of *Acacia longifolia*, also in *Viburnum tinus*, and *Clerodendron fragrans*. The cavities communicate with the surface of the leaves by means of canals.

Glands are occasionally sunk in the epidermis, so as merely to have the apex projecting; at other times they lie below the epidermal cells, as in the Myrtle, Orange, St. John's-wort, and Rue. In the latter case they are sometimes called *vesicular*, and are formed by cells surrounding cavities containing oil (fig. 89). When they occur in the leaves, they give rise, when viewed by transmitted light, to the appearance of transparent points or dots. *Verrucæ*, or warts, are collections of thickened cells on the surface of plants, assuming a rounded form, and containing starch or other matters. *Lenticels*, or *Lenticular glands*, are cellular projections on the surface of the bark, arising from its inner part.



64. **The Special Functions** of the Epidermis and its appendages, are to protect the parts beneath from various atmospheric and meteorological influences. In plants growing in dry climates, the epidermis

Fig. 87.—Gland from flower-stalk of *Dictamnus albus*, cut vertically, showing central cavity *l*, filled with greenish oil, and surrounded by a layer of cells *c*, which contain a red juice, and are connected with the epidermis *e*.

Fig. 88.—Gland from *Rosa centifolia*. *e*, The epidermis.

Fig. 89.—Vesicular gland from *Ruta graveolens*, or Common Rue. *g*, Gland formed by large transparent cells, surrounding a central lacuna, *l*. *e*, Epidermis from upper surface of the leaf. *uc*, *uc*, Cells filled with Chlorophyll.

is often very thick, and coated with a waxy secretion, to prevent too great transpiration or exudation of fluids. In those which inhabit humid places, the epidermis is thin and absorbent; while in submerged aquatics, there is no proper epidermal covering. The stomata regulate the transpiration, opening and closing according to the state of humidity and dryness of the atmosphere surrounding them. When a plant is growing vigorously, the constant passage of fluids keeps the regulating cells around the stomata in a distended state, and thus opens the orifice; whereas, when the circulation is languid and the fluids are exhausted, the cells collapse and close the opening. The opinion that the succulency of plants is a sort of dropsical condition, caused by the absence of stomata to carry off the fluids, has not been confirmed by observation. Hairs, according to their structure, serve various purposes. Lymphatic hairs protect the surface, and regulate evaporation. Plants thickly covered with hairs, as *Verbascum thapsus*, have been known to resist well an extended period of drought. Glandular hairs, and glands in general, form secretions which are employed in the economy of vegetation, or are thrown off as excretions no longer fitted for the use of the plant itself. Many of these secretions constitute important articles of *materia medica*. Lenticels keep up a connection between the air and the inner bark, and probably perform the function of stomata in the advanced period of the growth of the plant. They are considered, by Decandolle and others, as being the points where young roots are produced in certain circumstances, and on that account they have been called *Rhizogens* (ρίζα, a root, and γεννάειν, to produce). They are conspicuous in Willows, the young branches of which form roots very readily when placed in moist soil. Some hairs occurring on the style of plants are called collecting hairs, from the functions which they perform in taking up the pollen. In the species of *Campanula*, these hairs are so formed, that after the pollen has been discharged, their upper part is drawn within the lower. In many hairs a circulation of fluids takes place, connected apparently with their nutrition and development (fig. 86). In the moniliform purple hairs on the stamens of *Tradescantia*, or *Spiderwort*, this movement may be easily seen under the microscope. The subject of the circulation in hairs will be considered under *Rotation*.

STEM OR ASCENDING AXIS.

Forms of Stems.

65. The stem is that part of a plant which bears the leaves and flowers. It receives the name of *Caulis* in ordinary herbaceous plants which do not form a woody stem, *Truncus* in trees, *Caudex* in shrubs, *Culm* in grasses, and *Stipe* in Palms and Ferns. It is not always con-

spicuous. Plants with a distinct stem are called *caulescent*; those in which it is inconspicuous are *acaules*. Some plants are truly stemless, and consist only of expansions of cellular tissue, called a *Thallus*, and hence are denominated *Thallogens*, or *Thallophytes* (θάλλος, a frond, γένναιον, to produce, Φυτόν, a plant). They have no true vascular system, but are composed of cells of various sizes, which sometimes assume an elongated tubular form, as in *Chara*. The cells are sometimes united in one or several rows, forming simple filaments, as in *Confervæ*; or branched and interlaced filaments, as in some *Fungi*; or membranous expansions, as in *Lichens* and sea-weeds.

66. Stems have usually considerable firmness and solidity, but sometimes they are weak, and either lie prostrate on the ground, thus becoming *procumbent*, or climb on plants and rocks by means of suckers like the *Ivy*, being then called *scandent*, or twist round other plants in a spiral manner like *Woodbine*, becoming *volubile*. Twining plants turn either from right to left, as the *French bean*, *Convolvulus*, *Passion-flower*, and *Dodder*; or from left to right, as *Honeysuckle*, *Hop*, and *Tamus*. *Bryonia alba* twines from right to left, and from left to right, alternately. In warm climates, twining plants (*lianas*) often form thick woody stems; while in temperate regions they are generally herbaceous. Exceptions, however, occur in the case of the *Clematis*, *Honeysuckle*, and *Vine*, the woody stems of which have received the name of *Sarmentum* (*sarmentum*, a twig, or cutting of a vine). Some stems are developed more in diameter than in height, and present a peculiar shortened and thickened aspect, as *Testudinaria*, or *Tortoise-plant*, and *Cyclamen*.

67. Stems have a provision for a symmetrical arrangement of leaves and branches; *nodes* (*nodus*, a knot), or points whence leaf-buds are produced, being placed at regular intervals. No such provision occurs in roots which ramify irregularly, according to the nature of the soil. The intervals between nodes are called *internodes*. The mode in which branches come off from the nodes, gives rise to various forms of trees, such as pyramidal, spreading, or weeping; the angles formed with the stem being more or less acute or oblique. In the *Italian Poplar* and *Cypress* the branches are erect, forming acute angles with the upper part of the stem; in the *Oak* and *Cedar* they are spreading or patent, forming nearly a right angle; in the weeping *Ash* and *Elm* they come off at an oblique angle; while in the weeping *Willow* and *Birch*, they are pendulous from their flexibility. The comparative length of the upper and under branches, also gives rise to differences in the contour of trees, as seen in the conical form of *Spruce*, and the umbrella-like form of the *Italian Pine* (*Pinus Pinea*).

68. Plants which form permanent woody stems above ground, are denominated *trees* and *shrubs*, while those in which the stems die down to the ground, and are not persistent, are called *herbs*. The


term *tree* (*arbor*) is applied to those plants which have woody stems many times exceeding the height of a man, the lower part free from branches being the trunk: a *shrub* (*frutex*) has a stem about three times taller than a man, and branches from near the base: an *under-shrub* (*suffrutex*) does not exceed the length of the arm; while a *bush* (*dumus*) is a low diminutive shrub, with numerous branches near the base. The terms *arborescent*, *fruticose*, *suffruticose*, and *dumose*, are derived from these.

69. Stems have usually a round form; but they are sometimes compressed or flattened laterally, while at other times they are angular: being *triangular*, with three angles and three flat faces; *trigonus* (*τρεῖς*, three, and *γωνία*, an angle), with three convex faces; *triquetrous* (*triquetrum*, a triangle), with three concave faces; *quadrangular*, or square; *quinguangular*, or five-angled; *octangular*, or eight-angled, &c.

70. The stem has been called the *ascending axis*, from being developed in an upward direction. It does not, however, always ascend into the air; and hence stems have been divided into *aerial*, or stems which appear wholly or partially above ground; and *subterranean*, or those which are entirely under ground. The latter are often called roots, but they are distinguished by producing leaf-buds at regular intervals.

The following are some of the more important modifications of stems:—The *Crown of the root* is a shortened stem, often partially under ground, which remains in some plants after the leaves, branches, and flower-stalks have withered. In this case the internodes are very short, and the nodes are crowded together, so that the plant appears to be stemless.

It is seen in perennial plants, the leaves of which die down to the ground annually. A *Rhizome* or *root-stock* (fig. 90), is a stem which runs along the surface of the ground, being partially covered by the soil, sending out roots from its lower side, and leaf-buds from its upper. It occurs in Ferns, Iris, Hedychium, &c. By many the




term *rhizome* is applied to stems creeping horizontally, whether they are altogether or only partially subterranean. A *Pseudo-bulb* is an enlarged bulbous-like aerial stem, common in Orchidaceous plants. It


contains numerous spiral cells and vessels, and is persistent. In the Kohl-rabi, a peculiar thickened rhizome. A *Soboles* is a creeping under-ground

Convallaria Polygonatum. *a*, A bud in the progress of developing a branch at the extremity of the rhizome. *c c*, Cicatrices or scars of branches which have decayed.

stem, sending roots from one part and leaf-buds from another, as in Couch grass, *Carex arenaria*, and *Scirpus lacustris* (fig. 91). It is often called a creeping root. A *Tuber* is a thickened stem produced



by the approximation of the nodes and the swelling of the internodes, as in the potato (fig. 92 *t*). Tubers are sometimes aerial, occupying the place of branches, more especially when the potato has been made to grow in darkness. The *eyes* of the potato are leaf-buds. The



ordinary herbaceous stem of the potato, when cut into slips and planted, sometimes forms branches from its base which assume the form of tubers. These tubers occasionally become nodulated, or elongated, or curved in various ways. Arrow-root is derived from the scaly tubers of *Maranta arundinacea*. A *Corm* is a solid underground stem which does not spread by sending out shoots, but remains of a rounded form, and is covered by thin scales on the outside (fig. 93). It occurs in *Colchicum*, *Tulip*, *Crocus*, and *Gladiolus*. It is distinguished from a root by sending off annually buds in the form of small corms or thickened branches, either from the apex as in

Fig. 91.—Soboles, or Creeping subterranean stem, *r*, of *Scirpus lacustris*. *se*, *se*, Scales on the stem. *pa*, Aerial portion of the plant. *t*, *t*, Level of the earth.

Fig. 92.—Lower portion of a potato plant. *ss*, Level of earth. *pa*, *pa*, Aerial portion bearing leaves. *t*, Subterranean portion of stem or tubers. *T*, Tuber showing eyes or leaf-buds, covered by scales, *b*, which are equivalent to leaves.

Fig. 93.—Corm or under-ground stem of *Colchicum autumnale*. *r*, Roots. *f*, Leaf. *a'*, Ascending axis of preceding year, withered. *a''*, Axis of the year. *a'''*, Point where axis of next year would be formed.

Gladiolus, or from the side as Colchicum (fig. 93 α'). These buds feed on the original corm α' , and destroy it. It will be noticed afterwards, when leaf-buds and bulbs are considered.

Internal Structure of Stems.

71. Stems, according to their structure, have been divided into three classes:—*Exogenous* (ἐξω, outward, and γεννᾶειν, to produce), when the bundles of vascular tissue are produced regularly in succession externally, and go on increasing indefinitely in an outward direction. *Endogenous* (ἐνδον, within), when the bundles of vascular tissue are produced in definite bundles and converge towards the interior, additions being thus in the first instance made internally. *Acrogenous* (ἄκρος, summit), when the vascular bundles are developed at the same time and not in succession, the addition to the stem depending on the union of the base of the leaves and extension of the growing point or summit. The plants which exhibit these three kinds of stem, are distinguished also by the structure of their embryo. Thus exogenous stems are met with in plants having an embryo or germ which has two cotyledons or seed-lobes, hence they are called *Dicotyledonous* (δύς, twice, and κοτυληδών, a seed-lobe); plants with endogenous stems have only one cotyledon, and are called *Monocotyledonous* (μόνος, one); while plants with acrogenous stems have no cotyledons, and are called *Acotyledonous* (α, privative). The terms connected with the embryo will be afterwards fully explained.

Exogenous or Dicotyledonous Stem.

72. The Exogenous or Dicotyledonous stem characterizes the trees of this country. It consists of a cellular and vascular system: the former including the outer bark, medullary rays, and pith; the latter, the inner bark, woody layers, and medullary sheath. In the early stage of growth, the young dicotyledonous stem is entirely cellular; but ere long fusiform tubes appear, forming bundles, having the appearance of wedges (fig. 94





Fig. 94.—Young Dicotyledonous or Exogenous stem. *ww*, Vascular bundles in the form of wedges. *p*, Pith. *rrr*, Medullary rays.

Fig. 95.—Same stem further advanced; the letters as in fig. 94, *nnn*, new vascular wedges interposed between those first formed.

ww) arranged in a circle round a central cellular mass of *pith* (fig. 94 *p*), which is connected to the outer part or *bark*, by means of cellular processes called *medullary rays* (fig. 94 *rrr*). At first, the cellular portion is large,—the pith, bark, and rays occupying a large portion of the stem; but by degrees new vascular bundles are formed, which are deposited between the previous ones (fig. 95 *n n n*). By this means the pith is more circumscribed, the medullary rays become narrow, and the bark more defined. Such is the structure presented by an annual herbaceous dicotyledonous stem, consisting of pith, a circle of fibro-vascular and woody tissue, medullary rays, bark, and epidermis.


73. The stems of trees and shrubs in their young state exhibit an arrangement similar to that represented as occurring in the herbaceous stem (fig. 95), with this difference, that the vascular circle is more firm and solid. As ligneous stems continue to grow, further changes take place by which their diameter is increased, and they are rendered more dense. The shoots or young branches given out annually, however, are similar in structure to annual herbaceous stems; and in making successive sections from the apex of a branch, which is succulent and green, to the base of a trunk, which is comparatively dry and hard, the various changes which take place can be easily traced. Fig. 96 represents a thin horizontal or *transverse* section of the upper part of a young branch of *Acer campestre*. In the centre, *m*, is the pith, very large at this period of growth, and occupying at least one-half of the whole diameter, its cells diminishing in size as they approach the circumference. Immediately surrounding the pith is a layer of a




greenish hue, the medullary sheath, *em*, from which the medullary rays *rm*, proceed towards the circumference, dividing the vascular circle into numerous compact segments, which consist of woody vessels, *fb*, and of porous vessels, *vp*. These are surrounded by a moist layer of greenish cellular tissue, *c*, called the cambium layer, which is covered by three layers of bark, *fc*, *ec*, and *p*, with laticiferous vessels, *vl*, the whole being enclosed by the epidermis, *ep*. On making a thin

Fig. 96.—Horizontal section of young stem of *Acer campestre*, magnified twenty-six diameters. *m*, Pith. *em*, *em*, Medullary sheath. *fb*, *fb*, Woody bundles. *vp*, Porous vessels. *rm*, Medullary rays. *c*, Cambium. *fc*, Fibres of Endophloeum. *vl*, Laticiferous vessels. *ec*, Cellular envelope, Mesophloeum. *p*, Corky envelope, Epiphloeum. *ep*, Epidermis.

the microscope, the parts composing the different portions become more obvious (fig. 97 *a*). The pith, *m*, with its hexagonal cells decreasing in size outwards, surrounded by a narrow fibro-vascular zone; the medullary sheath, consisting chiefly of spiral vessels, *t*; the medullary ray, *rm*; the vascular zone, consisting of porous vessels, *vp*, of large diameter, and forming the large round apertures seen in a transverse section; the woody fibres, *fl*, with their thick walls and smaller apertures; the inner bark or liber, *fc*, with the layer of cambium cells, *c*; the second layer of bark, or the cellular envelope, *ec*, with the laticiferous vessels, *vl*; the outer or suberous layer of bark, *p*, with the thin layer of epidermis, *ep*, having hairs scattered over its surface. A transverse section of a bundle of vascular tissue of a dicotyledonous plant, magnified 230 times, is

97 *a*

represented in fig. 97 *b*. The arrow indicates the direction from within outwards. We here perceive the vascular bundle proper surrounded by a very large-celled tissue (*a á b e f*). The almost square cells, *a á*, form the epidermis on which follows the less dense cellular tissue of the bark. The latter surrounds a half-moon-shaped bundle of bast cells, *c*, which are separated in the direction towards the interior, by a layer of cambium, *d d' d''*, from the bundles of vascular tissue, consisting of vessels and longitudinal cells. The latter tissues may be distinguished in the transverse section by the thicker walls (*g g*), and by their greater breadth (*h h*).


97 *b*

74. Such is the structure of a young shoot during the first year of its growth. At the end of a second year the shoot is found to have increased in diameter by the formation of a zone of vessels consisting of porous and woody tissue, and a zone of fibrous bark, the medullary

Fig. 97 *a*.—Vertical section of the same stem more highly magnified. *t*, Tracheæ or spiral vessels. *fl*, *fl*, *fl*, Woody fibres. The other letters as in fig. 96.

Fig. 97 *b*.—Transverse section of a bundle of vascular tissue of a Dicotyledonous plant. *a á*, Epidermis. *b e f*, Large celled tissue of bark. *d d' d''*, Layer of cambium.

rays being at the same time continued from within outwards. This is represented in fig. 98, where (1, 1) indicates the section of the stem



of the first year's growth (the letters referring to the same part as in figs. 96, 97); and (2) shows the interposed zones of the second year, by which the diameter of the stem is increased.

75. **The Pith**, or the central part of a dicotyledonous stem, is composed of cellular tissue, which is developed in an upward direction, the cells diminishing in size towards the circumference, and being often hexagonal. In the young plant it occupies a large portion of the stem, and sends cellular processes outwards at regular intervals to join the medullary rays (figs. 94, 95, *p*). The pith has at first a greenish hue, and is full of fluid, but in process of time it becomes pale-coloured, dry, and full of air. These changes take place first in the central cells. Sometimes the pith is broken up into cavities, which have a regular arrangement, as in the Walnut and the Jessamine; it is then called *discoïd* or *disciform* (δίσκος, a disc, from the circular partitions). At other times, by the rapid growth of the outer part of the

Fig. 98.—Vertical section of a branch of *Acer campestre*, two years old, where (1, 1) indicates the portion formed the first year, and (2) that formed the second. The letters as in figs. 96 and 97.

Fig. 98 bis.—Certain parts of the preceding magnified in order to show the structure of the vessels and cells, as well as their form and direction. Fig. 98 ter.—A portion of a porous vessel magnified.

stem, the pith is ruptured irregularly, and forms large cavities, as in the fistular stem of Umbelliferous plants. Circumscribed cavities in the internal cellular portions of stems are by no means unfrequent, arising either from rupture or absorption of the cells. In some rare instances, vessels occur in pith, as in Elder, Pitcher-plant, and Ferula; and occasionally its cells are marked by pores indicating the formation of secondary deposits. The extent of pith varies in different plants, and in different parts of the same plant. In Ebony it is small, while in the Elder it is large. In the Rice-paper plant, a species of *Æschynomene*, the interior of the stem is occupied almost entirely by cellular tissue, which may be called pith; from this the paper is made by cutting thin sections in a circular manner. The same kind of tissue occurs in the Papyrus of the Nile. When the woody circle of the first year is completed, the pith remains stationary as regards its size, retaining its dimensions even in old trunks, and never becoming obliterated.


76. The Medullary Sheath is the fibro-vascular layer immediately surrounding the pith. It forms the inner layer of the vascular bundle of the first year (fig. 97 *t*), and consists chiefly of true spiral vessels, which continue to exercise their functions during the life of the plant, and which extend into the leaves. With the spiral vessels there are a few woody fibres intermingled. The processes from the pith are prolonged into the medullary rays between the vessels of the sheath.

77. Woody Layers. During the first year, the vascular circle consists of an internal layer of spiral vessels forming the medullary sheath, and external bundles of porous and ligneous vessels. In subsequent years the layer of spiral vessels is not repeated, but concentric zones of porous vessels (fig. 98 *ter.*), and pleurenchyma are formed, constituting what are commonly called the woody circles of trees. The vascular bundles, from their mode of development in an indefinite manner externally, have been called *Exogenous*; and, for the same reason, Schleiden has denominated them *Indefinite*. Exogenous plants have sometimes received the name of *Cyclogens* (κύκλος, a circle), in consequence of exhibiting concentric circles in their stems. On a transverse section, each zone or circle is usually seen to be separated from that next to it by a well-marked line of demarcation. This line, as in the Oak (figs. 99, 100), and in the Ash, is indicated by holes which are the openings of large porous vessels; the remainder of the tissue in the circle being formed by pleurenchyma, with thickened walls and of smaller calibre. In some trees, as the Lime, Hornbeam, and Maple, the line is by no means so well marked, as the openings are smaller and more generally diffused; but there is usually a deficiency of porous vessels towards the outer part of the circle. In cone-bearing plants, as the Fir, in which the woody layers consist entirely of punctated woody tissue (fig. 47), without any large porous


vessels, the line of separation is marked by the pleurenchyma becoming dense and often coloured. In some kinds of wood, as Sumach, the zones are separated by a marked development of cellular tissue. The separation between the zones is said to be owing to the interruption in the growth of the tree during autumn and winter, and hence it is well defined in trees of temperate and cold climates. But even in tropical trees, the lines, although often inconspicuous, are still visible; the dry season, during which many of them lose their leaves, being their season of repose.

78. The woody layers vary in their texture at different periods. At first the vessels are pervious and full of fluid, but by degrees thickening layers are deposited which contract their canal, and sometimes obliterate it. The first-formed layers are those which soonest become thus altered. In old trees, there is a marked division between the central *Heart-wood* or *Duramen* (*durus*, hard), and the external *Sap-wood* or *Alburnum* (*albus*, white); the former being hard and dense, and often coloured, with its

tubes dry and thickened; while the latter is less dense, is of a pale colour, and has its tubes permeable by fluids. The difference of colour between these two kinds of woods is often very visible. In the Ebony tree, the duramen, or perfect-wood, is black, and is the part used for furniture, &c.; the alburnum is pale: in the Beech, the heart-wood is light-brown: in the Oak, deep-brown: in Judas tree, yellow; in Guaiacum, greenish. The alteration in colour is frequent in tropical trees. In those of temperate climates, called *white-wood*, as the Willow



99



100

Fig. 99.—Horizontal section of the stem of an oak eight years old. *b*, Wood, showing concentric circles or zones, separated by points which correspond to the opening of the large porous vessels, or Bothrenchyma. *c*, Bark, showing also eight concentric circles, thinner and less distinct. The wood and bark are traversed by medullary rays, some of which extend from the bark to the pith, and others reach only a certain way inwards.

Fig. 100.—Horizontal section of two woody bundles of Cork-oak, separated from each other by the medullary ray, *r m'*. The two primary bundles are divided by secondary rays, *r m''*, *rm'''*, *r m'''*, which vary in extent according to the period when they originated. *m*, Pith. *ec*, Cellular envelope. *p*, Corky envelope, which is highly developed and exhibits several layers.

and Poplar, no change in colour takes place; this is also the case in the Chestnut and Bombax. The relative proportion of alburnum and duramen differs in different trees. Duhamel says that in the oak, six inches in diameter, the alburnum and duramen are of equal extent; in a trunk one foot in diameter, they are as two to seven; in a trunk two feet in diameter, as one to nine. The heart-wood is more useful than the sap-wood, and less liable to decay. The wood of different trees varies much in its durability. Pieces of wood, $2\frac{5}{8}$ inches square, were buried to the depth of one inch in the ground, and decayed in the following order:—Lime, American Birch, Alder, and Aspen, in three years; Willow, Horse-chestnut, and Plane, in four years; Maple, Red Beech, and Birch, in five years; Elm, Ash, Hornbeam, and Lombardy Poplar, in seven years; Robinia, Oak, Scotch Fir, Weymouth Pine, Silver Fir, were decayed to the depth of half an inch in seven years; while Larch, common Juniper, Virginian Juniper, and Arbor Vitæ, were uninjured at the end of that time.

79. From the mode in which the woody layers are formed, it is obvious that each vascular zone is moulded upon that which precedes it; and as in ordinary cases each woody circle is completed in the course of one year, it follows, that, by counting the concentric circles, the age of a tree may be ascertained. Thus fig. 99 represents an oak eight years old having eight woody layers, *b*. This computation can only be made in trees having marked separations between the circles. There are, however, many sources of fallacy. In some instances, by interruption to growth, several circles may be formed in one year, and thus lead to an erroneous estimate. Care must be taken to have a complete section from the bark to the pith, for the circles sometimes vary in diameter at different parts of their course, and a great error might occur from taking only a few rings or circles, and then estimating for the whole diameter of the tree. When by the action of severe frost, and other causes, injury has been done to the tender cells from which the young wood is developed, while, at the same time, the tree continues to live, so as to form perfect woody layers in subsequent years; the date of the injury may be ascertained by counting the number of layers which intervene between the imperfectly formed circle and the bark. In 1800, a Juniper was cut down in the forest of Fontainebleau, exhibiting near its centre a layer which had been affected by frost, and which was covered by ninety-one woody layers, showing that this had taken place in the winter of 1709. Inscriptions made in the wood become covered, and may be detected in after years when a tree is cut down; so also wires or nails driven into the wood. As the same development of woody layers takes place in the branches as in the stem of an Exogenous tree, the time when a branch was first given off may be computed by counting the circles on the stem and branch respectively. If there are fifty circles for instance in the trunk,

thirty in one branch, and ten in another, then the tree must have been twenty years old when it produced the first, and forty when it formed the other.

80. In Exogenous stems the pith is not always in the centre. The layers of wood on one side of a tree may be larger than those on the other, in consequence of more full exposure to light and air, or the nature of the nourishment conveyed, and thus the pith may become *excentric*. Zones vary in size in different kinds of trees, and at different periods of a plant's life. Soft wooded trees have usually broad zones, and old trees form smaller zones than young ones. There are certain periods of a plant's life when it seems to grow most vigorously, and to form the largest zones. This is said to occur in the oak between twenty and thirty years of age.

81. **Cambium.**—External to the woody layers, and between them and the bark, there is a layer of mucilaginous semifluid matter, which is particularly copious in spring, and to which the name of *Cambium* (*cambio*, to change, from the alterations that take place in it) has been given (figs. 96, 97 c). According to Schacht, this is the proper nourishing tissue. It constitutes the *thickening zone*, by means of which the stem is enlarged, those cambium cells situated most internal being subservient to the purposes of the wood formation, those most external originating the new bark. In this cells are formed, called cambium cells, of a delicate texture, in which the protoplasm and primary utricle are conspicuous. These cells undergo changes, so as to assume an elongated fusiform shape, and ultimately become thickened pleureenchyma. So long as the primary utricle can be detected, they appear to be in an active state, and capable of developing new cells. This cambium layer marks the separation between the wood and the bark, and may be regarded as constituting the active formative tissue of Dicotyledonous stems.

82. **Bark or Cortical (cortex, bark) System**, lies external to the wood, and, like it, consists of several layers. In the early state it is entirely cellular, and is in every respect similar to the pith; but, as the vascular bundles are developed, the bark and pith are separated, and the former gradually becomes altered by the formation of secondary deposits. The bark consists of a cellular and vascular system. In this respect it resembles the wood, but the position and relative proportion of these two systems is reversed. In the bark the cellular system is external, and is much developed; while the vascular is internal, and occupies comparatively a small space. The cellular portion of the bark consists of an external layer, or *Epiphlæum* (ἐπί, upon, on the outside, and φλοιός, bark), and the cellular envelope, or *Mesophlæum* (μεσός, middle; while the vascular system forms the internal portion called *Liber*, or *Endophlæum* (ἐνδόν, within).

83. The inner bark, or *endophlæum* (fig. 98 f c), is composed of

elongated pleurencyhyma mixed with laticiferous vessels and some cellular tissue. It is separated from the wood by the cambium layer. The pleurencyhymatous tubes are thickened by concentric deposits in their interior, and thus they acquire a great degree of tenacity. The liber of the Lime tree and of *Lepurandra saccidora* (a species of *Antiaris*?), are used to form mats, cordage, and sacks; and the toughness of the fibres of the inner bark of flax, hemp, and of many of the nettle and mallow tribe, render them fit for various manufacturing purposes. The liber is sometimes called the *bast-layer*, from its uses. Occasionally it is continuous and uninterrupted, as in the Vine and Horse-chestnut; at other times, as in the Oak, Ash, and Lime, the fibres are separated during the progress of growth, and form a sort of net-work, in the interstices of which the medullary rays are seen. The fibres of the lace-bark tree (*Lagetta linearia*) are thus formed. In fig. 101, is represented the bark of *Daphne Laureola*; *f* indicating the woody fibres of liber, and *r* the medullary rays. The endophlœm increases by layers on its inside, which are thin, and may be separated like the leaves of a book, and hence the application of the name *liber*. The term *liber* may be derived from the fact of the inner bark being used for writing upon.

84. The cellular envelope, or *mesophlœum*, lies immediately on the outside of the liber. It consists of polyhedral, often prismatical cells (fig. 98 *bis*, *e c*), usually having chlorophylle, or green colouring matter, in their interior, but sometimes being colourless and containing raphides. They are distinguished from those of the epiphlœum by their form and direction, by their thicker walls, their green colour, and the intercellular spaces which occur among them. This covering is usually less developed than the outer suberous layer, but sometimes, as in the Larch and common Fir, it becomes very thick, and separates like the epiphlœum. In the cellular envelope laticiferous vessels occur.

85. The *Epiphlœum* is the outer covering of the bark, consisting of cells which usually assume a cubical or flattened tabular form (fig. 98 *bis*, *p*). The cells have no chlorophylle in their interior, are placed close together, and are elongated in a horizontal direction; and thus they are distinguished from the cells of mesophlœum. In the progress of growth they become often of a brown colour. This covering may be composed of a single layer of tabular cells; but in some trees it consists of numerous layers, forming the substance called *cork*, which




Fig. 101.—Network formed by liber of *Daphne Laureola*. *f f*, Fibrous bundles. *r r*, Medullary rays.


is well seen in *Quercus suber*, the Cork-oak (fig. 100 p); hence the name *suberous*, or *corky layer*, which is given to it. The form of its cells varies in some instances, being cubical at one part, and more compressed or tabular at another, thus giving rise to the appearance of separate layers. After a certain period (sometimes eight or nine years), the corky portion becomes dead, and is thrown off in the form of thickish plates, leaving a layer of tabular cells or *periderm* below. On the exterior of the epiphloeum is situated the epidermis, which has already been described (§ 47). It is formed of a layer of cells, which in woody stems serve only a temporary purpose, becoming ultimately dry, and being thrown off in the form of plates or shreds.

86. The bark, in its increase, follows an order exactly the reverse of that which occurs in the woody layers. Its three portions increase by additions to their inside. The layers of liber owe their increase to the cambium cells, which, by their constant reproduction, mark the separation between the vascular bundles of the wood, and the fibres of the endophloeum. These layers are often so compressed and united together as to be counted with difficulty, while at other times they are separated by rings of cellular tissue, and thus remain conspicuous. In the case of the cellular portions of the bark, there are also successive additions, sometimes to a great extent, but they do not usually exhibit any marked divisions.

87. As the additions are made to the woody layers on the outside, and to the bark on the inside, there is a constant distension going on, by which the bark becomes compressed, its layers of liber are condensed, the fibres are often separated so as to form meshes (as in the lace-bark), its epidermis is thrown off, and the epiphloeum is either detached along with it, or, when thick, is ruptured in various ways, so as to give rise to the rugged appearance presented by such trees as the Elm and Cork-oak. In some instances the bark is very distensible, and its outer cellular covering is not much developed, so that the surface remains smooth, as in the Beech. The outer suberous layer sometimes separates with the epidermis, in thin plates or scales. In the Birch, these have a white and silvery aspect. There is thus a continual destruction and separation of different portions of the bark. The cellular envelope and liber may remain while the epiphloeum separates, or they also may be gradually pushed off—the parts which were at first internal becoming external. In the case of some Australian trees, both the cellular and fibrous portions are detached in the form of thin flakes, and occasionally each annual layer of liber throws off that which preceded it. The epidermis separates early, and no renewal of it takes place. There is, however, an internal covering, which is formed of various portions of the bark. To this covering the name *Periderm* (περί, around, and δερμα, skin) has been given by Mohl.

88. From the mode in which the outer layers of bark separate, it follows that inscriptions made on them, and not extending to the wood, gradually fall off and disappear. A nail driven into these layers ultimately falls out. In consequence of the continued distension of an exogenous stem, it is found that woody twining plants cause injury, by interrupting the passage of their fluids. A spiral groove may thus be formed on the surface of the stem, by the compression exercised by a twining plant, such as honeysuckle. From what has been stated relative to the changes which take place in the bark, it will be understood that it is often difficult to count its annual layers, so as to estimate the age of the tree by means of them. This may, however, be done in some cases, as shown at fig. 99, where there are eight layers of bark, *e*, corresponding to eight woody layers, *b*.

89. **Medullary Rays or Plates.**—While the bark and pith become gradually separated by the intervention of vascular bundles, the connection between them is kept up by means of processes called *medullary rays* (figs. 94, 95 *r*). These form the *silver grain* of carpenters; they communicate with the pith and the cellular envelope of the bark, and they consist of cellular tissue, which becomes compressed and flattened so as to assume a muriform appearance (fig. 102 *m r*). At first



they occupy a large space (fig. 94 *r*); but as the vascular bundles increase, they become more and more narrow, forming thin laminae or plates, which separate the woody layers. On making a transverse or horizontal section of a woody stem, the medullary rays present the


Fig. 102.—Vertical section of a one-year old branch of *Acer campestre* highly magnified, and extending from the pith to the bark, parallel to the medullary rays. *m r*, A medullary ray or plate extending from the pith, *p*, to the bark, *c e*, crossing tracheae, *t*, woody fibres, *l*, porous vessels, *b*, and cortical fibres, *c f*.

Fig. 103.—Vertical section of the same branch perpendicular to medullary rays. *l l*, Woody fibres which interlace, leaving spaces, *m r*, *m r*, *m r*, where the medullary rays pass.

aspect of narrow lines running from the centre to the circumference (figs. 99, 100 *r m*); and in making a vertical section of a similar stem through one of the rays, the appearance represented in fig. 102 will be observed, where a medullary ray, *m r*, composed of flattened muri-form cells passes from the pith, *p*, to the cellular envelope, *c e*, crossing the tracheæ of the medullary sheath, *t*, the ligneous tissue, *l*, the porous vessels of the wood, *b*, and the fibres of the liber, *c f*. The laminae do not by any means preserve an uninterrupted course from the apex to the base of the tree. They are broken up by the intervention of woody fibres, as seen in a vertical section of a woody stem (fig. 103), perpendicular to the medullary rays *m r*, *m r*, *m r*, which are separated by interlacing woody fibres, *l l*. The medullary rays are usually continuous from the pith to the bark, additions being made to them as they proceed outwards. But, occasionally, secondary rays arise from the outer cells, which pass only to a certain depth between the vascular bundles, as in the Cork-oak (fig. 100, *r m'''*, *r m''*, *r m'''*). Medullary rays are conspicuous in the Cork-oak, Hazel, Beech, Ivy, Clematis, Vine. They are not so well marked in the Lime, Chestnut, Birch, Yew.

Anomalies in the Structure of the Exogenous Stem.

90. The stems of Dicotyledonous plants occasionally present anomalous appearances in the structure and arrangement of their wood, bark, and medullary rays. In place of concentric circles, there are sometimes only a few rows of wedge-shaped vascular bundles produced during the life of the plant, additions being made by the interposition of bundles of a similar kind annually, resembling in this respect the formation of woody bundles in the early growth of herbaceous plants (fig. 95). In the Pepper tribe, Aristolochiaceæ, and Menispermaceæ, these anomalous stems occur. In *Gnetum* (fig. 104), the vascular



104

Fig. 104.—Horizontal section of stem of *Gnetum*. *m*, Pith. *e m*, Medullary sheath. *b b b b b b b*, Woody bundles forming seven concentric zones, each of which is the produce of several years. *l l l l l l l*, Fibres of liber forming interposed circles, equal in number to the woody zones.

bundles, *b b b b b b b*, form zones, which are each the produce of several years' growth, and are separated by layers, *l l l l l l l*, which may be considered as representing different zones of liber.

In some of the Menispermum tribe, the separating layers are of a cellular and not of a fibrous nature. In *Banisteria nigrescens* (fig. 105), the young stem (1) presents a

four-lobed surface; the lobes gradually deepen (2), and ultimately (3) the stem is divided into a number of separate portions, the central one of which alone exhibits pith and medullary rays. The portions are separated by interposed cortical layers.

Many of the Malpighiaceæ, Sapindaceæ, and Bignoniaceæ of Brazil, exhibit stems in which the woody layers are arranged in a very irregular manner. In the stem of *Calycanthus floridus*, and of some Brazilian Sapindaceæ, such as *Paullinia pinnata* (fig. 106), *Serjania triternata*, and *Selloviana*, there is a central woody mass with from three to ten small secondary ones around it. Each of the masses contains true pith, apparently derived from the cortical cellular tissue, or from the original medullary centre. Gaudichaud and Jussieu state that around these separate collections of pith, there is a medullary sheath and spiral vessels. No annual rings have been detected in the secondary masses, but medullary rays exist, usually in their outer portion (fig. 106). In these anomalous Sapindaceæ, the central and



105

lateral woody masses are enclosed in a common bark, with a continuous layer of liber. Some have supposed that the lateral masses are undeveloped branches united together under the bark; but Treviranus considers them as connected with the formation of leaves, and as depending on a peculiar tendency of the vascular bundles to be developed independently of each other round several centres.

In some Bignoniaceæ (fig. 107), the layers of wood are divided in

Fig. 105.—Horizontal section of stem of *Banisteria nigrescens* at different ages. 1. Stem presenting four superficial lobes. 2. Showing six deeper lobes, with intermediate divisions. 3. The lobes separated by cellular tissue, the middle one alone having pith and medullary sheath. The points indicate the orifices of porous vessels.


a crucial manner into four wedge-shaped portions by the intervention of plates differing in texture from the ordinary wood of the plant, and probably formed by introversion, or growing inwards of the liber. In some Guayaquil Bignonias, Gaudichaud perceived first four of these



106



107



108



109

plates, next eight, then sixteen, and finally thirty-two. In *Aspidospermum excelsum* of Guiana, and in *Heteropterys anomala* (fig. 108), the stem assumes a peculiar lobed and sinuous aspect, and in some woody climbing plants, pressure causes the stems to become flattened on the side next the tree on which they are supported, while from being twisted alternately in different directions, they present a remarkable zigzag form, having the woody layers developed only on one side

Fig. 106.—Horizontal section of the stem of *Paullinia pinnata*, one of the Sapindaceæ of Brazil, showing numerous secondary woody masses, surrounding a central one. Each of the separate masses has pith, often excentric, with a medullary sheath, containing spiral vessels, and a few medullary rays chiefly towards the circumference of the stem.

Fig. 107.—Horizontal section of the stem of *Bignonia capreolata*, showing the crucial division of the woody layers

Fig. 108.—Horizontal section of stem of *Heteropterys anomala*, one of the Brazilian Malpighiaceæ, showing an irregularly lobed surface. The dots indicate porous vessels.

Fig. 109.—Fragment of a stem of a climbing species of *Banisteria* (*B. scandens*) showing the effects of compression.


(fig. 109). In Firs the wood is occasionally produced in an oblique in place of a perpendicular manner, thus injuring the timber, and causing it to split in an unusual way. The young plants produced from the seed of such twisted-wooded firs, are said to inherit the peculiarity of their parents.

Endogenous or Monocotyledonous Stem.

91. This kind of stem is composed of cells and vessels which are differently arranged from those of the Exogenous stem. The vascular bundles are scattered through the cellular tissue, and there is no distinction between pith, wood, or bark. There are no medullary-rays, nor concentric circles (fig. 110). In the young state, the centre of the stem is occupied entirely by cells, which may be said to represent pith, and around this the vessels are seen increasing in number towards the circumference. The central cellular mass has no medullary sheath. In some cases its cells are ruptured, and disappear during the progress of growth, leaving a hollow cavity (fig. 111); but in general it remains permanent, and is gradually encroached upon by the development of the vascular system. The latter consists of vessels arranged in *definite* bundles, which do not increase by additions to their outside after being once formed, although they are developed in a progressive manner. These bundles may be considered as representing the vascular wedges, produced during the first year of an exogenous stem's growth (fig. 94). They consist of woody vessels enclosing some cellular tissue between them, with spiral and porous vessels. The outer



110



111


part of the stem is not formed by a separable bark, but consists of a dense mass of fibrous tissue, mixed with laticiferous vessels and cells. It is intimately connected to the inner part of the stem, without the intervention of medullary rays.

92. On making a transverse section of a young endogenous stem (fig. 112), there is observed a mass of cells or utricles, *u*, of various sizes, often small in the vicinity of the vascular bundles, spiral vessels or tracheæ, *t*, large porous vessels, *v p*, laticiferous vessels, *l*, and woody

Fig. 110.—Part of the stem of *Asparagus* cut transversely, showing the vessels as points distributed through the cellular tissue. *l*, Leaf in the form of a scale

Fig. 111.—Transverse section of stem of *Phragmites communis*, or common reed. The cellular tissue in the centre has disappeared, leaving a fistular or hollow stem, with a ring of cells and vessels, the latter indicated by dots. *n*, Node where the fibres cross, so as to form a solid partition.

fibres, *f*, resembling those of liber, thickened by internal deposits. A similar section of a further advanced endogenous stem, as of a Palm (fig. 113), shows numerous bundles of vessels dispersed irregularly in cellular tissue; those near the centre, *m*, being scattered at a distance from each other, while those towards the outside are densely aggregated, so as to form a darkish zone, *b*, and are succeeded at the circumference by a paler circle of less compact vessels, *l*, with some compressed cells, covered by an epidermis, *e*. The peripheral portion, *l*, differs from *true bark*, in not being separable from the rest of the tissue. It has received the name of *false bark*, and consists of the epidermal cells,



112



113

e, and what has been called the cortical integument, *l*. This portion of the stem is often very inconspicuous, but sometimes it is much developed as in *Testudinaria elephantipes*, in which it is rugged, and is formed of a substance resembling cork in many respects.


93. Mohl states that, in the stem of a Palm, there may be distinguished a central region, a fibrous layer, and a cortical region; and the same divisions are pointed out by Henfrey in the stem of *Sparganium ramosum* and other monocotyledons. The central portion, representing the pith of dicotyledons, consists in *Sparganium* of spherical cells, containing starch, while the cortical or outer portion is formed by irregular cells, which are usually destitute of starch.

94. It was at one time supposed that the woody portion of these stems was increased by additions to the centre, so that the first-formed

Fig. 112.—Horizontal section of a vascular bundle from the stem of a Palm (*Corypha frigida*). *t*, Tracheæ, or spiral vessels *vp*, Large porous vessels. *u*, Cells or utricles of various kinds surrounding the vessels, and forming the parenchyma. *l*, Laticiferous vessels. *f*, Fibres analogous to those of liber, thickened by concentric deposits.

Fig. 113.—Transverse section of part of the stem of a Palm (*Astrocaryum Murumura*). *m*, Central or medullary portion in which the woody bundles are distant and scattered. *b*, External woody portion, where the fibres are numerous and densely aggregated, so as to form a dark zone. *l*, Paler circle of more slender and less compact fibres, which may be considered as analogous to liber. *e*, Cellular epidermal portion.

fibres were gradually pushed towards the circumference by those which succeeded them, in the manner represented in fig. 114, 1: hence the term Endogenous (ἐνδον, within, and γεννάω, I produce), meaning internal growth. But Mohl has shown that this was not strictly correct. For although the fibres connected with the leaves, in the first instance, are directed towards the centre, and are therefore always internal to those previously formed, yet, when they are traced downwards, they are found not to continue in a parallel direction, but to arch outwards, so as ultimately to reach the circumference. Hence, the newly-formed fibres really become external at the base, although internal above. On making a vertical section of an endogenous stem, as of a Palm, there is observed an interlacing of fibres, similar to what is represented in fig. 114, 2, where the four vascular bundles, *a b c d*, are first directed towards the centre, and then curve outwards towards the circumference, so that those last formed ultimately become external. The term Endogenous, will, therefore, only apply strictly to the fibres at the early part of their course. On this account, the terms Endogenous and Exogenous have been recently discarded by many writers, the terms Monocotyledonous and Dicotyledonous being substituted. The true distinction between Exogenous and Endogenous stems consists in this, that in the former, the woody or vascular bundles increase indefinitely at their periphery, while in the latter, they are arrested in their transverse growth at a definite epoch. The investing bark of the former undergoes, as regards its external layers, a process of vital disin-



114

FIG. 114.—Figures to represent the arrangement of four pairs of vascular bundles (*a a*, *b b*, *c c*, *d d*), in endogenous stems. 1. According to the old idea of internal development throughout the stem. 2. According to the view of Mohl, who has shown that the fibres interlace, and that those which are at first internal, become external, lower down.

tegration which preserves its distensibility, and, in consequence, permits an unlimited extension of woody growth. The fibres on the surface of the latter maintain an intimate union with the general structure which precludes such an operation. Distensibility and vitality are, therefore, in the latter, so closely associated, that when one reaches its extreme the other proportionately declines.

95. The composition of the vascular bundles, in different parts of their course, varies. Thus, at the upper part, where they proceed from the leaves towards the centre, they contain spiral vessels, porous vessels with some cellular tissue, a few laticiferous vessels, and woody fibres resembling those of liber (fig. 112.) As the bundles descend, the spiral vessels disappear, then the porous vessels, and when they have reached the periphery, and have become incorporated with it, nothing but fibrous tissue, or pleurenchyma, remains, forming by its division a complicated anastomosis, or net-work. Thus, at the commencement the bundles are large, but as they descend they usually become more and more attenuated. In some instances, however, as in *Ceroxylon andicola*, they increase at different parts of their course, probably by interstitial growth, and give rise to irregular swellings of the stem. This distension takes place occasionally at the base of the stem, as in *Euterpe montana*.

96. There are many herbaceous plants in this country, as Lilies, Grasses, &c., having endogenous stems, in which the course of the vascular bundles may occasionally be traced, but there are no British endogenous plants with permanent aerial woody stems. All the British trees are exogenous. Illustrations of endogenous stems must therefore be taken from trees of foreign countries. Palms furnish the best examples. In them the stem forms a cylinder of nearly uniform diameter throughout. The leaves are produced from a single terminal and central bud, called a *Phyllophor*, or *Phyllogen*, (*φύλλον*, a leaf, and *φέρω*, I bear, or *γεννάω*, I produce). Connected with the leaves are the vascular bundles, and the bases of the leaves remain attached to the outer part of the stem, surrounded by the *matulla* or *reticulum* (§ 57). While the leaves produced by one bud decay, another bud is developed in the centre in a similar manner. As the definite vascular bundles are produced, the stem acquires increased thickness, but it is arrested in its transverse diameter at a certain epoch. The bundles, although developed progressively, do not multiply indefinitely; and thus a Palm-stem seldom becomes of great diameter.

97. In consequence of this mode of formation, the outer part of a Palm-stem is the hardest and densest, and after acquiring a certain degree of solidity, it resists all further distension, and frequently becomes so hard as to resist the blow of a hatchet. It has been already stated, that in the exogenous stem, provision is made for unlimited extension laterally, by the development of indefinite bundles of woody

fibres and vessels, and the formation of a separable bark which can be thrown off; but in the endogenous stem there is no such provision. Hence, when the first formed part of the stem has increased to a certain amount, its progress is stopped by the hard indistensible outer fibrous covering; and the same thing takes place with the other parts in succession, till at length all have acquired a comparatively uniform size, as is seen in the coco-nut palm (fig. 115, 1). In consequence of the small lateral increase of Palm-stems, a woody twining plant does less injury to them than to trees of exogenous growth.

98. The growth of endogenous stems may be said to resemble the upward growth of Exogens by terminal buds only, for there is no cambium layer, and no peripheral increase. Hence, in Palms, the terminal shoot is developed, but there are no annual rings. The hardening of the stem depends, in all probability, partly on internal changes in the woody fibres, similar to what takes place in the heart-wood of Exogens. Occasionally, at the upper part of a palm-stem there is an appearance of zones, but it does not continue throughout the stem. From the absence of concentric circles, the age of a Palm cannot be estimated in the same way as an exogenous tree. The elongation, however, of each species of Palm is pretty regular, and by it an opinion may be formed of the age. The rings on the surface of the stem are not indicative of yearly growth.

99. In Palms, there is in general no provision for lateral buds, and no branches are formed. Hence, destroying the central bud will kill the tree. In some Palms, however, as the Doom palm of Egypt (*Cucifera thebaica*), the stem divides in a forked or dichotomous (δίχα, two ways, and τέμνω, I cut,) manner. Gardner,



Fig. 115.—Two endogenous or monocotyledonous trees, belonging to two different families. 1. *Cocos nucifera*, or coco-nut, belonging to the Palm family. 2. *Pandanus odoratissimus*, or screw-pine, belonging to Pandanaceae. The first has a simple unbranched stem, with a cluster of leaves at the summit; the second has a branched stem, with numerous leafy clusters, and peculiar aerial roots, proceeding from different parts of the stem. Two human figures are given to indicate the height of the trees.

in his travels in Brazil, noticed a Palm in which the centre bud had been destroyed, and two side ones had been produced, so as to give it a forked appearance. Other plants with endogenous stems, also produce lateral buds. In fig. 115, 2, there is a representation of such a stem, in the case of the Screw-pine, (*Pandanus odoratissimus*), and examples are seen in Grasses, as the Bamboo, in Asparagus, Asphodels, and Dracænas. In these cases, the stem is conical, like that of Exogens, and the destruction of the terminal bud is not necessarily followed by the death of the plant. The development of lateral buds is accompanied often by an increased diameter of the stem. A Dracæna in the Canary Islands, has a hollow stem capable of holding several men; and the fact of its living in this state, is marked by Jussieu as an argument against the strict endogenous formation; for, if the centre were the youngest and newest part, its destruction would put an end to the existence of the tree, in the same way as the removal of the outer part of the wood would destroy an exogenous stem. The branches in such plants are formed on the same principle as the stems; but their fibres, when reaching the stem, do not proceed to the centre, but extend outside the previous layers, between them and the outer false bark (fig. 113 *le*), and thus it is that they give rise to lateral increase. In Grasses, the stem or culm is usually hollow or fistular (fig. 111), in consequence of the outer part, by its rapid increase, causing the rupture and ultimate disappearance of the internal cellular portion. The fibres in some Grasses cross from one side to the other, forming partitions, as in Bamboo, which add much to the strength of the stem. The age of a Palm may therefore be determined rather by the length of the stem and the number of cicatrices it presents.

100. In many Endogenous or Monocotyledonous plants, the stem remains below ground, developing shoots which are simple, as in Banana and Plantain, or branched, as in Asparagus. In the former, the stem above ground is an herbaceous shoot, composed of the sheaths of the leaves. It dies after fruiting, and is succeeded by other shoots from the subterranean stem. The shoots or buds from such stems occasionally remain below ground in the form of bulbs, as in Lilies.

101. In some instances, the aerial stem has the usual endogenous structure, while the under-ground stem has the vascular bundles developed in the form of wedges, with cellular tissue in the centre, thus resembling some Exogens. This structure has been remarked in the Smilax or Sarsaparilla family. Lindley calls these plants *Dictyogens* (*δίκτυον*, a net), from their netted leaves, a character by which they differ from other Endogens. Henfrey holds that the ring of woody fibres in Tamus and Smilax, is merely an alteration of the parenchymatous cells of the periphery, and is not produced, as some have supposed, in the same way as the zones of Dicotyledons. He considers

this ring as probably analogous to the liber, and not to the indefinite vascular bundles of Exogenous stems.

Acrogenous or Acotyledonous Stem.

102. This stem, in its general external aspect, resembles that of Endogens. It is unbranched, usually of small, nearly uniform diameter, and produces leaves at its summit. It is easily distinguished by its internal structure. Tree Ferns furnish the best example of this kind of stem. In them it is denominated a *Stipe*, or a *Rachis*, and often attains the height of 120 feet (fig. 116). A transverse section of the stem (fig. 117) exhibits a circle of vascular tissue composed of masses, *z l*, of various forms and sizes, situated near the circumference; the centre, *m*, being either hollow or formed of cellular tissue. On the outside of the vascular circle, cells exist, *p*, covered by an epidermal layer or cellular integument, *e*, often of hard and dense consistence, formed originally by the bases of the leaves, which remain for a long time attached to the stem.

103. The vascular bundles are formed *simultaneously*, and not progressively as in the stem already noticed; and additions are always made in an upward direction. The stem then is formed by additions to the summit, and by the elongation of vessels already formed; hence the name *Acrogenous* (*ἀκρος*, summit). The leaves unite by their bases to form the stem, and the arrangement of their vessels is traced into it. The vascular system is of greater density than the rest of the tissue, and is usually distinguished by the dark colour of the pleurenchyma




Fig. 116.—Tree fern (*Alseodora perrotetiana*), of the East Indies. Stem or stipe is cylindrical, unbranched, and presents at its base, *ra*, a conical enlargement, formed by a mass of adventitious roots. The leaves are terminal, and in the young state are rolled up in a circinate manner.

(fig. 117 *f*), which surrounds the paler vessels in the centre (fig. 117 *v v*). The vascular bundles do not follow a straight course, but unite



117

and separate, leaving spaces between them, similar to those seen in the liber of Exogens (fig. 101).
104. The acrogenous stem in the young state is solid, but it frequently becomes hollow in the progress of growth, by the rupture and absorption of the walls of the cells in the centre. The bases of the leaves remain long attached, but ultimately fall off, leaving marked scars which are at first close together, but often separate afterwards by interstitial growth. On these scars or *cicatrices* (*cicatrix*, a wound), the markings of the vessels are easily seen, arranged in the same manner as those of the stem with which they are continuous. The vascular system of ferns consists chiefly of scalariform vessels (fig. 62), mixed with annular (fig. 57), and porous vessels (fig. 98 *ter*). There are no true tracheæ with fibres which can be unrolled. In the stems of Lycopodiaceæ, closed tracheæ or ducts (§ 34) occur; and in Equisetaceæ, the rings of the annular vessels are closely united.

105. The stem of Ferns is generally of small diameter; it does not increase much laterally, after having been once formed, and it does not produce lateral buds. Sometimes it divides into two (fig. 118), by the formation of two buds at its growing point. This, however, is an actual division of the stem itself, and differs from the branching of the Exogenous and Endogenous stem. In the Ferns of this country, the stems usually creep along and under the ground, and the leaves which they produce die annually without giving origin to an elevated trunk. In the common Brake (*Pteris aquilina*), the arrangement of the vascular system may be seen by making a transverse section of the under-ground stem. The plant has received its name *aquilina*, from a supposed resemblance to a spread eagle, presented by the vessels when thus cut across.



118

Fig. 117.—Transverse section of the stem of a Tree fern (*Cyathea*). *m*, Cellular tissue, corresponding to pith, occupying the central part. *zl*, Vascular circle composed of numerous irregularly-formed masses. *f*, Dark-coloured woody or prosenchymatous fibres, forming the borders of the vascular masses. *v v*, Pale-coloured vessels, chiefly scalariform, occupying the centre of the masses. *p*, Parenchymatous or cellular external zone, often communicating with the central portion. *e*, Hard epidermal envelope, occupying the place of the bark.

Fig. 118.—Vertical section of part of the forked stem or stipe of *Alsophila perrotetiana*. *m*, Cellular central portion. *zl*, *zl*, Vascular zone, consisting of woody fibres and scalariform vessels. The forking is caused by an actual division of the stipe.

106. In some *Thallogens*, which have been noticed as being stemless, the thallus or frond is supported by a stalk, in which there are concentric circles, with divisions in the form of rays, and a sort of pith. These are all forms of cellular tissue, however, without any woody fibres. These appearances are presented by some large antarctic sea-weeds, species of *D'Urvillea*, and by some lichens, as *Usnea*.

107. There are thus three kinds of stems in the vegetable kingdom:—

1.—*Exogenous* or *Dicotyledonous*, having a separable bark; distinct concentric circles, composed of progressive indefinite vascular bundles, increasing at their periphery, the solidity diminishing from the centre towards the circumference; pith, enclosed in a longitudinal canal or medullary sheath, with cellular prolongations in the form of medullary rays.

2.—*Endogenous* or *Monocotyledonous*, having no separable bark; no distinct concentric circles; vascular bundles progressive and definite, not increasing at their periphery, the solidity diminishing from the circumference to the centre; no distinct pith, no medullary sheath nor medullary rays, the cellular tissue being interposed between the vascular bundles.

3.—*Acrogenous* or *Acotyledonous*, having no separable bark; no concentric circles; vascular bundles simultaneous, forming an irregular circle; additions being made to the summit; no distinct pith, no medullary sheath nor medullary rays; conspicuous scars left by the bases of the leaves.

Formation of the different parts of Stems, and their special Functions.

108. The stem bears the buds from which branches are developed, as also the leaves and flowers, exposes them to the atmosphere and light, conveys fluids and air, and receives secretions. Stems vary much in their size, both as regards height and diameter. Some oaks in Britain have a height of nearly 120 feet; forest trees in France have attained to 120 and 130 feet, and in America even to 150 feet; while palms are frequently still higher. The trunks of some Baobabs in Senegal (*Adansonia digitata*), are said to be 30 feet in diameter.

109. The *pith*, in its early state (fig. 94 *p*), is of a greenish colour, and contains much fluid, which is employed in the nourishment of the young plant. After serving a temporary nutritive purpose, it becomes dry, or disappears by rupture and absorption of the walls of the cells which enter into its composition. The *medullary sheath*, which is the first formed vascular layer (fig. 96 *e m*), keeps up a connection between the central parts of the stem and the leaves, by means of spiral vessels, which seem to be concerned partly in the conveyance of air. This is the only part of a *Dicotyledonous* stem in which these vessels ordinarily occur. The *medullary rays* (fig. 97 *r m*), preserve a com-

munication between the bark and the pith. The cells of which they are composed, are concerned in the production of leaf buds, and they assist in the elaboration and conveyance of secretions. They have a direct connection with the *cambium cells* (fig. 97 *c*), or the cells between the wood and bark, whose function is to aid in the formation of new wood. The size of those rays varies according to the age of the plant. In the young stem they are large and complete: in the more advanced stem they are seen as lines only. They constitute the *silver grain* conspicuous in many woods. In some instances they appear large and broad, while the ligneous wedges are comparatively small. In others, many rays seem to be imperfect and supplemental, extending but partially through the stem. A section of Clematis or Aristolochia will exemplify the first,—a transverse incision of the stem of the Cork oak, the second of those appearances. The *bark* (fig. 97 *f c, e c, p*), protects the tender wood, conveys the elaborated sap downwards from the leaves, and is the part in which many valuable products, such as gum, tannin, and bitter principles, are formed and deposited. The *vascular bundles* (fig. 97 *f l, v p*), convey the sap from the root to the leaves. This function is carried on during the life of the plant by the annular vessels and the porous vessels, as well as other kinds of spurious fibro-vascular tissue; but in the woody fibres it ceases at a certain epoch, in consequence of the tubes being filled up by secondary deposits, so as to form the perfect wood, which gives strength and stability to the stem.

110. Considerable differences of opinion have arisen on the subject of the formation of wood. All agree that it cannot be properly formed unless the leaves are exposed to air and light, but physiologists differ as to its mode of deposition. Some say that it is deposited in a horizontal, others, in a vertical direction. There seems to be no doubt, that the cambium cells perform an important part in the formation of wood, and that their activity depends on the proper development of leaves. These formative cells, although most easily detected in exogenous stems, appear also to be present in the other forms of stems which have been described.

111. The early physiologists made experiments on exogenous stems, as being most easily procured. They espoused the *horizontal* theory of deposition, and disputed as to the formation of cambium; some maintaining that it was formed by the cells of the bark; others, by the central cells of the stem; and others, by both united. Duhamel, by putting silver plates between the bark and wood; and Dr. Hope, by detaching portions of bark, endeavoured to show that the bark alone was concerned in the formation of wood; while Decandolle and others were led to the conclusion, that both were concerned in the process of forming cambium, by means of which a layer of liber and a layer of wood was annually produced.

112. Knight espoused what is called the *vertical* theory, considering the wood as developed in a downward direction by the leaves, and in this view he is supported by Petit-Thouars and Gaudichaud. These physiologists maintain that there are two vascular systems in plants, an ascending and descending; the one connected with the leaf formation, or the spiral vessels; the other connected with the production of roots, or the woody fibres—the cellular tissue being more especially concerned in horizontal development. Every bud is thus, according to them, an embryo plant fixed on the stem, sending leaves upwards, and roots downwards. In Palms, Dracænas, and other Endogenous stems, the peculiar manner in which the woody fibres interlace (fig. 114, 2), favours the opinion that they are developed like roots, by additions to their extremities; and this is also strengthened by the formation of adventitious or aerial roots, which burst through different parts of the stem in Screw-pines (fig. 115, 2), in the Banyan, and in the Fig tribe in general. In Vellozias and Tree Ferns, the surface of the stem is often covered with thin roots, protruding at various parts, and becoming so incorporated with the stem as to appear to be a part of it. In the Tree Fern, represented in fig. 116, the lower part of the stem is enlarged in a remarkable degree by these fibres, so as to give it a conical form. In exogenous stems, when ligatures are put round the stem, and when portions of bark are removed, a swelling takes place above the parts where the injury has been inflicted, thus apparently proving that the new matter is developed from above downwards.

113. Gaudichaud endeavours to account for various anomalous forms of stems (figs. 105—108), by considering them as depending on the arrangement of the leaves, and on the mode in which the woody fibres are sent down from them. Thus, the four secondary masses surrounding the central one in the stem of *Calycanthus floridus*, are traced to four vascular bundles from the leaves, penetrating the cellular tissue of the bark, distinct from the central wood and from each other, except at the nodes, where the cross bundles unite them so as to form a ring round the central mass. New fibres are formed on the inner side of these bundles, and by degrees they assume a crescentic shape, while the horns of the crescent ultimately unite on the outer side (centrifugally), and enclose a portion of the bark, which thus forms a kind of spurious excentric pith, with numerous woody layers on the inside, and a smaller number on the outside. Again, in Brazilian Sapindaceæ (fig. 106), with five, seven, nine, or ten woody masses, the same thing is said to occur, with this difference, that the pith of each of the masses is derived from the original medullary centre, portions of which are enclosed by the vascular bundles in a centripetal manner, or from without, inwards.

114. Treviranus states that the fibrous and vascular bundles descending from the leaves, are destined in general to unite around a

common centre, but that they retain a certain degree of independence, and may be developed separately in some instances, giving rise to anomalous fasciculated stems.

115. Gardner, from an examination of Brazilian Palms, adopts the vertical theory, and Lindley also supports it. It is strongly opposed by Schleiden, Mirbel, Naudin, Henfrey, and others, who consider the development of the vascular bundles, as proceeding from below upwards; in Dicotyledons, by peripheral production of woody and vascular tissue from cambium cells; and in Monocotyledons, by a definite formation of woody and vascular bundles by means of terminal buds; the hardening of the stem depending on the interstitial changes which take place afterwards in the woody fibres.

116. A consideration of all the observations made on the formation of woody stems, leads apparently to the conclusion, that there is an ascending and descending axis in plants, and that each plant consists of one or more individuals, or *phytons* (*φυτὸν*, a plant), as they are called by Gaudichaud and others, having both axes developed; the Exogenous stem being formed by the original formation of two opposite phytons, the Endogenous by one: and that woody fibres are produced from cells, which, in Exogens, are formed annually between the wood and bark, as cambium cells; and, in Endogens, are developed in the internal parts of the stem. Proof seems, however, still wanting of the direction in which the development of wood takes place in the former; while, in the latter, observations seem to be in favour of a vertical formation, or of additions of woody fibre being made in a downward direction, as in roots, thus following the course of the descending elaborated sap.* Mirbel, from his investigations respecting the growth of the Date, believes that its fibres extend from below upward. He states, that at the summit of Monocotyledonous plants the aggregated cellular tissue eventuates in the production of a phyllophore, and that the formation of a vascular system in the plant precedes the development of the leaves.† Trecul has also shown that woody fibres in Dicotyledonous plants are not necessarily dependent on the presence of leaves, since ligneous tissue has been developed in the central portion of stems from which the bark had been removed. He believes that local increase depends on local action, that ligneous tissue is locally developed from previously existing cells, and that, therefore, a descending action has no direct influence on such formation.‡

117. Notwithstanding these various opinions, the formation of wood has been found to depend mainly on the functions of the leaves being

* For full details relative to the formation of wood, see Gaudichaud, *Recherches sur l'Organographie*, &c., Paris, 1841; Mirbel, *Annales des Sciences Naturelles*, 2d series tom. xx., &c.; Naudin, *do.*, 3d series, tom. 1.; Henfrey, *Annals of Natural History*, 2d series, vol. 1.

† Mirbel, *Recherches sur la Structure du Dattier*-*Annales des Sciences, Nat.* 2d series, xx. 5.
‡ Trecul, *Sur l'accroissement endiamètre des Vegetans dicotylédones ligneux*, *Annales des Sc. Nat.* 3d series, xvii.

carried on properly, and this can only be effected by exposure to air and light. The more vigorously the plant grows, the better is the wood produced. Experiments made in the British dockyards proved that those oaks which had formed the thickest zones, yielded the best timber. Barlow's experiments at Woolwich, showed that a plank of quick-grown oak, bore a greater weight than a similar plank of slow-grown oak.

118. In order that trees may grow well, and that timber may be properly formed, great care should be taken in planting at proper distances and in soil fitted for the trees. Firs ought to be planted from 6 to 8 feet apart, and hardwood trees for a permanent plantation, 28 feet distant, the spaces being filled up with larch, spruce, or Scotch fir, according to soil and situation. Hardwood is of no value till it has attained some age, while larch and spruce may be applied to use in ten or twelve years; and thus judicious thinning may be practised. When trees are set too close, their leaves are interrupted in their functions; many of them fall off, leaving the stems bare; the wood is imperfectly formed, and the roots are not sent out vigorously. When such plantations are allowed to grow without being thinned, the trees are *drawn up* without having a hold of the ground; and when a portion of them is subsequently removed, the remainder is easily blown over by the wind. In thick plantations, it is only on the trees next the outside, where the leaves and branches are freely formed, that the wood and roots are properly developed. When a tree is fully exposed to air and light on one side only, it is frequently found that the woody zones on that side are largest. When trees are judiciously planted, there is a great saving both in the original outlay and in the subsequent treatment. Pruning or the shortening of branches and the removal of superfluous ones ought to be cautiously practised. It is only applicable to young branches and twigs, and is had recourse to chiefly in the case of fruit-trees when the object is to make the plants produce flowers and fruit. If forest trees are properly planted and thinned, little pruning is required.

ROOT OR DESCENDING AXIS.

Structure of Roots.

119. In the young state there is no distinction between stem and root, as regards structure; both being cellular, and an extension of each other in opposite directions. In stemless plants, as Thallogens, the root remains in a cellular state throughout the life of the plants. The root is afterwards distinguished from the stem, by the want of a provision for the development of leaf-buds, and by increasing from above downwards. Some plants, however, as the Moutan Pæony, the

Plum-tree, *Pyrus japonica*, and especially *Anemone japonica*, have a power of forming buds on their roots. The last-mentioned plant develops these buds on every part of its extensively ramifying roots, which may be chopped into numerous pieces, each capable of giving rise to a new plant. The part where the stem and root unite is the *collum* or *neck*. In woody plants, the fibres of the stem descend into the roots, and there is a similar internal arrangement of woody layers, as is seen in the stem itself.

120. Roots are usually subterranean and colourless. Externally, they have a cellular epidermal covering of a delicate texture, sometimes called *epiblema* (¶ 47), in which no stomata exist. Their internal structure consists partly of cells, and partly of vascular bundles, in which there are no vessels with fibres which can be unrolled. Roots do not exhibit true pith, nor a medullary sheath. The axis of the root gives off branches which divide into radicles or fibrils (fig. 119), the extremities of which are composed of loose sponge-like cellular tissue, and are called *spongioles* or *spongelets*. Over these a very thin layer of cells is extended, called, by Trecul, a *Pileorrhiza* (πίλος, a cup, and ρίζα, a root). This sometimes becomes thickened, and separates in the form of a cup, as in Screw-pines (fig. 115, 2), and in Lycopodiums. Occasionally the extremities of roots are enclosed in a sheath, or *ampulla*, as in Lemna. Cellular papillæ, and hairs are often seen in roots (fig. 77), but no true leaves. Those hairs



119

seem to consist of simple elongated cells which occur singly, and are conjoined by their extremities. Roots do not grow throughout the whole length like stems, but by additions to their extremities, which are constantly renewed, so that the minute fibrils serve only a temporary purpose, and represent deciduous leaves. Schleiden has, therefore, observed that the tissue in the extremities of roots is older and more dense than that immediately adjoining it. In this respect they differ from branches, whose summits are usually their most recent growth.

121. Roots, in some instances, in place of being subterranean, become aerial. Such roots occur in plants called *Epiphytes*, or air-plants (ἐπι, upon, and φυτόν, a plant, from growing on other plants), as in Orchidaceæ; also in the Screw-pine (fig. 115, 2), the Banyan (*Ficus indica*), and many other species of *Ficus*, where they assist in supporting the stem and branches, and have been called *adventitious* or

abnormal. In Screw-pines, these aerial roots follow a spiral order of development. In Mangrove trees, they often form the entire support of the stem, which has decayed at its lower part. The name of adventitious has also been applied to those roots which are formed where portions of stems and branches, as of the Willow and Poplar, are planted in moist soil. They appear first as cellular projections, into which the fibres of the stem are prolonged, and by some are said to proceed from lenticels (§ 63). They frequently arise from points where the epidermis has been injured. A Screw-pine in the palm-house of the Edinburgh Botanic Garden, had one of its branches injured close to its union with the stem. This branch was at the distance of several feet above the part where the aerial roots were in the course of formation. At the part, however, where the injury had been inflicted, a root soon appeared, which extended rapidly to the earth, and now the branch is firmly supported.

122. Green-coloured aerial roots are frequently met with in endogenous plants. Such roots possess stomata. In the Ivy, root-like processes are produced from the stem, by means of which it attaches itself to trees, rocks, and walls. Those processes are subservient to the purposes of support rather than nutrition. In parasites, or plants which derive nourishment from other plants, such as Dodder (*Cuscuta*), roots are sometimes produced in the form of *suckers*, which enter into the cellular tissue of the plant preyed upon.


123. When roots have been exposed to the air for some time, they occasionally assume the functions of stems, losing their fibrils, and developing abnormal buds. Duhamel proved this experimentally, by causing the branches of a willow to take root while attached to the stem, and ultimately raising the natural roots into the air.

Forms of Roots.

124. The forms of roots depend upon the mode in which the axis descends and branches. When the central axis goes deep into the ground in a tapering manner, without dividing, a *tap-root* is produced (fig. 119). This kind of root is sometimes shortened, and becomes succulent, forming the *conical* root of carrot, or the *fusiform*, or spindle-shaped root of radish, or the *napiform* root of turnip; or it ends abruptly, thus constituting the *præmorse* (*præmorsus*, bitten) root of *Scabiosa succisa*; or is *twisted*, as in the *contorted* root of *Bistort*.

125. When the descending axis is very short, and at once divides into thin, nearly equal fibrils, the root is called *fibrous*, as in many grasses; when the fibrils become short and succulent, the root is *fasciculated*, as in *Ranunculus Ficaria* and *Asphodelus luteus* (fig. 120); when the succulent fibrils are of uniform size, and arranged like coral, the root is *coralline*, as in *Corallorhiza innata*; when some of the

fibrils are developed in the form of tubercules containing starchy matter, as in *Orchis*, the root is *tubercular* (fig. 121); when the fibrils enlarge in certain parts only, the root is *nodulose*, as in *Spiræa Filipendula* (fig. 122), or *moniliform*, as in *Pelargonium triste* (fig. 123), or *annulated*, as in *Ipecacuanha*.




126. Root of Dicotyledonous or Exogenous Plants.—In these plants the root in its early state, or the *radicle* as it is then called, is a prolongation of the stem, and elongates directly by its extremity. It then continues to grow in a simple or branched state (fig. 119). From this mode of root development, these plants have been called *Exorhizal* (ἐξω, outwards, and ῥίζα, a root), by Richard. In their after progress, these roots follow the arrangement seen in the woody part of the stem. In some cases, as in the Walnut and Horse-chestnut, there is a prolongation of the pith into the root to a certain extent.

127. Root of Monocotyledonous or Endogenous Plants.—In these plants the young root or radicle pierces the lower part of the axis (fig. 124 *r*), is covered with a cellular sheath, *c*, and gives rise to numerous fibrils, *r' r' r' r'*, which are similarly developed. These plants are therefore called by Richard, *Endorhizal* (ἐνδον, within); and the sheath is denominated *Coleorhiza* (κολεός, a sheath). In their after

Fig. 120.—Fasciculated root of *Asphodelus luteus*.
 Fig. 121.—Tubercular root of *Orchis*. Several of the radical fibres retain their cylindrical form, while two are tubercles containing starchy matter.
 Fig. 122.—Nodulose root of *Spiræa Filipendula*.
 Fig. 123.—Moniliform root of *Pelargonium triste*.

progress, they usually retain their compound character, consisting of fibrils, most of which often remain unbranched (figs. 120, 121). The first-formed roots which surround the axis, if the plant is perennial, gradually die, and others are produced in succession farther from the central axis. In Endogenous roots, the same structure is observed as in the stem. Thus, fig. 125 represents a section of a Palm root, com-



posed of cellular tissue, porous vessels, *v p*, scalariform vessels, *v s*, fibrous or woody tissue, *f*, and laticiferous vessels, *l*. Roots are pushed out from various parts of the stems of many Palms, and ultimately appear as part of the external integument.

128. Root of Acotyledonous or Acrogenous Plants.—In these plants, the young root is a development of superficial cells from no fixed point, and they have been called *Heterorhizal* (ἑτεροῖς, diverse). In their subsequent progress, these roots present appearances similar to those seen in the stem. They frequently appear in the form of fibres on the outer part of the stem, giving rise, by their accumulation at the base, to the conical appearance represented in fig. 116 *r a*.

Functions of Roots.

129. Roots fix the plant, either in the soil or by attachment to other bodies. They absorb nourishment by a process of imbibition or endosmose (§ 27), through their spongioles or cellular extremities.

Fig. 124.—Grain of wheat germinating. *g*, The mass of the grain. *t*, The young stem beginning to shoot upwards. *r*, The principal root of axis. *r' r' r' r'*, Lateral roots, covered like the preceding, with small hairs or threads. *c c c*, Coleorhiza or sheath, with which each of the roots is covered at its base, while piercing the superficial layer of the embryo.

Fig. 125.—Transverse section of part of the root of a Palm (*Diplothemum maritimum*), to show the mode in which the cells and vessels are arranged. *v p*, Large porous vessels situated in the interior. *v s*, Scalariform vessels more external, and becoming smaller the farther they are from the centre. *f*, Fibrous tissue, or elongated cells, accompanying the vessels. *l*, Groups of laticiferous vessels of different sizes, the larger being inside.

The experiment of Duhamel and Senebier, conducted by inserting at one time the minute fibrils alone into fluid, and at another, the axis of the root alone, showed clearly that the cellular extremities were the chief absorbing parts of the roots. Hence the importance, in transplanting large trees, of cutting the roots some time before, in order that they may form young fibrils and spongelets, which are then easily taken up in an uninjured condition, ready to absorb nourishment.

130. The elongation of the roots by their extremities, enables them to accommodate themselves to the soil, and allows the spongioles to extend deeply without being injured. Roots in their lateral extension, bear usually a relation to the horizontal spreading of the branches, so as to fix the plant firmly, and to allow fluid nutritive substances to reach the spongioles more easily. It is of importance to permit the roots to extend easily in all directions. By restricting or cutting the roots, the growth of the plant is to a certain degree prevented, although it is sometimes made to flower and bear fruit sooner than it would otherwise have done. The system of restrictive potting, formerly practised in green-houses, often destroyed the natural appearance of the plants. The roots filled the pots completely, and even raised the plants in such a way as to make the upper part of the root appear above the soil.

131. To roots there are sometimes attached reservoirs of nourishment, in the form of tubercules, containing starch and gum (fig. 121), which are applied to the nourishment of the young plant. These are seen in the Dahlia and in terrestrial Orchids. In epiphytic Orchids, on the other hand, the roots are aerial, and the stems are much developed, forming pseudo-bulbs. Upon the roots of *Spondias tuberosa* there exist round black-coloured tubercules, about eight inches in diameter, consisting internally of a white cellular substance, which is full of water. These tubercules seem to be intended to supply water to the tree during the dry season. They are often dug by travellers, each of them yielding about a pint of fluid of excellent quality.

132. Roots also give off certain excretions, which differ in different species. These are given off by a process of exosmose (§ 27), and consist both of organic and inorganic matter. They were examined by Macaire and Decandolle, and at one time they were thought to be injurious to the plant, and by their accumulation to cause its deterioration. It was also supposed, that while they were prejudicial to the species of plant which yielded them, they were not so to others, and that hence a rotation of crops was necessary. Daubeney and Gyde have found by experiment, that these excretions are not injurious, and it is now shown, that the necessity for rotation depends on the want of certain nutritive matters in the soil.* In very rich and fertile land, the same crop may be grown successively for many years.

* This subject is considered when the sources whence plants derive their nourishment are treated of.

LEAVES AND THEIR APPENDAGES.


Structure of Leaves.

133. Leaves are expansions of the bark, developed in a symmetrical manner, as lateral appendages of the stem, and having a connection with the internal part of the ascending axis. They appear at first as small projections of cellular tissue, continuous with the bark, and closely applied to each other. In the early stages of their development they are undivided. These cellular papillæ, from which they derive their origin, gradually expand in various ways, acquire vascular tissue, and ultimately assume their permanent form and position on the axis. They may be divided into *aerial* and *submerged* leaves, the former being produced in the air, and the latter under water.

134. **Aerial Leaves.**—These leaves consist of vascular tissue in the form of *veins*, *ribs*, or *nerves*, of cellular tissue or *parenchyma* filling up the interstices between the veins, and of an *epidermal* covering.

135. The **Vascular System** of the leaf is continuous with that of the stem, those vessels which occupy the internal part of the stem becoming superior in the leaf, while the more external become inferior. Thus, in the upper part of the leaf, which may represent the woody layers, there are spiral vessels (fig. 126 *t*), annular reticulated or porous vessels, *v*, and woody fibres, *f*; whilst in the lower side, which may represent the bark, there are laticiferous vessels and fibres, resembling those of liber, *l*. There are usually two layers of fibro-vascular tissue in the leaf, which may be separated by maceration. They may be seen in what are called *skeleton leaves*, in which the cellular part is removed, and the fibro-vascular left. The vascular system of the leaf is distributed through the cellular tissue in the form of simple or branching veins.


126



136. The **Epidermis** (fig. 127 *e s, e i*), composed of cells more or less compressed, has usually a different structure and aspect on the two surfaces of the leaf. It is chiefly on the epidermis of the lower surface (fig. 128 *e i*), that stomata, *s s*, are produced, occupying spaces between the veins, and it is there also that hairs usually occur. In

Fig. 126.—Bundle of fibro-vascular tissue, passing from a branch, *b*, into a petiole, *p*. The vessels are first vertical, then nearly horizontal, but they continue to retain their relative position. Changes take place in the size of the cells at the articulation *a*. *tt*, Tracheae, in which the fibre can be unrolled. *vv*, Annular vessels. *ff*, Woody fibres. *ll*, Cortical fibres, or fibres of liber.

these respects, the lower epidermis resembles the outer bark of young stems, with which it may be said to correspond. The lower epidermis is often of a dull or pale-green colour, soft, and easily detached. The upper epidermis (figs. 127 and 128 *es*) is frequently smooth and



127



128

shining, and sometimes becomes very hard and dense. Many tropical plants present on the upper surface of their leaves several layers of compressed epidermal cells. These appear to be essential for the preservation of moisture in the leaf. In leaves which float upon the surface of water, as those of the water-lily, the upper epidermis alone possesses stomata (§ 56). On removing a strip of epidermis, part of the parietes of the cells below is often detached in the form of a green net-work (fig. 129 *p p*), and on examination under the microscope, the stomata, *ss*, are seen communicating with colourless spaces, *l l l*, surrounded by green matter.



129

137. The **Parenchyma** of the leaf is the cellular tissue surrounding the vessels, and enclosed within the epidermis (fig. 127 *ps*, *pi*). It has sometimes received the names of *Diachyma* (διά, in the midst, and χῶμα, tissue), or *Mesophyllum* (μέσος, middle, and φύλλον, a leaf), or *Diploë*, (διπλοῖς, a covering). It is formed of two distinct series of cells, each containing chlorophylle or green-coloured granules, but differing in their form and arrangement. This may be seen on making a vertical section of a leaf, as in fig. 127. Below the epidermis of the upper side of the leaf, there are one or two layers of oblong blunt cells, placed perpendicularly to the surface (fig. 127 *p s*), and


Fig. 127.—Thin vertical section of the leaf of a Lily, highly magnified. *es*, Epidermis of upper pagina or surface. *ei*, Epidermis of lower surface. *ps*, Parenchyma of upper portion of the leaf, composed of close vertically-placed cells. *pi*, Parenchyma of lower portion, composed of loose horizontal cells. *m*, Intercellular passages. *l l*, Lacunæ.

Fig. 128.—Similar section of the leaf of Balsam. The letters denote the same parts as in fig. 127. *ss*, Stomata.

Fig. 129.—Strip of the lower epidermis, *ee*, of the leaf of Balsam, showing a net-work formed by a portion of the parenchyma below, *pp*, being detached. The spaces of the net are lacunæ, *l l l*, often corresponding to stomata, *ss*.

applied so closely to each other as to leave only small intercellular spaces (fig. 127 *m*), except when stomata happen to be present. On the under side of the leaf, the cells are irregular, often branched, and are arranged more or less horizontally (fig. 127 *p i*), leaving cavities between them, *l l*, which often communicate with stomata (fig. 128 *s s*). On this account the tissue has received the name of *cavernous*. The form and arrangement of the cells, however, depend much on the nature of the plant, and its exposure to light and air. Sometimes the arrangement of the cells on both sides of the leaf is similar, as occurs in leaves which have their edges presented to the sky. In very succulent plants, the cells form a compact mass, and those in the centre are often colourless. In some cases the cellular tissue is deficient at certain points, giving rise to distinct holes in the leaf, as in *Dracontium pertusum*. In *Victoria regia* perforations in the leaf seem to be subservient to the purposes of nutrition, in permitting the air collected beneath the large expanded leaf to escape, and so allowing its under surface to be brought into immediate contact with the water.

138. Submerged Leaves.—Leaves which are developed under water differ in many points of structure from aerial leaves. They have no fibro-vascular system, but consist of a congeries of cells which sometimes become elongated and compressed so as to resemble veins. They have a layer of compact cells on their surface (fig. 130 *p*), but no true epidermis, and no stomata. The internal structure consists of cells, disposed irregularly, and sometimes leaving spaces which are filled with air for the purpose of floating the leaf (fig. 130 *l*). When exposed to the air, these leaves easily part with their moisture, and become shrivelled and dry. In some instances there is only a net-work of



130



131

filamentous-like cells formed, the spaces between which are not filled with parenchyma, giving a peculiar skeleton appearance to the leaf, as in *Hydrogeton* or *Ouvirandra fenestralis* (fig. 131). Such a leaf has been called *fenestrate* (*fenestra*, a window).

Fig. 130.—Perpendicular section through a small portion of the submerged leaf of *Potamogeton perfoliatus*. *p*, Parenchyma. *l*, Lacunae.

Fig. 131.—Fenestrate leaf of *Ouvirandra fenestralis*, composed of vascular tissue, without intervening cellular tissue or diachyma.

139. A leaf, in general, whether aerial or submerged, consists of a flat expanded portion (fig. 132 *l*), called the *blade*, *limb*, or *laminar merithal*, (*μέρος*, a part, and *θαλλός*, a frond), of a narrower portion called the *petiole*, (*petiolus*, a little foot or stalk), *stalk*, or *petiolar merithal*



132

(fig. 132 *p*), and sometimes of a portion at the base of the petiole, which forms a *sheath* or *vagina* (fig. 132 *g*), or is developed in the form of leaflets, called *stipules* (fig. 191 *s*). The sheathing portion or *vaginal merithal* is sometimes incorporated with the stem, and has been called *Tigellary* (*tige*, Fr., a stem or stalk), by Gaudichaud. These portions are not always present. The sheathing, or stipulary portion, is frequently wanting, and occasionally only one of the other two is developed.

When a leaf has a distinct stalk it is called *petiolate*; when it has none, it is *sessile* (*sessilis*, from *sedeo*, I sit). When sessile leaves embrace the stem, they are called *amplexicaul* (*amplector*, I embrace, and *caulis*, a stem). The part of the leaf next the petiole or the axis is its *base*, while the opposite extremity is the *apex*. The surfaces of the leaf are called the *paginae* (*pagina*, a flat page), and its edges or margins form the *circumscription* of the leaf. The leaf is usually horizontal, so that the upper pagina is directed towards the heavens, and the lower pagina towards the earth; but in many cases leaves are placed vertically, as in some Australian *Acacias*, *Eucalypti*, &c.; in other instances, as in *Alströmeria*, the leaf becomes twisted in its course, so that what is superior at one part becomes inferior at another.

140. The upper angle formed by the leaf with the stem is called its *axil* (*axilla*, arm-pit), and everything arising at that point is called *axillary*. It is there that leaf-buds (¶ 178) are usually developed. The leaf is sometimes articulated with the stem, and when it falls off a *scar* or *cicatricula* remains; at other times it is continuous with it, and then decays gradually, while still attached to the axis. In their early state all leaves are continuous with the stem, and it is only in their after-growth that articulations are formed. When leaves fall off annually, they are called *deciduous*; when they remain for two or more years, they are *evergreen*. The laminar portion of a leaf is occasionally articulated with the petiole, as in the *Orange* (fig. 185), and a joint at times exists between the vaginal or stipulary portion and the petiole.

Fig. 132.—Leaf of *Polygonum Hydropiper*, with a portion of the stem bearing it. *l*, Limb lamina, or blade. *p*, Petiole or leaf-stalk. *g*, Sheath or vagina, embracing the stem, and terminated by a fringe.

Distribution of the Veins, or Venation of Leaves.

141. The distribution of the veins has been called *Venation*, sometimes *Nervation*. In most leaves this can be easily traced, but in the case of succulent plants, as Hoya, Agave, and Mesembryanthemum, the veins are obscure, and the leaves are said to be *Hidden-veined* (figs. 171, 172). In the lower tribes of plants, as sea-weeds, and in submerged leaves, there are no true veins, but only condensations of elongated cellular tissue, and the term *Veinless* (*avenia*) is applied. In an ordinary leaf, as that of Lilac or Chestnut, there is observed a central vein larger than the rest, called the *midrib* (fig 133 *m*); this gives off veins laterally (*primary veins*), *ns ns ns*, which either end in a curvature within the margin, as in Lilac (fig. 133), or go directly to the edge of the leaf, as in Oak and Chestnut (fig. 134). If they are curved, then external veins and marginal veinlets are interspersed through the parenchyma external to the curvature. There are also other veins of less extent (*costal veins*) given off by the midrib, and these give origin to small *veinlets*. In some cases, as Sycamore and Cinnamon, in place of there being only a single central rib, there are several which diverge from the part where the blade



133



134



135

joins the petiole or stem. Thus, the primary veins give off secondary veins, and these in their turn give off tertiary veins, and so on, until a

Fig. 133.—Leaf of Belladonna. *p*, Petiole or leaf-stalk. *nm*, Midrib. *ns ns ns*, Primary veins, ending in curvatures at their extremities.

Fig. 134.—Leaf of Oak, pinnatifid or divided into lobes in a pinnate manner; feather-veined, the veins going directly to the margin.

Fig. 135.—Leaf of Banana, showing midrib and primary veins running parallel and in a curved manner to the margin. No reticulation. Plant monocotyledonous.

complete net-work of vessels is produced. To such a distribution of veins, the name of *Reticulated* or *Netted* venation has been applied.

142. In the leaves of some plants there exists a central rib or midrib, with veins running nearly parallel to it from the base to the apex of the leaf, as in grasses (fig. 194) and Fan palms; or with veins coming off from it throughout its whole course, and running parallel to each other in a straight or curved direction towards the margin of the leaf, as in Plantain and Banana (fig. 135). In these cases the veins are often united by cross veinlets, which do not, however, form an angular net-work. These are called *Parallel-veined*.

143. Leaves may thus be divided into two great classes, according to their venation—*Reticulated* or *netted* leaves, in which there is an angular net-work of vessels, as occurs generally in the leaves of exogenous or dicotyledonous plants; and *Parallel-veined*, in which the veins run in a straight or curved manner from base to apex, or from the midrib to the margin of leaf, and in which, if there is a union, it is effected by transverse veins which do not form an angular net-work. This kind of leaf occurs commonly in endogenous or monocotyledonous plants.

144.—TABULAR ARRANGEMENT OF VENATION.

A.—Reticulated Venation.

1. *Unicostate* (*unus*, one). A single rib or costa in the middle (midrib).
 1. Primary veins coming off at different points of the midrib.
 - a. Veins ending in curvatures within the margin (fig. 133), and forming what have been called *true netted* leaves (Lilac).
 - b. Veins going directly to the margin (fig. 134), and forming *feather-veined* leaves (Oak and Chestnut).
 2. Primary veins coming off along with the midrib (fig. 143) from the base of the leaf.
- II. *Multicostate* (*multus*, many). More than one rib. In such cases there are frequently three (*tricostate*), as in fig. 162; or five (*quinquecostate*), as in fig. 158. Authors usually give to these leaves the general name of *costate*, or *ribbed*.
 1. *Concostate* (*con*, together, *costa*, a rib). Ribs converging, running from base to apex in a curved manner, as in Cinnamon, *Laurus Cinnamomum* (fig. 158). There is occasionally an obscure rib running close to the edge of the leaf, and called *intramarginal*, as in the Myrtle.
 2. *Discostate* (*dis*, separate). Ribs diverging or proceeding in a radiating manner; this is called *radiating venation*, and is seen in Sycamore, Vine, Geranium (figs. 144, 146).


B.—Parallel Venation.—The term parallel is not strictly applicable, for the veins often proceed in a radiating manner, but it is difficult to find a comprehensive term. This venation may be characterized as *not reticulated*.

- I. Veins proceeding from midrib to margin, usually with convexity towards the midrib, as in *Musa* and *Canna* (fig. 135).
- II. Veins proceeding from base to apex.
 1. Veins more or less convergent (fig. 173), as in *Iris*, *Lillies*, *Grasses* (fig. 194).
 2. Veins more or less divergent, as in *Fan Palms*.

To this may be added the venation common in *Ferns* where the veins divide in a forked manner. This venation has been called *Furcate* (*furca*, a fork).

Forms of Leaves.

145. Leaves have been divided into *simple* and *compound*. The former have no articulation beyond the point of their insertion on the stem, or consist of one piece only, which, however, may be variously divided (figs. 136, 137, 138, &c.). The latter have one or more articu-



136



137



138

lations beyond the point of their insertion on the stem, or consist of one or more leaflets (*foliola*) separately attached to the petiole or leaf-stalk (fig. 141). In the earliest stage of growth all leaves are simple and undivided, and it is only during the subsequent development that divisions appear. The forms which the different kinds of simple and compound leaves assume, are traced to the character of the venation, and to the amount of parenchyma produced.

146. **Simple Leaves.**—When the parenchyma is developed symmetrically on each side of the midrib or stalk, the leaf is *equal* (fig. 149); if otherwise, the leaf is *unequal* or *oblique* (fig. 136), as in *Begonia*. If the margins are even and present no divisions, the leaf is *entire* (*integer*), as in figs. 149 and 150; if there are slight projections of cellular or vascular tissue beyond the margin, the leaf is not entire (fig. 136); when the projections are irregular and more or less pointed, the leaf is *dentate* or toothed (fig. 155); when they lie regularly over each other, like the teeth of a saw, the leaf is *serrate* (figs. 136, 154); when they are rounded, the leaf is *crenate* (fig. 159). If the divisions extend

Fig. 136.—Leaf of *Ulmus effusa*. Reticulated venation; primary veins going to the margin, which is serrated. Leaf unequal at the base.

Fig. 137.—Pinnatifid leaf of *Valeriana dioica*.

Fig. 138.—Bipinnatifid leaf of *Papaver Argemone*. Feather-veined.

more deeply than the margin, the leaf receives different names according to the nature of the segments: thus, when the divisions extend about half way down (figs. 134, 144), it is *cleft* (*fissus*), and its segments



are called *fissures* (*fissura*, a cleft); when the divisions extend nearly to the base or to the midrib (fig. 170), the leaf is *partite*, and its segments are called *partitions*.

147. These divisions take place in simple leaves exhibiting different kinds of venation, and thus give rise to marked forms. Thus, if they occur in a *feather-veined* leaf (fig. 137), it becomes either *pinnatifid* (*pinna*, a wing or leaflet, and *fissus*, cleft), when the segments extend to about the middle and are broad; or *pectinate* (*pecten*, a comb), when they are narrow; or *pinnatipartite*, when the divisions extend nearly to the midrib. These primary divisions may be again subdivided in a similar manner, and thus a *feather-veined* leaf will become *bipinnatifid* (fig. 138), or *bipinnatipartite*; and still further subdivisions give origin to *tripinnatifid* and *lacinated* leaves. If the divisions of a pinnatifid leaf are more or less triangular, and are pointed downwards towards the base, the extremity of the leaf being undivided and triangular, the leaf is *runcinate* (*runcina*, a large saw), as in the Dandelion. When the apex consists of a large rounded lobe, and the divisions, which are also more or less rounded, become gradually smaller towards the base (fig. 139), as in *Barbarea*, the leaf is called *lyrate*, from its resemblance to an ancient lyre.* When there is a concavity on each side of a leaf, so as to make it resemble a violin, as in *Rumex pulcher* (fig. 140), it is called *panduriform* (*παυδοῦρρα*, fiddle).

148. The same kind of divisions taking place in a simple leaf with *radiating* venation, gives origin to the terms *lobed* or *cleft* (figs. 174, 146), when the divisions extend about half-way through the leaves: and thus they may be *three-lobed*, *five-lobed*, *seven-lobed*, *many-lobed*; or, *trifid*, *quinquefid*, *septemfid*, *multifid*, according to the number of

Fig. 139.—Lyrate leaf of *Barbarea*.

Fig. 140.—Panduriform, a fiddle-shaped leaf of *Rumex pulcher*.

Fig. 141.—Compound leaf; ternate, the leaflets being obcordate.

Fig. 142.—Compound leaf; quaternate, the leaflets being rotundate-cuneiform, or wedge-shaped with rounded apices.

Fig. 143.—Two-lobed leaf, somewhat cordate at the base, emarginate, and mucronate.

Fig. 144.—Palmate leaf, the divisions acute and serrated at their margins. Radiating venation.

* Under the term lyrate, some include compound pinnate leaves in which the several pinnae are united at the apex of the leaf, and the others become gradually smaller towards the base.

divisions. The name of *palmate*, or *palmatifid* (fig. 144), is applied to leaves with radiating venation, in which there are several fissures united by a broad expansion of parenchyma, like the palm of the hand, as in *Passion-flower* and *Rheum palmatum*; while *digitate* (*digitus*, a finger), or *digitipartite*, includes leaves in which there are deeper partitions, five in number, like the fingers, as in *Janipha*; and *dissected* applies to leaves with radiating venation, having numerous narrow divisions, as in *Geranium dissectum*. When in a radiating leaf there are three primary partitions and two lateral ones spreading and forming divisions on their inner margin only, as in *Helleborus* (fig. 170), the leaf is called *pedate* or *pedatifid* (*pes*, a foot), from a fancied resemblance to the claw of a bird.

149. In all the instances already alluded to, the leaves have been considered as flat expansions in which the ribs or veins spread out on the same planes with the stalk. In some cases, however, the veins spread at right angles to the stalk. If they do so equally on all sides, and are united by parenchyma, so that the stalk occupies the centre (fig. 145), the leaf becomes *orbicular* (*orbis*, a circle), as in *Hydrocotyle*; if unequally, so that the stalk is not in the centre, the leaf is *peltate* (*pelta*, a buckler), as in the *Castor oil plant* (fig. 146). The edges or margins of orbicular and peltate leaves are often variously divided.

150. It has been observed that the order of the venation in the leaf bears a close analogy to the arrangement of the branches on the stem. That a certain unity so pervades vegetable organization, that the root, the stem, and the leaves may, in their ultimate arrangement, be regarded as being typical the one of the other. The peculiar formation of the leaf is not a mere act of local organic construction, but an advanced step in progressive vital development, consummating a pre-existing plan for symmetrical arrangement.* It is impossible to notice all the forms of leaves without exceeding due limits. The following are enumerated as the most important. When the veins do not spread out, but run from the base to the apex with a narrow strip of parenchyma, the leaf is *linear* or





Fig. 145.—Orbicular leaf of *Hydrocotyle vulgaris*. Radiating venation. *p*, Petiole. *l*, Lamina.
Fig. 146.—Peltate leaf of *Ricinus communis*, or *Castor oil plant*. Radiating venation. *p*, Petiole or leaf-stalk. *l*, Lamina or blade.

* M'Cosh on the plant morphologically considered. *Proceed. of the Edin. Bot. Soc.*, July, 1851. *Bot. Gazette*, September, 1851.

acicular (fig. 147), as in Pines and Firs. When the veins diverge, those in the middle being longest, and the leaf tapering at each end (fig. 166), it becomes *lanceolate* (*lancea*, a spear). If the middle veins only exceed the others slightly, and the ends are convex, the leaf is either *rounded* (*rotundatus*), as in fig. 164, *elliptical* (fig. 162), *oval* (fig. 149), or *oblong* (fig. 150). If the veins at the base are longest, the leaf is *ovate* or egg-shaped, as in Chick-week (fig. 152), and if those at the apex are longest, the leaf is *obovate*, or inversely egg-shaped. Leaves are *cuneate* (*cuneus*, a wedge) or wedge-shaped, in Saxifraga (fig. 155); *spatulate*, or spatula-like, having a broad rounded apex, and tapering down to the stalk in the Daisy (fig. 148); *subulate* (fig. 167), or narrow and tapering like an awl (*subula*); *acuminate*, or drawn out into a long point, as in Ficus religiosa (fig. 159), *mucronate*, with a hard stiff point or mucro at the apex (figs. 160 and 143). When the parenchyma



is deficient at the *apex* so as to form two rounded lobes, the leaf is *obcordate* or inversely heart-shaped; when the deficiency is very slight, the leaf is called *emarginate* (fig. 143) as having a portion taken out of the margin; when the apex is merely flattened or slightly depressed



Fig. 147.—Linear, or acicular leaf of Fir.

Fig. 148.—Spatulate leaf of Daisy.

Fig. 149.—Oval leaf.

Fig. 150.—Oblong leaf.

Fig. 151.—Petiolate, reticulated, somewhat oblong leaf, truncate at the base.

Fig. 152.—Ovate pointed leaf.

Fig. 153.—Cordate pointed leaf.

Fig. 154.—Ovato-lanceolate leaf, i. e. lanceolate in its general contour, but ovate at the base doubly serrated, or having large and small serratures alternately at the margin.

Fig. 155.—Cuneate or wedge-shaped leaf of Saxifraga, ending in an abrupt or truncate manner, and toothed or dentate at the apex.

Fig. 156.—Perfoliate leaf of Bupleurum, formed by lobes uniting at the base on the opposite side of the stem from that to which the leaf is attached.


Fig. 157.—Retuse leaf, i. e. slightly depressed at the apex. Margin slightly waved.

Fig. 158.—Ovate, five-ribbed leaf.

Fig. 159.—Rounded acuminate leaf of Ficus religiosa, with the margin crenate or slightly sinuous.

Fig. 160.—Sub-ovate, retuse, mucronate leaf.

(fig. 157), the leaf is *retuse* (*retusus*, blunt); and when the apex ends abruptly in a straight margin, as in the Tulip tree (fig. 163), the leaf is *truncate*. When the venation is prolonged downwards at an obtuse angle with the midrib, and rounded globes are formed, as in Dog-violet, the leaf is *cordate* or heart-shaped (fig. 153), or kidney-shaped (*reniform*) when the apex is rounded (fig. 161), as in Asarum. When the lobes are prolonged downwards and acute (fig. 165), the leaf is *sagittate* (*sagitta*, an arrow); when they proceed at right angles, as in Rumex Acetosella, the leaf is *hastate* (*hasta*, a halbert) or halbert-shaped. When a simple leaf is divided at the base into two leaf-like appendages (fig. 169), it is called *auriculate* (*auricula*, the ear). When the veins spread out in various planes, and there is a large development of cellular tissue, so as to produce a succulent leaf, such forms occur as *conical*, *prismatical*, *ensiform* or sword-like (*ensis*, a sword), *acinaciform* (*acinaces*, a scimitar), or scimitar-shaped (fig. 172), and *dolabriform* (*dolabra*, an axe) or axe-shaped (fig. 171). When the development of cells is such that they more than fill up the spaces between the veins, the margins become *wavy*, *crisp*, or *undulated*, as



in Rumex crispus and Rheum undulatum (fig. 174). By cultivation the cellular tissue is often much increased, giving rise to the *curled* leaves of Greens, Savoy, Cresses, Lettuce, &c.

151. **Compound Leaves** are those in which the divisions extend to

Fig. 161.—Reniform or kidney-shaped entire leaf of Asarum. Radiating venation.

Fig. 162.—Elliptical and somewhat lanceolate leaf; three-ribbed.

Fig. 163.—Three-lobed, truncate, or abrupt leaf of Liriodendron tulipifera.

Fig. 164.—Rounded entire leaf, ending in a short point.

Fig. 165.—Sagittate or arrow-shaped leaf of Sagittaria.

Fig. 166.—Lanceolate, acute leaf, with minute teeth or dentations at the margin.

Fig. 167.—Subulate or awl-shaped leaf.

Fig. 168.—Whorl or verticill of linear-ovate leaves.

Fig. 169.—Auriculate lanceolate leaf, oblique at the base, with minute toothings at the margin.

the midrib, or petiole (fig. 175), and receive the name of *foliola* or *leaflets*. The midrib, or petiole, has thus the appearance of a branch



with separate leaves attached to it, but it is considered properly as one leaf, because in its earliest state it arises from the axis as a single piece, and its subsequent divisions in the form of leaflets are all in one plane. When a compound leaf dies, it usually separates as one piece. The leaflets are either sessile (fig. 176), or have stalks, called



petiolules (fig. 175), according as the vascular bundles of the veins spread out or divaricate at once, or remain united for a certain length.

Fig. 170.—Pedate or pedatifid leaf of Hellebore. Radiating venation.

Fig. 171.—Dolabriform or axe-shaped fleshy succulent leaf. Hidden veined.

Fig. 172.—Acinaciform or scimitar-shaped succulent leaf. Hidden-veined.

Fig. 173.—Oval leaf with converging veins; not reticulated.

Fig. 174.—Palmately-lobed leaf, crisp or undulated at the margin. Radiating venation.

Fig. 175.—Leaf of *Robinia pseudo-acacia*, often called *Acacia*. The leaf is impari-pinnate, or alternately pinnate. The pinnae are supported on stalks or petiolules. *p*, Petiole or leaf-stalk.

l, Lamina or blade divided into separate leaflets or pinnae

Fig. 176.—Septenate leaf of *Æsculus Hippocastanum* or Horse Chestnut *p*, Petiole. *l*, Lamina, divided into seven separate leaflets.

152. Compound leaves have been classified according to the nature of the venation, and the development of parenchyma. In a feather-veined leaf, if the divisions extend to the midrib, and each of the primary veins spreads out or branches so as to become covered with parenchyma, and thus form separate leaflets, which are usually articulated to the petiole or midrib (fig. 177), the leaf is *pinnate* (*pinna*, a wing or feather). If the midrib and primary veins are not covered with parenchyma, while the *secondary* (or those coming off in a feather-like manner from the primary veins) are, and separate leaflets are thus formed which are usually articulated with the veins, the leaf is *bipinnate* (fig. 178). In this case the secondary veins form as it were partial petioles. A farther sub-division, in which the *tertiary* veins only are covered with parenchyma and have separate leaflets, gives *tripinnate* or *decompound*, in which case the tertiary veins form the partial petioles; and a leaf divided still more is called *supradecomound* (fig. 179).


153. When a pinnate leaf has one pair of leaflets, it is *unijugate*



177



178



179

(*unum*, one, and *jugum*, a yoke); when it has two pairs, it is *bijugate*; many pairs, *multijugate* (fig. 175). When a pinnate leaf ends in a pair of pinnæ (fig. 177,) it is *equally* or *abruptly pinnate* (*pari-pinnate*); when there is a single terminal leaflet (fig. 175), the leaf is *unequally pinnate* (*impari-pinnate*); when the leaflets or pinnæ are placed alternately on either side of the midrib, and not directly opposite to each


Fig. 177.—Pari-pinnate leaf with six pairs of pinnæ (*sexjugate*).

Fig. 178.—Bipinnate leaf, with sessile foliola or leaflets.

Fig. 179.—Part of the supradecomound leaf of *Laserpitium hirsutum*.

other, the leaf is *alternately pinnate* (fig. 175); and when the pinnæ are of different sizes, the leaf is *interruptedly pinnate* (fig. 180).

154. In the case of a leaf with radiating venation, if the ribs are separately covered with parenchyma, and each leaflet is articulated to the petiole, the leaf becomes *ternate* (figs. 141, 181) if there are three divisions; *quaternate*, if four (fig. 142); *quinate*, if five; *septenate*, if seven (fig. 176), and so on.* If the three ribs of a ternate leaf subdivide each into three primary veins, which become covered with parenchyma so as to be separate articulated leaflets, the leaf is *biterbate*; and if another three-fold division takes place, it is *triterbate* (fig. 182).



180

181

182

155. **Petiole or Leaf-stalk.**—This is the part which unites the limb or blade of the leaf to the stem (figs. 132, and 175 *p*). It is absent in *sessile* leaves, and in many sheathing leaves is not well defined. It consists of one or more bundles of vascular tissue, with a varying amount of parenchyma. The vessels are, spiral-vessels connected with the medullary sheath in Exogens and with the fibro-vascular bundles in Endogens, porous vessels and other forms of fibro-vascular tissue, woody tissue, and laticiferous vessels. These vessels are enclosed in an epidermal covering, with few stomata, and are more or less compressed. When the vascular bundles reach the base of the lamina, they separate and spread out in various ways, as already described under venation. A large vascular bundle is continued through the lamina to form the midrib (fig. 133, *nm*), and sometimes several large bundles form separate ribs (figs. 146, 162), whilst the ramifications of the smaller bundles constitute the veins.

* Some apply the term *digitate* to radiating compound leaves with five or seven leaflets.
Fig. 180.—Impari and alternately pinnate leaf. Leaflets or pinnæ sessile, and serrated at the margin.

Fig. 181.—Ternate leaf of Strawberry. Margin of leaflets, toothed or dentate. *p*, Petiole with projecting hairs. *l*, Lamina divided into three leaflets.


Fig. 182.—Triterbate leaf. Leaflets cordate.

156. At the place where the petiole joins the stem, there is frequently an articulation or a constriction with a tendency to disunion, and at the same time there exists a swelling (fig. 203 *p*), called *pulvinus* (*pulvinus*, a cushion), formed by a mass of cellular tissue. The fully developed leaf is thus seen, not sharply cut off from the stem, but running gradually into it. At other times the petiole is not articulated, but is either continuous with the stem, or forms a sheath around it. At the point where the petiole is united to the lamina, or where the midrib joins the leaflets of a compound leaf, there is occasionally a cellular dilatation called *struma* (*struma*, a swelling), with an articulation. This articulation or joint is by many considered as indicating a compound leaf, and hence the leaf of the orange is considered as such, although it consists of one undivided lamina (fig. 185). In articulated leaves, the pulvinus may be attached either to the petiole or to the axis, and may fall with the leaf, or remain attached to the stem. When articulated leaves drop, their place is marked by a cicatrix or scar, seen below the bud in fig. 203. In this scar, the remains of the vascular bundles, *c*, are seen; and its form furnishes characters by which particular kinds of trees may be known when not in leaf. Where no petiole exists, or where the leaf is not deciduous, when the cycle of vegetable life is perfected, it may be observed to wither on the stem.

157. The petiole varies in length, being usually shorter than the lamina, but sometimes much longer. In some palms it is fifteen or twenty feet long, and is so firm as to be used for poles or walking-sticks. In general,



183



184



185

the petiole is more or less rounded in its form, the upper surface

Fig. 183.—Leaf with a quadrangular toothed lamina or blade, *l*, and an inflated petiole, *p*, containing air cells.

Fig. 184.—Ascidium or pitcher of *Nepenthes*. *p*, Winged petiole which becomes narrowed, and then expands so as to form the pitcher by being folded on itself. *e*, The operculum or lid, formed by the blade of the leaf, and articulated to the pitcher.

Fig. 185.—Leaf of Orange, which some call compound. *p*, Dilated or winged petiole, united by an articulation to the blade. In such a leaf, if the vessels of the petiole were developed in a circular manner, so as to form a pitcher, the lamina or blade would form the jointed lid.

being flattened or grooved. Sometimes it is compressed laterally, as in the Aspen, and to this peculiarity the trembling of the leaves of this tree is attributed. In aquatic plants, the leaf-stalk is sometimes distended with air (fig. 183 *p*), as in *Pontederia* and *Trapa*, so as to float the leaf. At other times it is winged, or has a leaf-like appearance, as in the pitcher plant (fig. 184 *p*), orange (fig. 185 *p*), lemon, and *Dionæa* (fig. 186 *p*). In some Australian *Acacias*, and in some species of *Oxalis*, *Bupleurum*, &c., the petiole is flattened in a vertical direction, the vascular bundles separating immediately after quitting



the stem, and running nearly parallel from base to apex. This kind of petiole (fig. 188 *p*), has been called *Phyllodium* (φυλλον, a leaf, and ειδος, form). In these plants the laminae or blades of the leaves are pinnate, bipinnate, or ternate, and are produced at the extremities of the phyllodia in a horizontal direction (fig. 188 *l*); but in many instances they are not developed, and the phyllodium serves the purpose

Fig. 186.—Leaf of *Dionæa muscipula*, or 'Venus' Fly-trap. *p*, Dilated or winged petiole. *e*, Jointed blade, the two fringed halves of which fold on each other, when certain hairs on the upper surface are touched.

Fig. 187.—Ascidium, or Pitcher of *Sarracenia*, formed by the petiole of the leaf. The lid is not articulated to the pitcher as in *Nepenthes* (fig. 184).

Fig. 188.—Leaf of *Acacia heterophylla*. *p*, Phyllodium or enlarged petiole, with straight venation. *l*, Lamina or blade which is bipinnate. The blade is frequently wanting, and the phyllodium is the only part produced.

of a leaf. Hence, some *Acacias* are called leafless. These phyllodia, by their vertical position, and their peculiar form, give a remarkable aspect to vegetation. On the same *Acacia*, there occur leaves with the petiole and lamina perfect; others having the petiole slightly expanded or winged, and the lamina imperfectly developed; and others in which there is no lamina, and the petiole becomes large and broad. Some petioles, in place of ending in a lamina, form a *tendrill* or *cirrhus* (§ 201), so as to enable the plant to climb.

Stipules.


158. At the place where the petiole joins the axis, a sheath (*vagina*) is sometimes produced, which embraces the whole or part of the circumference of the stem (fig. 132 *g*). This sheath is formed by the divergence of the vascular bundles which separate so as to form a hollow cavity towards the stem. The sheath is occasionally developed to such a degree as to give a character to the plants. Thus, in the *Rhubarb* tribe, it is large and membranous, and has received the name of *ochrea* or boot (fig. 132 *g*); while in *Palms* it forms a kind of network, to which the name of *reticulum* has been given (§ 57); and in umbelliferous plants, it constitutes the *pericladium* (περίκλιον, around, and κλάδος, a branch). In place of a sheath, leaves are occasionally produced at the base of the petiole (fig. 189 *s s*), which have been denominated *stipules* (*stipula*, straw or husk). These stipules are often two in number, and they are important as supplying characters in certain natural orders. Thus they occur in the *Pea* and *Bean* family, in *Rosaceous* plants, and the *Cinchona* bark family. They are rarely met with in *Endogens*, or in *Exogens* with sheathing petioles, and they are not common in *Exogens* with opposite leaves. Plants having stipules, are *stipulate*; those having none, are *extipulate*.

159. Stipules are formed by some of the vascular bundles diverging as they leave the stem, and becoming covered with parenchyma so as to resemble true leaves. Like leaves they are large or small, entire or divided, deciduous or persistent, articulated or nonarticulated. Their lateral position at the base of the petiole, distinguishes them from true leaves. In the *Pansy*, the true leaves are stalked and crenate, while the stipules are large, sessile, and pinnatifid. In *Lathyrus aphaca*, and some other plants, the true pinnate leaves are abortive, the petiole forms a tendrill, and the stipules alone are developed, performing the office of leaves.



Fig. 189.—Portion of a branch, *r*, of *Salix aurita*, bearing a single petiolate leaf, *f*, which has been cut across. *s s*, Stipules. *b*, Bud in the axil of the leaf.


160. When stipules are attached separately to the stem at the base of the leaf, they are called *caulinary*. Thus, in fig. 189, *r* is a branch of *Salix aurita*, with a leaf, *f*, having a bud, *b*, in its axil, and two caulinary stipules, *s s*. When stipulate leaves are opposite to each other at the same height on the stem, it occasionally happens that the stipules at either side unite wholly or partially, so as to form an *interpetiolar* or *interfoliar* (*inter*, between) stipule, as in *Cinchona* (fig. 190 *s*). In the case of alternate leaves, the stipules at the base of



190



191



192



193

each leaf are sometimes united to the petiole and to each other, so as to form an *adnate*, *adherent*, or *petiolar* stipule, as in the Rose (fig. 191 *s*), or an *axillary* stipule, as in *Houttuynia cordata* (fig. 192 *s*).

Fig. 190.—Branch, *r*, and two leaves, *f f*, of *Cephalanthus occidentalis*. *s*, Interpetiolar or interfoliar stipule, formed by the partial union of two.

Fig. 191.—Portion of a branch, *r*, of *Rosa canina*, or dog-rose, bearing a single leaf, *f*, with its petiole, *p*, its petiolar or adnate stipules, *s*, its axillary bud, *b*, and its aculei or prickles, *a*.

Fig. 192.—Portion of a branch, *r*, of *Houttuynia cordata*, with a leaf, *f*, and an axillary stipule, *s*, formed by the union of two.

Fig. 193.—Branch, *r*, and portion of the leaf, *f*, of *Astragalus Onobrychis*, with a synochreate stipule, formed by the union of two stipules on the opposite side of the branch from that to which the leaf is attached. The leaf is pinnate, and in the figure three pairs of leaflets or pinnae are left.

In other instances, the stipules unite together on the side of the stem opposite the leaf, and become *synochreate* (συν, together), as in *Astragalus* (fig. 193 s). The union or adhesion of stipules is not an accidental occurrence taking place after they have been developed; but is intimately connected with the general law, in accordance with which the parts of the plants are formed.

161. Stipules are sometimes large, enveloping the leaves in the young state, and falling off in the progress of growth, as in *Ficus*, *Magnolia*, and *Potamogeton*; at other times they are so minute as to be scarcely distinguishable without the aid of a lens, and so fugaceous as to be visible only in the very young state of the leaf. In grasses, the sheath or sheathing petiole (fig. 194 g v) has a prolongation or folding of the epidermis* at its upper part, distinct from the leaf, to which the name of *ligule* (*ligula*, a small slip) has been given (fig. 194 g l). Some consider it as equivalent to a stipule. It is either long or short, acute or blunt, entire or divided, and thus gives rise to various characters. At the base of the leaflets or foliola of a compound leaf, small stipules are occasionally produced, to which some have given the name of *stipels*.



Anomalous Forms of Leaves and Petioles.

162. Variations in the structure and forms of leaves and leaf-stalks are produced by the increased development of cellular tissue, by the abortion or degeneration of parts, by the multiplication or repetition of parts, and by adhesion. When cellular tissue is developed to a great extent, leaves become succulent, and occasionally assume a crisp or curled appearance. Such changes take place naturally, but they are often increased by the art of the gardener; and the object of many horticultural operations is to increase the bulk and succulence of leaves. It is in this way that Cabbages and Greens are rendered more delicate and nutritious.

163. In some plants true leaves are not produced, their place being occupied by dilated petioles or phyllodia (§ 157), or by stipules (§ 159). In other instances scales are formed instead of leaves, as in

* See *Deduplication*, under the head of *Corolla*.

Fig. 194.—Portion of a leaf of *Phalaris arundinacea*, one of the grasses. *f*, Laminar merithal or blade of the leaf, with straight parallel venation. *g v*, Vaginal, or sheathing portion representing the petiole, ending in a membranous process or ligule, *g l*.

Orobanche, Lathræa, and young Asparagus (fig. 110 l). Divisions take place in leaves when there is a multiplication of their parts; and a union of two or more leaves, or of parts of leaves, occurs in many cases. When two lobes at the base of a leaf are prolonged beyond the stem and unite (fig. 156), the leaf is *perfoliate* (*per*, through, and *folium*, leaf), the stem appearing to pass through it, as in *Bupleurum perfoliatum*, and *Chlora perfoliata*; when two leaves unite by their bases they become *connate* (*con*, together, and *natus*, born), as in *Lonicera Caprifolium*; and when leaves adhere to the stem, forming a sort of winged or leafy appendage, they are *decurrent* (*decurro*, to run down or along), as in Thistles.

164. The vascular bundles and cellular tissue are sometimes developed in such a way as to form a circle, with a hollow in the centre, and thus give rise to what are called *fistular* (*fistula*, a pipe) or hollow leaves, and to *ascidia* (ασκίδιον, a small bag) or *pitchers*. Hollow leaves are well seen in the Onion. Pitchers are formed either by petioles or by laminæ, and they are composed of one or more leaves. In some *Convallarias*, two leaves unite to form a cavity. In *Sarracenia* (fig. 187) and *Heliamphora*, the pitcher is composed apparently of the petiole of the leaf. In *Nepenthes* (fig. 184) and perhaps in *Cephalotus*, while the folding of a winged petiole, *p*, forms the pitcher, *a*, the lid, *e*, which is united by an articulation, corresponds to the lamina. This kind of ascidium is called *calyptrimorphous* (καλύπτρα, a covering, and μορφή, form), and may be considered as formed by a leaf such as that of the Orange (fig. 185); the lamina, *e*, being articulated to the petiole, *p*, which, when folded, forms the pitcher. In *Dischidia Rafflesiana*, a climbing plant of India, the pitchers, according to Griffith, are formed by the lamina of the leaf, and have an open orifice into which the rootlets at the upper part of the plant enter. These pitchers would seem therefore to contain a supply of fluid for the nourishment of the upper branches of the plant. In *Utricularia*, the leaves form sacs called *ampullæ*. In *Bryophyllum Calycinum*, buds are produced along the margin of the leaf, which are capable of originating new plants when placed in circumstances favourable to their growth.

Structure and Form of Leaves in the Great Divisions of the Vegetable Kingdom.

165. **Exogenous or Dicotyledonous Leaves.**—In Exogens, the venation is reticulated, the veins coming off at various angles and forming an angular net-work of vessels (fig. 136), and the tracheæ communicating with the medullary sheath. They are frequently articulated, exhibit divisions at their margin, and become truly compound. There are no doubt instances in which the veins proceed in a parallel manner, but this will be found to occur chiefly in cases where the petiole may be considered as occupying the place of the leaf. Examples of

this kind are seen in *Acacias* (fig. 188), as well as in *Ranunculus gramineus*, and *Lingua*.

166. Endogenous or Monocotyledonous Leaves.—In Endogens, the leaves do not present an angular net-work of vessels, nor do they exhibit divisions on their margin. Their venation is generally parallel, and their margin entire (figs. 135, 194). Exceptions to this rule occur in some plants, as *Tamus* and *Dioscorea*, which have been called Dictyogens by Lindley, on account of their somewhat netted venation; and in Palms, in which although the leaves are entire at first, they afterwards become split into various lobes. Endogenous leaves are rarely stipulate, unless the ligule of grasses be considered as being a stipule. Their leaves are often sheathing, continuous with the stem (forming a spurious stem in Bananas), and do not fall off by an articulation. When there is only a slight divergence of their veins, they may be looked upon more as enlarged and flattened petioles than as true laminæ. This remark is illustrated by the leaves of *Typha* and *Iris*. In some Endogens, as in *Sagittaria sagittifolia*, the submerged and floating leaves are narrow, like petioles, while those growing erect above the water expand and assume an arrow-like shape (fig. 165).


167. Acrogenous or Acotyledonous Leaves.—In Acrogens, the leaves vary much; being entire or divided, petiolated or sessile, often feather-veined, occasionally with radiating venation, the extremities of the veins being forked. The fibro-vascular bundles of the leaves resemble those of the stem both in structure and arrangement. In Thallogens, the leaves when present have no vascular venation. In many of them, as Lichens, Fungi, and Algæ, there are no true leaves.

Phyllotaxis, or the Arrangement of the Leaves on the Axis.

168. Leaves occupy various positions on the stem and branches, and have received different names according to their situation. Thus, leaves arising from the crown of the root, as in the Primrose, are called *radical*; those on the stem are *cauline*; on the branches, *ramal*; on flower-stalks, *floral leaves*. The first leaves developed are denominated *seminal* (*semen*, a seed), or *cotyledons* (*κοτυληδών*, a name given to a plant); and those which succeed are *primordial* (*primus*, first, and *ordo*, rank).

169. The arrangement of the leaves on the axis and its appendages is called *phyllotaxis* (*φύλλον*, a leaf, and *τάξις*, order). In their arrangement leaves follow a definite order. It has been stated already (§ 67) that there are regular *nodes* or points on the stem (fig. 195 n) at which leaves appear, and that the part of the stem between the nodes is the *internode* or *merithal* (fig. 195 m). Each node is capable of giving origin to a leaf. Occasionally several nodes are approximated so as to form as it were one, and then several leaves may be produced at the same height on the stem. When two leaves are thus produced,

one on each side of the stem or axis, and at the same level, they are called *opposite* (fig. 196); when more than two are produced (figs. 168, 197), they are *verticillate* (*verto*, I turn), and the circle of leaves is then



called a *verticil* or *whorl*. When leaves are opposite, the pairs which are next each other, but separated by an internode, often cross at right angles (fig. 196 *a b*), or *decussate* (*decusso*, I cut crosswise), following thus a law of alternation. The same occurs in verticils, the leaves of each whorl being alternate with those of the whorl next to it; or, in other words, each leaf in a whorl occupying the space between two leaves of the whorl next to it. There are considerable irregularities, however, in this respect, and the number of leaves in different whorls is not always uniform, as may be seen in *Lysimachia vulgaris*.

170. When a single leaf is produced at a node, and the nodes are separated so that each leaf occurs at a different height on the stem, the leaves are *alternate* (fig. 198). The relative position of alternate leaves varies in different plants, although it is tolerably uniform in each

Fig. 195.—Portion of a branch of a Lime tree, with four leaves arranged in a distichous manner, or in two rows. *a*, The branch with the leaves numbered in their order, *n* being the node, and *m* the internode or merithal. *b*, Is a magnified representation of the branch, showing the cleatrics of the leaves and their spiral arrangement, which is expressed by $\frac{1}{4}$, or one turn of the spiral and two leaves.

Fig. 196.—Opposite, decussate leaves of *Pimelea decussata*. *a*, A pair of opposite leaves. *b*, Another pair placed at right angles.

Fig. 197.—Leaves of *Lysimachia vulgaris*, in verticils or whorls of three. The leaves of each verticil alternate with those of the verticils next it. In this plant the number of the leaves in a verticil often varies.

species. In fig. 195, leaf 1 arises from a node, *n*; leaf 2 is separated by an internode or merithal, *m*, and is placed to the right or left; while leaf 3 is situated directly above leaf 1. The arrangement in this case is *distichous* (δίς, twice, and *στίχος*, order), or the leaves are arranged in two rows. In fig. 199, on the other hand, the fourth leaf is that directly above the first, and the arrangement is *tristichous* (τρεῖς, three, and *στίχος*, order). The same arrangement continues throughout the stems, so that in fig. 199 the 7th leaf is above the 4th, the 10th above the 7th; also the 5th above the 2nd, the 6th above the 3rd, and so on. There is thus throughout a tendency to a spiral arrangement, the number of leaves in the spire or spiral cycle, and the number of turns varying in different plants. In plants whose leaves are close to each other, the spiral tendency is easily seen. In the Screw pine (*Pandanus odoratissimus*), in the Pine-apple family, and in some Palms, as *Corypha cerifera*, the screw-like arrangement of the leaves is obvious. This mode of development prevails in all parts of plants, and may be considered as depending on their manner of growth in an upward and at the same time in a lateral direction. Alternation is looked upon as the normal arrangement of all parts of plants. This arrangement is liable to be interrupted by many causes, so that its distinct existence cannot be always detected.


171. In a regularly-formed straight branch covered with leaves, if a thread is passed from one to the other, turning always in the same direction, a spiral is described, and a certain number of leaves and of complete turns occur before reaching the leaf directly above that from which the enumeration commenced. This arrangement has been reduced to mathematical precision,* and Braun has expressed it by a fraction, the numerator of which indicates the number of turns, and the denominator the number of leaves in the spiral cycle. Thus, in fig. 198 *a b*, the cycle consists of five leaves, the 6th leaf



Fig. 198.—Part of a branch of a Cherry with six leaves, the 6th being placed vertically over the first, after two turns of the spiral. This is expressed by $\frac{2}{5}$ or the quincunx. *a*, The branch, with the leaves numbered in order. *b*, A magnified representation of the branch, showing the cicatrices of the leaves or their points of insertion, and their spiral arrangement.

* For a full account of Phyllotaxis, see Bravais Mem. sur la Disposition Géométrique des Feuilles. Annales des Sciences Naturelles. Jan. and Feb. 1837.

being placed vertically over the first, the 7th over the 2nd, and so on; while the number of turns between the 1st and 6th leaf is two: hence, this arrangement is indicated by the fraction $\frac{2}{3}$. In other words, the distance or divergence between the first and second leaf, expressed in parts of a circle, is $\frac{2}{3}$ of a circle, or $360 \div \frac{3}{2} = 144^\circ$.




199

In fig. 195, *a b*, the spiral is $\frac{1}{2}$, i.e. one turn and two leaves; the third leaf being placed vertically over the first, and the divergence between the first and second leaf being one-half the circumference of a circle, $360 \div \frac{1}{2} = 180^\circ$. Again, in fig. 199, *a b*, the number is $\frac{1}{3}$, or one turn and three leaves, the angular divergence being 120° .

172. In cases where the internodes are very short, and the leaves are closely applied to each other, as in the House-leek, it is difficult to trace what has been called the *generating spiral*, or that which passes through every leaf of the cluster. Thus in fig. 200, there are thirteen leaves which are numbered in their order, and five turns of the spiral marked by circles in the centre ($\frac{5}{13}$ indicating the arrangement); but this could not be detected at once. So also in Fir cones (fig. 201), which are composed of scales or modified leaves, the generating spiral cannot be determined easily. In such cases, however, there are secondary spirals running parallel to each other, as is seen in fig. 201, where spiral lines pass through scales numbered 1, 6, 11, 16, &c., and 1, 9, 17, &c., and by counting those which run parallel in different directions, the number of scales intervening between every two in the same parallel coil may be ascertained. Thus, in fig. 201, it will be found that there are five secondary spirals running towards the right and parallel to each other, the first passing through the scales 1, 6, 11, 16, &c.; the second through 9, 14, 19, 24, &c.; the third through 17, 22, 27, 32, 37, &c.; the fourth through 30, 35, 40, 45, &c.; the fifth through 43, 48, 53, &c. The number of these secondary spirals indicates the number of scales intervening between every two scales in each of these spirals—the common difference being five. Again, it will be found on examination that there is a number of secondary spirals running to the left, in which the common difference between every two scales is eight, and that this corresponds to the number of

Fig. 199.—Young plant of *Cyperus esculentus*, with leaves in three rows, or tristichous, expressed by the fraction $\frac{1}{3}$, or one turn and three leaves. *a*, The plant, with its leaves numbered in their order. *b*, Magnified representation of the stem, showing the insertion of the leaves and their spiral arrangement.

secondary spirals, the first of which passes through the scales 1, 9, 17, &c.; the second through 6, 14, 22, 30, &c.; the third through 3, 11, 19, 27, 35, 43, and so on. Thus it is that, by counting the secondary spirals, all the scales may be numbered, and, by this means, the generating spiral may be discovered.



173. The primitive or generating spiral may pass either from right to left or from left to right. It sometimes follows a different direction in the branches from that pursued in the stem. When it follows the same course in the stem and branches, they are *homodromous* ($\delta\mu\omicron\iota\omicron\varsigma$, similar, and $\delta\rho\omicron\mu\omicron\varsigma$, a course); when the direction differs, they are *heterodromous* ($\epsilon\tau\epsilon\rho\omicron\varsigma$, another. In different species of the same genus the phyllotaxis frequently varies.

174. Considering alternation as the usual leaf-arrangement, some have supposed that opposite leaves are owing to the development of two spirals in opposite directions, while others look upon them as produced by two nodes coming close together without an internode. A verticil, in the latter view, will be the result of the non-development of more than one internode, and may occur on plants, the normal arrangement of whose leaves are alternate. Thus, in fig. 195, if the space between 1 and 2 were obliterated, or the internode, *m*, not developed,

Fig. 200.—Cycle of thirteen leaves placed closely together so as to form a rosette, as in *Sempervivum*. A, is the very short axis .o which the leaves are attached. The leaves are numbered in their order, from below upwards. The circles in the centre indicate the five turns of the spiral, and show the insertion of each of the leaves. The divergence is expressed by the fraction $\frac{5}{13}$.

Fig. 201.—Cone of *Pinus alba*, with the scales or modified leaves numbered in the order of their arrangement on the axis of the cone. The lines indicate a rectilinear series of scales, and two lateral secondary spirals, one turning from left to right, the other from right to left.

the leaves would be opposite. In fig. 198, if the spaces between each of the leaves were obliterated, there would be a verticil of five leaves. In many plants there is a law of arrestment of development, by which opposite and verticillate leaves are naturally produced : but in such cases the alternation is still seen in the arrangement of the different *clusters* of leaves.

175. In some cases, the effect of interruption of growth, in causing alternate leaves to become opposite and verticillate, can be distinctly shown, as for instance in *Rhododendron penticum*. In other cases, parts which are usually opposite or verticillate, become alternate by the vigorous development of the axis : and on different parts of the same stem, as in *Lysimachia vulgaris*, there may be seen alternate, opposite, and verticillate leaves. When the interruption to development takes place at the end of a branch, the leaves become *fasciculate* (*fasciculus*, a bundle) or clustered, as in the Larch. A remarkable instance of the shortening of internodes, and the clustering of leaves, occurred in the Palm house of the Botanic Garden of Edinburgh, in the case of a Bamboo which was exposed for many months to a low temperature, during the time that the roof of the house was being renewed. The plant had been growing rapidly, with its internodes of the usual length, but it was suddenly arrested near the summit, the internodes became gradually shortened, till the nodes were close to each other, and the leaves came off in bunches. All modifications of leaves follow the same laws of arrangement as true leaves—a fact which is of importance in a morphological point of view.

176. In Exogenous plants, the first leaves produced, or the cotyledons, are opposite. This arrangement often continues during the life of the plant, but at other times it changes. Some tribes of plants are distinguished by their opposite or verticillate, others by their alternate, leaves. Labiate plants have decussate leaves, while Boraginaceæ have alternate leaves, and Tiliaceæ have distichous leaves in general ; Cinchonaceæ have opposite leaves ; Galiaceæ, verticillate. Such arrangements as $\frac{2}{2}$, $\frac{3}{3}$, $\frac{5}{15}$, and $\frac{8}{17}$, are common in Exogens. The first of these, called *quincunx* (*quincunx*, an arrangement of five), is met with in the Apple, Pear, and Cherry (fig. 198) ; the second, in the Bay, Holly, *Plantago media* ; the third, in the cones of *Pinus alba* (fig. 201) ; and the fourth, in those of the *Pinus Picea*. In Endogenous plants, there is only one seed-leaf or cotyledon produced, and hence the arrangement is at first alternate ; and it generally continues so more or less. Such arrangements as $\frac{1}{2}$ $\frac{1}{3}$ (fig. 199), and $\frac{2}{3}$, are common in Endogens, as in Grasses, Sedges, and Lilies. In Acrogens, the leaves assume all kinds of arrangement, being opposite, alternate, and verticillate. It has been found in general that, while the number 5 occurs in the phyllotaxis of Exogens, 3 is common in that of Endogens.

177. Although there is thus, in the great divisions of the vegetable kingdom, a tendency to certain definite numerical arrangements, yet there are many exceptions. In speaking of Palms, which are endogenous plants, Martius states that the leaves of different species exhibit the following spirals— $\frac{1}{2}, \frac{3}{5}, \frac{5}{8}, \frac{7}{9}, \frac{8}{13}, \frac{13}{21}, \frac{21}{34}$. In the species of the genus *Pinus*, $\frac{2}{3}, \frac{5}{13}, \frac{13}{34}, \frac{8}{21}, \frac{21}{55}$, occur. Thus, while it has been shown that the phylloplastic (*φύλλον*, a leaf, and *πλαστικός*, formative), or leaf-formative power, moves in a spiral round the axis, it has been found impossible to apply phyllotaxis satisfactorily to the purposes of classification.

178. The spiral arrangement of the leaves allows all of them to be equally exposed to air and light, and thus enables them to carry on their functions with vigour. The form of the stem is also probably connected with the leaf arrangement. M. Cagnat has remarked that an analogy in arrangement of leaves and character of stem may be traced. The leaves of juniper are in verticils of three, and the pith is triangular; the leaves of cypress being opposite, the pith presents the form of a cross. When leaves are opposite and decussate, the stems are often square, as in Labiate plants. The ordinary rounded stem appears to be associated with a certain degree of alternation in the separate leaves, or in the different pairs of leaves when they are opposite.

179. The study of the structure, forms, and arrangement of leaves, is of great importance, when it is considered that all parts of plants are to be looked upon as leaf-formations variously modified, in order to serve special purposes in the economy of vegetation. The morphological relations of leaves, or the varied forms which they assume, will be illustrated during the consideration of the organs of reproduction, and of the doctrine of metamorphosis, as propounded by Goethe and others. It is only by looking upon all the organs of plants in their relation to the leaf as a type, that a philosophical view can be given of the great plan on which they have been formed.

Leaf-buds.

180. **Leaf-buds** contain the rudiments of branches, and are found in the *axil* of previously-formed leaves (fig. 202 *ba, ba, ba*); or, in other words, in the angle formed between the stem and leaf. They are hence called *axillary*, and may be either *terminal*, *bt*, or lateral, *ba*. In their commencement, they are cellular prolongations from the medullary rays bursting through the bark. The central cellular portion is surrounded by spiral vessels, and is covered with rudimentary leaves. In the progress of growth, vascular bundles are formed continuous with those of the stem; and, ultimately, branches are produced which in every respect resemble the axis whence the buds first sprung. The cellular portion in the centre remains as pith

with its medullary sheath, which is closed and not continuous with that of the parent stem. Thus, in the stem and branch, this sheath forms a canal which is closed at both extremities, and which sends prolongations of spiral vessels to the leaves. As the axis or central portion of the leaf-bud increases, cellular projections appear at regular intervals, which are the rudimentary leaves.




181. A leaf-bud may be removed in a young state from one plant and grafted upon another, by the process of *budding*, so as to continue to form its different parts; and it may even be made to grow in the soil, in some instances, immediately after removal. In certain cases, leaf-buds are naturally detached during the life of the parent, so as to form independent plants, and thus propagate the individual. Leaf-buds have on this account been called *fixed embryos*, by Petit-Thouars and others. They are embryo plants fixed to the axis, capable of sending stems and leaves in an upward direction, and woody fibres downwards, which, according to some, may be considered as roots. A tree may be said to consist of a series of leaf-buds, or *phytons* (*φυτόν*, a plant), attached to a common axis or trunk. In ordinary trees, in which there is provision made for the formation of numerous lateral leaf-buds, any injury done to a few branches is easily repaired; but in Palms, which only form central leaf-buds, and have no provision for a lateral formation of them, an injury inflicted on the bud in the axis is more likely to have a prejudicial effect on the future life of the plant.

182. In the trees of temperate and cold climates, the buds which are developed during one season lie dormant during the winter, ready to burst out under the genial warmth of spring. They are generally protected by external modified leaves in the form of *scales*, *tegmenta* or *perulæ* (*tegmenta*, coverings, *perulæ*, small bags), which are of a firmer and coarser texture than the leaves themselves. These scales or protective appendages of the bud, consist either of the altered laminæ, or of the enlarged petiolar sheath, or of stipules, as in the Fig and Magnolia, or of one or two of these parts combined. They serve a temporary purpose, and usually fall off sooner or later after the leaves are expanded. The bud is often protected by a coating of resinous matter, as in the Horse-chestnut and Balsam poplar, or by a thick downy covering, as in the Willow. Linnæus called leaf-buds *hibernacula*, or the *winter quarters* of the young branch.

183. In the bud of a common tree, as the Sycamore (fig. 203),

* Fig. 202.—Upper portion of a branch of *Lonicera nigra* in a state of hibernation, that is to say, after the fall of the leaves; covered with leaf-buds. *bt*, A terminal bud. *ba*, *ba*, *ba*, Axillary lateral buds. Below the buds, the cicatrix or scar left by the fallen leaves is seen.

there is seen the cicatrix left by the leaf of the previous year, *c*, with the pulvinus or swelling, *p*, then the scales, *ee*, arranged alternately in a spiral manner, and overlying each other in what is called an *imbricated* (*imbrex*, a roof tile) manner. On making a transverse section of the bud (fig. 204), the overlying scales, *eeee*, are distinctly seen surrounding the leaves, *f*, which are plaited or folded round the axis or growing point. In plants of warm climates, the buds are often formed by the ordinary leaves without any protecting appendages; such leaves are called *naked*.



184. **Vernation.**—The arrangement of the leaves in the bud has been denominated *vernation* (*ver*, spring), or *præfoliation* (*præ*, before, and *folium*, leaf), or *gemination* (*gemma*, a bud). This differs in different plants, but in each species it follows a regular law. The leaves in the bud are either placed simply in apposition, as in the Mistletoe, or they are folded or rolled up longitudinally or laterally, giving rise to different kinds of vernation, as delineated in fig. 205 *a*—*n*, where the *dot* represents the axis, and the folded or curved lines, the leaves with the thickened part indicating the midrib; figs. *a* and *g*, being vertical sections; *h*—*n*, horizontal.

185. The leaf taken individually, is either folded longitudinally from apex to base (fig. 205 *a*), as in the Tulip-tree, and called *reclinate*; or rolled up in a circular manner from apex to base, as in Ferns (fig. 205 *g*), and called *circinate* (*circino*, I turn round); or folded laterally, *conduplicate*, as in Oak (fig. 205 *b*); or it has several folds like a fan, *plicate* or *plaited*, as in Vine and Sycamore (figs. 204 *f*, 205 *c*), and in leaves with radiating vernation, where the ribs mark the foldings; or it is rolled upon itself, *convolute* or *supervolute*, as in Banana and Apricot (fig. 205 *d*); or its edges are rolled inwards, *involute*, as in Violet (fig. 205 *e*); or outwards, *revolute*, as in Rosemary (fig. 205 *f*). The different divisions of a cut leaf may be folded or rolled up separately, as in Ferns, while the entire leaf may have either the same or a different kind of vernation.

Fig. 203.—Leaf-bud of *Acer pseudo-platanus* covered with scales. *r*, The branch. *p*, Pulvinus or swelling at the base of the leaf which has fallen, leaving a scar or cicatrix, *c*, in which the remains of three vascular bundles are seen. *ee*, Imbricated scales of the bud.

Fig. 204.—Transverse section of the same leaf-bud. *eeee*, The scales arranged in an imbricated manner, like the tiles on a house. *f*, The leaves folded in a plaited manner, exhibiting plicate vernation.

186. Other kinds of vernation receive their names from the arrangement of the leaves in the bud, taken as a whole. Leaves in the



205

bud are opposite, alternate, or verticillate; and thus different kinds of vernation are produced. Sometimes they are nearly in a circle at the same level, remaining flat, or only slightly convex externally, and placed so as to touch each other by their edges; thus giving rise to *valvate* vernation (fig. 205 *h*). At other times they are at different levels, and are applied over each other, so as to be *imbricated*, as in Lilac, and in the outer scales of Sycamore (figs. 203, 204); and occasionally the margin of one leaf overlaps that of another, while it, in its turn, is overlapped by a third, so as to be *twisted* or *spiral* (fig. 205 *i*). When the leaves are more completely folded, they either touch at their extremities (fig. 205 *k*), or are folded inwards by their margin, and become *induplicate* (fig. 205 *l*); or a conduplicate leaf covers another similarly folded, while it covers a third, and thus the vernation is *equitant* (riding), as in Privet (fig. 205 *m*); or conduplicate leaves are placed, so that the half of the one covers the half of another, and thus they become *half-equitant* or *obvolute*, as in Sage (fig. 205 *n*). The scales of a bud sometimes exhibit one kind of vernation, and the leaves another (fig. 204). The same modes of arrangement occur in the flower-buds, as will be afterwards shown.

187. Leaf-buds, as has been stated, are either terminal or lateral. By the production of the former (fig. 202 *b t*), stems increase in length, while the latter (fig. 202 *b a*, *b a*, *b a*) give rise to branches, and add to the diameter of the stem. The terminal leaf-bud, after producing leaves, sometimes dies at the end of one season, and the

Fig. 205.—Figures to show the different kinds of vernation. *a*—*g*, The folding of individual leaves; *a* and *g* being vertical sections, *b c d e* and *f*, being horizontal. *a*, Reclinate. *b*, Conduplicate. *c*, Plicate. *d*, Convolute. *e*, Involute. *f*, Revolute. *g*, Circinate. *h*—*n*, Folding of leaves when united together in the leaf-bud. The sections are horizontal or transverse, and show the relative position of the leaves, and the mode in which each of them is folded. *h*, Valvate. *i*, Twisted or spiral. *k l*, Induplicate. *m*, Equitant. *n*, Obvolute or half-equitant. In all the figures, the thickened portion indicates the midrib of the leaf, and the dot marks the position of the axis.

whole plant, as in annuals, perishes; or part of the axis is persistent, and remains for two or more years, each of the leaves before its decay producing a leaf-bud in its axil. This leaf-bud continues the growth in spring.

188. In some trees of warm climates, as *Cycas*, Papaw-tree, Palms, and Tree ferns, the production of terminal buds is well seen. In these plants, the elongation of the stem is generally regular and uniform, so that the age of the plant may be estimated by its height. Such stems (often endogenous) may thus be considered as formed by a series of terminal buds, placed one over the other. From this mode of growth they do not attain a great diameter (fig. 115, 1). In other trees, especially *Exogens*, besides the terminal bud there are also lateral ones. These, by their development, give rise to *branches* (*rami*), from which others called *branchlets* or *twigs* (*ramuli*) arise. Such buds being always produced in the axils of leaves, are of course arranged in the same manner as the leaves are. By the continual production of lateral leaf-buds, the stem of exogenous plants acquires a great diameter.

189. Although provision is thus made for the regular formation of leaf-buds, there are often great irregularities in consequence of many being abortive, or remaining in a dormant state. Such buds are called *latent*, and are capable of being developed in cases where the terminal bud, or any of the branches, have been injured or destroyed. In some instances, as in *Firs*, the latent buds follow a regular system of alternation; and in plants with opposite leaves, it frequently happens that the bud in the axil of one of the leaves only is developed, and the different buds so produced are situated alternately on opposite sides of the stem.

190. When the terminal leaf is injured or arrested in its growth, the elongation of the main axis stops, and the lateral branches often acquire increased activity. By continually cutting off the terminal buds, a woody plant is made to assume a bushy appearance, and thus *pollard* trees are produced. Pruning has the effect of checking the growth of terminal buds, and of causing lateral ones to push forth. The peculiar bird-nest appearance often presented by the branches of the common *Birch*, depends on an arrestment in the terminal buds, a shortening of the internodes, and a consequent clustering or fasciculation of the twigs. In some plants there is a natural arrestment of the main axis after a certain time, giving rise to peculiar shortened stems. Thus the crown of the root (§ 70) is a stem of this nature, forming buds and roots. Such is also the case in the stem of *Cyclamen*, *Testudinaria elephantipes*, and in the tuber of the potato. The production of lateral in place of terminal buds, sometimes gives the stem a remarkable zigzag aspect.

191. In many plants with a shortened axis, the lateral buds produce

long branches. Thus the *flagellum* (*flagellum*, a whip or twig), or *runner* of the Strawberry and Ranunculus, is an elongated branch, developing buds as it runs along the ground; the *propagulum* (*propago*, a shoot), or *offset*, is a short thick branch produced laterally in fleshy plants from a shortened axis, and developing a bud at its extremity, which is capable of living when detached, as in Houseleek. Fig. 206 represents a strawberry plant in which a' is the primary axis, ending in a cluster of green leaves, r , and some rudimentary leaves, f , and not elongating; from the axil of one of the leaves proceeds a branch or runner, a'' , with a rudimentary leaf, f , about the middle, and another cluster of leaves, f and r , forming a young plant with roots; from this a third axis comes off, a''' , and so on. In many instances



206

the runner decays, and the young plant assumes an independent existence. Gardeners imitate this in the propagation of plants by the process of *layering*, which consists in bending a twig, fixing the central part of it into the ground, and, after the production of adventitious roots, cutting off its connection with the parent.



192. When the stem creeps along the surface of the ground, as in the Rhizome (fig. 90), or completely under ground, as in the Soboles or creeping stem (fig. 91), the terminal bud continues to elongate year after year, thus making additions to the axis in a horizontal manner. At the same time buds are annually produced on one side which send shoots upwards and roots downwards. Thus, in fig. 91 (soboles of a Rush), r is the extremity of the axis or terminal bud, f the leaves in the form of scales, p the aerial shoots or branches, t being the level of the ground. Again, in fig. 90 (rhizome of Solomon's seal), a is the terminal bud which has been formed subsequently to b , b the bud which has sent up leaves, and which has decayed, c being the scars left by the similar buds of previous seasons.

193. **Aerial and Subterranean Leaf-buds.**—According to the nature of the stems, leaf-buds are either *aerial* or *subterranean*; the former occurring in plants which have the stems above ground, the latter in those in which the stems are covered. In the case of Asparagus and other plants which have a perennial stem below ground, sub-

Fig. 206.—Flagellum, or Runner of the Strawberry. a' , One axis which has produced a cluster of leaves, the upper, r , green, the lower, f , rudimentary. From the axil of one of the latter, a second axis, a'' , arises, bearing about the middle a rudimentary leaf, f , and a cluster of leaves, r , partly green, and partly rudimentary, at its extremity. From the axil of one of the leaves of this cluster, a third axis, a''' , proceeds.

terranean buds are annually produced, which appear above ground as shoots or branches covered with scales at first (fig. 110 *l*), and ultimately with true leaves. The young shoot is called a *Turio* (*turio*, a young branch). These branches are herbaceous and perish annually, while the true stem remains below ground ready to send up fresh shoots next season. In Bananas and Plantains, the apparent aerial stem is a shoot or leaf-bud sent up by an underground stem, and perishes after ripening fruit. In some plants, several branches are sent up at once from the underground stem, in consequence of a rapid development of lateral as well as terminal buds; and in such cases the lateral ones may be separated as distinct plants in the form of *suckers* (*surculi*). The potato is a thickened stem or branch capable of developing leaf-buds, which in their turn form aerial and subterranean branches, the former of which decay annually, while the latter remain as *tubers* to propagate the plant. Thus, in fig. 92, *s s* is the surface of the soil, *p a* is the aerial portion of the potato covered with leaves, *t* is the subterranean stem or tuber covered with small scales or projections, as represented at *τ b*, from the axil of which leaf-buds are produced. This provision for a symmetrical development of axillary leaf-buds at once distinguishes the tuber of the potato from fleshy roots, like those of the *Dahlia*.

194. **Bulb.**—A good example of a subterranean bud occurs in the *Bulb*, as seen in the Hyacinth, Lily, and Onion. This is a subterranean



leaf-bud covered with scales, arising from a shortened axis. From the centre of the bulb a shoot or herbaceous stem is produced which dies

Fig. 207.—Tunicated bulb of *Allium Porrum*, or the Leek. *r*, Roots. *p*, A circular disc, or shortened stem intervening between the roots and the bulbous swelling. *e e*, Scales, or subterranean modified leaves. *f*, Upper leaves which become green.

Fig. 208.—Vertical section of the tunicated bulb of the Leek. The letters indicate the same parts as in the last figure. *b*, Bud situated in the axil of a scale, which, by its development, forms a new bulb.

Fig. 209.—Scaly or naked bulb of *Lillium album*. *r*, Roots. *e e e*, Scales or modified underground leaves. *i*, The stem cut.

down. New bulbs, or *cloves* as they are called, are produced from the subterranean axis. At the base of the scales there is a flattened disc, varying in thickness, which is formed by the base of the buds, and which has sometimes been called the stem. The parts of the bulb are seen in fig. 207, where *p* marks the disc or round flat portion formed by the bases of the lateral buds from which the fasciculated roots, *r*, proceed, *e* the scales or modified leaves, and *f* the true leaves. In the vertical section (fig. 208), *b* is the new bulb formed like a bud in the axil of a scale. The new bulb sometimes remains attached to the parent bulb, and sends up an axis and leaves; at other times it is detached in the course of growth, and forms an independent plant. The new bulbs feed on the parent one, and ultimately cause its absorption. The scales are sometimes all fleshy, as in the *scaly* or *naked bulb* of the white lily (fig 209 *e e e*), or the outer ones are thin and membranous, overlapping the internal fleshy ones, and forming a *tunicated bulb*, as in the Onion, Squill, and Leek (fig. 207).

195. The **Corm** (κορμός, a stump) has already been noticed under the head of subterranean stems (¶ 70, fig. 93). It may be considered as a bulb in which the central portion or axis is much enlarged, while the scales are reduced to thin membranes. Some have called it a *solid bulb*. A Corm may be generally distinguished from a Bulb, by a transverse section of the latter presenting a series of circles, equal in number to the fleshy scales arranged around its central axis. It is seen in the Tulip, Colchicum, Crocus, and Gladiolus. It produces either terminal buds, as in Gladiolus and Crocus, in which several annual additions to the corm remain attached together, and the newly produced corms come gradually nearer and nearer to the surface of the soil; or lateral buds, as in Colchicum, represented at fig. 93, where *r* indicates the roots, *f* the leaf, *a'* the stem or axis of the preceding year withered, *a''* the secondary axis, or the stem developed during the year, and taking the place of the old one, and which, in its turn, will give origin to a new axis, *a'''*, on the opposite side, according to the law of alternation. The new axes or corms being thus produced alternately at either side, there is very little change in the actual position of the plant from year to year. Bulbs and corms contain a store of starch and of other substances, for the nourishment of the young plants.

196. **Anomalies and Transformations of Leaf-buds.**—Leaf-buds arise from the medullary system of the plant, and in some instances they are found among the cellular tissue, without being in the axil of leaves. In this case they are *extra-axillary*, and have been called *adventitious* or *abnormal*. Such buds are produced after the stem and leaves have been formed, and in particular circumstances they are developed like normal buds. What have been called *embryo-buds*, are woody nodules seen in the bark of the Beech, Elm, and other trees. They are looked

upon as partially developed abnormal buds, in which the woody matter is pressed upon by the surrounding tissue, and thus acquires a very hard and firm texture. When a section is made, they present woody circles arranged around a central pith, and traversed by medullary rays (fig. 210). The nodules sometimes form *knots* on the surface of the stem, at other times they appear as large *excrescences*, and in some cases twigs and leaves are produced by them. Some consider embryo-buds as formed by layers of woody matter, which originate in the sap conveyed downward by the bark and cambium cells, and are deposited round a nucleus or central mass.



210

197. Leaf-buds sometimes become extra-axillary (fig. 211 *b*), in consequence of the non-appearance or abortion of one or more leaves, or on account of the adhesion of the young branch to the parent stem. In place of one leaf-bud, there are occasionally several accessory ones produced in the axil, giving origin to numerous branches (fig 212 *b*).



211



212


Such an occurrence is traced to the presence of latent or adventitious buds. Fig. 211 represents a branch *r*, of walnut, *p* the cut petiole, and *b* two buds, of which the upper is most developed, while fig. 212 exhibits a branch of *Lonicera tartarica*, with numerous buds, *b*, in the axil of the leaves, the lowest of which are most advanced. By the union of several such leaf-buds, branches are produced having a thickened or flattened appearance, as is seen in the Fir, Ash, and other trees. These *fasciated* (*fascia*, a band) branches, in some cases, however, are owing to the abnormal development of a single bud.

Fig. 210.—Vertical section of a nodule, *n*, or embryo-bud embedded in the bark of the Cedar. It forms a projection on the surface. The woody layers form zones round a kind of pith.

Fig. 211.—Portion of a branch, *r*, of the walnut, bearing the petiole, *p*, of a leaf which has been cut. In the axil of the leaf, several buds, *b*, are produced, the highest of which are most developed.

Fig. 212.—Portion of a branch, *r*, of *Lonicera tartarica*, bearing two opposite leaves, one of which has been cut, the other, *f*, being preserved. In the axil of the leaves, clusters of buds, *b*, are seen, the lowest of which are most developed.


198. In the axil of the leaves of *Lilium bulbiferum*, *Dentaria bulbifera*, and some other plants, small conical or rounded bodies are produced called *bulbils* or *bulblets* (fig. 213 *b b b*). They resemble bulbs in their aspect, and consist of a small number of thickened scales enclosing a growing point. These scales are frequently united closely together so as to form a solid mass. Bulbils are therefore transformed leaf-buds, which are easily detached, and are capable of producing young plants when placed in favourable circumstances.



199. Occasionally leaf-buds are produced naturally on the edges of leaves, as in *Bryophyllum calycinum* and *Malaxis paludosa* (fig. 214), and on the surface of leaves, as in *Ornithogalum thyrsoides* (fig. 215). These are capable of forming independent plants. Similar



214



215

buds are also made to appear on the leaves of *Gesnera*, *Gloxinia*, and *Achimenes*, by wounding various parts of them, and placing them in moist soil; this is the method often pursued by gardeners in their propagation. The cellular tissue near the surface of plants, seems therefore to have the power of developing abnormal leaf-buds in certain circumstances. Even roots, when long exposed to the air, may thus assume the functions of stems. Leaves bearing buds on their margin, are called *proliferous* (*proles*, offspring, and *fero*, I bear).

200. **Spines or Thorns.**—Branches are sometimes arrested in their development, and, in place of forming leaves, become transformed into *spines* and *tendrils*. Spines or thorns are undeveloped branches, ending in more or less pointed extremities, as in the Hawthorn. Plants which have spines in a wild state, as the Apple and Pear, often lose them when cultivated, in consequence of their being changed into branches; in some cases, as in *Prunus spinosa*, or the Sloe (fig. 217), a branch

Fig. 213.—Portion of the stem of *Lilium bulbiferum* with three alternate leaves, *f f f*, and three bulbils or bulblets, *b b b*, in their axils.


Fig. 214.—Extremity of a leaf, *l*, of *Malaxis paludosa*, the margin of which is covered with adventitious buds, *b b*; thus becoming proliferous.

Fig. 215.—Portion of the blade of a leaf, *f*, of *Ornithogalum thyrsoides*, on the surface of which are developed adventitious or abnormal buds, *b b b*, some of which are large.

bears leaves at its lower portions, and terminates in a spine. Leaves themselves often become spiny by the hardening of their midrib or primary veins, and the diminution or absence of parenchyma, as in *Astragalus massiliensis* (fig. 217 *r*), where the midrib becomes spiny after the fall of some of the leaflets; in the Holly, where all the veins are so; and in the Barberry (fig. 218), where some of the leaves, *fff*, are produced in the form of spiny branches, with scarcely any parenchyma. In place of producing a lamina or blade at its extremity, the petiole sometimes terminates in a spine. Stipules are occasionally transformed into spines, as in *Robinia pseudo-acacia* (fig. 219, *s s*), and such is also the case with the swelling or pulvinus at the base of the



216



217



218



219



220

leaf, as in *Ribes Uva-crispa* (fig. 220, *c c c*). Branches are sometimes

Fig. 216.—Branch of *Prunus spinosa*, or Sloe, with alternate leaves, and ending in a spine or thorn.

Fig. 217.—Pinnate leaf of *Astragalus massiliensis*, the midrib of which, *r*, ends in a spine. *s*, Petiolar stipules. *f*, Nine pairs of leaflets.

Fig. 218.—Branch of *Berberis vulgaris*, or Barberry, the leaves of which, *fff*, are transformed into branching spines. In the axil of each, a cluster, *r r r*, of regularly formed leaves is developed.

Fig. 219.—Base of the pinnate leaf of *Robinia pseudo-acacia*, the stipules of which, *s s*, are converted into spines or thorns. *b*, Branch. *r*, Petiole.

Fig. 220.—Branch of *Ribes Uva-crispa*, in which the pulvinus or swelling, *c c c*, at the base of each of the leaves, *fff*, is changed into a spine, which is either simple, or double, or triple. *b b*, Leaf-buds arising from the axil of the leaves.

arrested in their progress at an early state of their development, and do not appear beyond the surface of the stem; at other times, after having grown to a considerable size, they undergo decay. In both instances, the lower part of the branch becomes embedded and hardened among the woody layers of the stem, and forms a *knot*.

201. Tendrils.—A leaf-bud is sometimes developed as a slender spiral or twisted branch, called a *tendril* or *cirrhus* (*cirrus*, a curl), as in the Passion-flower, in which the lateral buds are thus altered with the view of enabling the plant to climb. When tendrils occupy the place of leaves, and appear as a continuation of the leaf-stalk, they are called *petiolar*y, as in *Lathyrus Aphaca*, in which the stipules perform the function of true leaves. In *Flagellaria indica*, *Methonica superba*, *Anthericum cirrhatum*, and *Albuca cirrhata*, the midrib of the leaf ends in a tendril; and in Vetches, the terminal leaflet, and some of the lateral ones at the extremity of their pinnate leaves, are changed, so as to form a branching tendril. In the Vine, the tendrils are looked upon as the terminations of separate axes, or as transformed terminal buds. In this plant there are no young buds seen in the angle between the stem and leaves, nor between the stem and tendrils; and the latter are not axillary. Fig. 221 represents the branch of a Vine, in which a' is the



221

primary or first formed axis, ending in v' , a tendril or altered terminal bud, and having a leaf, f' , on one side. Between this leaf and the tendril which represents the axis, a leaf-bud was formed at an early date producing the secondary axis, or branch, a'' , ending in a tendril, v'' , with a lateral leaf, f'' , from which a tertiary axis or branch, a''' , was developed, ending in a tendril v''' , and so on.

202. Tendrils twist in a spiral manner, and enable the plants to rise into the air by twining round other plants. The direction of the spiral frequently differs from

Fig. 221.—Portion of a branch of the Vine (*Vitis vinifera*). a' , First axis, terminated by a tendril or cirrhus, v' , which assumes a lateral position, and bears a leaf, f' . From the axil of this leaf, a second axis, a'' , comes off, which seems to be a continuation of the first, and is terminated also by a tendril, v'' , bearing a leaf, f'' . From the axil of this second leaf, a third axis, a''' , arises, terminated by a tendril, v''' , and bearing a leaf, f''' , from the axil of which a fourth axis, a'''' , arises.

that of the climbing stem, producing the tendril. In the Vine, the lower part of the stem is strong, and needs no additional support; the tendrils therefore occur only in the upper part where the branches are soft, and require aid to enable them to support the clusters of fruit. In *Vanilla aromatica*, the vanilla plant, tendrils are produced opposite the leaves, until the plant gains the top of the trees by which it is supported; the upper tendrils being then developed as leaves. The midrib is sometimes prolonged in a cup-like form: this is occasionally seen in the common cabbage, and seems to depend on the vascular bundles of the midrib spreading out at their extremity in a radiating manner, and becoming covered with parenchyma in such a way as to form a hollow cavity in the centre.

Special Functions of Leaves.

203. Leaves expose the fluids of the plant to the influence of air and light, and their spiral arrangement enables them to do so effectually. They are concerned in the elaboration of the various vegetable secretions, in the formation of wood, and in the absorption of fluid and gaseous matters. A plant, by being constantly deprived of its leaves, will ultimately be destroyed. On this principle, weeds, with creeping stems and vigorous roots, which are with difficulty eradicated, may be killed. In the cells of the leaves changes take place under the agency of light, by which oxygen is given off and carbon fixed. These will be considered under the head of vegetable *respiration*. The absorption of carbonic acid and of fluids is carried on by the leaves, chiefly through their stomata, according to Bonnet, who found that absorption was rapidly effected by the under surface of ordinary leaves in which the cuticle is thinnest, cellular tissue least condensed, and stomata most abundant; the upper surface of the leaf usually presenting a polished and dense epidermis, and having few stomata, taking little part in such a process. Hoffman* has carried these researches farther, and ascertained that leaves absorb fluids in large quantities; that during a fall of rain the vegetable fluids undergo from such a cause a process of dilution, leading to an immediate and more rapid descent of sap, which under such circumstances is capable of general diffusion throughout the several vegetable tissues. Some physiologists have expressed doubts as to absorption being carried on by the leaves in ordinary circumstances. Leaves, it has been shown, absorb gaseous matters. Saussure states that oxygen is absorbed by the leaves during night, the quantity varying according to the nature of the plant. Boussingault† has found that the leaves of the Vine detract carbonic acid from the air. Other experiments prove that

* Hoffman on Circulation of Sap, Scientific Memoir, Nat. Hist. i. 46.

† Boussingault, *Economie Rurale*, i. 66.

ammonia and nitrogen are similarly acted on. Leaves also give off gases and fluids by a process of exhalation or transpiration. Carbonic acid, to a moderate extent, is exhaled during darkness, and a large quantity of fluid is given off by transpiration. The number and size of the stomata regulate the transpiration of fluids, and it is modified by the nature of the epidermis. The absorbing power of leaves depending on similar causes, is capable of being increased by any process which removes either natural or imposed obstructions to the free action of their surface. It is thus that rain, while supplying the material for absorption, at the same time renders the leaf more capable of such action. In plants with a thick and hard epidermal covering, exhalation is less vigorous than in those where it is thin and soft. Some succulent plants of warm climates have a very thick covering. The peculiar character of the leaves or phyllodia of Australian plants, is probably connected with the dry nature of the climate. While heat acts in promoting evaporation, the process of transpiration is more under the influence of light. It assists the process of endosmose, by rendering the fluid in the cells thicker, and thus promotes the circulation of sap.

204. The quantity of fluid exhaled varies in amount in different plants. A Sunflower three feet high, gave off twenty ounces of watery fluid daily. Hales found that a Cabbage, with a surface of 2,736 square inches, transpired at an average nineteen ounces; a Vine, of 1,820 square inches, from five to six ounces. Experiments on exhalation may be made by taking a fresh leaf with a long petiole, putting it through a hole in a card which it exactly fits, and applying the card firmly and closely to a glass tumbler, about two-thirds full of water, so that the petiole is inserted into the water, then inverting an empty tumbler over the leaf, and exposing the whole to the sun, the fluid exhaled will be seen on the inside of the upper tumbler. The experiment may be varied by putting the apparatus in darkness, when no exhalation takes place, or in diffuse daylight, when it is less than in the sun's rays. This process of exhalation imparts moisture to the atmosphere, and hence the difference between the air of a wooded country and that of a country deprived of forests. The cells in the lower side of a leaf where stomata exist, are chiefly concerned in the aeration of the sap, whilst other assimilative processes go on in the upper cells.

205. Leaves, after performing their functions for a certain time, wither and die. In doing so, they frequently change colour, and hence arise the beautiful and varied tints of the autumnal foliage. This change of colour is chiefly occasioned by the diminished circulation in the leaves, and the higher degree of oxidation to which their chlorophyll has been submitted. Leaves which are articulated with the stem, as in the Walnut and Horse-chestnut, fall and leave a scar,

while those which are continuous with it remain attached for some time after they have lost their vitality, as in the Beech. Most of the trees of this country have deciduous leaves, their duration not extending over more than a few months; while in trees of warm climates, the leaves often remain for two or more years. In tropical countries, however, many trees lose their leaves in the dry season. This is seen in the forests of Brazil, called *Catingas*. The period of defoliation varies in different countries according to the nature of their climate. Trees which are called evergreen, as Pines and Evergreen oak, are always deprived of a certain number of leaves at intervals, sufficient being left, however, to preserve their green appearance. Various causes have been assigned for the fall of the leaf. In cold climates, the deficiency of light and heat in winter causes a cessation in the functions of the cells of the leaf; its fluid disappears by evaporation; its cells and vessels become contracted and diminished in their calibre; various inorganic matters accumulate in the texture; the whole leaf becomes dry; its parts lose their adherence; and it either falls by its own weight or is detached by the wind. In warm climates, the dry season gives rise to similar phenomena. In addition to these appreciable causes, we must include a process of preordained vital action, by which the cycle of vegetable life is preserved whether the leaves be deciduous, annual, or persistent.

SECTION II.—GENERAL VIEW OF THE FUNCTIONS OF THE NUTRITIVE ORGANS.

206. In order that plants may be nourished, food is required. This food, in a crude state, enters the roots by a process of *absorption* or *imbibition*; it is then transmitted from one part of the plant to another, by means of the *circulation* or *progressive movement of the sap*; it reaches the leaves, and is there submitted to the action of light and air, which constitutes the function of *respiration*; and thus the fluids are finally fitted for the process of *assimilation*, and form various vegetable *products* and *secretions*.

1.—FOOD OF PLANTS AND SOURCES WHENCE THEY DERIVE THEIR NOURISHMENT. CHEMICAL COMPOSITION OF PLANTS.

207. The nutriment of plants can only be ascertained when their chemical composition has been determined. The physiologist and chemist must unite in this inquiry, in order to arrive at satisfactory conclusions. Much has been done of late by Liebig, Mülder, Dumas, Boussingault, and other chemists, to aid the botanist in his investigations, and to place physiological science on a sound and firm basis.

It is true that many processes take place in plants which cannot as yet be explained by the chemist, and to these the name of *vital* has been applied. This term, however, must be considered as implying nothing more than that the function so called occurs in living bodies, and in the present state of our knowledge is not reducible to ordinary chemical or physical laws. A greater advance in science may clear up many difficulties in regard to some of the vital functions, while others may ever remain obscure.

208. Plants are composed of certain *chemical elements*, which are necessary for their growth. These are combined in various ways, so as to form what have been called *organic* and *inorganic* compounds. The former are composed of carbon, oxygen, hydrogen, and nitrogen, or azote, with a certain proportion of sulphur and phosphorus; while the latter consist of various metallic bases, combined with oxygen, metalloids, and acids. In all plants there is a greater or less proportion of water, the quantity of which is ascertained by drying at a temperature a little above that of boiling water. By burning the dried plant the organic constituents disappear, and the inorganic part or the ash is left. The relative proportion of these constituents varies in different species, as seen in the following table by Solly, in which the proportions are given in 10,000 parts of the fresh plants:—

	Water.	Organic Matter.	Inorganic.
Potato,	7713	2173	114
Turnip,	9308	588	104
Sea Kale,	9238	705	57
French Beans,	9317	619	64
Red Beet,	8501	1390	109
Asparagus,	9210	735	55
Water Cress,	9260	633	107
Sorrel,	9207	702	91
Parsley,	8430	1299	271
Fennel,	8761	1048	191
Salsafy,	7951	1929	120
Mustard,	9462	436	102

209. The analysis of 100 parts of Fruits gives the following results:—

	Water.	Organic.	Inorganic.
Strawberry,	90·22	9·37	0·41
Green Gage, whole fruit,	83·77	15·83	0·40
Cherry, do.	82·48	17·09	0·43
Pear, do.	83·55	16·04	0·41
Apple, do.	84·01	15·72	0·27
Gooseberry,	90·26	9·35	0·39

210. The following table, by Johnston, represents the constitution in 1000 parts of plants and seeds, taken in the state in which they are

given to cattle, or laid up for preservation, and dried at 230° Fahrenheit; the organic matter being indicated by the carbon, oxygen, hydrogen, and nitrogen; the inorganic by the ash:—

	Wheat.	Oats.	Peas.	Hay.	Turnips.	Potatoes.
Carbon	455	507	465	458	429	441
Hydrogen.....	57	64	61	50	56	58
Oxygen.....	430	367	401	387	422	439
Nitrogen.....	35	22	42	15	17	12
Ash	23	40	31	90	76	50

By the process of drying, the 1000 parts of these substances lost water in the following proportions:—

Wheat, 166	Peas, 86	Turnips, 925
Oats, 151	Hay, 158	Potatoes, 722

211. As plants have no power of locomotion, it follows that their food must be universally distributed. The atmosphere and the soil accordingly contain all the materials requisite for their nutrition. These materials must be supplied either in a gaseous or a fluid form, and hence the necessity for the various changes which are constantly going on in the soil, and which are aided by the efforts of man. Plants are capable of deriving all their nourishment from the mineral kingdom. The first created plants in all probability did so, but in the present day the decaying remains of other plants and of animals are also concerned in the support of vegetation.

Organic Constituents and their Sources.

212. **Carbon** (C) is the most abundant element in plants. It forms from 40 to 50 per cent. of all the plants usually cultivated for food. When plants are charred the carbon is left, and as it enters into all the tissues, although the weight of the plants is diminished by the process, their form still remains. When converted into coal (a form of carbon), plants are frequently so much altered by pressure as to lose their structure, but occasionally it can be detected under the microscope. Carbon is insoluble, and therefore cannot be absorbed in its uncombined state. When united to oxygen, however, in the form of carbonic acid, it is readily taken up either in its gaseous state by the leaves, or in combination with water by the roots. The soil contains carbon (*humus*), and in some soils, as those of a peaty nature, it exists in very large quantity. The carbon in the soil is converted into carbonic acid in order to be made available for the purpose of plant-growth. Carbon has the power of absorbing gases, and in this way, by enabling certain combinations to go on, it assists in the nourishment of plants. In the atmosphere, carbonic acid is always present, averaging about $\frac{1}{1000}$ part, arising from the respiration of man and animals,

combustion, and other processes. A certain atmospheric equilibrium is thus maintained, consequent on the difference in vegetable and animal respiration, the former giving out oxygen, which the latter consumes.

213. **Oxygen (O)** is another element of plants. Air contains about 21 per cent. of it. Every 9 lbs. of water contain 8 of oxygen, and it is combined with various elements, so as to form a great part of the solid rocks of the globe, as well as of the bodies of animals and man.

214. It is chiefly in its state of combination with **Hydrogen (H)**, so as to form water (HO), that oxygen is taken up by plants. Hydrogen is not found in a free state in nature, and with the exception of coal, it does not enter into the composition of the mineral masses of the globe. It forms $\frac{1}{8}$ of the weight of water, and it is present in the atmosphere in combination with nitrogen. Hydrogen is also furnished by sulphuretted hydrogen, and some compounds of carbon. Carbonic acid and water existing to a large extent in the atmosphere, vegetables can thence also derive carbon, oxygen, and hydrogen, which enter so extensively into their constitution.

215. **Nitrogen (N)** is another element found in plants. It forms 79 per cent. of the atmosphere, and abounds in animal tissues. It is therefore requisite for the purposes of animal life that nitrogen be furnished in food. Those vegetables containing the greatest quantity of nitrogenous matter are consequently the most nutritive. The latter, during their decay, give off nitrogen, combined with hydrogen, in the form of ammonia (NH^3), which is absorbed in large quantities by carbon, is very soluble in water, and seems to be the chief source whence plants derive nitrogen. In tropical countries where thunderstorms are frequent, the nitrogen and oxygen of the air are sometimes made to combine, so as to produce nitric acid, (NO^5) which, either in this state, or in combination with alkaline matters, furnishes a supply of nitrogen. Daubeny thinks that the ammonia and carbonic acid in the atmosphere are derived in part from volcanic actions going on in the interior of the globe. The continued fertility of the Terra del Lavoro, and other parts of Italy, is attributed by him to the disengagement of ammoniacal salts and carbonic acid by volcanic processes going on underneath; and to the same source he traces the abundance of gluten in the crops, as evidenced by the excellence of Italian macaroni.

216. Mülder maintains that the ammonia is not carried down from the atmosphere, but is produced in the soil by the combination between the nitrogen of the air, and the hydrogen of decomposing matters. The same thing takes place, as in natural saltpetre caverns of Ceylon, with this exception, that, by the subsequent action of oxygen, ulmic, humic, geic, apocrenic, and crenic acids, are formed in place of nitric acid. These acids consist of carbon, oxygen, and hydrogen, in

different proportions, and they form soluble salts with ammonia. By all porous substances like the soil, ammonia is produced, provided they are moist, and filled with atmospheric air, and are exposed to a certain temperature. It is thus, he states, that moist charcoal and humus become impregnated with ammonia.

217. These four elementary bodies then are supplied to plants, chiefly in the form of carbonic acid (CO^2), water (HO), and ammonia (NH^3). In these states of combination they exist in the atmosphere, and hence some plants can live suspended in the air, without any attachment to the soil. When a volcanic or a coral island appears above the waters of the ocean, the lichens which are developed on it are nourished in a great measure by the atmosphere, although they subsequently derive inorganic matter from the rocks, to which they are attached. Air plants, as Bromelias, Tillandsias, and Orchidaceæ, and many species of Ficus, can grow for a long time in the air. In the Botanic Garden of Edinburgh, a specimen of *Ficus australis* has lived in this condition for upwards of twenty years, receiving no supply of nourishment except that afforded by the atmosphere and common rain water, containing, of course, a certain quantity of inorganic matter. The following analysis was made of the leaves of this plant, in 1847, by my pupil, Mr. John Macadam:—

	Organic In 100 parts.	Inorganic In 100 parts.
Petiole of former year's growth, including midrib	82.98 ...	17.02
Three leaves of former year's growth.....	86.24 ...	13.76
Petiole of present year's growth, with midrib.....	92.65 ...	7.35
Seven leaves of present year's growth	92.28 ...	7.72

All were dried at 212° Fahrenheit.

In the experimental Garden of Edinburgh, Mr. James McNab has cultivated various plants, as *Strelitzia augusta*, currants, gooseberries, &c., without any addition of soil, and simply suspended in the air, with a supply of water kept up by the capillary action of a worsted thread. Some of the plants have flowered and ripened fruit. These experiments show that the atmosphere and rain water contain all the ingredients requisite for the life of some plants. Boussingault, from observations made on the cultivation of Trefoil, was led to the conclusion, that under the influence of air and water, in a soil absolutely devoid of organic matter, some plants acquire all the organic elements requisite for growth. Messrs. Wiegman and Polstorf took fine quartz sand, burnt it to destroy any organic matter, digested it for sixteen hours in strong nitro-muriatic acid, and then washed it with distilled water. Various kinds of seeds, as barley, oats, vetch, clover, and tobacco, were then sown in it, and watered with distilled water, and all grew more or less.

218. The elementary bodies already mentioned, in various states of combination, constitute the great bulk of plants. They occur in the

form of binary compounds, as water and oily matters; ternary, as starch, gum, sugar, and cellulose; quaternary, as gluten, albumen, caseine, and fibrine. The latter compounds seem to require for their composition, not merely the elements already noticed, in the form of a basis, called Proteine ($C^{40} H^{51} N^5 O^{12}$ according to Mülder, or $C^{48} H^{36} N^6 O^{14}$ according to Liebig), but certain proportions of sulphur and phosphorus in addition; thus, albumen = 10 Pr. + 1 P + 1 S; fibrine = 10 Pr. + 1 P + 2 S; caseine = 10 Pr. + 2 S. The tissues into the composition of which these proteine compounds enter, are tinged of a deep orange-yellow, by strong nitric acid. These compounds are highly important in an agricultural point of view, and the consideration of them will be resumed when treating of the application of manures.

Inorganic Constituents, and their Sources.

219. The consideration of the inorganic constituents of plants is no less important than the study of their organic elements. The organic substances formed by plants are decomposed by a moderately high temperature; they easily undergo putrefaction, especially when exposed to a moist and warm atmosphere, and they have not been formed by human art. Their inorganic constituents, on the other hand, are not so easily decomposed; they do not undergo putrefaction, and they have been formed artificially by the chemist.

220. The combustible or organic part of plants, even in a dried state, forms from 88 to 99 per cent. of their whole weight. Consequently, the ash or inorganic matter frequently constitutes a very small proportion of the vegetable tissue. It is not, however, on this account to be neglected, for it is found to be of great importance in the economy of vegetation, not merely on account of its entering directly into the constitution of various organs, but also from assisting in the production of certain organic compounds. Some of the lower tribes of cellular plants can exist apparently without any inorganic matter. Thus Mülder could not detect a particle of ash in *Mycoderma vini*, nor in moulds produced in large quantity by milk sugar. Deficiency of inorganic matters, however, in general injures the vigour of plants, and it will be found that, in an agricultural point of view, they require particular attention—a distinct relation subsisting between the kind and quality of the crop, and the nature and chemical composition of the soil in which it grows. It has been shown by careful and repeated experiments, that, when a plant is healthy and fairly ripens its seeds, the quantity and quality of the ash is nearly the same in whatever soil it is grown; and that, when two different species are grown in the same soil, the quantity and quality of the ash varies—the difference being greater the more remote the natural affinities of the plants are.

221. The inorganic elements of plants and their combinations, are thus given by Johnston:—

Chlorine (Cl.)	combined with metals forming	chlorides.
Iodine (I.) metals	iodides.
Bromine (Br.) metals	bromides.
	{ metals	... sulphurets.
Sulphur (S.)	{ hydrogen	... {sulphuretted hydrogen, or hydro-sulphuric acid.
	{ oxygen	... sulphuric acid.
Phosphorus (P.) oxygen	... phosphoric acid.
Potassium (K.)	{ oxygen	... potassa.
	{ chlorine	... chloride of potassium.
	{ oxygen	... soda.
Sodium (Na.)	{ chlorine	... {chloride of sodium. (common salt.)
Calcium (Ca.)	{ oxygen	... lime.
	{ chlorine	... chloride of calcium.
Magnesium (Mg.) oxygen	... magnesia,
Aluminum (Al.) oxygen	... alumina.
Silica (Si.) oxygen	... silica.
Iron (Fe.)	{ oxygen	{ oxides and
Manganese (Mn.)		
Copper (Cu.)		
 sulphur	... {sulphurets.

To these we may add Fluorine, whose presence in plants has been recently dwelt on by some writers. The extraordinary attraction of this element for Silica renders it a matter of impossibility to procure it in a separate state for examination. It is found only in those vegetable structures in which Silica abounds, in the stems of the Gramineæ, and Equisetaceæ. No combination of Fluorine and Oxygen has been yet discovered.

222. The quantity of inorganic matter or ash left by plants, varies in different species, and in different parts of the same plant. The dried leaves usually contain a large quantity. Saussure found that—

Dried bark of Oak gave.....	60	of ash in 1000
Dried leaves.....	53
Dried Alburnum.....	4
Dried duramen.....	2

The dried leaves of Elm contain more than 11 per cent. of inorganic matter, while the wood contains less than 2 per cent.; the leaves of the Willow, 8 per cent., wood, 0·45; leaves of Beech, 6·69, wood, 0·36; leaves of Pitch-pine, 3·5, wood, 0·25. Thus, the decaying leaves of trees restore a large quantity of inorganic matter to the soil.

223. The following tables show the relative proportion of inorganic compounds present in the ash of plants:—

According to Sprengel, 1000 lbs. of wheat leave 11·77 lbs., and of wheat straw 35·18 lbs. of ash, consisting of—

	Grain.	Straw.
Potash.....	2·25	0·20
Soda	2·40	0·29
Lime.....	0·96	2·40
Magnesia.....	0·90	0·32
Alumina with trace of Iron.....	0·26	0·90
Silica	4·00	28·70
Sulphuric acid.....	0·50	0·37
Phosphoric acid.....	0·40	1·70
Chlorine.....	0·10	0·30
	<hr/> 11·77 lbs.	<hr/> 35·18 lbs.

In 1000 lbs. of the grain of the Oat, are contained 25·80 lbs., and of the dry straw 57·40 lbs. of inorganic matter, consisting of—

	Grain.	Straw.
Potash.....	1·50	8·70
Soda.....	1·32	0·02
Lime.....	0·86	1·52
Magnesia.....	0·67	0·22
Alumina.....	0·14	0·06
Oxide of Iron.....	0·40	0·02
Oxide of Manganese.....	0·00	0·02
Silica	19·76	45·88
Sulphuric acid.....	0·35	0·79
Phosphoric acid.....	0·70	0·12
Chlorine.....	0·10	0·05
	<hr/> 25·80 lbs.	<hr/> 57·40 lbs.

In 1000 lbs. of the field Bean, field Pea, and Rye-grass hay, after being dried in the air, the following is the amount of ash, and its composition:—

	Field Bean.		Field Pea.		Rye-grass.
	Seed.	Straw.	Seed.	Straw.	Hay.
Potash.....	4·15 ..	16·56	8·10 ..	2·35	8·81
Soda.....	8·16 ..	0·50	7·39 ..	—	3·94
Lime.....	1·65 ..	6·24	0·58 ..	27·30	7·34
Magnesia.....	1·58 ..	2·09	1·36 ..	3·42	0·90
Alumina.....	0·34 ..	0·10	0·20 ..	0·60	0·31
Oxide of Iron.....	— ..	0·07	0·10 ..	0·20	—
Oxide of Manganese.....	— ..	0·05	— ..	0·07	—
Silica.....	1·26 ..	2·20	4·10 ..	9·96	27·72
Sulphuric acid.....	0·89 ..	0·34	0·53 ..	3·37	3·53
Phosphoric acid.....	2·92 ..	2·26	1·90 ..	2·40	0·25
Chlorine.....	0·41 ..	0·80	0·38 ..	0·04	0·06
	<hr/> 21·36	<hr/> 31·21	<hr/> 24·64	<hr/> 49·71	<hr/> 52·86

224. Dr. R. D. Thomson gives the following analysis of the inorganic matter in the stem and seeds of *Lolium perenne*:—

	Stem.	Seed.
Silica	64·57	42·28
Phosphoric acid	12·51	18·89
Sulphuric acid	—	3·12
Chlorine	—	trace.
Carbonic acid	—	3·61
Magnesia	4·01	5·31
Lime	6·50	18·55
Peroxide of Iron	0·36	2·10
Potash	8·03	4·80
Soda	2·17	1·38

225. These substances are variously combined in plants, in the form of sulphates, phosphates, silicates, and chlorides. Some plants, as Wheat, Oats, Barley, and Rye, contain a large quantity of Silica in their straw; others, such as Tobacco, Pea-straw, Meadow-clover, Potato-haulm, and Sainfoin, contain much lime; while-Turnips, Beet-root, Potatoes, Jerusalem-artichoke, and Maize-straw, have a large proportion of salts of potash and soda in their composition. Sulphates and phosphates are required to supply part of the material necessary for the composition of the nutritive proteine compounds found in grain.

226. **Silica** abounds in Grasses, in Equisetum, and other plants, giving firmness to their stems. The quantity contained in the Bamboo is very large, and it is occasionally found in the joints in the form of Tabasheer. Reeds, from the quantity of siliceous matter they contain, are said, during hurricanes in warm climates, to have actually caused conflagrations by striking against each other. In the species of Equisetum, the silica in the ash is as follows:—

	Ash.	Silica.
<i>Equisetum arvense</i>	13·84	6·38
— <i>limosum</i>	15·50	6·50
— <i>hyemale</i>	11·81	8·75
— <i>Telmateia</i>	23·61	12·00

The third of these furnishes Dutch Rush, used for polishing mahogany. The silica is deposited in a regular manner, forming an integral part of the structure of the plant. Many insoluble matters, as silica, seem to be deposited in cells by a process of decomposition. Thus, silicate of potash in a vegetable sap may be mixed with oxalic acid, by which oxalate of potash, and silicic acid will be produced, as in the cells of Grasses and Equisetum. *Chara translucens* has a covering of silicic acid, while *C. vulgaris* has one composed of silicic acid and carbonate of lime; and *Chara hispida* has a covering of carbonate of lime alone. Silica, the only known oxide of Silicon, contains 22·18 parts silicon, and 24 parts oxygen. It is in reality an acid, and a very powerful one. Its insolubility in water prevents the manifestation of its acid properties under ordinary circumstances. In those plants in which silica most abounds, Fluorine has also been discovered. The test for the presence of the latter rests in acting on the fluorine

in its combination with silicon, by concentrated sulphuric acid, and so producing hydrofluoric acid, which possesses the property of fixing etches on glass; the glass to be engraved being coated with etching ground or wax, and the design traced in the usual way with a pointed instrument.

227. **Lime** is found in all plants, and in some it exists in large quantity. It occurs sometimes in the form of carbonate on the surface of plants. Thus, many of the Characeæ have a calcareous encrustation. The crystals or raphides (¶ 18) found in the cells of plants, have lime in their composition. In the roots of Turkey and East India Rhubarb, the crystals of oxalate of lime have been estimated at about 25 per cent., while in those of the English plant the proportion is about 10 per cent. In the Cacti tribe crystals of the same character have been observed, the presence of which, in excessive quantity, impart in the old plant a brittleness to the stem.

228. **Soda and Potash** occur abundantly in plants. They are taken up from the soil in combination with acids. Those growing near the sea have a large proportion of soda in their composition, while those growing inland contain potash. Various species of *Salsola*, *Salicornia*, *Halimocnemum*, and *Kochia*, yield soda for commercial purposes, and are called Halophytes (ἅλς, salt, and φυτόν, plant). The young plants, according to Göbel, furnish more soda than the old ones. There are certain species, as *Armeria maritima*, *Cochlearia officinalis*, and *Plantago maritima*, which are found both on the sea-shore, and high on the mountains removed from the sea. In the former situation they contain much soda and some iodine; while in the latter, according to Dr. Dickie, potash prevails, and iodine disappears.

229. **Iron, Manganese, and Copper**, especially the two last, exist in small quantity in plants. Iron is met in the soil either as an oxide, a sulphuret, or a carbonate. The former is that most usual, peroxide of iron. Iron when held in solution as a proto-carbonate is capable of being absorbed into the vegetable tissues. Copper was detected, by Sarzean, in coffee.

230. All these inorganic matters are derived in a state of solution from the soil, and plants are said to have, as it were, a power of selection, certain matters being taken up by their roots in preference to others. Saussure made a series of experiments on this subject, and stated that when the roots and plants were put into solutions containing various saline matters in equal proportions, some substances were taken up by imbibition in larger proportion than others. Bouchardat doubts the accuracy of Saussure's conclusions on this point. He thinks that errors arose from the excretions of the plants and other causes. He performed similar experiments with plants of Mint, which had been growing for six months in water previous to experiment, and he found that in cases of mixed salts in water, the plant absorbed

all in equal proportions. Daubeney states, that if a particular salt is not present, the plant frequently takes up an isomorphous one.

231. The differences in the absorption of solutions depend, perhaps, on the relative densities alone, and not on any peculiar selecting power in roots, for it is well known that poisonous matters are absorbed as well as those which are wholesome. The following experiments show that poisonous matters in solutions, varying from half a grain to five grains to the ounce of water, are taken up by roots, and that some substances which are poisonous to animals do not appear to act energetically upon plants:—

Growing Plants.		
Chloride of zinc,	on beans,	} quickly destroyed.
Sulphate of zinc,	... cabbages and wheat,	
Sulphate of copper,	... beans,	
Nitrate of copper,	... beans,	
Acetate of copper,	... cabbages,	
Bichloride of mercury,	... { beans, wheat, cabbages,	} weak solutions did not destroy. destroyed in a few days. destroyed unless much diluted. destroyed in a few days.
Arsenious acid,	... cabbages and wheat,	
Arsenate of potash,	... barley and cabbages,	
Acetate of lead,	... beans,	
Bichromate of potash,	... cabbages, beans, barley,	
Nitrate & sulphate of iron,	... beans,	} quickly destroyed. plants uninjured, except solution strong. improved when very diluted. injured, and if strong destroyed. no injury when diluted.
Chloride of barium,	... beans,	
Nitrate of baryta,	... cabbages and wheat,	
Nitrate of strontia,	... beans,	
Muriate, sulphate, and nitrate of lime,	... beans,	
Sulphate and muriate of magnesia,	... beans and cabbages,	} no injury when diluted.
Phosphate of soda,	... beans and cabbages,	
Chloride of sodium,	... beans and cabbages,	

232. **Rotation of Crops.**—As the inorganic materials which enter into the composition of plants vary much in their nature and relative proportions, it is evident that a soil may contain those necessary for the growth of certain species, while it may be deficient in those required by others. It is on this principle that the rotation of crops proceeds; those plants succeeding each other in rotation which require different inorganic compounds for their growth. In ordinary cases, except in the case of very fertile virgin soil, a crop, by being constantly grown in successive years on the same field, will deteriorate in a marked degree. Dr. Daubeney has put this to the test of experiment, by causing plants to grow on the same and different plots in successive years, and noting the results:—

		Average of 5 yrs.
Potatoes,	{ in the same plot	72.9 lbs. tubers.
	{ in different plots.....	92.8 — —
Flax,	{ same	15.0 lbs.
	{ different.....	19.9
Beans,	{ same	32.8
	{ different.....	34.8
Barley,	{ same	30.0
	{ different.....	46.5
Turnips,	{ same	104.0
	{ different.....	173.0
Oats,	{ same	28.0
	{ different.....	32.4

This shows a manifest advantage in shifting crops, varying from 1 to 75 per cent. ; the deficiency of inorganic matter being the chief cause of difference. As this matter is shown to be of great importance to plants, it follows that the composition of soil is a subject requiring special notice.

CHEMICAL COMPOSITION OF SOILS.

233. Soils have been divided in the following way, according to the proportion of clay, sand, and lime, which they possess:—

1. Argillaceous soils, possessing little or no calcareous matter, and above 50 per cent of clay.
2. Loamy soils, containing from 20 to 50 per cent of clay.
3. Sandy soils, not more than 10 per cent. of clay.
4. Marly soils, 5 to 20 per cent. of calcareous matter.
5. Calcareous soils, more than 20 per cent. of carbonate of lime.
6. Humus soils, in which vegetable mould abounds.

Below the superficial soil there exists what is called *subsoil*, which varies in its composition, and often differs much from that on the surface. Into it the rain carries down various soluble inorganic matters, which, when brought to the surface by agricultural operations, as trenching and subsoil ploughing, may materially promote the growth of crops. The advantages of subsoil ploughing are dependent on the nature of the soil. Such a process increases the aptitude for damp of badly drained localities, promotes the retention of wet, and in light gravelly soils, renders the surface more porous and less capable of affording nutrition. A knowledge of the chemical as well as mechanical nature of soils, affords to the agriculturist a certain guidance by which to direct his operations; since in the judicious application of manures, certain deficiencies may be supplied, and by the admixture of soils they may be rendered the more suitable for the purposes of vegetation.

234. *Humus*, or decaying woody fibre, exists in soils to a certain

amount. This has been called also *ulmine*, or *coal of humus*. In a soluble state it forms humic and ulmic acid. Humus absorbs ammonia, and it is slowly acted upon by the atmosphere, so as to form carbonic acid by combination with oxygen. Peaty soils contain much of this substance. When peroxide of iron is present in such soils, it loses part of its oxygen, and is converted into the protoxide.

235. **Silica** in greater or less quantity, is found in all soils; but it abounds in sandy soils. In its ordinary state it is insoluble, and it is only when acted upon by alkaline matter in the soil that it forms compounds which can be absorbed by plants. Silica, in a soluble state, exists in minute quantities in soils; the proportion, according to Johnston, varying from 0.16 to 0.84 in 100 parts, while the insoluble siliceous matter varies from 60.47 to 83.31 in 100 parts. Wiegman and Polstorf found that plants took up the silica from a soil composed entirely of quartz sand, from which everything organic and soluble had been removed. The following table shows the plants which germinated, the height to which they grew previously to being analyzed, the quantity of silica they contained when planted, and the increase:—

	Height. inches	Silica in the ash.			Silica had increased.
		Seed.	Plant.		
Barley.....15	0.034	0.355	10 times.
Oats.....18	—	0.064	0.354	5½ —
Buckwheat...18	—	0.004	0.075	18 —
Vetch.....10	—	0.013	0.135	10 —
Clover..... 3½	—	0.009	0.091	10 —
Tobacco..... 5	—	0.001	0.549	500 —

236. **Alumina** exists abundantly in clayey soils, but it does not enter largely into the composition of plants. It has the power of absorbing ammonia, and may prove beneficial in this way.

237. **Lime** is an essential ingredient in all fertile soils. In 1000 lbs. of such soil, there are, according to Johnston, 56 lbs. of lime; while barren soil contains only 4 lbs. The presence of phosphoric acid in soils, in the form of phosphates of potass, soda, and lime, is essential for the production of certain azotised compounds in plants; and sulphuric acid, similarly combined, is required for the formation of others. In wet low lands the addition of lime is found to be of the greatest advantage. It forms a union with such acids as may be generated from the partial decomposition of vegetable remains, and in those localities promotes the formation of carbonic acid by rendering the organized matter in the soil thoroughly decomposable.

238. A rough way of estimating the general nature of a soil, is thus given by Professor Johnston:—

1. Weigh a given portion of soil, heat it and dry it. The loss is water.
2. Burn what remains. The loss is chiefly vegetable matter.

3. Add muriatic acid to residue, and thus the quantity of lime may be determined.
4. Wash a fresh portion of soil to determine the quantity of insoluble siliceous sand.

Such an analysis, however, is by no means sufficient for the purposes of the farmer.

239. The chemical composition of a plant being known, conclusions can be drawn as to the soil most suitable for its growth. This is a matter of great importance both to the farmer and to the planter. In order that a plant may thrive, even in a suitable soil, exposure and altitude must also be taken into account. It is only by attention to these particulars that agricultural and foresting operations can be successful. As regards trees, the following practical observations are given as an illustration of what has been stated. The Scotch Fir thrives best in a healthy soil, incumbent on a pervious subsoil, and at a high altitude; Larch in loam, with a dry subsoil, in a high situation, and on sloping banks; Spruce and silver firs, in soft loam or peaty soil, in a low moist situation, but they will also grow in a dry soil, and in a pretty high altitude; Oak in any soil and situation under 800 feet above the level of the sea, but it thrives best in clayey loam, on a rather retentive subsoil, and on gently sloping ground; Ash and Elm, on a gravelly loam, on gravel or sand, at an altitude under 500 feet above the level of the sea; Sycamore, at 100 feet higher than the ash or elm, and in a more retentive soil and subsoil; Beech, on a dry gravelly soil, and in a rather high situation, but it is often luxuriant on strong retentive clay, and in a low damp situation.

APPLICATION OF MANURE.

240. If the soil does not contain the ingredients required for a crop, they must be added in the form of manure. The principle of manuring is to supply what the plant cannot obtain from the soil, and to render certain matters already in the soil available for nutrition. In order that this may be properly practised, there must be an analysis of the soil, of the plant, and of the manure. Hence the importance of agricultural chemistry to the farmer.

Various kinds of Manure.

241. **Natural Manures**, as farm-yard dung, are more valuable than *simple manures*; inasmuch as the former furnish all the substances required for the growth of plants, while the latter only supply a particular ingredient. Natural manures may be regarded as conferring on the soil the most lasting advantage, as from the slowness of

their decomposition their beneficial effects are not so readily exhausted. The plant itself, in a soluble state, would be the best manure. In ordinary farm-yard manure, the straw is again made available for the purpose of the plant. The whole crop of wheat and oats, however, cannot be returned to the soil, as part must be retained for food. A substitute, therefore, must be found for the portion thus taken away. This contains both azotised and unazotised matters, the former consisting of proteine compounds which supply nitrogen for the muscular tissue of man and animals; the latter of starchy, mucilaginous and saccharine matters, which furnish carbon as a material for respiration and fat. The object of manuring is chiefly to increase the former, and hence those manures are most valuable which contain soluble nitrogenous compounds.

242. The value of manures is often estimated by the quantity of gluten which is produced by their application. Hermbstaedt sowed equal quantities of the same wheat on equal plots of the same ground, and manured them with equal weights of different manures, and from 100 parts of each sample of grain produced, he obtained gluten and starch in the following proportions:—

	Gluten.		Starch.
Without manure.....	9·2	66·7
Cow dung.....	12·0	62·3
Pigeons' do.....	12·2	63·2
Horse do.....	13·7	61·6
Goats' do.....	32·9	42·4
Sheep do.....	32·9	42·8
Dried night soil.....	33·1	41·4
Dried Ox blood.....	34·2	41·3

243. Manures containing ammonia, owe their excellent qualities to the nitrogen which enters into their composition; hence the value of sulphate of ammonia, ammoniacal liquor of gas-works, and urine. The value of guano, or the dung of sea-fowl, depends chiefly on the ammoniacal salts, and the phosphates which it contains; thus supplying the nitrogen and phosphorus requisite for the proteine compounds which contain the elements of flesh and blood. The guano, which is imported, is the excrement of numerous sea-fowl which frequent the shores of South America and Africa. It often contains beautiful specimens of infusoria, as *Campylodiscus*, *Coscinodiscus*, &c. The guano found in caves on the coasts of Malacca and Cochin-China, is the produce of frugivorous and insectivorous bats, and of a species of swallow—the last being the best.

244. The following analyses by Dr. Colquhoun of Glasgow, which are the result of an examination of a large number of samples, give a general idea of the composition of guano. The term *ammoniacal matter* includes urate of ammonia and other ammoniacal salts, as

oxalate, phosphate, and muriate, as well as decayed organic matter of animal origin. The term *bone earth*, includes phosphate of lime (always the principal ingredient), phosphate of magnesia (always in small amount), oxalate of lime; and in African guano, a minute quantity of carbonate of lime, and from $\frac{1}{2}$ to 2 per cent. of fragments of sea-shells. The *fixed alkaline salts*, are various salts of soda, as muriate, phosphate, and sulphate; a little of a potash salt has been detected.

South American Guano.

	Fine Chincha.		Middling.		Inferior.		Low Qualities.	
Ammoniacal matter.....	62	42	28	12	... 15
Bone earth.....	20	24	30	50	... 37
Fixed alkaline salts.....	10	14	21	10	... 5
Rock, sand, earth.....	0.5	5	3	15	... 34
Water.....	7.5	15	18	13	... 9
	<hr/> 100.0		<hr/> 100		<hr/> 100		<hr/> 100	

African Guano.

	Best Ichaboe.		Inferior.		Low Quality.	
Ammoniacal matter.....	45	28	20
Bone earth.....	20	21	17
Fixed alkaline salts.....	12	16	14
Rock, sand, earth.....	1	3	25
Water.....	22	32	24
	<hr/> 100		<hr/> 100		<hr/> 100	

245. The guano from the islands on the British coasts, contains the same ingredients, but the soluble salts are generally washed out by the action of rain. The following is the analysis, by Dr. R. D. Thomson, of guano gathered on Ailsa Craig:—

Water.....	50.30
Organic matter and ammoniacal salts, containing 3.47 per cent. of ammonia.....	12.50
Phosphates of lime and magnesia.....	12.10
Oxalate of lime.....	1.50
Sulphate and phosphate of potash, and chloride of potassium....	1.00
Earthy matter and sand.....	15.00

246. **Simple Manures** supply only one or two of the materials required for the growth and nourishment of plants. The *ammoniacal liquor of gas-works*, in a very diluted state, has been advantageously applied to the soil, on account of the nitrogen which it supplies. *Soot* has also been used, from furnishing salts of ammonia. *Nitrates of potash and soda* have been recommended not only on account of the alkalies, but also on account of the nitrogen which they contain, in the

form of nitric acid. The quantity of gluten is said to be increased by the use of nitrates. *Carbonate of potash and soda* and *chloride of sodium*, are frequently used as manures. The latter is especially useful in the case of plants cultivated inland, which were originally natives of the sea-shore, as Cabbage, Asparagus, and Sea-kale. As *lime* is found in all plants, the salts containing it are of great importance. It may be used in the caustic state with the view of decomposing vegetable matter, and aiding in the formation of carbonic acid. It also neutralizes any acid previously in the soil, as is said to occur occasionally in boggy and marshy land, abounding in species of *Juncus*, *Carex*, and *Eriophorum*, with some *Calluna vulgaris*. Lime also combines with certain elements of the soil, and sets potash free, which reacts on the silica, and renders it soluble. Lime is sometimes washed down into the subsoil; and in such cases, trenching improves the land. *Phosphate of lime* is a valuable manure, both on account of the lime, and of the phosphorus which it contains. Without the presence of phosphates, gluten, and the proteine compounds of plants, cannot be formed. Phosphate of lime exists abundantly in animal tissues; and hence it must be furnished by plants. The use of bone-dust as a manure, depends in a great measure on the phosphate of lime which it contains. Besides phosphate of lime, bones contain about 3 per cent. of phosphate of magnesia, carbonate of lime, and salts of soda. The gelatine of bones also seems to act beneficially, by forming carbonic acid and ammonia. Bones are best applied mixed with sulphuric acid, so as to give rise to the formation of soluble phosphates by decomposition. They are broken into pieces, and mixed with half their weight of boiling water, and then with half their weight of sulphuric acid. The mixture is applied to the soil, either in a dry state by the drill, with saw-dust and charcoal added, or in a liquid state diluted with 100 to 200 waters. Phosphates and other inorganic matters, sometimes exist *potentially* in the soil, but in a dormant state, requiring the addition of something to render them soluble. Allowing the ground to lie fallow, and stirring and pulverizing it, are methods by which air and moisture are admitted, and time is allowed for the decomposition of the materials, which are thus rendered available for plants. *Sulphur* exists in considerable quantity in some plants, as *Cruciferæ*, and it forms an element in albumen; hence the use of sulphuric acid and of sulphates as manures. Sulphate of lime or gypsum, is well fitted as a manure for clover. It acts in supplying sulphur and lime, and in absorbing ammonia. *Charcoal* in a solid state, has been applied with advantage as a manure. It acts partly by taking up ammonia in large quantities, and partly in combining slowly with oxygen, so as to form carbonic acid. The effects of carbonic acid on vegetation are said to be remarkably conspicuous in some volcanic countries, in which this gas is evolved from the bottom

of lakes. When it accumulates in large quantities, however, it destroys plants as well as animals.

247. **Manuring with Green Crops** is sometimes practised. The mode adopted is to sow certain green crops, the roots of which extend deeply into the soil; and when the plants have advanced considerably in growth, to plough them in, and sow a crop of some kind of grain. In this way the nutritive matter from the deeper part of the soil is brought within reach of the roots of the grain crop. Manuring with *sea-weeds* is also resorted to in cases where they are accessible. They supply abundance of carbonate, phosphate, and sulphate of lime, besides chloride of sodium. There are considerable differences in their chemical composition; thus, while in *Laminaria saccharina*, alkaline carbonates, potash, and iodine, predominate; in *Fucus vesiculosus* and *serratus*, sulphates and soda are in excess, and iodine is less abundant. In the cultivation of the Coco-nut Palm, Mr. M'Nab finds that sea-weeds act very beneficially.

248. **Liquid Manures** have of late years been much employed, and the formation of tanks for their reception has been strongly recommended, in which the ammonia is fixed by the addition of sulphuric acid or charcoal. They can be applied after vegetation has advanced, and they are in a state to be made at once available to the crop. More recently some have advocated a system of *steeping* seeds and grains in certain solutions before sowing them. Professor Johnston suggests a mixture of phosphate of soda, sulphate of magnesia, nitrate of potash, common salt, and sulphate of ammonia (1 lb. of each), in ten gallons of water, to steep 300 lbs. of seeds, which are to be afterwards dried with gypsum or quicklime.

249. The following experiment, conducted by Mr. Wilson, at Knock, near Largs, shows the mode of estimating the effects of manures. The land was a piece of three-year old pasture, of uniform quality. It was divided into ten lots, and these were treated with different kinds of manure. The quantity of well-made hay is given in lbs. :—

	Produce per Lot.	Rate per Acre.
Lot 1.—Left untouched.....	420	3360
" 2.— 2½ barrels Irish quicklime.....	602	4816
" 3.— 20 cwt. Lime of gas-works.....	651	5208
" 4.— 4½ cwt Wood charcoal powder.....	665	5320
" 5.— 2 bushels Bone-dust.....	693	5544
" 6.— 18 lbs. Nitrate of potash.....	742	5936
" 7.— 20 lbs. Nitrate of soda.....	784	6272
" 8.— 2½ bolls Soot.....	819	6552
" 9.— 28 lbs. Sulphate of ammonia.....	874	6776
" 10.—100 gallons Ammoniacal liquor of gas-works. } 5° Tweddell's hydrometer... }	945	7560

The value of each application was the same, all were applied at the same time, and the grass also was cut at the same time.

250. Plants are thus employed to form from the atmosphere and soil those organic products which are requisite for the nourishment of man and animals. Nutrition derivable from the atmosphere being generally diffused, is accessible to all plants, and is perpetually renewed. Nutrition derivable from the soil being fixed to certain localities, requires that those elements contributing to it, as they become exhausted, be mechanically supplied. While an animal consumes carbon so as to form carbonic acid, gives off ammonia in various excretions, transforms organized into mineral matters, and restores its elements to air and earth; a plant, on the other hand, fixes carbon in its substance and gives off oxygen, forms from ammonia solid compounds, transforms mineral into organized matters, and derives its elements from the air and earth. Thus, says Dumas, what the atmosphere and soil yield to plants, plants yield to animals, and animals return to the air and earth, a constant round in which matter merely changes its place and form.*

EPIPHYTIC AND PARASITIC PLANTS.

251. Some plants grow without any attachment to the soil, and are able to derive in a great measure, from the atmosphere, all the materials required for their growth. Such plants are called *Epiphytes* (ἐπι, upon, and φυτόν, a plant), or *air-plants*, and may be illustrated by the Tillandsias, Bromelias, and Orchids of warm climates. Such plants, when attached to the surface of trees, may perhaps derive some nourishment from the inorganic matter in the decaying bark; but they do not become incorporated, so to speak, with the trees. Orchidaceous plants, which are always perennial, are found in the greatest variety and profusion in those regions where heat and moisture abound. Extremes of cold or dryness are the least favourable to their growth. Tillandsias and Bromelias flourish in dry hot air without any contact with the earth.

252. There are other plants, however, which are true *Parasites* (παρά, beside, and σίτος, food, deriving food from another), sending prolongations of their tissue into other plants, and preying upon them. Many Fungi, for instance, develop their spores (seeds) and spawn (mycelium) in the interior of living or dead plants, and thus cause rapid decay. The disease of corn, called smut and rust, and the dry rot in wood, are due to the attacks of these parasitic Fungi. The minute dust or powder produced by these plants, consists of millions of germs which are easily carried about in the atmosphere, ready to fix themselves on any plants where they can find a nidus. There are also flowering plants which

* For fuller particulars as to the food of plants, analyses of plants, soils, manures, and rotation of crops, see Johnston's Lectures on Agricultural Chemistry; Liebig's Works; Dumas on Organic Nature; Davy's Agricultural Chemistry, by Shier; Mülder's Chemistry of Organic Bodies, translated by Fromberg; and various Papers in the Quarterly Journal of Agriculture, 1844—46; Saussure's Works; Daubeny on Rotation of Crops, Phil. Trans. 1845; Boussingault, Economie Rurale.

grow parasitically, and they may be divided into two classes; 1. Those which are of a pale or brownish colour, and have scales in place of leaves; and 2. Those which are of a green colour, and have leaves. The former, including *Orobanche* or broom-ropes, *Lathræa* or toothwort, *Cuscuta* or dodder, derive their nourishment entirely from the plant to which they are united, and seem to have little power of elaborating a peculiar sap; while the latter, as *Loranthus*, *Viscum* or mistletoe, *Myzodendron*, *Thesium*, *Euphrasia*, *Melampyrum*, and *Buchnera*, expose the sap to the action of air and light in their leaves, and thus allow certain changes to take place in it. The Mistletoe, from its power of elaboration, is able to grow on different species of plants, as on the apple, beech, oak, &c. Some of these parasites are attached to the roots of plants by means of suckers, as in the case of Broom-ropes, Toothwort, and *Thesium*; while others, as Dodder, Mistletoe, &c., feed upon the stems. The plants to which the parasites are attached give origin frequently to their specific names. The species of *Cuscuta* or dodder, inhabit all the temperate and warm parts of the globe, and are peculiarly destructive to clover and lint. They are produced from seed which at first germinates in the soil like other plants; but after the stem has coiled closely round another plant, and becomes attached to it by means of suckers then all connection with the soil ceases, and the Dodder continues its life as a parasite. A remarkable tribe of parasites, called *Rafflesias*, has been found in Sumatra and Java. They are leafless, and produce brown-coloured flowers, which are sometimes three feet in diameter. On account of their only producing a flower and root, they are denominated *Rhizanthus* (ρίζα, a root, and ἄνθος, a flower). No example of a green colour is known among this tribe.

2.—ABSORPTION AND CIRCULATION OF FLUIDS.

253. While the leaves and other aerial organs of plants have the power of absorbing fluids, it is chiefly in the roots that this process takes place. The cells of the spongioles or fibrils of the roots, are covered by a very delicate membrane (§ 120), which allows the imbibition of fluids to proceed rapidly; and as additions are made to their extremities, they are constantly placed in circumstances favourable for the reception of fresh nutriment. This nutriment is supplied to the plant. Animals having the power of locomotion are enabled, as they exhaust one locality of its nutrition, to remove to another. Plants being fixed, should speedily perish from want of nourishment in the surrounding soil, had not nature, while supplying by many sources the deficiencies their consumption occasion, at the same time enabled them to send roots to considerable distances in search of the requisite food. The nutritive materials in the soil, partly derived from the decomposition of organic and inorganic materials, and partly from the atmosphere,

are supplied to the roots in a state of solution ; and as the substances in the cells of plants are usually denser than the external fluid matters, a process of endosmose takes place by which the latter pass through the cell-membrane in large quantities, while a small portion of the former is given off or excreted by exosmose. These movements have already been alluded to as taking place between fluids of different densities, when separated by an animal or vegetable membrane (§ 27). They are referred by some to electrical agency, and they perform an important part in the motions of vegetable fluids.

254. Endosmose and Exosmose, then, are the names given to the phenomena of mixture through a membrane accompanied with change of volume ; the former being given when the volume increases by an in-going strong current, the latter when the volume diminishes by an out-going weak current. In most cases, but not all, the dense fluid increases. The rapidity of the mixture depends on the position which the denser fluid occupies being quicker when it is uppermost. In fig. 222 is represented the mode of showing endosmose by means of a bladder full of syrup, which is attached to the end of a tube and immersed in water. In this case the water passes rapidly into the bladder by endosmose, so that the fluid rises in the tube, while a portion of the thicker fluid passes out by exosmose. The force of this endosmose may be measured by a graduated tube, as in the figure, or by a tube with a double curvature, as fig. 224, the lower part of which is filled with mercury. In the latter case, the mercury is pushed upwards into a graduated tube, and thus an endosmosmeter (*μέτρον, α* measure), or measure of the force of endosmose, is formed.



255. Dutrochet found that with a membrane of 40 millimetres* in diameter, a tube of 2 millimetres, and a solution of sugar, the density of which was 1·083, the fluid rose 39 millimetres in the space of an hour and a-half ; with syrup, of density 1·145, the rise was 68 millimetres ; and with syrup, of density 1·228, the rise was 106 millimetres. Syrup of density 1·3, produced a current capable of raising a column of mercury of 127 inches, which is equal to a pressure of $4\frac{1}{2}$ atmospheres. Thus the velocity and force of the rise depends on the excess of density of the interior liquor over that of the water outside. Different substances act with various intensity in producing endosmose. The following ratio expresses the variable intensity of endosmose, in different

* A millimetre is about 1-25th of an English inch.

Fig. 222.—Instrument to show Endosmose and Exosmose, consisting of a bladder containing syrup attached to a tube, and plunged in a vessel of water. The inward motion of the water (endosmose) exceeds the outward movement of the syrup (exosmose).

cases in which the density of the solution was the same:—Solution of gelatine, 3; of gum, 5·17; of sugar, 11; of albumen, 12. In order that endosmose and exosmose may take place, the liquids must have an affinity for the interposed membrane, and an affinity for each other, and be miscible. According to Matteucci and Cima, the interposed membrane, whether animal or vegetable, is very actively concerned in the intensity and direction of the endosmotic current. The different surfaces of membranes also act variously, and it is probable that the physiological condition of the membrane has an important effect.

256. The fluid matters absorbed by the roots are carried upwards through the cells and vessels of the stem, under the form of *ascending* or *crude sap*; they pass into the leaves, where they are exposed to the influence of air and light, and afterwards return through the bark in the form of *descending* or *elaborated sap*, and a portion of them ultimately reaches the root, where it is either excreted or mixed with the new fluid entering from the soil. The presence of light is essential for this elaboration of the descending sap. Vegetable growth cannot progress without the vegetable circulation be perfectly accomplished. This act of vegetable vitality may, however, be effected while the plant is removed from the action of light, but the oxygenation of the juices cannot be perfected without their free exposure to its influence.

257. **Ascending or Crude Sap.**—In order to show the course of the fluids in exogenous stems, numerous experiments have been performed by Walker, De la Baisse, Burnett, Schultz, and others. These consisted in making incisions or notches in the bark and wood of trees at different heights, and noting the points where the sap made its appearance at different periods of the year, more especially in spring; also in plunging plants with their roots entire into certain coloured solutions, and marking the course of the coloured fluids. These experiments led to the conclusion that the sap ascends chiefly through the alburnum or newer wood, proceeds along the upper side of the leaves, and returns by their lower side to the bark and root. If incisions are made into the trunk of a tree at different heights early in spring, it is found that the discharge of sap takes place, first from the lower parts of the incisions, and chiefly from the alburnum; while at a later period of the year the discharge, or the bleeding, occurs on both sides of the incision, chiefly from the new wood on the lower side, and from the bark on the upper side. If a plant be plunged into a weak solution of acetate of lead (which is capable of being absorbed), the metal may be detected, first in the new wood, next in the leaves, and then in the bark. In cellular plants, where this division of structures does not exist, the sap appears to equally permeate each cell, and to generally diffuse itself throughout the plants.

258. From the minuteness of the tissue, and the difficulty of examining the circulation in a living plant, it is not easy to determine the vessels through which the sap moves. In its upward course, it appears to pass through the intercellular spaces of the recent woody tissue and the porous vessels, and in its downward course through the laticiferous vessels and cellular tissue of the bark, being also transmitted laterally through the cells of the medullary rays. In some cases, when the bark has been removed, and the tree continues to live, the descent or fall of the sap takes place by the cells of the medullary rays. In the course of this circulation, the sap nourishes the different organs, its carbonic acid and water are partly decomposed, combinations take place with nitrogen, protoplasm or formative matter is produced, and various secretions are formed in the cells and intercellular passages.

259. Gaseous matters are taken up by the roots of plants and circulated along with the sap, as well as in the spiral vessels. These usually consist of common air, carbonic acid, and oxygen. Hales showed the existence of a large quantity of air in the vessels of the Vine, and Geiger and Proust have proved that the sap of this plant contains much carbonic acid. In some aquatic plants, as *Pontederia* and *Trapa*, there is a quantity of air contained in the vessels or intercellular spaces, with the view of floating them. In *Vallisneria*, the large cells in the centre of the leaves are surrounded by air cavities, which are seen as dark lines under the microscope. When cut, the air comes out in bubbles, and this escape will continue under water for several days, from the part of the leaf attached to the plant, when exposed to the light. An ounce of air has been collected from two leaves of the plant in six days. This air, as well as that contained in sea-weeds, does not enter by stomata, for none exist, but must be taken up by the cells probably in solution.

260. Changes take place in the composition and density of the sap in its upward course, but the chief alterations take place in the leaves. There it is exposed to the influence of light and air, by means of which, as will afterwards be seen, carbon and hydrogen are fixed, oxygen is given off, and an exhalation of watery fluids takes place. The sap becomes denser, and consequently the process of endosmose is promoted, so that the fluids pass from cell to cell along the upper surface of the leaf, and are gradually propelled into the lower cells, where they are acted upon by the air through the stomata, and are ultimately sent into the vascular and cellular tissue of the bark, where further changes take place. Walker, from his experiments, has concluded that until after the development of the leaves no descent takes place.*

261. **Elaborated or Descending Sap.**—The elaborated sap is sometimes clear and transparent, at other times it is milky or variously coloured and opaque. By Schultz it has been called *latex*, and the

* Walker's Exp. on Motion of Sap in Trees. Trans. Roy. Soc. Edin. i. 3.

vessels transmitting it have been denominated *laticiferous* (§ 38). The latex contains granules, which exhibit certain movements under the microscope. These were first noticed by Schultz, who has written a very elaborate treatise on the subject.* Carpenter considers such movements as analogous to those observable in the capillary circulation of animals. On account of these movements in the latex, the laticiferous vessels have been denominated *Cinenchymatous* (κίνηω, I move), and the movements themselves are included under the name *Clycosis* (κύκλος, a circle.) Schultz looks upon the latex as a fluid of vital importance, and similar to the blood in animals. His views are opposed by Mohl, Tristan, and Treviranus, who consider the latex as a granular fluid containing oil, resin, and caoutchouc, which exhibits molecular movements only when injury is done to the vessels containing it.

262. The plants in which the movements are best observed, are those in which the latex is milky or coloured, such as various species of *Ficus*, *Euphorbia*, and *Chelidonium*. In fig. 223 there is represented

a small fragment of a leaf of *Chelidonium majus* (*celandine*), which shows the current of orange granules in the laticiferous vessels, their direction being indicated by arrows. Observation seems to confirm the truth of Schultz's statements. It is true, as Mohl remarks, that any injury done to the part examined causes peculiar oscillatory movements, which speedily cease. Thus if the young unexpanded sepal of the *Celandine* is removed from the plant, and put under the microscope, or if the inner lining of the young stipule of *Ficus elastica* be treated in a similar manner, very obvious motion is seen in the granular contents of the vessels, and this motion is affected by pricking the vessels or by pressure. In order to avoid fallacy, however, if the microscope be applied to the stipules of *Ficus elastica*, while still attached to the plant and uninjured, it will be remarked that, while



223

* Nova acta Academiae Cæsar. Leopold-Carol. Naturæ Curios. tom. xviii.

Fig. 223.—Small portions of the leaf of *Chelidonium majus* or *Celandine*, (highly magnified), showing a net-work of laticiferous vessels. The direction of the currents in the vessels is indicated by the arrows.

pressure with any blunt object on the stipule causes a marked oscillation in the vessels showing their continuity, there will, nevertheless, be observed a regular movement from the apex towards the base, independent of external influences, when the stipule is simply allowed to lie on the field of the microscope without any pressure or injury whatever. This movement has been observed to continue for at least twenty minutes, and there is no doubt it might have been observed longer. It is of importance to distinguish between those molecular movements which are caused by injury and pressure, and those which depend on processes going on in the interior of the living plant. Those experiments, though by no means complete, lead at present to the adoption of Schultz's opinion relative to the existence of cyclosis.

263. The elaborated sap descends partly by the vessels of latex, and partly by those of the liber. It has been said that there is sometimes a difference in the sap contained in these two kinds of vessels. Occasionally, as in *Euphorbia canariensis*, the elaborated sap has acrid properties, while the ascending sap is bland and wholesome. The elaborated sap contributes to the formation of the cambium, which is produced between the bark and wood of exogens.

264. It appears, then, that in the case of Exogenous plants, the fluid matter in the soil, containing different substances in solution, is absorbed by the extremities of the roots, ascends to the stem, passes through the woody tissue, porous vessels and cells, dissolving and appropriating various new substances. Proceeding upwards and outwards, this sap reaches the leaves and the bark, where it is exposed to the air, and is elaborated by the function of respiration. It then returns, or descends chiefly through the bark, either directly or in a circuitous manner, communicating with the central parts by the medullary rays, depositing various secretions, more especially in the bark, and giving origin to substances which are destined to nourish and form new tissues. Finally, it reaches the extremity of the root, where absorption had commenced; a small portion is there excreted, while the remainder mixes with the newly-absorbed fluids, and again circulates in the sap. The celerity with which the sap ascends is thus dependent on the endosmotic property of the cells in the roots, and the density of the fluids presented for their transmission. In the ratio of the transpiration from the leaves will be the absorption by the roots, and the quickness of the ascending fluid. In the spring season, the vegetable circulation is at its greatest activity, for at such a period the roots are most freely supplied with moisture, and the foliation thoroughly under the influence of air, light, and heat. In winter, though the foliation in many plants be persistent, the circulation is scarce perceptible, owing to the suspension of the active agency which those elements exercise.

265. In the case of Endogenous plants, observations are still wanting by which to determine the exact course of their fluids. The vascular bundles contain woody vessels, which probably are concerned in the ascent of the sap, and vessels equivalent to those of the bark and of the latex, which serve for the descent of it. The cellular tissue is also probably concerned in the movements. Cambium is produced in these plants in the neighbourhood of the vascular bundles, and is thus generally diffused through the texture of the stem. In acrogenous stems, it is likely that the sap follows the same course as in Endogens, although, in regard to both, experiments are still wanting. In cellular plants, transmission of the sap takes place from one cell to another; and, as their texture is often delicate, the movements are rapid. Many of these, as sea-weeds, when plunged into water, after having been dried by evaporation, imbibe the fluid with very great rapidity.

266. The **Cause of the Progression of the Sap** has been investigated by Malpighi, Hales, Dutrochet, Draper, Brücke, and Liebig. While the capillarity of the vessels in the higher plants operates to a certain degree, it would appear that the process of endosmose is that by which the continued imbibition and movement of fluids is chiefly carried on. From the loss of its watery contents, by exhalation, and the metamorphoses going on during the process of nutrition and secretion, the sap becomes gradually more and more dense, and thus, throughout the whole plant there is a forcible endosmotic transmission of the thinner fluids, and a constant change in the contents of the cells and vessels. These movements will of course take place with greater vigour and rapidity according to the activity of the processes going on in the leaves, which thus tend to keep up the circulation. While the ascending movement of the sap is powerfully promoted by the active operation at the surface of the leaves, its lateral movements are no less influenced by the individual relations of each distinct cell, since the different functions of separate cells when actively exercised, call into action those vital agencies by which a transmission of the cellular contents is effected.

267. Draper attributes the movement of the sap to capillary attraction, which he considers as an electrical phenomenon. This attraction takes place when a fluid moistens a capillary tube, and there can be no flow unless a portion of this fluid is removed from the upper extremity; for capillarity will not of itself raise a fluid beyond the end of the tube. Evaporation and transpiration, which take place in the leaves, remove a portion of the vegetable fluids, and thus they promote the capillary action of the vessels. When two fluids of different kinds come into contact in a tube on different sides of a membrane (which membrane being porous, may be considered as made up of numerous short capillary tubes), that will pass the fastest which wets it most completely, or has the greatest affinity for it.

Hence, Draper explains the phenomena of endosmose and exosmose by referring them to capillary attraction, aided by transpiration.

268. Liebig adopts a somewhat similar view of the phenomena. He states that the accurate experiments of Hales have shown the effects of evaporation and transpiration on the movements of sap. Transpiration takes place chiefly in clear and dry weather, and consequently is regulated by the hygrometric state of the atmosphere. When the weather is cloudy and the atmosphere moist, transpiration is checked, and stagnation of the juices takes place. The greater the transpiration, the greater the supply of fluid necessary. Hence, plants kept in the dry atmosphere of rooms fade from want of a due supply to compensate for transpiration; and hence the importance of pruning plants before transplanting them, so as to diminish the evaporating surface, and of performing the operation in dull and moist weather, so as to allow the absorption of fluids to keep pace with the transpiration. This process of transpiration, therefore, by forming a vacuum, assists capillary attraction and the atmospheric pressure, and thus the fluids rise. As the process of endosmose and exosmose depends on the chemical affinity between the fluids on each side of a membrane, the porosity of the membrane, and the attraction existing between it and either of the fluids, it follows that the nature of the parietes of the cells and vessels of plants must have a marked effect on their contents and secretions.

269. The observations of physiologists and chemists thus lead to the conclusion, that the movement of the sap in plants is due partly to the changes effected in the leaves and other green parts, by light and air; partly to capillary attraction, the continuous influence of which is kept up by the constant loss of fluids; and partly to endosmose and exosmose. It may be said that there is a *vis a tergo*, without the presence of leaves, as shown by the experiments of Hales (fig. 224), combined with *vis a fronte*, depending on the suction-power of the leaves.

270. When cut twigs or flowers are put into water, their functions are kept up for some time by endosmose and capillarity. The latter power has great influence in such a case, and hence the cleaner the cut the better, so that no lacerated or ragged edge may interrupt its operation. In these circumstances also small solid particles and colouring matters will enter the tubes. Boucherie found that felled trees, the extremities of which were immediately immersed in various solutions, continued to imbibe them with great force and rapidity for many days. A Poplar, 92 feet high, absorbed in six days nearly sixty-six gallons of a solution of pyrolignite of iron.

271. Heat and light have a powerful influence on the movements of the sap, by promoting transpiration and the action of the cells.

After the winter's repose, the first genial sunshine of spring stimulates the sap to activity, and after the leaves are expanded, the circulation goes on with vigour. The effect of leaf-buds in promoting the movement of sap, may be exhibited by introducing a single branch of a vine growing in the open air into a hot-house during winter, thus exposing it to heat and light. In this case the leaves are developed, and the fluids are set in motion from the roots upwards, so as to supply this single branch, although in the other branches there is no circulation.

272. In spring, the first effect of light and warmth is to stimulate the leaf-buds. These enlarge, and the endosmotic process commences in their cells. This is communicated to other cells, and gradually extends to the root, which draws up a continued supply of fluids from the soil. The matter stored up during the winter undergoes changes; certain substances are dissolved, and thus the sap is thickened, so that the endosmotic process is powerfully increased, and the whole plant exhibits an active and vigorous circulation. The starch deposited in the previous season becomes converted into sugar and dextrine, it is thus readily acted on by the ascending fluids, and in a state of solution admits of being generally diffused. Towards the latter part of the season, when the heat and light decrease, the leaves perform their functions more languidly, and there is a near approach to equilibrium in the density of the fluids, and ultimately there is a cessation of the circulation.

273. Liebig thinks that in the case of the vine, in which, according to Brücke, the specific gravity of the sap in spring is very little more than that of water, the rise of the sap does not at this season depend on endosmose, but on the disengagement of gas, which was shown by Hales to be given off in large quantities, when the vessels were cut. The gas is conjectured to be carbonic acid gas, judging from the experiments of Geiger and Proust, who showed that the sap of the vine contains much of this acid. In hollow stems the disruption and absorption of the medullary cells renders their cavities vast receptacles for air. In the leaves of many aquatics, air has been observed to form distension, and thereby to render them more buoyant.

274. The height to which the sap rises in the case of lofty trees, with spreading roots, is very great. The force with which it ascends has been measured by Hales, and is found to vary according to the state of the weather and the vigour of the plant. By fastening a bent tube, containing mercury, on the stem of the vine, he found in one of his experiments that the sap raised the mercury upwards of thirty inches. The apparatus used by Hales is similar to that used by Dutrochet, to measure endosmose, as is represented at fig. 224, where *c* is the stem of a vine cut, *t* is a bent glass tube fitted to the

cut extremity of the vine by a copper ring, *v*, carefully luted and secured by a bit of bladder, *m*; *n n* represents the level of the mercury in the two branches of the lower curvature, before the experiment, and *n' n'* the level at the conclusion of it. He calculated that the force of the sap in the vine, in some of his experiments, was five times greater than that of the blood in the crural artery of the horse.

275. Special Movements of Fluids.—

Besides this general circulation of the sap, special movements have been observed in the individual cells of plants, which have been included under the name of *Rotation* (*rota*, a wheel), or *Gyration* (*gyrus*, a circuit or circle). These motions have been detected in the cells of many aquatic plants, especially species of *Chara* and *Vallisneria*, and in the hairs of *Tradescantia*. The currents proceed in a more or less spiral direction, and are rendered visible by the granules of chlorophyll which they carry along with them. There exist also other granules in the fluids, which are coloured yellow by iodine, and are probably of a nitrogenous nature.

276. The species of *Chara*, in which rotation has been observed, are aquatic plants growing in stagnant ponds, and are composed of a series of cylindrical cells, placed end to end. Sometimes the plant consists of a single central cell; at other times there are several smaller ones surrounding it, which require to be scraped off in order to see the movements. Many of the species are incrustated with calcareous matter, so as become opaque, while others, as *Chara flexilis*, included




Fig. 224.—Experiment by Hales, to show the force of ascent of the sap. *c*, Stock of a vine cut. *t*, A glass tube with a double curvature attached to the upper part of the vine-stock, by means of a copper cap, *v*, which is secured by means of a lute and piece of a bladder, *m*. *n n*, Level of the column of mercury in the two branches of the tube at the commencement of the experiment. *n' n'*, Level at the conclusion of the experiment.


under the division Nitella, have no incrustation, and are transparent. Those plants with unencrusted tubular cells best exhibit such movements. In these plants the movements take place between the two membranes of which the cell-wall is composed. They are not interrupted should a division of the cell be effected by ligature, as a separate motion may then be observed in either section. Some granules, of a green colour, are attached to the cell-wall, while others are carried with the current, which passes along one side and returns by the other, following an elongated spiral direction. The descending current in the branches is next to the axis.

277. In *Vallisneria spiralis* (which includes *V. Micheliana* and *Jacquiniiana*), the cells in all parts of the plant, as in the leaf, root, flower-stalk, and calyx, contain numerous green granules, and an occasional cytoblast or nucleus, which, under certain circumstances, are carried, with the juices of the plant, in continual revolution round the walls of each cell. Although in different cells the currents proceed often in different directions, still, in any given cell the rotation is uniform; for if stopped by cold it resumes the same direction. Rotation will continue in detached portions of the plant for several days, or even for three or four weeks. The best way of showing these motions is to take a small portion of a young leaf and divide it in halves, by making a very oblique section on the plane of the leaf, by which means a transparent end is obtained. This should be done at least an hour before it is put under the microscope. The part is to be viewed in water, between two pieces of glass; and a little heat is sometimes useful in causing the movements to commence. In *Vallisneria* the motion ceases at about 45° Fahr., while in *Chara*, it goes on at a lower temperature; if the temperature be raised above 150° the motion ceases.

278. A similar intra-cellular circulation is seen in species of *Potamogeton*, *Hydrocharis*, and many aquatics, as well as in the moniliform purple hairs on the filaments, and in the calycine hairs of *Tradescantia virginica*. In the examination of these hairs a higher microscopic power is required than in the case of the plants previously mentioned. The nucleus in the cells of these hairs is usually fixed to the walls, and the movements take place to and from it, and appear to be confined between a double cell-wall. Fig. 225 shows a calycine hair, *p*, of *Tradescantia virginica*, with a small portion of the epidermis, *e e*, on which a stoma, *s*, is seen. In each of the cells, both of the epidermis and the hair, there is a nucleus, *n*, and rotatory currents, the direction of which is indicated by that of the arrows. In each cell, as seen in *a*, there are several currents, which cross each other at the point where the nucleus is situated, thus giving rise to the appearance of an irregular net-work. The hairs of many other

flowering plants exhibit rotation (fig. 86), and it is probable that in all young cells there are currents or streams radiating from the nucleus. The fluid circulating is a mucilaginous protoplasm or formative matter, and in *Chara* and *Vallisneria* it forms a uniform investing layer on the inner surface of the cell. The motions would appear to be connected in some way with the nutrition of cells and the formation of new ones; and, while they continue throughout life in aquatics, they often cease in plants living in air, after they have attained a certain development. Mohl's experiments have shown that at the temperature of 66° Fahrenheit, the quickest motion was 1-125th of a Parisian line,* the slowest, 1-600th, and the mean, 1-185th.

279. Some of these movements, especially in hairs, were looked upon by Schultz as occurring in minute vessels, and therefore he included them under cyclosis. Schleiden maintains that in the *Vallisneria* cells it is not the cellular sap that is in motion, but a mucilaginous fluid, with which the chlorophylle granules and the nucleus are connected, and which flows in an uninterrupted manner along the cell-walls, but on account of its transparency and slight thickness, is not easily seen. In *Chara*, also, he states it is not the cell-sap which moves, but a denser fluid, present in large quantity, and occupying the outer part of the cell-cavity. Mohl thinks that a homogeneous protoplasm fills these cells at first completely, but that during growth it becomes hollowed out into one or more cavities, and that around these the mucilaginous matter circulates. In *Vallisneria*, there is only one cavity, while in other plants there are several, giving rise to the



225

* Parisian line = 0.88815 of inch.

Fig. 225.—Hair, *p*, taken from the calyx of *Tradescantia virginica*, with a small portion of the epidermis, *e.e.* on which there is a stoma, *s*. In each of the epidermal cells there is a nucleus, *n*, and currents (rotation), the direction of which are indicated by the arrows. In each cell there are several currents moving to and from the nucleus, as is well seen at *a*. In the elongated cells of the hair, the nucleus, *n*, is carried along with the currents.

appearance of mucilaginous streams or lines running from the nucleus to the cell-wall. These mucilaginous lines, he says, occasionally after the circulation has ceased, remain permanently on the cell-wall. The existence of spiral fibres in cells has been traced to currents of this kind. Amici is of opinion that in chara the chlorophylle-granules lining the walls of the cell exercise a galvanic action upon the sap, and that the action is thereby produced.

280. The velocity of the currents in various plants, at 66° to 68° Fahrenheit, is thus given by Mohl:—

Filamental hairs of *Tradescantia virginica*,— $\frac{1}{300}$ to $\frac{1}{500}$ of a Parisian line in a second; mean, $\frac{1}{500}$.

Leaves of *Vallisneria spiralis*—quickest, $\frac{1}{125}$; slowest, $\frac{1}{600}$; mean, $\frac{1}{163}$ of a line in a second.

Stinging hairs of *Utrica baccifera*—quickest, $\frac{1}{27}$; slowest, $\frac{1}{875}$; mean, $\frac{1}{50}$.

Cellular tissue of young shoot of *Sagittaria sagittifolia*, $\frac{1}{125}$, to $\frac{1}{1000}$; mean, $\frac{1}{55}$.

“ “ leaf of do., $\frac{1}{170}$, to $\frac{1}{1300}$; mean, $\frac{1}{255}$.

Hairs of *Cucurbita Pepo*—quickest, $\frac{1}{70}$; slowest, $\frac{1}{600}$; mean, $\frac{1}{87}$.

The measurements were made by noting the passage of the globules across the field of a micrometer, fixed in the ocular of the microscope, and counting the strokes of a second's pendulum. These movements appear more rapid to the observer; but then it must be recollected that the parts are seen in a highly magnified state.

281. **The Cause of Rotation** has not been satisfactorily explained. Some attribute it to electrical or magnetic currents causing attraction and repulsion of the granular contents of cells. The different contents of the cells, according to them, mutually act and react on each other, and thus give rise to movements similar to those which take place on the surface of water when oily or resinous matters are added, and which have been called *epipolic* (ἐπιπολῆς, on the surface). Recent observations, by Dutrochet, seem to show that the magnetic force exercises no influence over the movements in Chara. Others believe that while heat, and electricity, and physical agents, stimulate these movements, they are not the cause of them. Some trace the movements to the presence of the nucleus, and look upon them as connected with the period of growth when new cells are being formed, and as ceasing after the nucleus has disappeared. Mechanical explanations fail to determine the nature of those movements in the variety of circumstances under which they are observable. They are equally inexplicable if regarded as purely vital actions. They must be considered as analogous to those other physico-vital phenomena identified with the active exercise of vegetable vitality.

3.—RESPIRATION OF PLANTS.

282. The changes which are produced in the atmosphere by living plants have been included under the title of *Vegetable Respiration*. The experiments of Priestley, in 1771, showed that plants when put

into an atmosphere containing a considerable proportion of carbonic acid, and exposed to light, purified the air by removing carbon and producing oxygen. Air in which animals had died, was thus rendered again fit for breathing. Percival has confirmed those observations. Scheele made a series of experiments with nitrogen in place of carbonic acid, and he found that plants did not purify an atmosphere composed of nitrogen alone. The *foul* air, then, in his experiments, differed completely from that in Priestley's experiments, and hence the difference of results. Ingenhouz and Senebier performed numerous experiments, which proved that during the day, plants gave out oxygen gas, while during darkness, this process was suspended. The former has shown that the green portions of all vegetables, irrespective of their specific properties, are equally available for such operations; that it is from the under surface of the matured leaves oxygen is chiefly given off, that those recently developed are not so active in such a process, which in plants situated in the shade is inadequate to prevent deterioration of the air. Saussure stated, that during the night, oxygen gas was absorbed in different quantities by plants. Fleshy plants absorbed least; next came evergreen trees, and then deciduous trees and shrubs. This absorption of oxygen is attended with the formation of carbonic and other acids. It has been said that some leaves, on account of this process of oxidation, are acid in the morning, and become tasteless during the day. Decandolle, Ellis, Daubeney, and numerous other observers, have confirmed the conclusions drawn by the early experimenters. The results of all these observations are, that plants, more especially their leaves and green parts, have the power of decomposing carbonic acid under the influence of solar light, and of evolving oxygen. While in darkness, no such decomposition takes place, oxygen is absorbed in moderate quantity, and some carbonic acid is given off. The former process caused by the deoxidizing power of plants, much exceeds the latter in amount.

283. Burnett endeavoured to show that there are two processes constantly going on in plants, one being what he calls *digestion*, consisting in the fixation of carbon and the evolution of oxygen, and only carried on during the day; the other being what he calls *proper respiration*, consisting in the evolution of carbonic acid gas, and carried on at all periods of a plant's growth. He thinks that his experiments prove the disengagement of carbonic acid from the leaves of plants, both during night and day. Carpenter entertains similar opinions. He considers that under all circumstances vegetable respiration is a process continued throughout and essential for vegetable life, that it consists of the elimination from the system of the superfluous carbon, either by its entering into combination with the oxygen of the air, or by giving off carbonic acid to replace the oxygen absorbed.*

* Carpenter, Prin. Phy. p. 730.

Mr. Pepys is of opinion that the evolution of carbonic acid indicates an abnormal condition of the leaf, which, in the process of healthy active vegetation, absorbs carbonic acid and disengages oxygen. He believes that the action of light leads to the greater perfection of this function which, though less energetically performed, is not wholly suspended during the night.* What is generally called vegetable respiration, may be regarded as equivalent to digestion, consisting, as it does, of the decomposition of certain matters, and the fixation of others by a process of assimilation; but there is no evidence of the constant exhalation of carbonic acid from healthy leaves in the same way as occurs in animal respiration. It would appear to be more correct to consider the processes in animals and vegetables as opposed; respiration in the former being the elimination of carbon, while in the latter it is the elimination of oxygen.

284. The changes produced in the atmosphere are caused chiefly by the superficial green parts of plants. It was long ago supposed that the spiral vessels from their structure were to be looked upon as true wind-pipes or tracheæ, conveying air from the stomata or pores in the leaves. But although they contain aeriform matters, they have been shown to be not directly concerned with the changes in the atmosphere, and to have no immediate connection with the stomata. The oxygen evolved by plants appears to be derived from the carbonic acid (CO_2), the carbon of which is appropriated, and from water (HO), the hydrogen of which is assimilated. Light is necessary for these decompositions, and it is probable that the alkalies taken up by the roots aid the process.

285. If the leaves of a plant are bent under an inverted tumbler of water, in a pneumatic trough, and exposed to the sun, bubbles of gas will soon be given off, which are found to be pure oxygen; and if the water contains carbonic acid, there will be a diminution in its quantity. The same leaves in darkness will not evolve any oxygen, light being essential for the process. The oxygen derived from the carbonic acid may be all evolved, or part of it may in its nascent state enter into certain combinations within the plant. The brighter and longer continued the light, the more oxygen is given off, and the greater the quantity of carbon added to the plant. If a healthy plant is covered by a bell jar, and exposed to light for twelve hours, oxygen will be formed, and if carbonic acid be added to the air, it will gradually diminish, while the oxygen will increase. During the night the action is reversed, and if the plant is left twelve hours in darkness, the oxygen will decrease, while carbonic acid will increase. Daubeny, from his experiments respecting the action of plants on a known amount of atmospheric air—effected by placing plants beneath large glasses,—

* Pepys on the Respiration of Leaves of Plants. Phil. Trans. 1843, p. 339.

states that leaves are requisite for the purification of the air, that the action of light causes their active operation for the emission of oxygen, and the decomposition of carbonic acid, that for the elimination of oxygen the presence of carbonic acid is requisite, and that the greatest amount of oxygen which can, by vegetable respiration, be added to air confined within a jar is 18 per cent.*

286. The fixation of carbon probably takes place gradually, giving rise at different stages to the formation of various organized compounds. Thus, two atoms of carbonic acid, by losing one of oxygen, become oxalic acid; this oxalic acid, with the aid of water, may yield other acids, from which by the elimination of oxygen, and the addition of the elements of water, various unazotised matters, as starch, gum, and sugar, may be derived; these changes being promoted by the presence of alkalies. The fixation of carbon and hydrogen from the decomposition of carbonic acid and water, gives rise to the formation of the various secretions found in the bark and external cells, as chlorophylle, resins, oils, caoutchouc, and wax.

287. Carbonic acid, as has been already noticed, is taken up in large quantity by the roots of plants from the soil, and it is also probably absorbed from the atmosphere by the leaves. In the interior of plants it is changed in various ways, but it is in the leaves more especially that its decomposition takes place. At night it is given off unchanged, by what Liebig considers as a mere process of exosmose, in consequence of the dissolved acid being no longer assimilated by the action of light. Others say that carbonic acid is not produced by exhalation only, but is also derived from the direct union between the oxygen of the air and the carbon of the plant. This may occur in some plants without leaves, as Fungi, where a direct process of oxidation takes place in the organic matters which have been assimilated. The quantity of this acid given off during the night, is by no means equal to that which is absorbed by the plant during the day.

288. The parts of plants which are not green, seem to absorb oxygen. Thus, roots and subterranean organs act in this way, and the presence of oxygen seems to be necessary for their growth. There are also certain periods in the life of a plant when carbonic acid is given off in large quantity, even during the day, depending on a chemical change taking place in the starch of the plant, by which it is converted into sugar. These periods are germination, flowering, and fruiting. The changes produced will be alluded to when these subjects are considered. When plants are decaying, or are in an unhealthy state, they undergo chemical changes, by which carbonic acid is formed. This was found by Burnett to have effected the results of some of Mr. Ellis's early experiments.

289. Certain plants have a great power of decomposing carbonic

* Daubeny on the Action of Plants on Atmosphere. Phil. Trans. 1836, p. 163.

acid under the action of light. This is particularly the case with aquatics. It is thus that they keep up the purity of the pools and ponds in which they grow. *Pistia Stratiotes* has this effect in the Batavian ponds, and Sir H. Davy notices the great vigour of aquatic plants in the lake Solfatara, where carbonic acid was constantly bubbling up on the surface. The oxygenation of the water by aquatics has also been observed by Morren of Geneva.

290. Experiments have been made as to the effect of the different rays of the spectrum in aiding the decomposition of carbonic acid, by the green parts of plants. Draper states that the light-giving rays, or those nearest the yellow, have the greatest effect; while the heat-giving and the tithonic, or chemical rays, had scarcely any influence. The experiments of Hunt also lead to the conclusion, that the yellow rays have most effect in the fixation of carbon, and in the production of woody matter.

291. While the breathing of man and animals, and the various processes of combustion, are constantly abstracting oxygen from the atmosphere, and substituting carbonic acid, plants are decomposing this noxious gas, and restoring the oxygen. In tropical countries, where the vegetation is luxuriant and the light intense, the fixation of carbon and evolution of oxygen goes on with great vigour, thus furnishing a supply to those regions where, during certain periods of the year, both vegetation and heat are deficient.

Effects of certain Gases on living Plants.

292. It has been already stated that plants can live in an atmosphere containing a considerable proportion of carbonic acid, provided they are exposed to the light. Thus, an atmosphere which could not be breathed by man and animals is capable of supporting vegetable life. The experiments of Priestley, Percival, and Saussure, show, however, that plants will not continue to exercise their functions in pure carbonic acid gas, but that in all cases a certain quantity of free oxygen must be present. It has been found that though plants do not thrive in pure nitrogen, nor in hydrogen gas, yet their vitality is not destroyed by their presence. Saussure observed that vegetable life was maintained by a plant of *Lythrum Salicaria* for five weeks, in an atmosphere of hydrogen gas. Nitrogen has been proved to be apparently innocuous. These gases seem of themselves to have no directly injurious effects, but to act chiefly by depriving the plants of carbon and oxygen.

293. There are certain gases which have very prejudicial effects on plants, as proved by the experiments of Turner and Christison.*

* See Edinburgh Medical and Surgical Journal, vol. xxviii. p. 356.

Some of them act as irritant poisons, causing local disorganization ; others as narcotic poisons, inducing a drooping and decay of the entire plant. To the former class belong sulphurous acid gas, hydrochloric or muriatic acid gas, chlorine and nitrous acid gas ; while under the latter are classed sulphuretted hydrogen, cyanogen, carbonic, oxide, and ammoniacal gas.

294. **Sulphurous Acid Gas** is highly injurious to plants. It produces greyish-yellow dry-looking spots, on the leaves, which gradually extend until the leaves are destroyed and fall. The effect resembles much the ordinary decay of the leaves in autumn. The proportion of gas, in some experiments, was only 1 in 9,000 or 10,000 parts of air, and the quantity $\frac{1}{2}$ of a cubic inch ; and yet the whole unfolded leaves of a mignonette plant were destroyed in forty-eight hours. This proportion of the gas is hardly or not at all discoverable by the smell.

295. **Muriatic Acid Gas** produced effects similar and scarcely inferior to those of the last-mentioned gas. When $\frac{1}{2}$ of an inch was diluted with 10,000 parts of air, it acted destructively on Laburnum and Larch, destroying the whole vegetation in less than two days. Even when in quantity not perceptible by the smell, it still acts as an irritant poison.

296. **Sulphuretted Hydrogen** acted in a different way from the acid gases. The latter attacked the leaves at the tips first, and gradually extended their operation to the leaf-stalks. When in considerable proportion, their effects began in a few minutes ; and, if diluted, the parts not attacked generally survived if the plants were removed into the air. But in the case of sulphuretted hydrogen, the leaves, without being injured in texture or colour, became flaccid and drooping, and the plant did not recover when removed into the air. It required a larger quantity of this gas to produce the effects stated. When six inches were added to sixty times their volume of air, the drooping began in ten hours. This gas then acts like a narcotic poison, by destroying vegetable life throughout the whole plant at once.

297. These observations point out the great injury which is caused to plants by the gases given off during the combustion of coal, and more especially by certain chemical works. In the vicinity of the latter, the vegetation, for a considerable distance around, is often destroyed, particularly in the direction of the prevailing winds of the locality. The atmosphere of large manufacturing towns, in which fuliginous matter and sulphurous gases abound, is peculiarly hurtful to vegetable life. In order to protect plants from such prejudicial influences, Mr. N. B. Ward has invented close glass Cases, in which plants can be made to grow independently of the noxious atmosphere around.* These Cases consist of a trough containing soil, and a frame of glass, which is accurately fitted upon it. The soil is well supplied

* See Ward on the growth of plants in closed Cases.

with water at first, and after the plants are put in they are kept exposed to the light. In these circumstances, they will continue to thrive for a long time, even for years, without any fresh supply of moisture or any direct exposure to the air. They are peculiarly fitted for rooms where the dryness of the atmosphere interferes with the vigour of plants, by causing greater exhalation than can be compensated by the absorption of moisture by the roots. Some tribes of plants, as Ferns, requiring a humid atmosphere, thrive well in such Cases. The windows of houses may be converted by this means into conservatories. Those who wish to see the effects thus produced, ought to visit Mr. Ward's house, in Wellclose square, London. Nothing can exceed the beauty and luxuriance of his Ferns.

298. But it is not merely as matters of luxury and curiosity that these Cases deserve notice. They serve as a most important means of transporting plants, in a living state, to and from foreign climates; and they are in constant use for that purpose. Plants have thus been brought to this country which could not have retained their vitality in the form of seed, and which would have been destroyed by exposure to the sea-breeze and to the vicissitudes of climate experienced during their transport. By this means Tea and Coffee plants have been transported to these countries without injury, and many valuable specimens, amongst which we may include the *Cinchona Calisaya*, have been forwarded in return to the Botanical gardens in the East. The stillness of the atmosphere in the Case contributes materially to prevent injurious consequences. In June 1833, Mr. Ward filled two Cases with Ferns, Grasses, &c., and sent them to Sydney, where they arrived in January 1834. The plants were taken out in good condition, and the Cases were refilled at Sydney, in February 1834, the thermometer then being between 90° and 100° Fahrenheit. In their passage to England, they encountered very varying temperatures. The thermometer fell to 20° on rounding Cape Horn, and the decks were covered a foot with snow. In crossing the line, the thermometer rose to 120°, and fell to 40° on their arrival in the British channel in the beginning of November, eight months after they had been enclosed. The plants were not once watered during the voyage, and received no protection by day or by night, but were taken out at Loddiges in a most healthy and vigorous condition.

299. It is a mistake to suppose that the air in the Cases is not changed. They are not hermetically sealed; and by the law of diffusion of gases there is a constant although gradual mixture of the external air, free however from many impurities, with that inside. Plants will continue to grow for a long time, even in Cases hermetically sealed, if supplied at first with abundance of good soil and water. By the united action of the plant and light, the air undergoes constant changes, and thus continues fit for vegetable life.

4.—PRODUCTS AND SECRETIONS OF PLANTS.

300. The sap in its progress through the cells and vessels, and especially in its passage through the leaves, is converted into organizable products, from which the vegetable tissues and the secretions contained in them are elaborated. Light, by enabling plants to fix carbon, has an important influence over these secretions. When plants are kept in darkness they become etiolated or blanched, and do not form their proper secretions. Gardeners resort to the practice of blanching when they wish to diminish or destroy certain secretions, and to render plants fit for food; a familiar example of which may be seen in their culture of the *Apium graveolens* or Celery. In speaking of the contents of cells and vessels, allusion has already been made to some of the more important organizable products. It is proposed in this place to take a general view of those vegetable secretions which are connected with the nutrition of plants, or which are important on account of their medical or commercial uses. Some of these occur in small quantity, and are limited to certain plants only; others are abundant, and more universal in their distribution. Thus, while quinine and morphine, the active ingredients of Peruvian bark and opium, are circumscribed, both as regards quantity and distribution, starch, gum, sugar, woody matter, and certain nitrogenous compounds are more abundant, and more generally diffused over the vegetable kingdom. The latter substances therefore demand especial attention. If a plant is macerated in water, and all its soluble parts removed, lignine or woody fibre is left, and the water in which it has been macerated, gradually deposits starch. If the liquid is boiled, a scum coagulates, formed of albumen and some azotised matters, while gum and sugar remain in solution.

301. **Starch** is a general product, being laid up as a store of nourishment, and undergoing changes at certain periods of a plant's life, which fit it for further uses in the economy of vegetation. It is not found in animal cells. It consists of $C^{12}H^{10}O^{10}$, and occurs in the form of grains of various sizes and forms, having an external membrane, enclosing a soluble substance. By boiling in water, the pellicle bursts, and the contents are dissolved, becoming gelatinous on cooling. The consistence of this jelly is due to the mutual adhesion of the swollen hydrated integuments of the Starch grains. The circular markings and striæ seen on the grains, and the part called the hilum, have already been noticed (§ 17). Some plants, such as potato, arrow-root, and wheat, contain a large quantity of starch, which varies, however, in quantity according to the period of growth. Thus, while starch abounds towards the latter part of the season in the potato, it decreases when

the tubers begin to germinate in spring. It was found that 240 lbs. of potatoes, left in the ground, contained of starch:—

In August.....	23 to 25 lbs., or	9·6 to 10·4 per cent.
“ September.....	32 “ 38 “	“ 13·3 “ 16 “
“ October.....	32 “ 40 “	“ 13·3 “ 16·6 “
“ November.....	38 “ 45 “	“ 16 “ 18·7 “
“ April.....	38 “ 28 “	“ 16 “ 11·6 “
“ May.....	28 “ 20 “	“ 11·6 “ 8·3 “

The quantity of starch remained the same during the dormant state in winter, but decreased whenever the plant began to grow, and to require a supply of nourishment.

302. Starch is stored up in many seeds. It exists in roots, especially in those which are fleshy; in stems; in the receptacles of flowers; and in pulpy fruits. The seed-lobes of the Bean and Pea, and many other leguminous plants; the roots and the under-ground stem of *Maranta arundinacea* or Arrow-root, and of *Canna coccinea* or Tous-les-mois, *Canna Achiras* and *C. edulis*; the stem of the Sago Palm (*Sagus Rumphii* and *farinifera*), and of the *Cycas* tribe; the receptacle of the artichoke, and the pulp of the apple, are familiar instances of parts in which starch abounds. The grains of potato-starch are pearly or sparkling in their appearance, of large size, having one or more hila, and often cracks on the surface. Those of arrow-root are dull, white, and small, while those of Tous-les-mois, present a glistening appearance like potato-starch, and are larger. In some cases, starch is associated with poisonous or acid juices, as in *Jatropha Manihot*, which yields Cassava and Tapioca, and in *Arum maculatum*, the under-ground stem of which furnishes Portland sago. Inuline is a substance analogous to starch, to which Iodine communicates a brown colour. It is found in the roots and tubers of *Inula Helenium* (Elecampane), *Dahlia variabilis*, and *Helianthus tuberosus* (Jerusalem artichoke); while Lichenin is a variety of starch occurring in *Cetraria islandica* (Iceland moss). Lichenin or lichen starch consists of $C^{12} H^{10} O^{10}$, and is deposited in the primary cell-wall of the plant, in the form of an encrusting layer. By the action of malt or of sulphuric acid upon starch, by long boiling in water, or by heating in a tray in an oven whose temperature has been raised to 300° Fahrenheit, a gummy matter is produced called *dextrin*,* or soluble starch composed of $C^{12} H^{10} O^{10}$. Some consider this to be the substance contained in the interior of the starch grains. When dried, it constitutes British gum. It is one of the steps in the process of the conversion of starch into sugar.

303. **Gum** is one of the substances which are produced abundantly in the vegetable kingdom. Its composition is $C^{12} H^{11} O^{11}$, the same as that of Cane-sugar. It exists in many seeds, exudes from the stems

* Dextrin is so called from possessing the property of effecting the right-handed rotation of the plane of polarization of a ray of light.

and twigs of many trees, and is contained in the juices of others from which it does not exude. It is one of the forms through which organic matter passes during the growth of plants. The different kinds of gums have been divided into those which are soluble in cold water (Arabine, mucilage), and those which only swell up into a gelatinous matter (Bassorine or Tragacanth, Cerasine and Pectine). Arabine is familiarly known by the name of gum-arabic or gum-senegal, and is the produce of various species of *Acacia*, chiefly natives of Arabia, Egypt, Nubia, and Senegambia, such as *Acacia Ehrenbergii*, *tortolis*, *Seyal*, *arabica*, *vera*, and *albida*. From the bark of these plants it exudes in the form of a thick juice, which afterwards concretes into tears. Ehrenberg has shown that the characters of gum from the same species of plant are liable to considerable variation; that the same tree may yield a transparent or an opaque, a light or a dark coloured gum. Old stunted trees, in hot and dry seasons, yield the most gum. Arabine exists with cerasine in the gum of the Cherry and Plum. *Mucilage* is present in many of the Mallow tribe, as *Malva sylvestris*, *Althæa officinalis* or marsh mallow, and in Linseed. In *Sphærococcus crispus*, mucilage is present, of which the formula is $C^{24} H^{19} O^{19}$. *Bassorine* forms the chief part of gum-tragacanth, the produce of several species of *Astragalus*, and of gum-bassora. It exists in Salep, procured from the tubercles of *Orchis mascula*. *Cerasine* is that part of the gum of the Cherry (*Cerasus*), Plum, and Almond trees, which is insoluble in cold water. *Pectine* is a substance procured from pulpy fruits, as the apple and pear. It forms a jelly with water, and when dried, resembles gum or isinglass. It is changed by alkalies into pectic acid, which is found in many fruits and esculent roots.

304. **Sugar.**—This substance, which forms an important article of diet, exists in many species of plants. Sugars have been divided into those which undergo vinous fermentation, as Cane and Grape sugar, and those which are not fermentescible, as Mannite. *Cane sugar*, $C^{12} H^9 O^9 + 2 HO$, is procured from *Saccharum officinarum* (sugar-cane), *Beta vulgaris* (beet-root), *Acer saccharinum* (sugar-maple), and many other plants. It has been conjectured that the *Calamus* or sweet cane mentioned in the Old Testament, may be the sugar cane. At all events, the plant was known as early as the commencement of the Christian era. In the East and West Indies, at the present time, numerous varieties of cane are cultivated, such as Country cane, Ribbon cane, Bourbon cane, Violet or Batavian cane, which are distinguished by their size, form, the position and colour of their joints, their foliage, and their glumes. Bourbon cane is richest in saccharine matter. Canes demand a fertile soil, and for their perfect maturation they require from twelve to fourteen months. Those which are grown from planted slips, are *plant-canes*, those which sprout up from the old stems, are *rattoons*. After being cut, the canes are crushed (the pressed canes

being called *begass*), the saccharine juice is extracted, evaporated, and crystallized, as Raw or Muscovado sugar, which is afterwards refined in vacuo, so as to form *loaf* sugar.

305. In 1844, the gross amount of sugar entered for consumption in the United Kingdom was 4,139,994 cwt. The quantity of sugar produced from the sugar cane in different parts of the world, in 1839, has been thus estimated:—

British Sugar Colonies.....	3,571,378 cwt.
British India.....	519,126 —
Danish West Indies.....	450,000 —
Dutch West Indies.....	260,000 —
French Sugar Colonies.....	2,160,000 —
United States of America.....	900,000 —
Brazil.....	2,400,000 —
Java.....	4,481,342 —

306. *Maple Sugar* is much used in America. It is procured from the sugar maple by making perforations in the stem, and allowing the sweet sap to flow out. Two or three holes, at the height of eighteen or twenty inches from the ground, are said to be sufficient for an ordinary tree. The season of collecting is from the beginning of February to the middle of April. *Beet sugar* is the produce of the root of *Beta vulgaris*, and is extensively manufactured in many parts of the continent. In the year 1841, there were 142,518 acres in France planted with beet-root for sugar, and the quantity of sugar produced was 31,621,923 kilogrammes (one kilogramme being equal to about $2\frac{1}{2}$ lbs). In Ireland much attention has been directed to the culture of this root with the view of manufacturing sugar. A public company has been formed to carry out the undertaking. Though the sugar obtained has been of a superior quality, it does not appear to have been hitherto afforded in sufficient quantity to render its manufacture profitable as a speculation. *Mannu sugar*, or *Mannite*, differs from the others in not being fermentescible. Its composition is $C^6 H^7 O^6$. It is the chief ingredient of Manna, which exudes from the *Ornus europæa* and *rotundifolia*. From Sicily and Calabria it is exported under the name of flake-manna. Mannite is found in the juices of Mushroom, in Celery, and in *Laminaria saccharina*, and *Eucalyptus mannifera*. Dr. Stenhouse has determined the quantity of Mannite in some sea-weeds as follows:—

<i>Laminaria saccharina</i>	12 to 15 per cent. of Mannite.
<i>Halydris siliquosa</i>	5 to 6 per cent. —
<i>Laminaria digitata</i>	4 to 5 per cent. —
<i>Fucus serratus</i>	rather less. —
<i>Alaria esculenta</i>	about the same. —
<i>Rhodomenia palmata</i>	2 to 3 per cent. —
<i>Fucus vesiculosus</i>	1 to 2 per cent. —
<i>Fucus nodosus</i>	nearly same. —

Knop and Schnederman have detected Mannite in *Agaricus piperatus*, and other chemists have found it in *Cantharellus esculentus*, and *Clavaria coralloides*.

307. *Grape sugar*, called also *Starch sugar*, or *Glucose*, is composed of $C^{12} H^{14} O^{14}$. It occurs in the juices of many plants, and is a product of the metamorphoses of starch, cane sugar, and woody fibre. It may be extracted from dry grapes, and may be prepared from starch by the action of an infusion of malt, or of a substance called *Diastase* (§ 310). It is less soluble and less sweet than cane-sugar. It gives sweetness to gooseberries, currants, apples, pears, plums, apricots, and most other fruits. It is also the sweet substance of the chestnut, of the brewer's wort, and of all fermented liquors.

308. **Lignine** is the substance which gives hardness and solidity to the cells and vessels of plants. It exists abundantly in woody fibre, which may be said to be composed of cellulose forming the parietes, and lignine forming the encrusting matter in the interior of the *Sclerogen* of Payen. The latter dissolves in strong nitric acid, forming oxalic acid, while the former is left undissolved. Lignine is said to be composed of $C^{35} H^{24} O^{20}$. According to Müllder, the formula for the ligneous matter of ordinary wood is $C^{40} H^{28} O^{26}$. When a portion of the stem of a herbaceous plant, or of newly cut wood, is reduced to small pieces and boiled in successive portions of water, alcohol, ether, diluted acids and alkalies, until everything soluble in these menstrua is removed, a white fibrous mass remains, to which the name of woody fibre is given. It varies slightly in its composition in different trees, thus:—

	Oak.		Beech.		Pine.		Willow.
Carbon.....	52.53	51.45	50.	49.8
Hydrogen.....	5.69	5.82	5.55	5.58
Oxygen.....	41.78	42.73	44.45	44.62
Iron wood contains 53.44 per cent. of carbon.							

This woody fibre exists in linen and paper; and these substances, when subjected to the action of sulphuric acid, are converted into grape sugar. Lignine gives support to the vegetable texture, and is often deposited in concentric layers. It occurs in large quantity in the wood of trees, and is also present in the stem of herbaceous plants. In some cellular plants it is absent, and the object of many horticultural operations, as blanching, is to prevent its formation. Beet-root and white turnips contain only 3 per cent. Lignine is not coloured by iodine.

309. All these organic substances, consisting of carbon united with the elements of water, are easily convertible into each other by the action of sulphuric acid and heat. Similar changes are induced during

the growth and development of plants, as will be noticed under the head of flowering, fruiting, and germination. In many unazotised matters the proportion of the elements is the same, or they are *isomeric*. Thus, cellulose and starch have the same composition, and the difference in their qualities seems to depend on the mode in which the elements are united. Their form is altered by a change in the molecular arrangement. The unazotised products which have been noticed, supply carbon for the respiration of man and animals, and probably assist in the formation of fat. It is impossible to notice all the compounds of carbon, oxygen, and hydrogen, found in plants. For example, Salicine, a bitter neutral crystalline substance, is procured from the bark of *Salix alba*, *Helix*, *purpurea*, *viminalis*, *pentandra*, &c.; and Phloridzine, an analogous substance, occurs in the bark of the roots of the apple, pear, and plum.

310. Azotised Products.—There are certain azotised products which exist in greater or less quantity in plants, and which are particularly abundant in grains and seeds. The nutritive matter of wheat consists of starch or unazotised matter, separable by washing, and of azotised matter or gluten. *Gluten* is composed of certain proteine compounds (Fibrine, Caseine, Albumen, Emulsine), containing carbon, oxygen, hydrogen, and nitrogen, with some phosphorus and sulphur. *Vegetable fibrine* is the essential part of the gluten of wheat, and of the cereal grains. It may be procured by treating with ether the glutinous mass left after kneading wheat flour in linen bags under water. *Vegetable caseine* or *legumine* is an essential part of the seeds of Leguminous plants, and also of oily seeds. It may be procured in solution from kidney beans and peas, by bruising them in a mortar with cold water, and straining. *Vegetable albumen* occurs in a soluble form associated with caseine. It forms a small proportion of cereal grains. Wheat is said to contain $\frac{1}{4}$ to $1\frac{1}{2}$ per cent.; Rye, 2 to $3\frac{1}{2}$ per cent.; Barley, $\frac{1}{10}$ to $\frac{1}{2}$ per cent.; and Oats, $\frac{1}{3}$ to $\frac{1}{2}$ per cent. It is distinguished by its coagulation at a temperature of 140° to 160° , and by not being precipitated by acetic acid. These three compounds dissolve in a solution of caustic potash; and if to the solution acetic acid is added, the same precipitate is obtained whichever of the three is employed. This precipitate is called Proteine (*πρωτεύω*, I have the first place). Its formula is $C^{48} H^{30} N^6 O^{14}$. Fibrine is $\text{proteine} + S. + Ph.$ Albumen is $\text{proteine} + S^2 + Ph.$ Caseine is $\text{proteine} + S.$ *Emulsine*, or *synaptase*, is a nitrogenous compound found in certain oily seeds, as in almonds. It exists in the milky emulsion which these seeds form in water, and it is coagulated by acetic acid, and by heat. In bitter almonds, it is associated with a substance called *amygdaline*, on which it acts in a peculiar manner, producing hydrocyanic acid. *Diastase* is an azotised substance procured from malt, and developed during

the germination of plants. It is probably fibrine in an altered state, and it has the power of promoting the conversion of starch into sugar.

311. The azotised products of plants have a similar composition with blood and muscular fibre, and hence their value in the food of man and animals. The following table gives a general view of the quantity of azotised and unazotised matters occurring in certain plants, with the amount of water and inorganic matter:—

	Water.	Azotised matter.	Carbonaceous matter.	Ashes.
Peas.....	16	29	52	3
Beans.....	14	31	52	3
Lentils.....	16	33	48	3
Oats.....	18	11	68	3
Barley.....	16	14	69	2
Potatoes.....	72	2	25	1
Turnips.....	89	1	9	1

312. The following arrangement is given by Fromberg of the comparative value of various plants as articles of food, taking into account the proteine compounds, and the starch, gum, and saccharine matter which they contain, the highest value being 100:—

Beans.....	100	Rye.....	55
Peas.....	80	Barley.....	50
Oats.....	75	Potatoes.....	45
Wheat.....	70	Rice.....	35
Maize.....	60		

313. As regards the produce of different crops per acre, Johnston gives the following estimate of the nutritive products which they yield:—

	Average produce per acre of tubers and grain.	No. of lbs. of true nutriment in pro- duce of an acre.
Beet, Mangel-wurzel, and Turnip	30 tons	672 lbs.
Beans.....	30 bushels, or 1980 lbs.	594 —
Potatoes.....	8 tons	358 —
Peas.....	20 bushels, or 1160 lbs.	348 —
Barley.....	36 bushels, or 1872 lbs.	243 —
Jerusalem Artichokes.....	10 tons	224 —
Wheat.....	25 bushels, or 1500 lbs.	180 —
Oats.....	30 bushels, or 1200 lbs.	132 —

314. **Fixed Oils** are found in the cells and intercellular spaces of the fruit, leaves, and other parts. Some of these are drying oils, as Linseed oil, from *Linum usitatissimum*; others are fat oils, as that from Olives (fruit of *Olea europæa*); while others are solid, as Palm oil. The solid oils or fats procured from plants, are Butter of Cacao, from *Theobroma Cacao*; of Cinnamon, from *Cinnamomum*

zeylanicum; of Nutmeg, from *Myristica moschata*; of Coco-nut, from *Cocos nucifera*; of Laurel, from *Laurus nobilis*; Palm oil, from *Elais guineensis*; Shea butter, from *Bassia Parkii*; Galam butter, from *Bassia butyracea*; and Vegetable tallow, from *Stillingia sebifera* in China, from *Vateria indica* in India, and from *Pentadesma butyracea* in Sierra Leone. These oils contain a large amount of stearine, and are used as substitutes for fat. Castor Oil, from the seeds of *Ricinus communis*, differs from other fixed oils in its composition.

315. Decandolle gives the following table to show the quantity of oils got from seeds:—

Hazel-nut.....60	per cent. in weight.	White Mustard...36	per cent. in weight.
Garden Cress.....57	— —	Tobacco.....34	— —
Olive.....50	— —	Plum.....33	— —
Walnut... ..50	— —	Woad.....30	— —
Poppy.....48	— —	Hemp.....25	— —
Almond.....46	— —	Flax.....22	— —
Euphorbia Lath-		Sunflower.....15	— —
yris.....41	— —	Buckwheat.....14	— —
Colza.....39	— —	Grapes.....12	— —

316. **Vegetable Wax** is a peculiar fatty matter sometimes found in the stem and fruit of plants. It is procured from several species of Palms, as *Ceroxylon andicola*, and *Corypha cerifera*, and from the fruit of *Myrica cerifera* or candle-berry myrtle, and *Myrica cordifolia*. By boiling these plants in water and compressing them the wax exudes, floats on the water, and may be collected and re-melted. It is of a greenish-yellow colour. By saponification yields stearic, margaric, and oleic acids, along with glycerine. It therefore more closely approximates the character of fat rather than wax. Waxy matter also occurs on the exterior of fruits, giving rise to the *bloom* of grapes, plums, &c., on the outer surface of the bracts of *Musa paradisiaca*, and on the leaves of many species of *Encephalartos*. In Cork there exists a fatty body which, when acted upon by nitric acid, yields suberic acid. Chlorophylle, or the green colouring matter of leaves, is allied to wax in its nature, being soluble in ether and alcohol, but insoluble in water.

317. **Volatile or Essential Oils** occur in the stem, leaves, flowers, and fruit of many odoriferous plants, and are procured by distillation along with water. They are called *essences*, and contain the concentrated odour of the plant. They usually exist ready-formed, but occasionally they are formed by a kind of fermentation, as oil of bitter almonds, and oil of mustard. Some of them consist of carbon and hydrogen only, as oil of turpentine, procured from various species of *Pinus* and *Abies*; oil of juniper, from *Juniperus communis*; oil of Savin, from *Juniperus Sabina*; oil of lemons and oranges, from the rind of the fruit; and oil of neroli, from orange flowers. A second series contain oxygen in addition, as oil of cinnamon, from *Cinnamo-*

mum zeylaicum; otto or attar of roses, from various species of Rose, especially *Rosa centifolia*; oil of peppermint, from *Mentha viridis*; oil of caraway, from *Carum carui*; oil of cloves, from *Caryophyllus aromaticus*. Oils of this kind are procured from many Labiatae, as species of *Lavandula*, *Origanum*, *Rosmarinus*, *Thymus*; and from the fruit of Umbelliferae, as species of *Anethum*, *Fœniculum*, *Coriandrum*, *Cuminum*, *Petroselinum*, *Pimpinella*; and from some Compositae, as species of *Anthemis*, *Pyrethrum*, and *Artemisia*. A third series have also sulphur in their composition, and have a peculiar pungent, often alliaceous smell, with an acrid burning taste, as oil of garlic, and of onion, procured from the bulbs of *Allium sativum* and *Cepa*; oil of assafœtida, from *Narthex Assafœtida*; and oil of Mustard, which is obtained from the seeds of *Sinapis nigra*, by a kind of fermentation induced by the action of a nitrogenous body, myrosine, on a substance called myronic acid, or myronate of potash, when macerated in water. A similar oil exists in many Cruciferae, as in *Erysimum Alliaria*, *Armoracia rusticana*, and *Cochlearia officinalis*, and in several Umbelliferae, yielding gum-resin, as *Opoponax*, *Ferula*, *Galbanum*, &c. Many of the essential oils deposit a solid crystalline matter, called Stearoptene, allied to *camphor*. This latter substance, which consists of carbon, oxygen, and hydrogen, is procured from *Camphora officinarum*, a native of Japan and India. There is also another kind of camphor, produced in Borneo, by *Dryobalanops Camphora*.

318. **Resinous Products.**—The milky and coloured juices of plants contain frequently resins mixed with volatile oils, in the form of balsams, besides a quantity of caoutchouc. The resinous substances found in plants, are either fluid or solid. The former may be illustrated by Balsam of Tolu, procured from *Myrospermum toluiferum*; Balsam of Peru, from *Myrospermum peruiferum*; Balsam of Copaiva, from various species of *Copaifera*, especially *Copaifera officinalis*; Carpathian Balsam, from *Pinus Pineae*; Strasburg turpentine, from *Abies pectinata*, or silver fir; Bourdeaux turpentine, from *Pinus pinaster*; Canada Balsam, from *Abies balsamea*, or Balm of Gilead fir; Chian turpentine, from *Pistacia Terebinthus*, &c. The latter may be illustrated by common resin or Colophony, and Burgundy pitch, from *Pinus sylvestris*; Mastich, from *Pistacia Lentiscus*; Sandarach, from *Callitris quadrivalvis*; Elemi, from several species of *Amyris*; Guaiac, from *Guaiacum officinale*; Dragon's-blood, from *Dracæna Draco*, and *Calamus Draco*; Dammar, from *Dammara australis* and *orientalis*; Labdanum, from *Cistus creticus*, and others; Tacamahaca, from *Calophyllum Cadaba*, and from *Elaphrium tomentosum*; Resin of Jalap, from *Exogonium Purga*; Storax, from *Styrax officinale*; Benzoin, from *Styrax Benzoin*; Copal, from *Vateria indica*, &c.; Lac, from various species of *Ficus*, as *Ficus indica*, and *benghalensis*, after

attacks of Cocci, and from *Aleurites laccifera*, and *Erythrina monosperma*; *Euphorbium*, from *Euphorbia officinarum*, *antiquorum*, and *canariensis*.

319. **Caoutchouc** is in some respects analogous to essential oils. It is found associated with them and resinous matters, in the milky juice of plants. It is procured from various species of *Ficus*, as *Ficus elastica*, *Radula elliptica*, and *prinoides*, from *Urceola elastica*, *Siphonia elastica*, and *Vahea gummifera*, by wounding the plants. A kind of caoutchouc, called *gutta percha*, imported from Singapore and Borneo, is procured from *Isonandra Gutta*, one of the *Sapotaceæ*. The milky juice of many plants, as of *Euphorbiaceæ*, *Asclepiadaceæ*, *Apocynaceæ*, *Artocarpaceæ*, and *Papayaceæ*, contain caotitchouc or gum elastic. Some of these coloured juices are bland, as that produced by the Cow-tree (*Galactodendron utile*); others are narcotic, as those of Poppy and *Chelidonium*; others are purgative, as *Gamboge*; others diuretic, as *Taraxacum*.

320. **Organic Acids** are produced by processes going on in living plants, and exist in vegetable juices combined often with peculiar bases and alkaloids. Thus Citric acid occurs in the fruit of the orange, lemon, lime, red currant, &c.; Tartaric acid, in the juice of the grape, and in combination with potash in tamarinds; Malic acid, in the fruit of the apple, gooseberry, mountain ash; Tannic acid or Tannine, in oak bark and nut-galls; Gallic acid, in the seeds of Mango; Meconic acid, in the juice of *Papaver somniferum*; Kinic acid, in the bark of various species of *Cinchona*. Besides these, there are numerous others, which are characteristic of certain species or genera. To these may be added Hydrocyanic acid, as found in *Prunus Laurocerasus*, &c., and Oxalic acid, which exists in combination with potash in *Rumex acetosa*, and *Acetosella*, *Oxyria reniformis*, *Oxalis Acetosella*, in the fluid in the pitcher of *Nepenthes distillatoria*; and in combination with lime in *Rhubarb*, and many species of *Parmelia* and *Variolaria*.

521. **Alkaloids or Organic bases** are azotised compounds found in living plants, and generally containing their active principles. They occur usually in combination with organic acids. Quinine and Cinchonine exist in the bark of *Cinchona*, the former predominating in yellow bark, the latter in pale bark; Morphine, Narcotine, Codeine, Thebaine, and Narceine, occur in the juice of *Papaver somniferum*; Solanine is an alkaloid found in many species of *Solanum*, as *Solanum tuberosum*, *nigrum*, and *Dulcamara*; Veratrine exists in *Veratrum Sabadilla* and *album*; Aconitine in *Aconitum Napellus*; Strychnine in *Strychnos Nux-vomica*, *Sancti Ignatii*, *Colubrina* and *Tieuté*; Brucine also in *Nux-vomica*, or False *Angustura* bark; Atropine in *Atropa Belladonna*; Bebeerine in *Neetandra Rodiei*; Piperine in *Piper longum* and *nigrum*; Emetine in *Cephuëlis Ipecacuanha*; Caffeine (Theine and Guaranine) in *Coffea arabica*, *Thea Bohea* and *viridis*, *Paullinia*

sorbilis, and *Ilex paraguayensis*; Theobromine in the seeds of *Theobroma Cacao* or chocolate; besides numerous others of less importance. These Alkaloids are often found in plants having poisonous properties.

322. **Colouring matters** are furnished by many plants, either directly or by a process of fermentation. *Yellow* colouring matters are procured from the roots of *Curcuma longa* or Turmeric, from the pulp surrounding the seeds of *Bixa orellana* (arnotto), from the stem of the Gamboge plant (*Hebradendron Cambogioides*), and various species of *Garcinia*, as *Garcinia Cambogia* and *elliptica*, from the flowers of *Carthamus tinctorius* (safflower), from the stigmata of *Crocus sativus* (saffron), from a kind of Mulberry (*Morus tinctoria*), from *Reseda Luteola* (weld), and from some Lichens, as *Parmelia parietina* (parietin or chrysophanic acid). *Red* colouring matters are produced from the root of *Anchusa tinctoria* (alkanet), from *Pterocarpus santalinus*, *Dracaena Draco* (Dragon's-blood), the root of *Rubia tinctorum* or madder (alizarine), the root of *Morinda citrifolia* (Sooranjee), *Hæmatoxylon campechianum* (logwood), *Cæsalpinia braziliæna* (Brazil wood), from Camwood, also from *Carthamus tinctorius* (Carthamine), and from some Lichens, as *Rocella tinctoria* (Archil and Litmus). *Blue* colouring matters are furnished by the flowers and fruits of many plants, and from the leaves of some, by chemical action. *Indigo*, a most valuable dye, is procured by fermentation from various species of *Indigofera*, as *Indigofera tinctoria*, Anil, *cœrulea* and *argentea*, as well as from *Wrightia tinctoria*, *Marsdenia tinctoria*, *Nerium tinctorium*, and *Gymnema tingens*, &c. The plants in full flower are cut and put into vats with water, fermentation takes place, and a peculiar substance is formed, which, by absorption of oxygen, becomes blue. The best and the largest quantity of Indigo is produced in the Delta of the Ganges. Several Lichens yield nitrogenous colouring matters, which give blue and purple colours with alkalies, &c. *Lecanora tartarea* yields Cudbear.

SECTION III.—ORGANS OF REPRODUCTION.

STRUCTURE, ARRANGEMENT, AND FUNCTIONS.

323. The reproductive organs consist of the flower and its appendages, the *essential* parts being the stamens and pistil. When the flower, or at least the essential organs, are conspicuous, the plants are called *Phanerogamous* (*φανερὸς*, conspicuous, and *γάμος*, union or marriage), or Flowering plants; when they are inconspicuous, the plants are *Cryptogamous* (*κρυπτός*, concealed, and *γάμος*, union or marriage), or Flowerless plants. The former include Exogens and Endogens, the latter Acrogens and Cellular plants. On careful examination it

will be found that the organs of reproduction and nutrition are modifications of each other. The parts of the flower, as regards their development, structure, and arrangement, may all be referred to the leaf as a type. They commence like leaves in cellular projections, in which fibro-vascular tissue is ultimately formed; they are arranged in a more or less spiral manner, and they are often partially or entirely converted into leaves.

1.—INFLORESCENCE, OR THE ARRANGEMENT OF THE FLOWERS ON THE AXIS.

324. The arrangement of the flowers on the axis, or the ramification of the floral axis, is called *Inflorescence* or *Anthotaxis* (ἄνθος, a flower, and τάξις, order). Flower-buds, like leaf-buds, are produced in the axil of leaves, which are called floral leaves or bracts. A flower-bud has not in ordinary circumstances any power of extension by the de-

velopment of its central cellular portion. In this respect it differs from a leaf-bud. In some cases, however, of monstrosity, especially seen in the Rose (fig. 226) and Geum, the central part, A, is prolonged, and bears leaves or flowers. In such cases the flowers are usually abortive, the essential organs being so altered as to unfit them for their functions. Such metamorphoses confirm Goethe's doctrine, that all the parts of the flower are altered leaves.



226

325. The general axis of inflorescence, is sometimes called *rachis* (ῥαχίς, the spine); the stalk supporting a flower, or a cluster of flowers, is a *peduncle* (*pes*, a foot) (fig. 231 a'); and if small branches are given off by it, they are called *pedicels* (fig. 231 a''). A flower having a stalk is called *pedunculate* or *pedicellate* (fig. 231); one having no stalk is *sessile* (fig. 237). In describing a branching inflorescence, it is common to speak of the

Rachis as the *primary* floral axis, its branches as the *secondary* floral axes, their divisions as the *tertiary* floral axes, and so on; thus avoid-

Fig. 226.—Proliferous or monstrous Rose, showing the prolongation of the axis beyond the flowers. c, Calyx transformed into leaves. p, Petals multiplied at the expense of the stamens, which are reduced in number. f, Coloured leaves representing abortive carpels. a, Axis prolonged, bearing an imperfect flower at its apex.

ing any confusion that might arise from the use of the terms *rachis*, *peduncle*, and *pedicel*.

326. The **Peduncle** assumes various forms. It is cylindrical, compressed, and grooved; simple, bearing a single flower, as in Primrose; and branched, as in London-pride. It is sometimes large and succulent, as in the Cashew (fig. 227 *p*), in which the peduncle forms the large coloured expansion supporting the nut; spiral, as in *Cyclamen* and *Vallisneria* (fig. 228); spiny, as in *Alyssum spinosum*. Sometimes the floral axis is shortened, assuming a flattened, convex, or concave form, and bearing numerous flowers, as in the Artichoke, Daisy, and Fig. In these cases it is called a *Receptacle* or *Phoranthium* (φορέω, I bear, and ἄνθος, flower), or *Clinanthium* (κλίνη, a bed, and ἄνθος, flower).

327. The *Floral axis* sometimes assumes a leaf-like or *phylloid* (φύλλον, a leaf, and ἔδος, form) appearance, bearing numerous flowers at its margin, as in *Xylophylla longifolia* (fig. 229), and in *Ruscus*; or it appears as if formed by several peduncles united together so as to become a fasciated axis, as in the Cockscomb (fig. 230), in which the flowers form a peculiar crest at the apex of the flattened peduncles. Adhesions take place between the peduncle and the bracts or leaves of the plant, as in the Lime tree, *Helwingia*, *Chaillertia*, several species of *Hibiscus*, and in *Zostera*. The adhesion of the peduncles to the stem accounts for the extra-axillary position of flowers, as in many *Solanaceæ*; when this union extends for a considerable length along the stem, several leaves may be interposed between the part where the peduncle becomes free, and the leaf whence it originated, and it may be difficult to trace the connection.




Fig. 227.—Fruit of Cashew (*Anacardium occidentale*). *p*, Enlarged peduncle. *a*, Fruit or nut.

Fig. 228.—Pistilliferous plant of *Vallisneria spiralis*, showing spiral peduncles or flower-stalks, by the uncoiling of which the flowers reach the surface of the water, previous to fertilization.

Fig. 229.—Leaf-like (*phylloid*) flattened peduncle, *r*, of *Xylophylla longifolia*. *fff*, Clusters of flowers developed in a centrifugal or cymose manner.

328. The peduncle occasionally becomes abortive, and in place of bearing a flower, is transformed into a tendril (§ 201); at other times it is hollowed at the apex, so as apparently to form the lower part of the outer floral envelope, as in *Eschscholtzia*.



329. The termination of the peduncle, or the part on which the whorls of the flowers are arranged, is called the *Thalamus* or *Torus*. It is the termination of the floral axis. The term *receptacle* is also sometimes applied to this, whether expanded so as to bear several flowers, or narrowed so as to bear one. It may be considered as the growing point of the axis, which usually is arrested by the production of the flowers, but which


sometimes becomes enlarged and expanded. Thus, in the *Geranium*, it is prolonged beyond the flower in the form of a beak; in the *Arum*, it is a club-shaped fleshy column (fig. 239, 2, a); in the *Strawberry*, it becomes succulent and enlarged, bearing the seed-vessels; while in *Nelumbium* it envelops them in the form of a truncated tabular expansion. In some cases it bears the seeds. In some monstrous flowers, as in *Rose* and *Geum*, it is prolonged as a branch bearing leaves (fig. 226).

330. There are two kinds of *inflorescence*—one in which flowers are produced in the axil of leaves, while the axis continues to be elongated beyond them, and to bear other leaves and flowers; the other in which the axis ends in a single terminal flower. In the former, the flowers are axillary, the axis extends in an indefinite manner, and the flowers, as they successively expand, spring from floral leaves placed higher on the axis than the leaf from which the first flower was developed. In the latter, the single solitary flower terminates and defines the axis, and the flowers developed subsequently, arise from floral leaves below this central flower, and therefore farther removed from the centre.

331. The first is *Indeterminate*, *Indefinite*, or *Axillary* inflorescence, in which the axis is either elongated, continuing to produce flower-buds as it grows, the lowermost expanding first; or it is flattened and depressed, and the outermost flowers expand first. The expansion of the flowers is thus *centripetal*, that is, from base to apex, or from circumference to centre. When this kind of inflorescence produces many flowers, it is *simple*, and proceeds from the development of the flower-buds of a single branch. This kind of inflorescence is shown in fig. 231, where the leaf from which the cluster of flowers is produced, *f*, represents the bract or floral leaf. The *rachis*, or primary axis of the flower, is *a'*; this produces small leaflets, *b*, which bear

Fig. 230.—Upper part of flattened or fasciated flowering stem of *Celosia cristata* (*Cockscomb*), having the form of a crest, covered with pointed bracts, and supporting flowers on its summit.

smaller flower-leaves or bractlets, from which peduncles or secondary axes spring, bearing each single flowers. The whole inflorescence is the product of one branch, the lower flowers having expanded first, and bearing fruit, while the upper are in bud, and the middle are in full bloom. In fig. 232, the same kind of inflorescence is shown on a shortened axis, the outer flowers expanding first, and those in the centre last.



231



232

332. The second is *Determinate*, *Definite*, or *Terminal* inflorescence, in which the axis is either elongated and ends in a solitary flower, which thus terminates the axis, and if other flowers are produced, they are secondary, and farther from the centre; or the axis is shortened, and produces at once a number of flower-buds, but of these the central flower expands first, being in fact the termination of the axis, while the other flowers are developed in succession farther from the centre. The expansion of the flowers is in this case *centrifugal*, that is, from apex to base, or from centre to circumference. When this inflorescence produces many flowers, it is *compound*, and proceeds from the development of the buds of several branches. It is illustrated in fig. 233, where a representation is given of a plant of *Ranunculus bulbosus*; a' is the primary axis swollen at the base in a bulb-like manner, b , and with roots proceeding from it. From the leaves which are radical proceeds the axis ending in a solitary terminal flower, f' . About the middle of this axis there is a leaf or bract from which a secondary floral axis, a'' , is produced, ending in a single flower, f'' , less advanced than the flower, f' . This secondary axis bears a leaf also from which a tertiary floral axis is produced, a''' , bearing an unexpanded solitary flower, f''' . From this tertiary axis a fourth is in progress of forma-

Fig. 231.—Raceme of Barberry (*Berberis vulgaris*), produced in the axil of a leaf or bract, f , which has been transformed into a spine, with two stipules, s , at its base. a' , Primary floral axis, bearing small alternate bracts, b , in the axil of which the secondary axes, a'' a''' , are produced, each terminated by a flower. The expansion of the flowers is centripetal, or from base to apex: the lower flowers have passed into the state of fruit, the middle are fully expanded, and those at the top are still in bud. Indeterminate simple inflorescence.

Fig. 232.—Head of flowers or glomerulus of *Scabiosa atro-purpurea*. The inflorescence is simple and indeterminate, and the expansion of the flowers centripetal, those at the circumference opening first.

tion. Here f' is the real termination of the axis, and this flower then expands first, the other flowers being developed centrifugally on separate axes. It is a compound inflorescence.

333. **Indefinite Inflorescence.**—The simplest form of this inflorescence is when single flowers are produced in the axils of the ordinary leaves of the plant, the axis of the plant elongating beyond them, as in *Veronica hederifolia*, *Vinca minor*, and *Lysimachia nemorum*. The ordinary leaves in this case become floral leaves, by producing flower-buds in place of leaf-buds. In place of solitary flowers there is often an elongated floral axis or peduncle arising from a more or less altered leaf or bract, and bearing numerous leaflets, called *bracteoles* or *bract-*



233




234

lets, from which smaller peduncles are produced, and those in their turn may be branched in a similar way. According to the nature of the subdivision, and the origin and length of the flower-stalks, there


Fig. 233.—Plant of *Ranunculus bulbosus*, showing determinate compound inflorescence. α' , Primary floral axis dilated at its base, so as to form a sort of bulb, b , whence the roots and radical leaves proceed. f' , Solitary flower, terminating the primary axis. About the middle of the axis a leaf is developed which gives origin to a secondary axis, α'' , ending in a solitary flower, f'' , which is not so advanced as f' . On the secondary axis a leaf is formed, from the axil of which a tertiary axis, α''' , proceeds, ending in a flower, f''' , which is still in bud. On this axis another floral leaf and bud is in the progress of formation.

Fig. 234.—Panicle or branching raceme of *Yucca gloriosa*. α' , Primary axis or rachis. α'' , Secondary axes or smaller peduncles. α''' , Tertiary axes or pedicels bearing flowers. $b b b b$, Bracts and bracteoles, in the axil of which the axes are produced. The inflorescence is indeterminate, and consists of a series of racemes on a common axis, α' . The expansion of the whole inflorescence is centripetal, and such is also the case with each of the racemes forming it, the flowers at the base of the axes opening first.

arise numerous varieties of floral arrangements. When the primary peduncle or floral axis, as in fig. 231 *a*, is elongated, and gives off pedicels, *a'*, of nearly equal length ending in single flowers, a *raceme* or *cluster* is produced, as in Currant, Hyacinth, and Barberry. If the secondary floral axis gives rise to tertiary ones, the raceme is branching, and forms a *panicle*. In fig. 234 is represented a panicle of *Yucca gloriosa*, *a'* being the primary axis or rachis with bracts, giving off numerous secondary axes, *a''*, which in their turn develop tertiary axes, *a'''*, the development in each of the secondary axes being centripetal, and *b b b b* being the bracts from which the separate axes are produced. If the peduncles in the middle of a dense panicle are longer than those at the extremities, a *thyrsus* is produced, as in Lilac. If in a raceme the lower flower-stalks are elongated, and come to nearly a level with the upper, a *corymb* is formed, which may be



235



236

simple, as in fig. 235, where the primary axis, *a'*, divides into secondary axes, *a'' a''*, which end in single flowers; or *compound*, as in fig. 236, where the secondary axes again subdivide.

334. If the peduncles or secondary axes are very short or awanting, so that the flowers are sessile, a *spike* is produced, as in *Plantago* and *Verbena officinalis* (fig. 237). The spike sometimes bears unisexual flowers, usually stamiferous, the whole falling off by an articulation, as in Willow or Hazel (fig. 238), and then it is called an *amentum* or *catkin*; at other times it becomes succulent, bearing

Fig. 235.—Corymb of *Cerasus Mahaleb*, produced in the axil of a leaf which has fallen, and terminating an abortive branch, at the base of which are modified leaves in the form of scales, *e*. *a'*, Primary axis, or peduncle, or rachis, producing alternate bracts, *b b*, from the axil of which secondary axes or pedicels, *a'' a''*, arise, each bearing a single flower. The evolution or expansion of the flowers is centripetal.

Fig. 236.—Compound or branching corymb of *Pyrus torminalis*. *a'*, Primary axis. *a'' a''*, Secondary axes. *a''' a'''*, Tertiary axes or pedicels bearing the flowers directly. *b b b*, Bracts.

numerous flowers surrounded by a sheathing bract or spathe, and then it constitutes a *spadix*, which may be simple, as in *Arum maculatum* (fig. 239), or branching, as in Palms. A spike bearing female flowers only, and covered with scales, is either a *strobilus*, as in the Hop; or a *cone*, as in the Fir (fig. 201). In grasses, there are usually numerous sessile flowers arranged in small spikes, called *Locustæ* or *spikelets*, and these clusters are either set closely along a central axis or rachis, or they are produced on a branched panicle.

335. If the primary axis, in place of being elongated, is depressed or flattened, it gives rise to other forms of indefinite inflorescence.



When the axis is so shortened that the secondary axes or peduncles arise from a common point, and spread out like *radii* of nearly equal length, each ending in a single flower, or dividing again in a similar

Fig. 237.—Spike of *Verbena officinalis*, showing sessile flowers on a common rachis; the inflorescence indeterminate, and the evolution of the flower centripetal. The flowers at the lower part of the spike have passed into fruit, those towards the middle are in full bloom, and those at the top are only in bud.

Fig. 238.—Amentum or catkin of Hazel (*Corylus Avellana*), consisting of an axis or rachis covered with bracts in the form of scales (*squamæ*), each of which covers a male flower, the stamens of which are seen projecting beyond the scale. The catkin falls off in a mass, separating from the branch by an articulation.

Fig. 239.—Spadix or succulent spike of *Arum maculatum*. 1, Exhibits the sagittate leaf, the spathe or sheathing bract, *b*, rolled round the spadix, the apex of which, *a*, is seen projecting. 2, Shows the spathe, *b*, cut longitudinally, so as to display the spadix, *a*. *f*, Female flowers at the base. *m*, Male flowers. On the spadix there are numerous abortive flowers indicated by hair-like projections.

radiating manner, an *Umbel* is produced, as in figs. 240 and 241. In fig. 240 the floral axes, a', a', a' , end in simple umbels, o', o', o' , and the umbels are called *stipitate* or *stalked*; while in fig. 241 the primary floral axis, a' , is very short, and the secondary axes, a', a' , come off from it in a radiating or umbrella-like manner, and end in small umbels, o', o' , which are called *partial umbels* or *umbellules*, to distinguish them from the *general umbel* arising from the primary axis. This inflorescence is seen in Hemlock, and other allied plants, which are hence called Umbelliferous.



241



240



242


336. If there are numerous flowers on a flattened convex or slightly concave receptacle, having either very short pedicels or none, a *capitulum* (a head) or *anthodium* (ἄνθος, a flower, ὁδός, a way or method), or *calathium* (καλάθιον, a small cup), is formed, as in Dandelion,

Fig. 240.—Several umbels, $o' o' o'$, of *Aralia racemosa*. a , General axis or the apex of the branch terminated by a single umbel farther advanced than the rest $a' a' a'$, Axes arising from it, which are secondary as respects the general axis, a ; each of them bears an umbel, and as regards this inflorescence they are primary. $a'' a'' a''$, Secondary axes, or the radii of the umbel. $b b b$, Bracts placed alternately on the general axis. d , Shows a double bud proceeding from the axil of one of these bracts, and thus giving rise to two-stalked or stipitate umbels. $i i i$, Verticillate bracts, forming involucre at the base of the radii of the umbels.

Fig. 241.—Compound umbel of Carrot (*Daucus Carota*). a' , Primary axis shortened and depressed, so as to present a convex surface. $a'' a''$, Secondary axes or radii of the general umbel, each ending in a partial umbel or umbellule, $o' o' o' o'$. $a''' a'''$, Tertiary axes or radii of the partial umbels or umbellules. f , Pinnatifid bracts, forming the general involucre. $i' i'$, Simple bracts, forming the partial involucre or involucl.

Fig. 242.—Capitulum, Anthodium, or Head of flowers of *Scorzonera hispanica*. b , Imbricated bracts, forming an involucre. f , Florets or small flowers on the receptacle, having a centripetal evolution.


Daisy, and other composite plants (figs. 242 and 243); or a *glomerulus** (a ball), as in *Scabiosa* (fig. 232), and in *Dipsacus* (fig. 244). Such a receptacle or flattened peduncle may some-



243

times be folded so as to enclose partially or completely a number of flowers (generally unisexual), giving rise to the peculiar inflorescence of *Dorstenia* (fig. 245), or to that of the Fig (fig. 246), where *f* indicates the flowers placed on the inner surface of the receptacle, and provided with bracteoles. This inflorescence has been called *Hypanthodium* (ὑπό, under, and ἄθος, a flower).


337. On reviewing these different kinds of inflorescence, it will be observed that the elongation or shortening of the axis, and the presence or absence of stalks to the flowers, determine the different varieties. Thus, a spike is a raceme in which the flowers are not stalked, the umbel a raceme in which the primary



244



245



246

axis is shortened, the capitulum or head a spike in which the same shortening has taken place.

338. **Definite Inflorescence.**—The simplest form of this inflorescence

* By some this term is applied to the centrifugal inflorescence of certain *Urticaceæ*, *Chenopodiaceæ*, and *Juncaceæ*.

Fig. 243.—The same Capitulum cut vertically. *r*, The Receptacle, Pboranthium, or the flattened and depressed apex of the peduncle, bearing the florets, *f*, which are surrounded by bracts, *b*.

Fig. 244.—Inflorescence of *Dipsacus sylvestris*. Glomerulus, or head of flowers, each of which is separated by long pointed bracts. The flowers are evolved in a centripetal manner. *e i*, The first expanded, followed by those at *e m*, while those at the apex, *e s*, are in bud.

Fig. 245.—Inflorescence of *Dorstenia Contrayerva*, consisting of a broad slightly concave receptacle, *r*, on which numerous male and female flowers, *f*, are placed.

Fig. 246.—Inflorescence of Fig (*Ficus Carica*), showing the hollow receptacle, *r*, or peduncle, which forms the fruit covered with flowers, *f*, of various kinds.

is seen in *Anemone nemorosa* (fig. 247), or in *Gentiana acaulis* (*Gentianella*), where the axis terminates in a single flower; and if other flowers are produced, they arise from the leaves below the first-formed flower. When numerous flowers are produced, and the axes are much shortened, it is sometimes difficult to understand this mode of in-



florescence. It may be distinctly traced, however, in plants with opposite leaves, in which the different axes are clearly developed. In fig. 248 is represented the flowering branch of *Erythræa Centaurium*. Here the primary axis, *a'*, ends in a flower, *f'*, which has passed

FIG. 247.—*Anemone nemorosa*. *a*, Subterranean stem. *f*, Leaf. *d*, Floral axis producing bracts, *b*, which form a three-leaved involucre. *e*, Solitary flower terminating the axis. Inflorescence determinate.

into the state of fruit. At its base two leaves are produced, each of which is capable of developing buds. In the *Gentiana acaulis* these leaves rarely produce buds, but in the present plant they generally do. The buds so produced are flower-buds, and constitute secondary axes, $a' a'$, ending in single flowers, $f' f'$, which thus are terminal and solitary; and at the base of these axes a pair of opposite leaves is produced, giving rise to tertiary axes, $a'' a'' a''$, ending in single flowers, $f'' f'' f''$, and so on. The divisions in this case always take place by two, or in a *dichotomous* (δίχα, in two ways, and τέμνω, I cut) manner. Had there been a whorl of three leaves in place of two, the division would have been by three, or *trichotomous* (τρίχα, in three ways).

339. When the leaves become very small, and are transformed into true bracts, this whole system forms a single inflorescence, and has received the name of *cyme*. As the definite inflorescence occurs in a marked degree in the cyme, it has hence been called *cymose*; and the cyme itself, according to its divisions, has been characterized as *dichotomous* or *trichotomous*. In figs. 249 and 250, the cyme is represented in two species of *Cerastium*, belonging to the natural order Caryophyllaceæ, in which cymose inflorescence is of general occurrence. The leaves in the figures are small bracts giving origin to flower-buds in the same way as in fig. 248; the flowers at $a' a'$ being the termination of the primary axis and expanding first, the others being subsequently developed in a centrifugal order. In some of the Pink tribe, as *Dianthus barbatus*, *Carthusianorum*, &c, in which the peduncles are short, and the flowers closely approximated, with a centrifugal expansion, the inflorescence has received the name of *fascicle*. A similar inflorescence is seen in such plants as *Xylophylla longifolia* (fig. 229). When the axes become very much shortened, the arrangement is more complicated in appearance, and the nature of the inflorescence is indicated by the order of opening of the flowers. In *Labiata*, as in the Dead nettle (*Lamium*), the flowers are produced at the axil

Fig. 248.—Flowering branch of *Erythraea centaurium*. a' , Primary axis. $a'' a''$, Two secondary axes. $a''' a''' a''' a'''$, Tertiary axes, four in number. $a'''' a'''' a'''' a''''$, Quaternary axes, eight in number. f' , The flower in various stages of development. f' , Solitary flower which has passed into fruit, terminating the primary axis. f'' , Flowers less advanced, terminating the secondary axes. f''' , Flowers in bud at the extremity of the tertiary axes, and so on. Inflorescence definite or determinate. Evolution of flowers centrifugal.

of each of the leaves, and might be looked upon as ordinary whorls, but on examination it is found that the central flower expands first, and that the expansion is thus centrifugal. The inflorescence is there-




249



250

fore truly cymose, the flowers being sessile, or nearly so, and the clusters are called *verticillasters* (*verticillus*, a kind of screw).

340. Sometimes the bract on one side of a dichotomous cyme, especially towards the summit of the inflorescence, does not give origin to buds, as seen in the upper flower of fig. 250. When a single bract only is produced, in place of two, there is often an anomalous cymose inflorescence produced, resembling a raceme. Thus, in *Alströmeria*, as represented at fig. 251, the axis, a' , ends in a flower, which has been cut off, and a leaf. From the axil of this leaf, that is, between it and the primary axis, a' , a secondary axis, a'' , is formed, ending in a flower f'' , and producing a leaf about the middle. From the axil of this leaf, a tertiary floral axis, a''' , ending in a flower, f''' ,



251

Fig. 249.—Inflorescence of *Cerastium grandiflorum*. $b\ b\ b$, Opposite bracts produced at each of the branchings. The axes are indicated as in last figure. The primary axis, a' , ends in a flower which has passed into fruit. Inflorescence determinate. Evolution of flowers centrifugal.

Fig. 250.—Inflorescence of *Cerastium tetrandrum*. Letters have the same meaning as in the last two figures. In the quaternary axes, a''' , the inflorescence becomes lateral by the non-development of the flower-buds on one side.

Fig. 251.—False raceme of a species of *Alströmeria*. $a' a'' a''' a''''$, Separate axes successively developed, which appear to form a simple continuous raceme, of which the axes form the internodes. It is a compound determinate inflorescence, however, with centrifugal evolution. Each of the axes is produced in the axil of a leaf, and is terminated by a flower, $f' f'' f''' f''''$, opposite to that leaf.

is developed, and so on. Sometimes the bract on the opposite side shows itself, as at *a*. This inflorescence, therefore, although it appears simple and racemose, is truly compound and cymose, consisting of a series of separate axes, with a centrifugal expansion. The flowering branch often exhibits, in such cases, a series of curvations. In certain orders of plants, especially Boraginaceæ, the bracts being alternate, give rise to an inflorescence of this kind. Thus, in fig. 252, *a* is a primary axis, ending in a flower, producing another, *b*, and that, a third, *c*, a fourth, *d*, &c., all on the same side. In such a case there is usually a remarkable curvature resembling the



252



253

tail of a scorpion, and the cyme is called *scorpioid* or *helicoid* (ἑλικοῖς, a spiral, and ἰδιός, form) or *gyrate*.* It is seen in the forget-me-not (fig. 253).


341. Instances of both kinds of floral expansion occur occasionally on the same plant. Thus, in Compositæ, the *heads* of flowers taken as a whole, are developed centrifugally, the terminal head first; while the *florets*, or small flowers on the receptacle, open centripetally, those at the circumference first. So also in Labiatae, the different whorls of inflorescence are developed centripetally, while the florets of the verticillaster are centrifugal. Sometimes this mixed character presents difficulties in cases such as Labiatae, where the leaves, in place of retaining their ordinary form, become bracts, and thus might lead to the supposition of all being a single inflorescence. In such, cases the cymes are described as spiked, racemose, or paniced, according to circumstances. Fig. 254 represents a paniced cyme of Privet, in which the primary axis, *a'*, gives off secondary axes, *a'' a''*, whence arise tertiary, *a''' a'''*, ending dichotomously, and producing three-flowered cymes, *c c*, in which the central flower expands first. Fig. 255 is a racemose cyme of Campanula, developed in a very irregular manner, and giving rise to a peculiar mixed inflorescence; *a' a'* is the primary axis, ending in a flower, *f'*, which has withered, and giving off secondary axes, *a'' a''*, each terminated by a flower, and developed centripetally, the lowest being most expanded. These are anomalous cases, and not

* Schleiden says that this inflorescence is simply a unilateral raceme, having centripetal expansion.

Fig. 252.—Figure to show the formation of a scorpioid or helicoid cyme, consisting of separate axes, *a b c d e*.

Fig. 253.—Scorpioid, gyrate, or helicoid cyme of Forget-me-not (*Myosotis palustris*).

easily explained. Such mixed inflorescences, partly definite and partly indefinite, are by no means uncommon.



254



255

342. Sometimes flowers proceed from what are called radical leaves ; that is, from an axis which is so shortened as to bring the leaves close together in the form of a cluster. From such stems, floral axes are pushed upwards occasionally, bearing single flowers, or flowers in umbels and racemes, as in Primrose, Auricula, Hyacinth, &c. In these cases, the name of *scape* is applied to the flowering stem.

343.—TABULAR ARRANGEMENT OF INFLORESCENCE OR ANTHOTAXIS.

A. Flowers Sessile.

I. Floral Axis elongated.

1. Axis permanent.

Spike (Plantago), Locusta or Spikelet (Lolium), Spadix (Arum), Cone (Fir), Strobilus (Hop).

Fig. 254.—Cymose panicle, or Panicle cyme of Privet (*Ligustrum vulgare*). The primary axis, *a'*, gives off secondary axes, *a'' a''*, which are opposite to each other, and produce tertiary axes, *a''' a'''*, which are dichotomous, and consequently end in small three-flowered cymes, *c c*. Of the three flowers terminating these tertiary axes, the central one expands first, the evolution of the others being centrifugal.

Fig. 255.—Racemose cyme, or Cymose raceme of Campanula. *a' a'*, Primary axis, terminated by a flower, *f'*, which has already withered, and is beginning to pass into the state of fruit. *a'' a''*, Secondary axes, each terminated by flowers, *f'' f''*, which are more advanced the lower they are in their position.

2. Axis deciduous.
Catkin or Amentum (Willow), Compound Catkin (Male flowers of some Palms).
- II. Floral Axis shortened or depressed (a Receptacle).
Capitulum, Anthodium or Calathium (Dandelion).
- . Flowers Pedicellate.
- I. Floral Axis elongated.
 1. Peduncles simple.
 - a. of equal or nearly equal length.
Raceme (Currant, Hyacinth).
 - b. lowermost longest.
Centripetal expansion—Corymb (Ornithogalum).
Centrifugal expansion—Fascicle (Pink).
 2. Peduncles branched.
Panicle (Poa), Thyrsus (Lilac), Anthela (Luzula), Compound Raceme (Plane-tree), Compound Corymb (Milfoil).
- II. Floral Axis shortened or depressed.
 1. Expansion centripetal.
 - a. Peduncles very short, Flowers forming a close head.
Glomerulus (Armeria, Scabious).
 - b. Peduncles nearly equal, radiating from a common centre.
Peduncles simple—Simple Umbel (Astrantia, Ramsons).
Peduncles branched—Compound Umbel (Hemlock).
 2. Expansion centrifugal.
 - a. Peduncles simple—Verticillaster (Lamium).
 - b. Peduncles branched—Cyme (Elder).

2.—BRACTS OR FLORAL LEAVES.

344. Flowers, with the exception of the terminal flower, arise from the axil of leaves, called *Bractæ*, *bracts* or *floral leaves*. The term bract is properly applied to the leaf, from which the primary floral axis, whether simple or branched, arises, while the leaves which arise on the axis between the bract and the outer envelope of the flower are *bracteoles* or *bractlets*. The two are distinct, but are often used indiscriminately in ordinary descriptions. Bracts sometimes do not differ from the ordinary leaves, and are then called *leafy*, as in *Ajuga*. Like leaves, they are either entire or divided. In general, as regards their form and appearance, they differ from the ordinary leaves of the plants, this difference being greater in the upper than in the lower branches of an inflorescence. They are distinguished by their position at the base of the flower or flower-stalk. When the flower is sessile, the bracts are often applied closely to the calyx, and may thus be confounded with it; intermediate between the leaf and the petal, and at times partaking of the characters of each. *Bracts* appear in many instances to be intended by nature as protecting organs, within or beneath which the young flowers in their earliest stage of growth may be secured.

345. When bracts become coloured, as in *Amherstia nobilis*, *Euphorbia splendens*, and *Salvia splendens*, they may be mistaken for

parts of the corolla. They are sometimes mere scales or threads, and at other times they are abortive, and remain undeveloped, giving rise to the *ebracteated* inflorescence of Cruciferae and some Boraginaceae. Sometimes flower-buds are not produced in their axil, and then they are *empty*. A series of empty coloured bracts terminates the inflorescence in *Salvia Horminum*. The smaller bracts or bracteoles, which occur among the subdivisions of a branching inflorescence, often produce no flower-buds, and thus anomalies occur in the floral arrangements.

346. Bracts are occasionally persistent, remaining long attached to the base of the peduncles, but more usually they are deciduous, falling off early by an articulation. In some instances they form part of the fruit, becoming incorporated with other organs. Thus, the cones of Firs (fig. 201) and the strobili of the Hop, are composed of a series of bracts arranged in a spiral manner, and covering fertile flowers; and the scales on the fruit of the Pine-apple (fig. 256 *a*), are of the same nature. In Amenta or catkins (fig. 238), the bracts are called *squamæ* or scales. As regards their arrangement, they follow the same law as leaves; being alternate, opposite, or verticillate.

347. At the base of the general umbel in umbelliferous plants, a whorl of bracts often exists called a *general involucre* (fig. 241 *i'*), and at the base of the smaller umbels or umbellules, there is a similar leafy whorl called *involucel* or *partial involucre* (fig. 241 *i''*). In the case of Compositæ, the name involucre is also applied to the leaves or scales surrounding the head of flowers (fig. 242 *b*), as in Dandelion, Daisy, and Artichoke. This involucre is often composed of several rows of leaflets, which are either of the same or different forms and lengths, and often lie over each other in an imbricated manner. When the bracts are arranged in two rows, and the outer row is perceptibly smaller than the inner, the involucre is sometimes said to be *caliculate*, as in *Senecio*. The leaves of the involucre are spiny in Thistles and in *Dipsacus* (fig. 244 *e i'*), and hooked in Burdock. Such whorled or verticillate bracts may either remain separate (*polyphyllous*), or be united by adhesion (*gamophyllous*), as in many species of *Bupleurum*, and in *Lavatera*. In the acorn they form the *cupula* or cup (fig. 257 *c*), and they also



Fig. 256.—Fruit of Pine-apple (*Ananassa sativa*), composed of numerous flowers united into one mass; the scales, *a*, being modified bracts or floral leaves. The crown, *b*, consists of a prolongation of the axis bearing leaves, which may be considered as a series of empty bracts, *i, e*, bracts not producing flowers in their axil.

form the husky covering of the Hazel-nut. Schleiden considers the cupula of the yew to be a late development of the primine of the ovule.



257


348. When bracts become united together, and overlie each other in several rows, it often happens that the outer ones do not produce flowers or become empty or sterile. In the artichoke, the outer imbricated scales or bracts are in this condition, and it is from the membranous white scales or bracts (*paleæ*) forming the choke attached to the edible receptacle, that the flowers are produced. The sterile bracts of the Daisy occasionally produce *capitula*, and give rise to the Hen-and-Chicken Daisy. In place of developing flower-buds, bracts

may, in certain circumstances, as in proliferous or viviparous flowers, produce leaf-buds.

349. A sheathing bract enclosing one or several flowers, is called a *spatha* or *spathe*. It is common among Endogens, as Narcissus, Arum (fig. 239 *b*), and Palms. Link considers the *spathe* as a modification of the petiole. It is often associated with the spadix, and may be coloured as in Calla or Richardia æthiopica. When the spadix is compound or branching, as in Palms, there are smaller spathes surrounding separate parts of the inflorescence, to which the name *spathellæ* has sometimes been given. The spathe protects the flowers in their young state, and often falls off after they are developed, or hangs down in a withered form, as in some Palms, Typha, and Pothos. In grasses, the outer scales have been considered as sterile bracts, and have received the name of *glumes*, and in Cyperaceæ bracts enclose the organs of reproduction.

3.—THE FLOWER AND ITS APPENDAGES.

350. **The Flower** consists of whorled leaves placed on an axis, the internodes of which are not developed. This shortened axis is the *Thalamus* or *torus*. There



258

are usually four of these whorls or verticils:—
1. The outer one called the *calyx*. 2. The *corolla*. 3. The *stamens*. 4. The most internal one, the *pistil*. Each of these consist normally of several parts, which, like leaves, follow a law of alternation. Thus, the flower of *Crassula rubens* (fig. 258) presents a calyx, *c c*, composed of five equal pieces arranged in a whorl; a corolla, *p p*, also five parts,


Fig. 257.—Acorn, or Fruit of the Oak. *c*, Cupula or cup, formed by the union of numerous bracts or floral leaves, the free points of which are seen arranged in a spiral manner.

Fig. 258.—Flower of *Crassula rubens*. *c c*, Foliola of calyx or sepals. *p p*, Petals. *e e*, Stamens. *o o*, Carpels, each of them having a small scale-like appendage, *a*, at their base.

placed in a whorl within the former, and occupying the intervals between the five parts of the calyx; five stamens, *e e e*, in the space between the parts of the corolla, and consequently opposite those of the calyx; and five parts of the pistil, *o o*, which follow the same law of arrangement. Again, in *Scilla italica*, the parts are arranged in sets of three in place of five, as shown in fig. 259, where *p' p' p'* are three parts of the external whorl; *p'' p'' p''*, three of the next whorl; *e'*, an outer row of stamens; *e''*, an inner row; *o*, the pistil formed of three parts. It is distinctly seen in these instances, that the parts of the flower are to be regarded as leaves arranged on a depressed or shortened axis.

351. When all the parts of the flower are separate, and normally developed, there is no difficulty in tracing this arrangement; but in many cases it is by no means an easy matter to do so, on account of changes produced by the union of one part to another, by degeneration, by the abortion or non-development of some portions, and by the multiplication or folding of others.* Of the four whorls noticed, the two outer (calyx and corolla), are called *floral envelopes*; the two inner (stamens and pistil), are called *essential organs*. When both calyx and corolla are present, the plants are *Dichlamydeous* (δις, twice, and χλαμύς, a covering); occasionally one or both become abortive, and then the flower is either *Monochlamydeous* (μόνος, single), having a calyx only, or *Achlamydeous* (α, privative) or naked, having only the essential organs, and no floral envelope.

352. The **Floral Envelopes** consist of the calyx and corolla. In most cases, especially in Exogens, these two whorls are easily distinguishable, the first being external and green, the latter internal, and more or less highly coloured. If there is only one whorl, then, whatever its colour



259



260

* At the conclusion of the remarks on the organs of reproduction, notice is taken of various metamorphoses produced in flowers by the causes above specified.

Fig. 259.—Flower of *Scilla italica*. *p' p' p'*, Three external leaflets, or divisions of the perianth or Perigone. *p'' p'' p''*, The three internal leaflets. *e'*, Stamens, opposite to the first or external leaflets. *e''*, Stamens, opposite the second or internal leaflets. *o*, Ovaries united together into one. *s*, Three styles, consolidated so as to form one.


Fig. 260.—Flower of White Lily (*Lilium album*). *p*, Perianth or Perigone, having three parts exterior, *p e*, alternating with three interior, *p i*. *e*, Stamens, having versatile anthers attached to the top of the filaments. *s*, Stigma at the apex of the style.

or degree of development, it is the *calyx*. Sometimes, as in many Endogens, the calyx and corolla both display rich colouring, and are apt to be confounded. In such cases, the term *Perianth* (περί, around, and ἄθος, flower), or *Perigone* (περί and γυνή, pistil) has been applied to avoid ambiguity. Thus, in the Tulip, Crocus, Lily, Hyacinth, in place of calyx and corolla, authors speak of the parts of the perianth, although in these plants, an outer whorl (calyx), may be detected, of three parts, and an inner (corolla), of a similar number arranged alternately. Thus, the perianth of the White Lily (fig. 260 p), consists of three outer parts, *p e*, alternating with three internal parts, *p i*, surrounding the essential organs, *e*, the stamens, and *s*, the pistil.

353. The term perianth then is usually confined to the flowers of Endogens, whatever colour they present, whether green, as in Asparagus, or coloured, as in Tulip. Some use the term perianth as a general one, and restrict the use of perigone to cases where a pistil is present, not applying it to unisexual flowers, in which stamens only are produced. In some plants, as *Nymphaea alba* (fig. 310), it is not easy to say where the calyx ends, and the corolla begins; as these two whorls pass insensibly into each other.

354. **Flower-bud.**—To the flower-bud, the name *alabastrus* (meaning *rose-bud*) is sometimes given, and its period of opening has been called *anthesis* (ἄνθησις, flower opening), whilst the manner in which the parts are arranged with respect to each other before opening, is the *æstivation* (æstivus, belonging to summer), or *præfloration* (præ, before, and flos, flower). The latter terms bear the same relation to the flower-bud, that veneration does to the leaf-bud, and distinctive names have also been given to the different arrangements which it exhibits. When the parts of a whorl are placed in an exact circle, and are applied to each other by their edges only, without overlapping or being folded, thus resembling the valves of a seed-vessel, the æstivation is *valvate*, as in calyx of *Guazuma ulmifolia* (fig. 261 c). The edges of each of the parts may be turned either inwards or outwards; in the former case, the æstivation is *induplicate*, as in corolla of *Guazuma ulmifolia* (fig. 261 p), in the latter *reduplicate*, as in calyx of *Althæa rosea* (figs. 262 c, 263 c). When the parts of a single whorl are placed in a circle, but each of them exhibits a torsion of its axis, so that by one of its sides it overlaps its neighbour, whilst its side is overlapped in like manner by that standing next to it, the æstivation is *twisted* or *contorted*, as in Corolla of *Althæa rosea* (figs. 262 p, 264 p). This arrangement is characteristic of the flower-buds of the Malvaceæ and Apocynaceæ, and it is also seen in the Convolvulaceæ and some Caryophyllaceæ. When the flower expands, the traces of twisting often disappear, but sometimes, as in Apocynaceæ, it remains.

355. In these instances of æstivation, the parts of the verticils are considered as being placed regularly in a circle, and about the same



height, and they are included under *circular* æstivation. But there are other cases in which there is a slight difference of level, and then the true spiral arrangement exhibits itself. This is well seen in the leaves of the calyx of *Camellia japonica* (fig. 265 c), which cover each other partially like tiles on a house. This æstivation is *imbricated*. At other times, as in the petals of *Camellia* (fig. 265 p), the parts envelop each other completely, so as to become *convolute*. This is also seen in a transverse section of the calyx of *Magnolia grandiflora* (fig. 267), where each of the three leaves embraces that within it. When the parts of a whorl are five, as occurs in many *Exogens*, and the imbrication is such, that there are two parts external, two internal, and a fifth which partially covers one of the internal parts by its margin, and is in its turn partially covered by one of the external parts, the æstivation is *quincuncial* (fig. 266). This quincuncx is common in the corolla of *Rosaceæ*. Fig. 266 is a horizontal section of the calyx in the flower-bud of *Convolvulus sepium*, in which the parts are numbered according to their arrangement in the spiral cycle, and the course of the spiral is indicated by a line of points. In fig. 268, a section is given of the bud of *Antirrhinum majus*, showing the imbricated spiral arrangement. In this case it will be seen, when contrasted with fig. 266, that the part marked 2, by a slight change



Fig. 261.—Diagram of calyx, c, and corolla, p, in the bud of *Guazuma ulmifolia*. Æstivation of calyx valvate, of petals induplicate.

Fig. 262.—Diagram of calyx, c, and corolla, p, in the flower-bud of *Althæa rosea*. Æstivation of calyx reduplicate, of petals contorted or twisted.

Fig. 263.—Flower-bud of *Althæa rosea* in a young state, showing calyx, c, still completely enveloping the other parts, and the edges of its divisions touching each other.

Fig. 264.—The same in a more advanced state, where the calycine divisions, c, are separated so as to allow the expansion of the corolla, the petals of which, p, are contorted in æstivation.

Fig. 265.—Flower-bud of *Camellia japonica*. c, Imbricated sepals of the calyx. p, Petals with convolute æstivation.

in position, has become overlapped by 4. In flowers, such as those of the Pea (§ 379, fig. 292), one of the parts, the vexillum, is often large and folded over the others, giving rise to *vexillary* æstivation, or the carina may perform a similar part, and then the æstivation is *carinal*.

356. The different verticils often differ in their mode of æstivation. Thus, in Malvaceæ, the corolla is contorted, and the calyx valvate, or reduplicate (fig. 264). In Convolvulaceæ, while the corolla is twisted, and has its parts arranged in a circle, the calyx is imbricated and exhibits a spiral arrangement (fig. 266). In Guazuma (fig. 261), the calyx is valvate, and the corolla induplicate. The circular æstivation is generally associated with a regular calyx and corolla; while the spiral æstivations are connected with irregular as well as regular forms.

357. The different parts of the flower, besides having a certain position as regards each other, bear also definite relations with respect to the floral axis whence they arise. An individual part of a flower may be turned to the one or the other side of the axis, to the right or to the left, in its normal state, and the same will be the case with the corresponding parts of the other flowers. This law often holds good with whole groups of plants, and a means is thus given of characterizing them. If a whorl of the flower consists of four parts, that which is turned towards the floral axis is called *superior* or *posterior*, that next the bract whence the pedicel arises is *inferior* or *anterior*, while the other two are *lateral*. If again, there are five parts of the whorl, then two are inferior, two lateral, and one superior. In plants having blossoms like the Pea, the vexillum, or odd petal, is the superior part; whilst in the calyx, the odd part, by the law of alternation, is inferior. Sometimes the twisting of parts makes an *apparent* change in their position.

External Floral Whorls, or Floral Envelopes.

358. **Calyx.**—The calyx is the external envelope of the flower, and consists of verticillate leaves, called *sepals*, *foliola*, or *phylla*, (*folium*, and φύλλον, a leaf). These calycine leaves are sometimes separate from each other, at other times they are united to a greater or less extent; in the former case, the calyx is *polysepalous* or *polyphyllous* (πολύς,


Fig. 266.—Horizontal section of calyx in flower-bud of *Convolvulus sepium*. Calyx consists of five sepals corresponding to the numbers in the figure, and the line of points indicates the direction of the spiral according to which they are arranged.

Fig. 267.—Arrangement of the three outer leaflets (calyx) in the bud of *Magnolia grandiflora*, cut transversely; æstivation convolute.

Fig. 268.—Arrangement of the parts of the calyx in the flower of Frogmouth (*Antirrhinum majus*). The arrangement differs from that in fig. 266, on account of a slight twisting and overlapping of the parts.

many), in the latter *gamosepalous* or *gamophyllous*, *monosepalous* or *monophyllous* (γάμος, union, or μόνος, one). The divisions of the calyx present usually all the characters of leaves, and in some cases of monstrosity they are converted into the ordinary leaves of the plant. This is frequently seen in the Rose (fig. 226 c), Pæony, &c. Their structure consists of cellular tissue or parenchyma, traversed by vascular bundles, in the form of ribs and veins, containing spiral vessels, which can be unrolled, delicate woody fibres, and other vessels,—the whole being enclosed in an epidermal covering, having stomata, and often hairs on its outer surface, which corresponds to the under side of the leaf.

359. The venation of the calyx in the great divisions of the vegetable kingdom, is similar to that of their leaves; parallel in Endogens, reticulated in Exogens. The leaves of the calyx are usually entire (fig. 269), but occasionally they are cut in various ways, as in the Rose (fig. 270 c f), and they are sometimes hooked at their margins, as in *Rumex uncatus* (fig. 271 c i). In the last-named plant, there are two



whorls of calycine leaves, the outer of which, c e, are entire, and there are also swellings, g, in the form of grains or tubercles on the back of the inner hooked sepals. The outer leaves, c e, may be looked upon in such cases as bracts occupying an intermediate place between leaves and sepals. It is rare to find the leaves of the calyx stalked. They usually consist of sessile leaves, in which the laminar portion is only slightly developed, and frequently the vaginal part is alone present. Sepals are generally of a more or less oval, elliptical, or oblong form, with the extremity either blunt or acute. In their direction they are erect or reflexed (with their apices downwards), spreading out-

Fig. 269.—Pentaphyllous or pentasepalous calyx of *Stellaria Holostea*; sepals entire.

Fig. 270.—Flower of the Rose, cut vertically. c t, Tube of the calyx. c f, Limb of calyx divided into leaflets. e e, Stamens. o o, Ovaries, each having a style which reaches beyond the tube of the calyx, and ending in a stigma, s.

Fig. 271.—Calyx of *Rumex uncatus*, composed of two verticils or whorls; the outer, c e, having short and entire divisions; the inner, c i, having larger divisions, which exhibit at the margin narrow hooked projections, and have at their base a granular swelling, g.

wards (*divergent* or *patulous*), or arched inwards (*connivent*). They are usually of a greenish colour, and are called *foliaceous* or *herbaceous*; but sometimes they are coloured, as in the Fuchsia, Tropæolum, and Pomegranate, and they are then called *petaloid*. Whatever be its colour, the external envelope of the flower must be considered as the calyx.

360. The nature of the hairs on the calyx gives rise to terms similar to those already mentioned as applied to the surfaces of plants (§ 60).



The vascular bundles sometimes form a prominent rib (figs. 272, 273), which indicates the middle of the sepal, at other times they form several (fig. 274). The venation is useful in pointing out the number of leaves which form a gamosepalous calyx. At the part where two sepals unite, there is occasionally a prominent line, formed by the union of the vessels of each

(fig. 274), which divides near the apex into two branches, each following the course of their respective sepals.

361. In a polysepalous calyx, the number of the parts is marked by Greek numerals prefixed. Thus, a *trisepalous* calyx has three sepals, *pentasepalous* or *pentaphyllous*, five, as in *Stellaria Holostea* (fig. 269), and so on. The sepals occasionally are of different forms and sizes. Thus, in *Aconite*, one of them has a peculiar helmet shape, and has been called *galeate* (*galea*, a helmet); some authors regard this as a petal, but it seems to belong to the outer whorl, and is consequently a part of the calyx. In *Calcophyllum*, one of the sepals enlarges after the corolla falls, and becomes of a fine pink colour.

362. In a monophyllous or gamosepalous calyx, the sepals adhere in various ways, sometimes very slightly, as in *Oenothera*; and their number is marked by the divisions at the apex. These divisions are either simple projections in the form of acute or obtuse teeth (fig. 273), or they extend about half way, as fissures, the calyx being *trifid* (three-cleft), *quinquefid* (five-cleft), as in *Primula elatior* (fig. 272), according to circumstances; or they reach to near the base in the form of partitions, the calyx being *tripartite*, *quadripartite*, *quinquepartite*, &c. The adhesion or union of the parts may be complete, and the calyx may be quite entire or *truncate*, as in some *Correas*, the venation being the chief indication of the different parts. The adhesion is sometimes irregular, some parts uniting to a greater extent than others, thus

Fig. 272.—Quinquefid or five-cleft calyx of *Primula elatior* or the oxlip.

Fig. 273.—Five-toothed calyx of *Silene inflata*.

Fig. 274.—Calyx, c, of *Hibiscus*, with its caliculus or epicalyx, b.

forming a two-lipped or *labiate* calyx, which, when the upper lip is arched, becomes *ringent*. The upper lip is often composed of three parts, which are thus posterior or next the axis, while the lower has two, which are anterior. The part formed by the union of the sepals is called the *tube* of the calyx; the upper portion where the sepals are free is the *limb*. Sometimes a monophyllous calyx assumes an angular or prismatic form, as in *Lamium* and *Primula*, and then the angles are marked by the midribs of the sepals which form it. Occasionally the calyx has a *globular* form, at other times it is *bell-shaped*, *funnel-shaped*, *turbinate* (like a top), or *inflated*.

363. Occasionally, certain parts of the sepals undergo marked enlargement. In the Violet, the calycine segments (*laciniæ*) are prolonged downwards beyond their insertions, and in the Indian Cress (*Tropæolum*) this prolongation is in the form of a spur (*calcar*), formed by three sepals (fig. 275 *e*); in Delphinium it is formed by one. When one or more sepals are thus enlarged, the calyx is *calcarate* or *spurred*. In the Pelargonium, the spur from one of the sepals is adherent to the flower-stalk.




364. In some plants, as the Mallow tribe, the flower appears to be provided with a double calyx, which has been denominated *caliculate*, the outer calyx being the *epicalyx*. In fig. 274, *c* represents the calyx of *Hibiscus*, and *b* the smaller calyx or epicalyx outside; and in fig. 276, the same thing is shown in *Potentilla verna*. Many authors look upon the outer calyx as a collection of whorled bractlets, or an involucre placed immediately below the flower. In some cases the projecting teeth between the divisions of the calyx, as in *Rosaceæ*, are to be traced to the transformed stipules of the calycine leaves. Degenerations take place in the calyx, so that it becomes dry, scaly, and glumaceous (like the glumes of grasses), as in the Rush tribe; hairy, as in *Compositæ*; and a mere rim, as in some *Umbelliferæ* and *Acanthaceæ*, when it is called *obsolete* or *marginate*.



365. In *Compositæ*, *Dipsacæ*, and *Valerianaceæ*, the tube of the calyx adheres to the pistil, and the limb is developed in the form of hairs, called *pappus*. This pappus is either simple (*pilose*) (fig. 278), or feathery (*plumose*) (fig. 279). In cases where, to the naked eye, the hairs appear to be simple, the application of a lens sometimes exhibits distinct tooth-like projections often irregularly scattered. In

Fig. 275.—Calcarate calyx of *Tropæolum* or Indian cress. *e*, Spur or calcar. *p*, Pedicel.
Fig. 276.—Calyx, *c*, of *Potentilla verna*, with its epicalyx or caliculus, *b*.

figs. 277, 278, and 279, there are examples of calyces, *c*, the tubes of which, *t*, are united to the pistil, while the limbs, *l*, show the transition from the narrowed thread-like form in *Catananche cærulea* (fig. 277), to the pilose in *Scabiosa atro-purpurea* (fig. 278), and thence to the plumose in *Pterocephalus palæstinus* (fig. 279). In *Valeriana*, the limb of the calyx at first seems to be an obsolete rim, but as the fruit ripens, it is shown to consist of hairs rolled inwards, which expand so as to waft the fruit.



366. The calyx sometimes falls off before the flower expands, as in Poppies, and is *caducous*; or along with the corolla, as in *Ranunculus*, and is *deciduous*; or it remains after flowering, as in *Labiatae*, *Scrophulariaceæ*, and *Boraginaceæ*; or its base only is persistent, as in *Datura Stramonium*. In *Eschscholtzia* and *Eucalyptus*, the parts of the calyx remain united at the upper part, and become disarticulated at the base or middle, so as to come off in the form of a lid or funnel. Such a calyx is *operculate* (*operculum*, a lid), or *calyptrate* (*καλύπτρα*, a covering). The existence or non-existence of an articulation determines the deciduous or persistent nature of the calyx. In the case of *Eschscholtzia*, the axis seems to be prolonged so as to form a sort of tube, from which the calyx separates. In *Eucalyptus*, the calyx consists of leaves, the laminae or petioles of which are articulated like those of the Orange, and the separation between the parts occurs at this articulation.

367. The tube of the calyx is sometimes united to the pistil, and enlarges subsequently, so as to form a part of the fruit, as in the Apple, Pear, Pomegranate, Gooseberry, &c. In these fruits the limb of the calyx is seen at the apex. Sometimes a persistent calyx increases much after flowering, without being incorporated with the fruit, becoming

Figs. 277–279.—Examples of calyces, the limbs of which, *l*, gradually pass into the state of hairs or pappus. *c*, Calyx, the tube of which, *t*, is united to the ovary, and forms a narrow column above it, in figs. 278, 279, the limb, *l*, consisting of numerous simple or feathery hairs.

i, Involucre or gamosepalous bracts cut vertically.

Fig. 277.—Calyx of *Catananche cærulea*.

Fig. 278.—Calyx of *Scabiosa atro-purpurea*.

Fig. 279.—Calyx of *Pterocephalus palæstinus*.

accrescent (*accresco*, to increase), as in various species of *Physalis* (fig. 280); at other times it remains in a withered or *marcescent* (*marcesco*, I decay) form; sometimes it becomes *inflated* or *vesicular*. In *Trifolium fragiferum*, the union of the inflated calyces causes the strawberry-like appearance of the head of flowers when in fruit.

368. **Corolla.**—The corolla is the more or less coloured inner floral envelope, forming the whorl of leaves between the calyx and the stamens. It is generally the most conspicuous whorl. The gay colours and fragrant odours of flowers are resident in it. It is present in the greater number of Exogens. It is composed of parts which are usually disposed in one or more verticillate rows, and which are called *petals* (πέταλον, a leaf). The petals sometimes form a continuous spiral with the calycine segments, but in general they are disposed in a circle, and alternate with the sepals.

369. Petals differ more from leaves than sepals do, and are much more nearly allied to the next whorl or the stamens. In some cases, however, they are transformed into leaves like the calyx, and occasionally leaf-buds are developed in their axil. They are seldom green, although occasionally this colour is met with, as in some *Cobæas*, *Hoya viridiflora*, *Gonolobus viridiflorus*, and *Pentatropis spiralis*. They are generally white, red, blue, or yellow, or exhibit some colour produced by an intermixture of these. The colouring matter is contained in cells, and differs in its nature from the chlorophylle of the leaves. As regards their structure, petals consist of cellular tissue traversed by true spiral vessels, and thin-walled tubes. In delicate flowers, as *Convolvulus* and *Anagallis*, these vessels are easily seen under the microscope. Petals do not usually present numerous layers of cells like the leaves, neither is the epidermis always distinct, although in some instances it may be detached, especially from the surface next the calyx. The cuticle of the petal of a *Pelargonium*, when viewed with a $\frac{1}{2}$ or $\frac{1}{4}$ inch object glass, shows beautiful hexagons, the boundaries of which are ornamented with several inflected loops in the sides of the cells.

370. On the outer surface of petals, corresponding to the lower side of leaves, stomata are sometimes found. Petals are generally glabrous or smooth; but, in some instances, hairs are produced on their surface. Petaline hairs, though sparse and scattered, present occasionally the same arrangement as those which occur on the leaves: thus, in




Fig. 280.—Accrescent calyx, c, connected with the fruit of *Physalis alkekengi*.

Bombaceæ they are stellate. Coloured hairs are seen on the petals of *Menyanthes*, and on the segments of the perianth of the *Iris*. Although petals are usually very thin and delicate in their texture, they occasionally become thick and fleshy, as in *Stapelia* and *Rafflesia*; or dry, as in *Heaths*; or hard and stiff, as in *Xylopia*. A petal often consists of two portions—the lower narrow, resembling the petiole of a leaf, and called the *unguis* or *claw*; the upper broader, like the blade of a leaf, and called the *lamina* or *limb*. These parts are seen in the petals of the *Pink* (fig. 281), where *o* is the claw, and *l* the limb. The claw is often wanting, as in the *Rose*, and the petals are then *sessile*. Those having a claw are *unguiculate*.

371. Petals, properly so called, belong to Exogenous plants, for in Endogens the flowers consist of a perianth or perigone, which is referred to the calycine envelope. Hence the venation of petals resembles that of exogenous leaves. In the claw the vessels are approximated, as in the petiole, and in the limb they expand. There may be a medium vein whence lateral veins go off, at the same or different heights, forming reticulations; or there may be several primary veins diverging from the base of the limb, and forming a sort of fan-shaped venation. At other times the median vein divides into two.

372. According to the development of veins, and the growth of cellular tissue, petals present varieties similar to those already noticed in the case of leaves. Thus the margin is either entire or divided into lobes or teeth. These teeth sometimes form a regular fringe round the margin, and the petal becomes *fimbriated* (*fimbriæ*, a fringe), as in the *Pink* (fig. 281); or *laciniated*, as in *Lychnis Flos cuculi*; or *crested*, as in *Polygala*. The median vein is occasionally prolonged beyond the summit of the petals in the form of a long process, as in *Strophanthus hispidus*, where it extends for seven inches; and at other times it ends in a free point or *cuspidis*, and thus becomes *cuspidate*; or the prolonged extremity is folded downwards or inflexed, as in the *Umbelliferae* (fig 282), so that the point approaches to the base. If the median vein divides into two, the space between the divisions may be filled up so as to leave only a



281



282



283

Fig. 281.—An unguiculate petal of *Dianthus monspessulanus*. *o*, Unguis or claw. *l*, Limb which is fimbriated, or has a fringed margin.

Fig. 282.—A petal of *Eryngium campestre*, with the apex inflexed or turned down towards the base.

Fig. 283.—A bipartite petal of *Alsine Media*, or common Chickweed. *l*, The limb split into two. *o*, The claw.

slight deficiency, and thus the petal becomes *emarginate*; or the deficiency may be greater, while the limb gradually expands from below upwards, and its extremity becomes two-lobed, so that the petal becomes *obcordate*. If the separation extends to the middle, it is *bifid*; if to near the base, *bipartite*, as in Chickweed (fig. 283 *l*). In the same way as in leaves, the venation of the petals is sometimes unequal, and the cellular tissue is developed more on one side than on the other, thus giving rise to an *oblique* petal.

373. According as the veins proceed in a straight or curved direction, so may the limb of the petal be flat or concave, or hollowed like a boat, *cymbiform* or *navicular* (*cymba*, a boat, and *navis*, a ship), or like a spoon, *cochleariform* (*cochleare*, a spoon). In the case of the navicular petal, the median vein forms a marked keel. In Hellebore, the petals become folded in a tubular form, resembling a horn; in Aconite (fig. 284), some of the petals, *p*, resemble a hollow curved horn, supported on a grooved stalk, while in Columbine (fig. 285), Violet, Snapdragon, and Centranthus, one or all of them are prolonged in the form of a spur, and are *calcarate* (*calcar*, a spur). In Valeriana, Antirrhinum, and Corydalis, the spur is very short, and the corolla or petal is said to be *gibbous* (*gibbus*, a bunch or swelling), or *saccate* at the base. In some Boraginaceæ (fig. 297), there are foldings at the upper part of the tube of the corolla, *r*, causing hollow projections, open on the outside, which might be considered as small internal spurs.

374. When a petal continues narrow, as if formed by a prolongation of the claw, it is called *linear*; when the limb is prolonged below, so as to form two rounded lobes, it is *cordate*, as in the petal of *Genista candicans* (fig. 286), and when the lobes are acute, it may be *sagittate* or *hastate*. The meaning of the epithets applied to the forms of petals, will be understood by considering those applied to leaves. In general,




Fig. 284.—Part of the flower of *Aconitum Napellus*, showing two irregular horn-like petals, *p*, supported on grooved stalks, *o*. These used to be called nectaries. *s*, The whorl of stamens inserted on the thalamus, and surrounding the pistil.

Fig. 285.—Single spurred petal of *Aquilegia vulgaris*, or common Columbine, formed by a folding of the margins.

Fig. 286.—Cordate or cordiform petal of *Genista candicans*. *o*, The claw. *l*, The limb.

it may be stated, that the terms refer to the limb of the petal, which is frequently the only portion developed. In the Poppy, the petals have a puckered or corrugated appearance, arising from their delicacy, and the mode in which they are folded in æstivation. Other petals have a *crisp* or *wavy* margin.

375. A corolla rarely consists of one petal, and when this occurs, as in *Amorpha*, it depends on the abortion or non-development of others. Such a corolla is *unipetalous* (*unus*, one), a term quite distinct from monopetalous (§ 376). In general, the corolla consists of several petals, equalling the sepals in number, or being some multiple of them. When this is the case, the floral envelopes are said to be *symmetrical*; when, however, by the abortion of some of the petals the numbers do not correspond, then the flower becomes *unsymmetrical*. When alluding to the general symmetry of the flower, the various changes produced by some petals being undeveloped will be considered. A corolla is *dipetalous*, *tripetalous*, *tetrapetalous*, or *pentapetalous*, according as it has two, three, four, or five separate petals.

376. The general name of *polypetalous* (πολύς, many), is given to corollas having separate petals, while *monopetalous* or *gamopetalous* (μόνος, one, and γάμος, union) is applied to those in which the petals are united. This union generally takes place at the base, and extends more or less towards the apex; in *Phyteuma*, the petals are united at their apices also. In some polypetalous corollas, as that of the Vine,

in which the petals are separate at the base, they adhere by their apices. That a monopetalous corolla consists of several petals united, is shown in such cases as *Phlox amœna*, some specimens of which have the petals more or less completely disunited, while others exhibit the normal form of coherent petals. When the petals are equal as regards their development and size, the corolla is *regular*; when unequal it is *irregular*. Even although the separate petals are oblique, still, if they are all equally so, as in many *Malvaceæ* with twisted æstivation, the corolla is regular. The size of the corolla as compared with the calyx, the number, direction, and form of its parts, and their relation to the axis of the plant, require attention.

377. When a corolla is monopetalous, it usually happens that the claws, or the lower parts of the petals, are united into a tube (figs. 287, t,



Fig. 287.—Regular monopetalous or gamopetalous tubular corolla of *Spigelia marylandica*. c, Calyx. t, Tube of the corolla. l, Limb of the corolla. s, Summit of style and stigma.
Fig. 288.—Irregular gamopetalous or monopetalous corolla of *Digitalis purpurea*, or Foxglove. c, Calyx. p, Corolla. t, Tube. l, Limb.

288 *t*), while the upper parts are either free or partially united, so as to form a common limb (fig. 287 *l*), the two portions being separated by the *faux* or *throat*, which often exhibits a distinct contraction or expansion. The number of parts forming such a corolla can be determined by the divisions, as by the number of teeth, crenations, fissures, or partitions; or if, as rarely happens, the corolla is entire, by the venation. The union may be equal among the parts, or some may unite more than others, leaving gaps between the united portions. Sometimes the tubular portion is bent, as in *Lycopsis*; at other times the limb is curved at its apex, as in *Lamium*.



289



290

378. **Regular Polypetalous Corollas.**—Among them may be noticed the *rosaceous* corolla, in which there are five spreading petals, having no claws, and arranged as in the single Rose (fig. 289) and *Potentilla*; the *caryophyllaceous* corolla, in which there are five petals with long narrow tapering claws, as in many of the Pink tribe (figs. 290, 281); the *alsinaceous*, where the claw is less narrow, and there are distinct spaces between the petals, as in some species of Chickweed; *cruciform*, having four petals, often unguiculate, placed opposite in the form of a cross, as seen in Wallflower (fig. 291), and other plants called *cruciferous* (*crux*, a cross, and *fero*, to bear).




291

Fig. 289.—Polypetalous flower of *Rosa rubiginosa*, or the Sweet-brier. *b*, Bract or floral leaf. *c* *t*, Tube of calyx, which forms the conspicuous part of what is commonly called the fruit. *c* *f*, *c* *f*, *c* *f*, *c* *f*, *c* *f*, Sepals or foliola of the calyx. *p* *p* *p* *p*, Petals without a claw. *e*, Stamens attached to the calyx.

Fig. 290.—Polypetalous flower of *Dianthus monspessulanus*. *b*, Bracts. *c*, Calyx. *p* *p*, Petals with their claws, *o*, approximated so as to form a tube.

Fig. 291.—Cruciferous flower of *Chieranthus Cheiri*, or Wallflower. *c*, Lobes of the sepals; the two external sepals being prolonged inferiorly, so as to form a sort of spur or swelling. *p* *p*, The four petals arranged like a cross. *e*, The four longer stamens, the summits of the anthers being visible.

379 Irregular Polypetalous Corollas.—The most marked of these is the *papilionaceous* (fig. 292), in which there are five petals; one superior (or posterior), turned next to the axis, usually larger than the rest, *e*, and folded over them in æstivation, called the *vexillum* or *standard*; two lateral, *a*, the *alæ* or wings; two inferior (or anterior), partially or completely covered by the *alæ*, and often united slightly by their lower margins, so as to form a single keel-like piece, *b*, called *carina*, or keel, which embraces the essential organs. This corolla occurs in the Leguminous plants of Britain, or those plants which have flowers like the pea. Among the irregular polypetalous corollas might be included the *orchidaceous*, although it is, properly speaking, the perianth of an Endogen. This perianth consists of three outer portions equivalent to the calyx, and three inner alternating with them, constituting the petals. The latter are often very irregular, some being spurred, others hooded, &c.; and there is always one, called the *labellum* or lip, which presents a remarkable development, and gives rise to many of the anomalous forms exhibited by these flowers.



380. Regular Monopetalous or Gamopetalous Corollas.—These are sometimes *campanulate* or *bell-shaped*, as in *Campanula rotundifolia*

Fig. 292.—Irregular polypetalous corolla in the papilionaceous flower of *Lathyrus odoratus*, or Sweet-pea. *c*, Calyx. *e*, Vexillum or standard. *a*, Two *alæ* or wings. *b*, Carina or keel, formed of two petals.


Fig. 293.—Regular monopetalous or gamopetalous campanulate or bell-shaped corolla of *Campanula rotundifolia*. *c*, Calyx. *l*, Limb of corolla. *s*, Stigma.

Fig. 294.—Regular monopetalous or gamopetalous infundibuliform corolla of *Nicotiana Tabacum*, or Tobacco. The letters as in fig. 293.

Fig. 295.—Regular monopetalous or gamopetalous hypocrateriform corolla of *Primula elatior*, or Oxlip. *c*, Calyx. *p*, Corolla. *t*, Tube. *l*, Limb. *a*, Anthers.

Fig. 296.—Regular gamopetalous tubular corolla of *Symphytum officinale*, or common Comfrey. *c*, Calyx. *l*, Tube of corolla. *l*, Limb. *s*, Stigma. *r*, External depressed surface of folds which project into the tube of the corolla.

(fig. 293); *infundibuliform* or *funnel-shaped*, when the tube is like an inverted cone, and the limb becomes more expanded, at the apex, as in Tobacco (fig. 294)); *hypocrateriform* or *salver-shaped*, when there is a straight tube surmounted by a flat spreading limb, as in Primula (fig. 295); *tubular*, having a long cylindrical tube, appearing continuous with the limb, as in Spigelia (fig. 287), and Comfrey (fig. 296); *rotate* or *wheel-shaped*, when the tube is very short, and the limb flat and spreading, as in Myosotis (fig. 297); when the divisions of the rotate corolla are very acute, as in Galium, it is sometimes called *stellate* or *star-like*; *urceolate* or *urn-shaped*, when there is scarcely any limb, and the tube is narrow at both ends, and expanded in the middle, as in Erica cinerea (fig. 298). Some of these forms may become irregular in consequence of certain parts being more developed than others. Thus, in Veronica, the rotate corolla has one division much smaller than the rest, and in Digitalis there is a slightly irregular campanulate corolla (fig. 288), which some have called *digitaliform*.



381. Irregular Monopetalous or Gamopetalous Corollas.—Among these may be remarked the *labiate* or *lipped* (fig. 299), having two divisions of the limb in the form of what are called *labia* or lips (from a fancied resemblance to a mouth), the upper one composed usually of two pieces, and the lower of three, separated by a *hiatus* or gap, *l*. In such cases the tube varies in length, and the parts of the calyx follow the reverse order in their union, two sepals being united in the lower lip, and three in the upper. When the upper lip of a labiate corolla is much arched, and the lips separated by a distinct gap, it is called *ringent* (*ringor*, to grin). The Labiate corolla

Fig. 297.—Regular gamopetalous rotate corolla of *Myosotis palustris*, or Forget-me-not. *c*, Calyx. *p*, Corolla. *r*, Folds of the corolla, forming projections at the upper part of the tube, which are opposite to the lobes of the corolla.

Fig. 298.—Regular gamopetalous urceolate or urn-shaped corolla of *Erica cinerea*, or cross-leaved Heath. Letters as in fig. 295.

Fig. 299.—Irregular gamopetalous labiate or lipped corolla of *Salvia pratensis*. *c*, Calyx. *t*, Tube of corolla. *l*, Limb, forming two lips, having a gap or hiatus between them. *s*, Summit of style.

Fig. 300.—Irregular gamopetalous personate or masked corolla of *Antirrhinum majus*, or Frog-mouth. *c*, Calyx. *t*, Tube of corolla, having a gibbosity or swelling, *a*, at its base. *l*, Limb of corolla. *g*, The faux or mouth closed by a projection of the lower lip, *p*.

characterizes the natural order Labiatae. In *Lobelia*, there is a Labiate corolla, the upper lip of which becomes convex superiorly, and is split to near the base. When the lower lip is pressed against the upper, so as to leave only a chink or *rictus* between them, the corolla is said to be *personate* or *masked* (*persona*, a mask), as in Frogmouth, Snapdragon, and some other Scrophulariaceae (fig. 300), and the projecting portion, *p*, of the lower lip is called the *palate*. In some corollas the two lips become hollowed out in a remarkable manner, as in *Calceolaria*, assuming a slipper-like appearance, similar to what occurs in the labellum of some Orchids, as *Cypripedium*. Such *calceolate* (*calceolus*, a slipper) corollas may be considered as consisting of two slipper-like lips.

382. When a tubular corolla is split in such a way as to form a strap-like process on one side with several tooth-like projections at its apex, it becomes *ligulate* (*ligula*, a little tongue), or *strap-shaped* (fig. 301). This corolla occurs in many composite plants, as in the florets of Dandelion, Daisy, and Chicory. The number of divisions at the apex indicates the number of united petals, some of which, however, may be abortive. Occasionally, some of the petals become more united than others, and then this corolla assumes a *bilabiate* or two-lipped form. In Compositae there are often two kinds of florets associated in the same head. Thus, in the Daisy, there are irregular ligulate white florets on the outside or in the ray, while there are regular tubular yellow florets in the centre or disc. In *Scævola* and in Honeysuckle, the corolla is split down to its base, so as to resemble somewhat the ligulate form.

383. **Nectaries and Anomalies in Petals.**—Certain abnormal appearances occur in the petals of some flowers, which received in former days the name of *nectaries*. The term nectary was very vaguely applied by Linnæus to any part of the flower which presented an unusual aspect, as the crown of *Narcissus*, the processes of the Passion-flower, &c. If the name is retained, it ought properly to include only those portions which secrete a honey-like matter, as the glandular depression at the base of the perianth of the Fritillary (fig 302 *r*), or on the petal of *Ranunculus*, or on the stamens of Rutaceae. Some say that in all flowers there is an apparatus for secreting honey connected with the essential organs of the flower, and in some way concerned in fertilization, and the nourishment of the



301

Fig. 301.—Irregular gamopetalous ligulate floret of *Catananche cærulea*. *c*, Calyx, with a quinquefid limb united inferiorly with the ovary, *o*. *e*, Stamens with united anthers, *a*, (*syngontherous* or *syngonous*), surrounding the style, *s*, with its bifid stigma.

young seeds. This opinion is particularly supported by Vaucher and Bravais. The sap of *Zea Mais* is said to contain much saccharine matter before flowering, which ultimately passes to the flower, and disappears from the rest of the plant. What have usually, however, been called Nectaries, are mere modifications of some part of the flower, produced either by degeneration, or by a process of *dilamination* (*dis*, separate, and *lamina*, a blade), or *chorization* (*χωρίζω*, I separate). This process, called *unlining* by Lindley, and *deduplication* by Henfrey, consists in the separation of a layer from the inner side of a petal, either presenting a peculiar form, or resembling the part from which it is derived. The parts thus produced are not alternate with the petals or the segments of the corolla, but opposite to them. In these cases, the petals at the lower part consist of one piece, but where the limb and claw separate, or where the tube ends, the vascular layer splits into two, and thus two laminæ are formed, posteriorly and anteriorly, one of which is generally less developed than the other. These dilaminated scales are well seen in *Lychnis* (fig. 303 *a*), *Silene*, *Cynoglossum*, and *Ranunculus*, and may be considered as formed in the same way as the ligule of grasses (§ 161). Corollas having these scaly appendages are sometimes denominated *appendiculate*. In other cases, as in *Cuscuta*, the scales are alternate with the petals, and are not traced to dilamination. This system of dilamination has been applied by the French botanists to all cases in which the parts of whorls become opposite in place of alternate. Lindley and others, however, believing that the law of alternation is the normal one, refer such cases in general to an abortion of a whorl, or to some peculiar arrestment in development, as will be shown under the section of Morphology and Symmetry.

384. In general, the parts called Nectaries, are to be considered as merely modifications of the corolla or stamens. Thus, the horn-like nectaries under the galeate sepal of *Aconite* (fig. 284 *p*), are modified petals, so also the tubular nectaries of *Hellebore*. The nectaries of *Menyanthes* and of *Iris*, consist of hairs developed on the petals. Those of *Parnassia* (fig. 304 *n*), and of the Passion-flower, *Stapelia*, *Asclepias*, and *Canna*, are fringes, rays, and processes, which are apparently



302



303

Fig. 302.—One of the segments, *s*, of the perianth of *Fritillaria imperialis*, or Crown Imperial, with a pit or depression, *r*, at its base, containing honey-like matter. The cavity is coloured differently from the rest of the segment, and it is often called a nectary, or a nectariferous gland.

Fig. 303.—Petal of *Lychnis fulgens*, seen on its inner side. *o*, Claw. *l*, Limb. *a*, An appendage formed by dilamination or chorization. This appendage has been called a nectary.

modifications of stamens, and some consider the crown of *Narcissus* as consisting of a membrane similar to that which unites the stamens in *Paneratium*. It is sometimes difficult to say whether these nectaries are to be referred to the row of the corolla or of the stamens. The paraphyses of the Passion-flower, the crown of *Narcissus*, and the coronet of *Stapelia*, are referred sometimes to the one and sometimes to the other. In general, they may be said to belong to that series with which they are immediately connected. Some have attempted to give names according to the parts of which they are modifications, by prefixing the term *para* (παρά, beside, or close to), and speaking of *paracorolla* and *parastemones*.

385. Petals are attached to the axis usually by a narrow base, but occasionally the base is larger than the limb, as in the Orange flower. When this attachment takes place by an articulation, the petals fall off either immediately after expansion (*caducous*), or after fertilization (*deciduous*). A corolla or petals which are continuous with the axis and not articulated, as *Campanula*, *Heaths*, &c., may be persistent, and remain in a withered or marcescent state while the fruit is forming. A gamopetalous corolla always falls off in one piece. Sometimes the base of the corolla remains persistent, as in *Rhinanthus* and *Orobanche*.

386. **Development of Floral Envelopes.**—The floral envelopes, when monosepalous and monopetalous, first appear in the form of a ring, whence various cellular projections arise, constituting the sepals and petals; when they are polysepalous and polypetalous, the ring is wanting. Even when the parts become ultimately unequal, as in *Digitalis* (fig. 288), they form equal cellular papillæ when first developed (fig. 305). Barneoud has shown this in the irregular *Ranunculaceæ*, *Violaceæ*, *Orchidaceæ*, *Labiataæ*, *Scrophulariaceæ*, *Leguminosæ*, and *Polygalaceæ*.

387. In *Begoniaceæ*, the floral envelope at first appears as a continuous ring, having five very equal small segments; some of these, especially in the male flowers, disappear entirely or become atrophied. All the observations of Barneoud confirm Decandolle's statement, that irregular flowers are to be referred to *regular* types, from which they seem to have degenerated. There appear to be three principal kinds of irregularity among corollas:—1. Irregularity by simple inequality of development of the several segments, often along with adhesion or atrophy, or arrest of growth:



Fig. 304.—Petal, *p*, of *Parnassia palustris*, or grass of Parnassus, with a nectary, *n*, which appears to be an abortive state of some of the stamens.

Fig. 305.—Bud of the Irregular gamopetalous flower of *Digitalis purpurea*. *c c*, Calyx. *p*, Corolla, which in its early development is regular. *e*, The stamens at first projecting beyond the corolla.

this is the most common kind. 2. Irregularity of deviation, when the segments, though equal, turn all to the same side, as in ligulate florets. 3. Irregularity by simple metamorphosis of stamina, as in *Canna*. The irregular corollas of *Acanthaceæ*, *Bignoniaceæ*, *Gesneraceæ*, *Lobeliaceæ*, and *Scrophulariaceæ*, are formed at first in a regular manner by equal projections from a sort of cup or ring. Even in *Calceolaria*, there is at first a scooped-out cup, with four regular and very minute teeth, which are ultimately developed as the corolla; the nascent calyx having also four divisions.

Inner Floral Whorls, or the Essential Organs of Reproduction.

388. These organs are the *stamens* and the *pistil*, the latter containing the seeds or germs of young plants, and corresponding to the female, while the former produces a powder necessary for fecundation, and is looked upon as performing the part of the male. The presence of both is required in order that perfect seed may be produced. A flower may have a calyx and corolla, and yet be imperfect if the essential organs are not present. The name of *hermaphrodite* is given to flowers in which both these organs are found; that of *unisexual* (one sex), or *diclinous* (δῖς, twice, and κλῖν, a bed), to those in which only one of these organs appears,—those bearing stamens only, being *stamiferous* (*stamen*, a stamen, and *fero*, I bear), or male; those having the pistil only, *pistilliferous* (*pistillum*, a pistil, and *fero*, I bear), or female.

389. The absence of one of the organs is due to *abortion* or *non-development*. When in the same plant there are unisexual flowers, both male and female, the plant is said to be *monoecious* (μόνος, one, and οἶκον, habitation), as in the Hazel and Castor oil plant; when the male and female flowers of a species are found on separate plants, the term *dioecious* (δῖς, twice) is applied, as in *Mercurialis* and Hemp; and when a species has male, female, and hermaphrodite flowers on the same or different plants, it is *polygamous* (πολύς, many, and γάμος, marriage). The term *agamous* (α, privative, and γάμος, marriage) has sometimes been applied to Cryptogamic plants, from the supposed absence of any bodies truly representing the stamens and pistil.

390. **Stamens.**—The stamens (*stamina*) arise from the *thalamus* or *torus* within the petals, with which they alternate, forming one or more verticils or whorls, which collectively constitute the *androecium* (ἀνδρ, a male, and οἶκον, habitation), or the male organs of the plant, as distinguished from the *gynæcium* (γυνή, a female, and οἶκον, habitation), or female organs of plant. Their normal position is below the inner whorl or the pistil, and when they are so placed (fig. 306 e), they are *hypogynous*; (ὑπο, under, γυνή, female or pistil). Sometimes they become united to the petals or *epipetalous* (ἐπὶ, upon, and πέταλον, a

leaf), and the insertion of both is looked upon as similar, so that they are still hypogynous, provided they are independent of the calyx and the pistil. In fig. 307, the stamens, *e*, and the petals, *p*, are both below the pistil or ovary, *o*, and separate both from it and the calyx, *c*, and are therefore *hypogynous*; when the stamens are inserted on the calyx, that is, become united to it to a greater or less height above the base of the pistil, then they become lateral as it were in regard to it, and are *perigynous*, ($\pi\epsilon\gamma\acute{\iota}$, around). This is shown in the flower of



almond (fig. 308), in which the petals, *p*, and the stamens, *e*, are united to the calyx *c*, while the pistil is free. When the union of the parts of the flower is such that the stamens are inserted upon the ovary, they are *epigynous* ($\epsilon\pi\acute{\iota}$, upon or above). In this case, the whorls are usually so incorporated, that the stamens appear also to come from the calyx. In the Orchis tribe, where the stamens adhere completely to the pistil, and when united extend above the ovary, they are said to be *gynandrous*. In *Aralia spinosa* (fig. 309), all the whorls,



Fig. 306.—Central part of the flower of *Liriodendron tulipifera*, the tulip-tree, composed of carpels, *c c*, which together form the pistil. They cover the upper part of the axis, *a*, and below them are inserted numerous stamens, some of which are seen, *e e*. These stamens are hypogynous and extrorse.

Fig. 307.—Section of a flower of *Geranium robertianum*. *c c*, Calyx. *p*, Petals. *e*, Stamens. Pistil composed of ovary, *o*, and style and stigmata, *s*. *t*, Torus or thalamus. The petals and stamens are hypogynous, and the latter are monadelphous.

Fig. 308.—Section of the flower of the Almond-tree. The letters indicate the same parts as in the last figure. The petals and stamens are perigynous. The pistil is free.

Fig. 309.—Section of the flower of *Aralia spinosa*. Letters as in last figure. The petals and stamens are epigynous, attached to a large disk, *d*, which covers the summit of the ovary. The ovary is adherent to the calyx, and has been laid open to show its loculaments and pendulous ovules.

observed to bear anthers, while at the same time their petaloid appearance seemed but little changed.

392. When there is only one whorl, the stamens are usually equal in number to the sepals or petals, and are arranged opposite to the former, and alternate with the latter. The flower is then *isostemonous* (ἴσος, equal, and στήμων, a stamen). When the stamens are not equal in number to the sepals or petals, the flower is *anisostemonous* (ἀνίσος, unequal). When there is more than one whorl of stamens, then the parts of each successive whorl are alternate with those preceding it. The staminal row is more liable to multiplication of parts than the outer whorls. If the stamens are double the sepals or petals as regards number, the flower is *diplostemonous* (διπλός, double); if more than double, *polystemonous* (πολύς, many). In general, when the stamens are normally developed, and are more numerous than the sepals and petals, they will be found arranged in several whorls, and their parts multiples of the floral envelopes. Thus, if a flower has five sepals, five petals, and twenty stamens, the latter are arranged in four alternate rows, having five in each. Although this is the usual law, yet various changes take place by abortion, arrestment of development, and other circumstances leading to abnormal growth. In this way the stamens may neither be equal to, nor a multiple of, the floral envelopes, and they may even be less numerous, so that the flower is *miostemonous* (μείων, less). In Cruciferous plants, while the petals and sepals are equal in number, (four,) and alternate in arrangement, the stamens are six in number, four long and two short, this imparity of numbers has been supposed to result from the splitting of the long stamens by lateral chorisis or separation, a presumption favoured by the fact, that partial union frequently exists between the apposed stamens, that teeth are found on but the one side of those long stamens, those sides which are free being adjoining, and that in many cruciferæ but four stamens exist.

393. In certain cases, as in *Primula*, the row of stamens is opposite to the petals forming the gamopetalous corolla. This opposition is by many looked upon as caused by the non-appearance of an outer row of stamens; by others it is considered as produced by chorization or separation of laminæ from the petals, which become altered so as to form stamens, a view which is thought to be confirmed by their development taking place before the petals; by a third party, each petal is looked upon when fully developed as formed by the halves of two contiguous petals, and thus the stamens are considered as being really alternate with the original petals.

394. When the stamens are under twenty, they are called *definite*, and the flower is *oligandrous* (ὀλίγος, few, and ἀνήρ, male or stamen); when above twenty, they are *indefinite* or *polyandrous* (πολύς, many), and are marked ∞. The number of stamens is indicated by the Greek

numerals prefixed to the term *androus*: a flower with one stamen being *monandrous* (μόνος, one); with two, *diandrous* (δίς, twice); with three, *triandrous* (τρεῖς, three); with four, *tetrandrous* (τετράς, four); with five, *pentandrous* (πέντε, five); with six, *hexandrous* (ἕξ, six); with seven, *heptandrous* (ἑπτὰ, seven); with eight, *octandrous* (ὀκτώ, eight); with nine, *enneandrous* (ἐννεά, nine); with ten, *decandrous* (δέκα, ten); with twelve, *dodecandrous* (δωδεκά, twelve). These terms will be referred to when treating of the Linnæan system of classification.

395. A stamen consists of two parts—a contracted portion, usually thread-like, equivalent to the petiole of the leaf, and termed the *filament* (*filum*, a thread); and a broader portion, representing the folded blade of the leaf, termed the *anther* (ἀνθηρός, belonging to a flower), which contains a powdery matter, called *pollen*. The filament is no more essential to the stamen than the petiole is to the leaf, or the claw to the petal. If the *anther* is absent, the stamen is abortive, and cannot perform its functions. The anther is developed before the filament, and when the latter is not produced the anther is sessile (*sessilis*, sitting), or has no stalk, as in the Mistletoe.

396. **The Filament**, when structurally considered, is found to consist of a thin epidermis, on which occasionally stomata and hairs occur, and of a layer of cellular tissue enclosing a bundle of spiral vessels, which traverses its whole length, and terminates at the union between the filament and the anther. The filaments of *Callitriche verna* are said to have no vessels. The filament is usually, as its name imports, filiform or thread-like, cylindrical, or slightly tapering towards its summit. It is often, however, thickened, compressed, and flattened in various ways. It sometimes assumes the appearance of a petal, or becomes *petaloid* (πέταλον, a leaf or petal, and ἔδος, form), as in *Canna*, *Maranta*, *Nymphæa alba* (fig. 310, 2); occasionally it is *subulate* (*subula*, an awl), or slightly broadened at the base, and drawn out into a point like an awl, as in *Butomus umbellatus*; and at other times it is *clavate* (*clava*, a club), or narrow below and broad above, like the club of Hercules, as in *Thalictrum*. In place of tapering, it happens, in some instances, as in *Tamarix gallica* (fig. 311), *Peganum Harmala*, and *Campanula*, that the base of the filament is dilated much, and ends suddenly in a narrow thread-like portion. In these cases, the base may represent the sheath or vagina of the petiole, and, like it, may give off stipulatory processes in a lateral direction. Sometimes the filament is forked, or divided at the apex into branches or teeth. In *Allium* there are three teeth, the central one of which bears the anther.



Fig. 311.—Three out of the ten stamens of *Tamarix gallica*, united together by the dilated bases of their filaments.

397. The filament varies much in length and in firmness. The length bears a relation to that of the pistil, and to the position of the flower, whether erect or drooping; the object being to bring the anther into more or less immediate contact with the upper part of the pistil, so as to allow the pollen to be scattered on it. The filament is usually of sufficient solidity to support the anther in an erect position; but sometimes, as in Grasses, *Littorella*, and *Plantago*, it is very delicate and capillary (*capillus*, a hair), or hair-like, so that the anther is pendulous. The filament is usually continuous from one end to the other, but in some cases it is bent or jointed, becoming *geniculate* (*genu*, a knee); at other times, as in the Pellitory, it is spiral. It is frequently colourless; but, in many instances, it exhibits different colours. In *Fuchsia* and *Poinciana*, it is red; in *Adamia* and *Tradescantia virginica*, blue; in *Oenothera* and *Ranunculus acris*, yellow.

398. Hairs, scales, teeth, or processes of different kinds are sometimes developed on the filament. In *Tradescantia virginica*, or Spiderwort, the hairs are beautifully coloured, and moniliform (*monile*, a necklace) or necklace-like. These hairs exhibit movements of rotation (¶ 278). Such a filament is bearded or stupose (*stupa*, tow). At the base of the filament, certain glandular or scaly appendages are occasionally produced, either on its internal or external surface. These may be either parts of a whorl, to be afterwards noticed under the name of the Disk, or separate prolongations from the filament itself. In fig. 313, *a* represents such a staminiferous appendage found on

the inner side of the base of the filament, *f*, which is hence called *appendiculate*, or sometimes *strumose* (*struma*, a swelling). The processes noticed in the Boraginaceæ as modified petals (fig. 312 *a*), may be considered external appendages of the filaments, the stamen being regarded as the lamina of a petal.

399. Filaments are usually articulated to the torus, and the stamen falls off after fertilization; but in *Campanula* and other plants, they are continuous with the torus, and the stamen remains persistent, although in a withered state. Certain changes are produced in the whorl of stamens by adhesion of the filaments to a greater or less extent,


while the anthers remain free; thus, all the filaments of the Androecium may unite, forming a tube round the pistil (fig. 307 *e*), or

Fig. 312.—Stamen of *Borago officinalis*. *f*, Appendiculate filament. *a*, Appendage prolonged in the form of a horn-like process. *l*, Lobes of the anther.

Fig. 313.—Stamen of *Zygophyllum fabago*. *f*, Filament, connected with a broad scaly appendage, *a*.



a central bundle when the pistil is abortive (fig. 314, 1), the stamens becoming *monadelphous* (μόνος, one, and ἀδελφός, brother), as occurs in *Geranium* (fig. 307), *Malva*, *Hibiscus*, and *Jatropha Curcas* (fig. 314, 1); or they may unite so as to form two bundles, the stamens being *diadelphous*, (δύς, twice), as in *Polygala*, *Fumaria*, and *Pea*; in this case the bundles may be equal or unequal. It frequently happens, especially in *Papilionaceous* flowers, that out of ten stamens, nine are united by their filaments, while one (the posterior one) is free. When filaments form three or more bundles, the stamens are *triadelphous* (τρεῖς, three), as in *Hypericum ægyptiacum* (fig. 315), or *polyadelphous*



(πολύς, many), as in *Luhea paniculata* (fig. 316, 1), or in *Ricinus communis* (fig. 317, 1). The union of the filaments takes place sometimes at the base only, as in *Tamarix gallica* (fig. 311); at other times it extends throughout their whole length, so that the bundles assume a columnar form. In certain cases, the cohesion extends to near the apex, forming what Mirbel calls an *androphore* (ἀνθή, male or stamen, and φέρω, I bear,), or a column which divides into terminal branches, each bearing an anther (fig. 315, *f e*). Occasionally some filaments are united higher up than others, and thus a kind of compound branching

Fig. 314.—Male or staminiferous flower (1), and female or pistilliferous flower (2), of *Jatropha Curcas*. *c*, Calyx. *p*, Corolla. *e*, Stamens united by filaments occupying the centre in flower 1, in consequence of the suppression of the pistil. *p*, Pistil in flower 2, composed of ovary, *o*, with three bifid styles at its summit. *a*, Small glandular appendages alternating with the divisions of the corolla. Above each of the flowers is a diagram representing the order in which the different parts of the flower are arranged. In diagram 1 are represented five parts of the calyx, five of the corolla, two rows of stamens, five in each. In diagram 2, the staminal rows are abortive, and there are three carpels, forming the pistil, in the centre.

Fig. 315.—Triadelphous stamens of *Hypericum ægyptiacum* surrounding the pistil, *o*, *f f*, United filaments forming columns. *e e*, Anthers, free. The outer envelope of the flower has been removed, the essential organs alone being left.

is produced (fig. 317, 2). In *Pancratium*, the filaments are united by a membrane, which may be considered as corresponding to the crown of *Narcissus*.



400. Filaments sometimes are united with the pistil, forming a *columna* or column, as in *Stylidium*, *Asclepiadaceæ*, *Rafflesia*, and *Orchidaceæ*. The column is called *gynostemium* (γυνή, pistil, and στήμων, stamen), and the flowers are denominated *gyandrous* (γυνή, pistil, and ἀνής, male or stamen).

401. In the case of certain Achlamydeous (§ 351) flowers, as *Euphorbia*, with only one stamen developed, there is the appearance of a jointed filament bearing one anther. This, however, is not a true filament, but a peduncle with a single stamen attached to it, as proved by the fact, that in some species of *Euphorbia* one or more verticils are produced at the joint. Thus the so-called anther is in reality a single flower supported on a stalk, all the parts being abortive, except a solitary stamen.


402. **The Anther** corresponds to the blade of the leaf, and consists of lobes or cavities containing minute powdery matter, called *pollen*, which, when mature, is discharged by a fissure or opening of some sort. The anther-lobes may be considered as formed by the two halves of the lamina, their back corresponding to the under surface, and their face to the upper surface, united by the midrib, the pollen being cellular tissue, and the fissure of the anther taking place at the margin, which, however, is often turned towards the face. In this view, the two cavities which are found to exist in each lobe, may correspond with the upper and under layer of cells, separated by a septum equivalent to the fibro-vascular layer of the leaf. Others view the

Fig. 316.—1. Flower of *Luhea paniculata*. cccc, Segments of calyx. pp, Petals. ee, Stamens grouped in bundles, which alternate with the petals. s, Stigma, composed of five parts, indicating the union of five carpels. 2. One of the staminal bundles magnified, showing all the filaments united into a single mass at the base, but separating superiorly. fa, The larger internal filaments, each ending in an anther. fs, The shorter outer ones, sterile and abortive.



Fig. 317.—1. Male flower of *Ricinus communis*, or Castor oil plant, consisting of a calyx, c, composed of five reflexed sepals, and of stamens, e, united by their filaments so as to form many bundles, thus being polyadelphous. 2. One of the staminal bundles, f, branching above, so as to leave the anthers free and separate.

anther as formed by each half of the lamina being folded upon itself, so that the outer surface of both face and back corresponds to the lower side of the leaf, and the septum dividing each cavity into two is formed by the upper surfaces of the folded half united.

403. There is a double covering of the anther—the outer, or *exothecium* (ἐξω, outwards, and ἔνχιον, a covering), resembles the epidermis, and often presents stomata and projections of different kinds (fig. 318 *c e*); the inner, or *endothecium* (ἐνδον, within), is formed by a layer or layers of fibro-cellular tissue (fig. 318 *c f*), the cells of which have a spiral (fig. 23), annular (fig. 24), or reticulated (fig. 25) fibre in their interior. This internal lining varies in thickness, generally becoming thinner towards the part where the anther opens, and there disappears entirely. The membrane of the cells is frequently absorbed, so that when the anther attains maturity the fibres are alone left, and these by their elasticity assist in discharging the pollen.



404. The anther is developed before the filament, and is always sessile in the first instance. In many examples it continues permanently so. It appears in the form of a small cellular projection, containing a mass of mucilaginous cells (fig. 319). In the progress of growth, certain grooves and markings appear on its surface, and its interior becomes hollowed out into two marked cavities, containing a mucilaginous matter (figs. 320, 321). In these cavities cells make their appearance—the outer small (figs. 320, 321, *c p*), forming ultimately the *endothecium* (fig. 318 *c f*); the interior layer forming cells in which the pollen is produced (figs. 320, 321, *u p*). As the




cavities become larger, the layer of cells (figs. 320, 321, *c i*), between

Fig. 318.—Transverse section of a portion of the covering of the anther of *Cobæa scandens* at the period of dehiscence. *ce*, Exothecium, or external layer, consisting of epidermal cells. *cf*, Endothecium, or inner layer, composed of spiral cells or inenchyma.

Fig. 319.—Transverse section of an anther of *Cucurbita Pepo*, or Gourd, taken from a bud about two millimetres, or 1-12th of an English inch, in length.

Fig. 320.—Similar horizontal section from a bud in a more advanced state. *ce*, Outer layer of cellules (*Exothecium*) forming the epidermis. *ci*, Intermediate layer of cellules in several layers, most of which are ultimately absorbed. *cp*, Internal layer of cells (*Endothecium*). *up*, Anther-cavities filled with large cells, which constitute the first state of the pollen-utricles.

the endothecium, *c p*, and exothecium, *c e*, is gradually absorbed more or less completely, forming at first septa in the cavities; and ultimately the anther assumes its mature form, consisting of two lobes with their membranous coverings (fig. 322, *l*).



321



322

405. In the young state there are usually four cavities produced, two for each anther-lobe, separated by the *connective*, and each divided by the septum, which sometimes remains permanently complete, and thus forms a *quadrilocular* (*quatuor*, four, and *loculus*, a pouch or box),

or *tetrathecal* (*τέτρας*, four, and *θήκη*, a sac) anther. The four cavities are sometimes placed in apposition, as in *Poranthera* (fig. 323) and *Tetratheca juncea* (fig. 324), and at other times two are placed above and two below, as in *Persea gratissima* (fig. 325 *l l*). In general, however, only two cavities remain in the anther, in consequence of the more or less complete removal of the septum, in which case the anther is said to be *bilocular* (*bis*, twice), or *dithecal* (*δύς*, twice), as seen in figs. 322, 326. Sometimes the anther has a single cavity, and becomes *unilocular* (*unus*, one), or



323



324



325

Fig. 321.—Similar section in a still more advanced state. The letters have the same meaning as in the last figure.

Fig. 322.—Anther of the Almond-tree. 'l', Seen in front. "!!", Seen behind. *f f*, Filament attached to the connective, *c*, by a point. *l l*, Anther-lobes containing pollen.

Fig. 323.—Quadrilocular anther, *l*, of *Poranthera*, attached to the filament, *f*, and opening at the summit by four pores, *p*.

Fig. 324.—Quadrilocular anther of *Tetratheca juncea*. 1. The anther entire, with its four loculements ending in one opening. 2. Anther cut transversely, showing the four loculements.

Fig. 325.—Anther of *Persea gratissima*, composed of four cavities or loculements, *l l*, united in pairs, one above the other, and opening each by a valve, *v*. At the base of the filament, *f*, are two glands, *g g*, which seem to be abortive stamens.

monotheal (μόνος, one), either by the disappearance of the partition between the two lobes, or by the abortion of one of its lobes, as in *Styphelia læta* (fig. 327), and *Althæa officinalis* (fig. 328). Occasionally, there are numerous cavities in the anther, as in *Viscum* and *Rafflesia*. The number of loculi or cavities is only seen when the anther opens.

406. The form of the anther-lobes varies. They are generally of a more or less oval or elliptical form (figs. 322, 329 *l*). Sometimes they are globular, as in *Mercurialis annua* (fig. 326); at other times linear or clavate (fig. 330), curved (fig. 331), flexuose, sinuose, or *anfractuose* (*anfractus*, winding), as in *Bryony* and *Gourd* (fig. 332). The lobes of the anther are sometimes in contact throughout their whole length (fig. 329), at other times they are separate (figs. 326, 333). In the former case their extremities may be rounded, forming a cordate anther (fig. 322), or the apex may be acute (figs. 312, 313); in the latter case the lobes may divide at the base only, and end in a sagittate or arrow-like manner (fig. 334 *l*); or at the apex, so as to be *bifurcate* or forked (fig. 335 *p*); or *quadrifurcate*, doubly forked (fig. 336 *l*); or at both base and apex, so as to be forked at each extremity, as in *Grasses* (fig. 337). The cavities of the anther are occasionally elongated so as to end in points (fig. 336 *l*). Sometimes the lower part of the antherine cavities is obliterated, and they degenerate into flattened appendages (fig. 338 *a*). It happens at times that the surface of the anther presents excrescences in the form of warts, awl-shaped pointed bodies (fig. 335 *a*), or crests (fig. 339 *a*).

407. That part of the anther to which the filament is attached, and which is generally towards the petals, is the *back*, the opposite being the *face*. The division between the lobes is marked on the face of the anther by a groove or *furrow*, and there is usually on the face, a *suture*, indicating the line where the membranous coverings open to discharge the pollen. The suture is often towards one side in consequence of the valves being unequal.

408. The anther-lobes are united either by a direct prolongation of




Fig. 326.—Pendulous anther lobes, *ll*, of *Mercurialis annua*, supported on the filament, *f*, and united by the connective, *c*.

Fig. 327.—Unilocular or monotheal anther of *Styphelia læta*, one of the *Epacridaceæ*, seen in front, *f*, and behind, *b*. *f*, Filament. *l*, Anther.

Fig. 328.—Unilocular anther of *Althæa officinalis*, or Marsh mallow. One of the lobes of the anther, *l*, abortive. *f*, Filament.

the filament, or more generally by a body called the *connective*, consisting of a mass of cellular tissue different from that contained in the filament. In this tissue the spiral vessels of the latter terminate. From the connective a partition or *septum* extends across each antherine



loculus, dividing it either partially or completely. The septum sometimes reaches the suture. When the filament is continuous with the connective, and is prolonged so that the anther-lobes appear to be united to it throughout their whole length, and lie in apposition and

Fig. 329.—Adnate or adherent anther of *Begonia manicata*, opening by longitudinal dehiscence. *l*, Anther-lobes. *f*, Filament.

Fig. 330.—Forked or bifurcate anther, *l*, of *Acalypha alopecuroides*, in the expanded flower.

Fig. 331.—Same anther in the bud, exhibiting a curved form.

Fig. 332.—Sinuous anther, *l*, of *Bryonia dioica*. *f*, Filament.

Fig. 333.—Anther of *Salvia officinalis*. *lf*, Fertile lobe full of pollen. *ls*, Barren lobe without pollen. *c*, Distractile connective.

Fig. 334.—Anther of *Nerium Oleander*, with its lobes, *ll*, sagittate at the base, and ending at the apex in a long feathery prolongation.

Fig. 335.—Anther, *l*, of *Vaccinium uliginosum*. *l*, Lobes ending in two pointed extremities, which open by pores. *a*, Appendages to the lobes.

Fig. 336.—Quadrifurcate anther of *Gualtheria procumbens*. *l*, Lobes ending in four points.


Fig. 337.—Versatile anther of *Poa compressa*. *f*, Filament. *l*, Lobes separating at each end.

Fig. 338.—Anther, *l*, of *Erica cinerea*. *f*, Filament. *r*, Lobes split partially downwards. *a*, Scale-like prolongations at the base.

Fig. 339.—Anther of *Pterandra pyroidea*. 1. Entire anther, seen laterally. 2. Lower half after having been cut transversely. *aaa*, Antherine appendages. *ll*, Anther-lobes. *cc*, Connective.

on either side of it, the anther is said to be *adnate* or *adherent* (fig. 329); when the filament ends at the base of the anther, then the latter is *innate* or *erect*. In these cases the anther is to a greater or less degree fixed. When, however, the attachment is very narrow, and an articulation exists, the anthers are then moveable, and easily turned by the wind. This is well seen in what are called *versatile* (*verto*, I turn) anthers, as in *Tritonia*, Grasses, &c. (figs. 260, 337), where the filament is attached only to the middle of the connective; and it may occur also in cases where it is attached to the apex, as in pendulous anthers (fig. 340).

409. The connective may unite the anther-lobes completely, or only partially. It is sometimes very short, and is reduced to a mere point, (fig. 326), so that the lobes are separate or free. At other times it is prolonged upwards beyond the lobes in the form of a point, as in *Acalypha* (fig. 331 c); or of a feathery awn, as in *Nerium*, *Oleander* (fig. 334); or of a conical or tongue-like process (figs 341, 342 c); or of a membranous expansion (fig. 343 c); or it is extended backwards



and downwards, in the form of a spur, as in fig. 343 a; or downwards, as in the case of the flaky appendage in *Ticorea febrifuga*. In *Salvia officinalis* (fig. 333), the connective is attached to the filament in a horizontal manner, so as to separate the two anther-lobes, and then it is called *distractile* (*dis*, separate, and *traho*, I draw). In *Stachys*, the connective is expanded laterally, so as to unite the bases of the antherine lobes, and bring them into a horizontal line.

410. The opening of the anthers to discharge their contents is denominated *dehiscence* (*dehisco*, I open). This takes place either by


Fig. 340.—Pendulous anther, *l*, of *Pyrola rotundifolia*. The anther is suspended from the summit of the filament, *f*, and opens at its apex by two pores, *p*.

Fig. 341.—Anther of *Humiria balsamifera*. *l l*, Anther lobes. *f*, Filament, ciliated or fringed with glandular teeth. *c*, Conical appendage, which seems to be a prolongation of the connective.

Fig. 342.—Anther of *Byrsonima bicorniculata*. *f*, Filament. *l*, Anther-lobes. The empty lobes at the summit are detached in the form of two small horn-like projections. *c*, A linguiform or tongue-like appendage prolonged from the connective.

Fig. 343.—Sessile anther of *Viola odorata*, or sweet violet. 1. Seen in front. 2. Seen behind. *l*, Anther-lobes. *a*, Spur-like appendage from the connective. *c*, Membranous expansion at the apex of anther-lobes.

clefts, by hinges, or by pores. When the anther-lobes are erect, the cleft takes place lengthwise along the line of the suture, constituting *longitudinal dehiscence* (figs. 322, 329, 342). At other times, the slit takes place in a horizontal manner, from the connective to the side, as in *Alchemilla arvensis*, and in *Lemna*, where the dehiscence is *transverse*. When the anther-lobes are rendered horizontal by the enlargement of the connective (figs. 328, 344 *a q*), then what is really longitudinal dehiscence may appear to be transverse. In other cases (fig. 344 *a g*), when the lobes are united at the base, the fissure in each of them may be continuous, and the two lobes may appear as one.



411. The cleft does not always proceed the whole length of the anther-lobe at once, but often for a time it extends only partially (figs. 343, 2, 338). In other instances the opening is confined to the base or apex, each loculament (*loculus*) opening by a single pore, as in *Pyrola* (fig. 340), *Vaccinium* (fig. 335), and *Solanum*, where there are two, and *Poranthera* (fig. 323), where there are four. In *Tetradlea juncea*, the four cavities (fig. 324, 2) open into a single pore at the apex (fig. 334, 1); and in the Mistletoe, the anther has numerous pores for the discharge of the pollen. Another mode of dehiscence is called *hinged*. In the Barberry, each lobe opens by a valve on the outer side of the suture, separately rolling up from base to apex; while in some of the Laurel tribe (fig. 325 *v*), there are two such separating valves for each lobe, or four in all. This may be called a combination of transverse and hinged dehiscence. In some *Guttiferæ*, as *Hebradendron cambogioides* (the Gamboge plant), the anther opens by a lid separating from the apex, or by what is called *circumscissile* (*circum*, around, and *scindo*, I cut) dehiscence. In the last-mentioned dehiscence, the anther may be considered as formed of jointed leaves like those of the Orange, the blades of which separate at the joint.

412. The anthers open at various periods of flowering; sometimes in the bud, but more commonly when the pistil is fully developed, and the flower is expanded. They either open simultaneously or in succession. In the latter case, individual stamens may move towards the pistil and discharge their contents, as in *Parnassia palustris*, or the outer or the inner stamens may open first, following thus a centripetal or centrifugal order. The anthers are called *introrse* (*introrsum*, inwardly), or *anticae* (*anticus*, the fore part), when they open on the

Fig. 344.—Corolla of *Digitalis purpurea*, cut in order to show the didynamous stamens (two long and two short) which are attached to it. *t*, Tube. *f*, Filaments which are united to the corolla at *t*, and run along its inner surface, having formed a marked adhesion. *a g*, Anthers of the long stamens. *a q*, Anthers of the short stamens.

surface next to the centre of the flower (fig. 345); they are *extrorse* (outwardly), or *posticæ* (*posticus*, behind), when they open on the outer surface; when they open on the sides, as in Iris, and some grasses, they are called *laterally* dehiscent (fig. 337). Sometimes anthers originally introrse, from their versatile nature, become extrorse, as in the Passion-flower and Oxalis. The attachment of the filament either on the outer or inner side, and the position of the anther in the young state, assist in determining the direction of the dehiscence when the anthers open by pores, or are versatile.

413. The usual colour of anthers is yellow, but they present a great variety in this respect. They are red in the Peach, dark purple in the Poppy and Tulip, orange in Eschscholtzia, &c. The colour and appearance of the anthers often change after they have discharged their functions.

414. Sometimes a flower consists of a single stamen, as already stated in regard to Euphorbia (§ 401). It is said also, that in the Coniferæ, as in the Fir, and in the Cycadaceæ, the stamens are to be regarded as single male flowers, supported on scales; being either a single stamen with bilocular anthers, as in Pinus, or unilocular, as in Abies, or several stamens united in an androphore, as in Taxus.

415. Stamens occasionally become sterile by the degeneration or non-development of the anthers, which, in consequence of containing pollen, are essential for fertilization; such stamens receive the name of *staminodia*, or rudimentary stamens. In Scrophularia (fig. 346), the fifth stamen, *s*, appears in the form of a scale; and in many Pentstemons it is reduced to a filament with hairs, or a shrivelled membrane at the apex. In other cases, as in double flowers, the stamens are converted into petals. In Persea gratissima (fig. 325), two glands, *g*, are produced at the base of the filament in the form of stamens, the anthers of which are abortive. Sometimes only one of the anther-lobes becomes abortive. In many unilocular anthers, the non-development of one lobe is indicated by the lateral production of a cellular mass resembling the connective. In Salvias, where the connective is distractile, one of the lobes only is perfect or fertile (fig. 333 *l f*), containing pollen, the other (fig. 333 *l s*) is imperfectly developed and sterile. In Canna, in place of one of the lobes, a petaloid appendage is produced.

Fig. 345.—Tetradynamous stamens (two long and two short) of Cheiranthus Cheiri. *p*, Top of the peduncle. *c*, Cicatrices left by the sepals of calyx which have been removed. *e g*, Two pairs of long stamens. *e p*, The short stamens. *t*, Torus or thalamus to which the stamens are attached.

Fig. 346.—Irregular corolla of Scrophularia with a staminodium, *s*, or abortive stamen in the form of a scale.



416. It has been already stated, that the term nectary has been sometimes applied to modified stamens presenting abnormal appearances. Thus, in *Parnassia palustris*, the so-called nectaries are clusters of abnormal stamens (fig. 304 *n*), united by a membrane at the base, and ending in glandular bodies like anthers. Staminodia were also called nectaries (fig. 346 *s*). When treating of the disk, other modifications of stamens will be considered.



417. The stamens, in place of being free and separate, may become united by their filaments (§ 399). They may also unite by their anthers, and become *syngenesious* or *synantherous* (σύν, together, and γενεαίς, origin, or ἀνθήρα, anther). This union occurs in Composite flowers, and in *Lobelia*, *Jasione*, *Viola*, &c.

418. Stamens vary in length as regards the corolla. Some are enclosed within the tube of the flower, as in *Cinchona*, and are called *included* (figs. 287, 288, 344); others are *exserted*, or extend beyond the flower, as in *Littorella*, *Plantago*, and *Exostemma*. Sometimes the stamens in the early state of the flower project beyond the petals, and in the progress of growth become included, as in *Geranium striatum* (fig. 347). Stamens also vary in their relative lengths as respects each other. When there is more than one row or whorl of stamens in a flower, those on the outside are sometimes longest, as in *Rosaceæ* (fig. 308); at other times those in the interior, as in *Luhea* (fig. 316, 2 *f a*). When the stamens are in two rows, those opposite the petals are usually shorter than those which alternate with the petals.

419. It sometimes happens that a single stamen is longer than all the rest. In some cases there exists a definite relation, as regards number, between the long and the short stamens. Thus, some flowers are *didynamous* (δύς, twice, and δύναμις, power or superiority), having only four out of five stamens developed, and the two corresponding to the upper part of the flower longer than the two lateral ones. This occurs in *Labiatae* and *Scrophulariaceæ* (figs. 344, 346). Again, in other cases there are six stamens, whereof four long ones are arranged in pairs opposite to each other, and alternate with two isolated short ones (fig. 345), and give rise to *tetradynamous* (τετράς, four, and δύναμις, power or superiority) flowers, as in *Cruciferae*.

420. Stamens, as regards their direction, may be erect, turned inwards, outwards, or to one side. In the last-mentioned case they are called *declinate* (*declino*, I bend to one side), as in *Amaryllis*, *Horsechestnut*, and *Fraxinella*.

421. **The Pollen.**—The pollen or powdery matter contained in the anther, consists of small cells developed in the interior of other cells.

Fig. 347.—Bud of polypetalous corolla of *Geranium striatum*, exhibiting the stamens, *ee*, at first longer than the petals, *pp*.

The cavities formed in the anther (fig. 321), are surrounded by a fibro-cellular envelope, *c p*, and within this are produced large cells, *u p*, containing a granular mass (fig 348, 1), which divides into four minute cells (fig. 348, 2), around which a membrane is developed, so that the original cell, or the parent *pollen-utricle*, becomes resolved by a merismatic division (¶ 24) into four parts (fig. 348, 3), each of which forms a granule of pollen. The four cells continue to increase (fig. 348, 4), distending the parent cell, and ultimately causing its absorption and disappearance. They then assume the form of perfect pollen grains, and either remain united in fours or multiples of four, as in some *Acacias*, *Periploca græca* (fig. 349), and *Inga anomala* (fig. 354), or separate into individual grains (fig. 348, 5), which by degrees become mature pollen, (figs. 348, 6, 351, 352). In *Acacia ringens*,



348



349

350



351

352

there are eight pollen-grains united; in *Acacia decipiens*, twelve; and in *Acacia linearis*, sixteen. Occasionally the membrane of the parent pollen-cell is not completely absorbed, and traces of it are detected in a viscous matter, surrounding the pollen-grains, as in *Onagrariaceæ*. In *Orchidaceous* plants, the pollen-grains are united into masses or

Fig. 348.—Development of the pollen of *Viscum album*, or the Mistletoe. 1. Two pollen-cells or pollinary utricles filled with granular matter. 2. Four nuclei produced in this matter. 3. Separation into four masses, each corresponding to a nucleus or a new utricule. 4. Pollenic or pollinary utricule containing three separate vesicles in its anterior. 5. Two of the latter, or the young pollen-grains, removed from the mother-cell or utricule. 6. The grains of pollen in their perfect state.

Fig. 349.—Pollen of *Periploca græca*, showing four grains agglutinated together.

Fig. 350.—Pollen of *Inga anomala*. The grains united in multiples of four.

Fig. 351.—Pollen-grain showing the extine covered with small punctuations.

Fig. 352.—Pollen-grain with the extine covered with granulations.

pollinia, by means of viscid matter. In *Asclepiadaceæ* (fig. 353), the pollinia, *p*, seems to have a special cellular covering, derived from a layer of reproductive pollen-cells, or from the endothecium. Pollinia in different plants vary from two to eight. Thus, there are usually two in *Orchis*, four in *Cattleya*, and eight in *Lælia*. The two pollinia in *Orchis Morio*, according to Amici, contain each about 200 secondary smaller masses. These small masses, when bruised, divide into grains which are united in fours. In Orchids, each of the pollen-masses has a prolongation or stalk, called a *caudicle* (*cauda*, a tail), which often adheres to a prolongation at the base of the anther, called *rostellum* (*rostellum*, a beak), by means of a viscid tenacious matter secreted by cells, and denominated *retinacula*, (*retinaculum*, a band or rein). Lindley considers the caudicle as derived from the stigma, and not from the pollinary tissue. The term *clinandrum* (*κλίνη*, a bed, and *ἄνθη*, a stamen), is sometimes applied to the part of the column in Orchids, where the stamens are situated.

422. When mature, the pollen-grain is a cellular body having an external covering, *extine* (*exto*, to stand out, or on the outside), and an internal, *intine* (*intus*, within). Fritzsche states that he has detected, in some cases, other two coverings, which he calls *intextine* and *exintine*. They occur between the extine and intine, and are probably formed by foldings of these membranes. In some aquatics, as *Zostera marina*, *Zannichellia pedunculata*, *Naias minor*, &c., only one covering exists, and that is said to be the intine. The extine is a firm membrane, which defines the figure of the pollen-grain, and gives colour to it. It is either smooth or covered with numerous projections, granules, points,




353



354



355



356

Fig. 353.—Flower of *Asclepias*, showing the pollinia or pollen-masses, *p*, attached to the stigma.
Fig. 354.—Pollen-grain of *Passiflora* before bursting. *o o o*, Opercula or lids formed by the extine, which open to allow the protrusion of the intine in the form of pollen-tubes.

Fig. 355.—Pollen-grain of *Cucurbita Pepo*, or Gourd, at the moment of its dehiscence or rupture. *o o*, Opercula or lids separated from the extine by the protrusion of the pollen-tubes, *t t*.

Fig. 356.—Pollen-grain of *Ipomœa*, with a reticulated extine.

minute hairs, or crested reticulations (fig. 356). The colour is generally yellow, and the surface is often covered with a viscid or oily matter. The intine is uniform in different kinds of pollen, thin and transparent, and possesses great power of extension. It is said to be the first envelope formed, the other being subsequently deposited while enclosed in the parent cell.


423. Within these coverings a granular semifluid matter called *fovilla*, is contained, along with some oily particles, and occasionally starch. The fovilla contains small spherical granules, sometimes the $\frac{1}{300000}$ of an inch in diameter (fig. 357), and larger ellipsoidal or elongated corpuscles (fig. 358), which are said to exhibit movements under the microscope similar to those seen in some Infusoria, and in some Algæ, to be afterwards noticed. These movements generally cease long before maturation, except in *Zostera marina* and some other plants.

424. Pollen-grains vary from $\frac{1}{3000}$ to $\frac{1}{7000}$ of an inch or less in diameter. Their form is much diversified. The most common form is ellipsoidal (figs. 358, 359), more or less narrow at the extremities, which are called its poles, in contradistinction to a line at *e*, equidistant from either extremity, and which is its equator. In figs. 359, 360, 1 and 2, the two surfaces of the pollen-grains of *Allium fistulosum* and *Convolvulus tricolor* are represented with their poles, *p*, their equator, *e*, and the longitudinal folds in their membrane; while at 3,




357

358



359



360

are shown transverse sections at the equators, with a single fold in one case, and three folds in the other. Pollen-grains are also of a spherical, triangular, trigonal (fig. 362), or polyhedral figure (fig. 364). In the latter case, when there are markings on their surface, those at the

Fig. 357.—Pollen-grain of *Amygdalus nana*, the intine or internal membrane of which is protruding at three pores under the form of as many ampullæ or sacs, *ttt*. One of these is open at the extremity, and from it is discharged the fovilla, *f*, composed of variously-sized granules.

Fig. 358.—Large granules of fovilla of *Hibiscus palustris*.

Fig. 359.—Pollen of *Allium fistulosum*. *p*, Pole. *e*, Equator. 1. Pollen-grain seen on the face. 2. On the opposite side or back. 3. Transverse section through its equatorial line.

Fig. 360.—Pollen of *Convolvulus tricolor*. The letters and numbers have the same signification as in fig. 359.

poles, *p*, sometimes differ from those at the equator, *e*. In *Tradescantia virginica*, the pollen is cylindrical, and becomes curved; it is polyhedral in *Dipsacæ* and *Compositæ*; nearly triangular in *Proteacæ* and *Onagrariacæ*. The surface of the pollen-grain is either uniform and



homogeneous, or it is marked by folds dipping in towards the centre, and formed by thinnings of the membrane. In Endogenous plants there is usually a single fold (fig. 359); in Exogens, often three (fig. 360). Two, four, six, and even twelve folds are also met with.

425. There are also pores or rounded portions of the membrane visible in the pollen-grain. These vary in number from one to fifty.



367

In Endogens, as in Grasses, there is often only one (fig. 365); while in Exogens, they number from three upwards. When numerous, the pores are either scattered irregularly (fig. 366), or in a regular order, frequently forming a circle round the equatorial surface (fig. 361). Sometimes at the place where the pores exist, the outer membrane, in place of being thin and transparent, is separated in the form of a lid, thus becoming *operculate* (*operculum*, a lid), as in the Passion-flower (fig. 354) and Gourd (fig. 355). Grains of pollen have sometimes both folds and pores. There may be a single pore in each fold, either in the middle (fig. 367), or at the extremities; or folds with pores may

alternate with others without pores; or finally, the pores and folds may be separate.

Fig. 361.—Grain of pollen of *Cannabis sativa*, or common Hemp. *e*, Equator. *pp*, Poles.

Fig. 362.—Pollen-grain of *Oenothera biennis* entire, with three angles, where tubes are produced.

Fig. 363.—The same, with one of its angles giving origin to a pollen-tube, which is formed by the intine. When the tube protrudes, the extine is ruptured.

Fig. 364.—Polyhedral pollen-grain of *Cichorium Intybus*, or Chicory.

Fig. 365.—Pollen-grain of *Daelylis glomerata*, or Cocks-foot grass.

Fig. 366.—Pollen-grain of *Fumaria capreolata*.

Fig. 367.—Grain of pollen of *Lythrum Salicaria*, showing six folds, three of which are perforated by a pore in their middle, and three alternating with them have no pores. *pp*, Poles. *e e*, Equator. 1. The grain in a dry state. 2. The grain swollen in water, so as to take a globular form and display its folds. The intine or internal membrane begins to protrude through the pores.

426. The form of the pollen-grains is much altered by the application of moisture. Thus, in fig. 367, 1, the pollen grain of *Lythrum Salicaria*, when dry, has an ellipsoidal form, but when swollen by the application of water, it assumes a globular form (fig. 367, 2). This change of form is due to endosmose, and depends on the fovilla being denser than the water. If the grains are retained in water, the distension becomes so great as to rupture the extine irregularly if it is homogeneous, or to cause projections and final rupture at the folds or pores when they exist. The intine, from its distensibility, is not so liable to rupture, and it is often forced through the ruptured extine, or through the pores, in the form of small sac-like projections (figs. 367, 2, 362). This effect is produced more fully by adding a little nitric acid to the water. The internal membrane ultimately gives way, and allows the granular fovilla to escape (fig. 357 *f*). If the fluid is applied only to one side of the pollen-grain, as when the pollen is applied to the pistil, the distension goes on more slowly, and the intine is prolonged outwards like a hernia, and forms an elongated tube called a *pollen-tube* (fig. 363). This tube, at its base, is often covered by the ruptured extine, and probably also by some of the coverings mentioned by Fritzsche as intervening between it and the intine. It contains in its interior fovilla-granules, and its functions will be particularly noticed under fertilization. The number of pollen-tubes which may be produced depends on the number of pores. In some pollinia, the number of tubes which are found is enormous. Thus, Amici calculates that the two pollen-masses of *Orchis Morio* may give out 120,000 tubes.

427. In **Cryptogamic Plants** there are certain organs which are supposed by some to be equivalent to stamens. On that account they were denominated by Hedwig *antheridia*, by others *pollinaria*. They consist of closed sacs of different forms, rounded, ovate, oblong, clavate, flask-like, &c., developed in different parts of the plants, containing a number of corpuscles immersed in a mucilaginous fluid, which at a certain period of growth are discharged through an opening at the surface. Sometimes the antheridium is a simple cell, at other times it is composed of a number of cells, as in *Hypnum triquetrum* (fig. 368, 1). It either appears on the surface of the plant, or is concealed within its tissue. Antheridia are sometimes confined to particular parts of the plant, at other times they are more



Fig. 368.—1. Antheridium or pollinarium, *a*, of a moss called *Hypnum triquetrum*, at the moment when its apex is rupturing to discharge the contents, *f*. 2. Four utricle of the contents containing each a phytozoon or moving corpuscle rolled up in a circular manner. 3. Single phytozoon separated.

generally diffused. Their contents are small utricles or cellules, varying, like pollen-grains, in the different orders of cryptogamic plants, and enclosing in place of fovilla, peculiar bodies called *phytozoa* (φύτον, a plant, and ζῷον, an animal) (fig. 368, 2), which are often rolled up in a circular or spiral manner, as in *Hepaticæ* and *Mosses* (fig. 368, 3). These exhibit active movements at certain periods of their existence, and resemble in this respect animalcules. In *Chara vulgaris* (fig. 369), the antheridium or *globule*, as it is called, contains cells, 1, from which proceed numerous *septate* (*septum*, a division) tubes, *t*. In each of the divisions of these tubes, 2, there is a phytozoon which escapes in a spiral form, leaving the division empty, 3, and ultimately becomes unrolled, 4,



369

exhibiting two vibratile cilia (*cilium*, an eyelash,) to which the movements are referred.

428. **The Disk.**—The term *disk* is applied to whatever intervenes between the stamens and the pistil, and is one of those organs to which the name of *nectary* was applied by old authors. It presents great varieties of form, such as scales, glands, hairs, petaloid appendages, &c., and in the progress of growth it often contains saccharine matter, thus becoming truly nectariferous. The degeneration and transformation of the stamens frequently form the disk. It may consist of processes rising from the torus, alternating with the stamens, and thus representing an abortive whorl; or it may be opposite to the stamens, and then formed by chorization (§ 383), as in *Crasula rubens* (fig. 258 a). In some flowers,



370

as *Jatropha Curcas*, in which the stamens are not developed, their place is occupied by glandular bodies forming the disk (fig. 314, 2, a).

Fig. 369.—1. Portion of antheridium or globule of *Chara vulgaris*. Several septate or partitioned tubes, *t*, attached to a utricle or vesicle. A mass of similar utricles, forming the bases of a large number of tubes, fills the cavity of the antheridium. 2. Extremity of one of these tubes, composed of several cellules, in each of which is a phytozoon. One of the phytozoa is represented half detached from the cellule. 3. Extremity of a tube from which the phytozoa have escaped, with the exception of the terminal cellule. 4. One of the phytozoa separated.

Fig. 370.—Disk, *d*, of *Pæonia Moutan*, or Tree Pæony, covering the ovary, and interposed between the whorl of stamens, *s*, and the pistil, *p*.

In Gesneraceæ and Cruciferæ the disk consists of tooth-like scales at the base of the stamens (fig. 345, *t*). The parts forming the disk sometimes unite and form a glandular ring, as in the Orange; or a dark-red lamina covering the pistil, as in *Pæonia Moutan* (fig. 370, *d*); or a waxy lining of the calyx tube, as in the Rose (fig. 270 *c t*); or a swelling at the top of the ovary, as in Umbelliferæ.

429. **The Pistil.**—The pistil occupies the centre or axis of the flower, and is surrounded by the stamens and floral envelopes, when these are present. It constitutes the innermost whorl, and is the female organ of the plant, which after flowering is changed into the fruit, and contains the seeds. It sometimes receives the name of *gynæcium* (γυνή, pistil, and οἶκος, habitation). It consists essentially of two parts, the *ovary* or *germen*, containing ovules or young seeds, and the *stigma*, a cellular secreting body, which is either seated immediately on the ovary, and is then called *sessile*, as in the Tulip and Poppy, (fig. 409), or is elevated on a stalk called the *style*, interposed between the ovary and stigma. The style is not necessary for the perfection of the pistil. Sometimes it becomes blended with other parts, as with the filaments of the anthers in the column of Orchidaceæ.

430. Like the other organs, the pistil consists of one or more modified leaves, which in this instance are called *carpels* (καρπός, fruit). The analogy of carpels to leaves may be deduced from their similarity in texture, and in venation, from the presence of stomata, hairs, and glands; from their resemblance to leaves in their nascent state; from their occasional conversion into true leaves, as in *Lathyrus latifolius*; and from the ovules corresponding in situation to the germs or buds found in some leaves, as those of *Bryophyllum calycinum*. When a pistil consists of a single carpel it is *simple*, a state usually depending on the non-development of other carpels; when it is composed of several carpels, more or less united, it is *compound*. In the first-mentioned case, the terms carpel and pistil are synonymous. Each carpel has its own ovary, style (when present), and stigma, and is formed by a folded leaf, the upper surface of which is turned inwards towards the axis, and the lower outwards; while at its margins are developed one or more buds called *ovules*. That this is the true nature of the pistil may be seen by examining the flower of the double-flowering Cherry. In it no fruit is produced, and the pistil consists usually of sessile leaves (fig. 371), the limb of each being green and folded, with a narrow prolongation upwards, *s*, as if from the midrib, *n*, and



Fig. 371.—Carpellary leaf of the double-flowering Cherry. In this plant the pistil is composed distinctly of one or more leaves folded inwards. *l*, Lamina or blade of the leaf or carpel. *s*, Prolongation of the midrib, *n*, representing the style, and ending in a circular thickened portion equivalent to the stigma.

ending in a thickened portion. When the single-flowering Cherry is examined, it is found that, in place of folded leaves, there is a single body (figs. 372, 373), the lower part of which is enlarged, forming the ovary, *o*, and containing a single ovule, *g*, attached to its walls, with a bundle of vessels, *fn*, entering it, a cylindrical prolongation, *t*, forming the style, and a terminal expansion, *s*, the stigma. It will be seen that in this case two carpellary leaves have become succulent and have united together, so as to form a compound pistil, with a single cavity containing one young seed.

431. The *Ovary* then represents the limb or lamina of the leaf, and is composed of cellular tissue with fibro-vascular bundles, and an epidermal covering. The cellular tissue, or parenchyma, often becomes much developed, as will be seen particularly when fleshy fruits are considered. The outer epidermis corresponds to the


lower side of the leaf, exhibiting stomata, and sometimes hairs; the inner surface represents the upper side of the leaf, being usually very delicate and pale, and forming a layer called sometimes *epithelium* (*ἐπί*, upon, and *θήλυς*, tender), which does not exhibit stomata. The vascular bundles correspond with the veins of the leaf, and consist of spiral, annular, and other vessels.

432. The *Style* has usually a cylindrical form, consists of cellular and vascular tissue, and when carefully examined is found to be traversed by a narrow canal (fig. 373 *c*), in which there are some loose projecting cells (figs. 374, 375), forming what is called the conducting tissue. A transverse section of the style of Crown Imperial (fig. 374), shows three vascular bundles, *v v v*, corresponding to three styles which are united into one, and *p*, loose cells in the canal of the style. This canal is bounded by cellular tissue (fig. 375 *c c*), traversed by spiral vessels, *v v*, and in its interior, besides the loose cells, *p p*, there are, especially at the period of fecundation, elongated tubes, *f f*, which in part fill up the canal. The name, *conducting tissue*, is given to that found in the canal of the style, on account of the part which it plays in conveying the influence of the pollen to the ovules, as will be explained under fertilization. Lindley has shown that in some instances the style seems to derive its origin from the placenta. The presence of the style is by no means essential to the perfection of the pistil. It varies in its shape and position, being usually apicular, but from altera-

Fig. 372.—Pistil or carpel of the single-flowering Cherry in its normal state. *o*, Ovary. *t*, Style. *s*, Stigma.

Fig. 373.—The same cut vertically, to show the central cavity of the ovary, *o*, with the ovule, *g*, suspended from its wall at a point where a bundle of nourishing vessels, *fn*, terminates. *t*, Style traversed by a canal, *c*, which runs from the stigma, *s*, to the cavity of the ovary.

tion in the direction of the central axis, it occasionally seems to be lateral. Its form and appearance also varies; under ordinary circumstances it is rounded in shape, but occasionally becomes flattened. In the *Clematis* it is furnished with downy hair, in the *Euphorbia* it becomes forked.



374



375

433. The *Stigma* is a continuation of the cellular tissue in the centre of the style, and it may be either terminal, when the canal opens at the top only (figs. 373 s, 376, 1), or lateral, when the splitting of the canal takes place on one side (fig. 377 s), or on both sides (fig. 378 s s). The stigma sometimes extends along the whole length of the style. In other instances the style is absent, when the stigma is said to be *sessile*. In Orchidaceous plants, it is placed on a part of the column called the *gynizus* (γυνίς, pistil, and ἰζώ, I sit). It is composed of cellular tissue more or less lax, and often having projecting cellules in the form of papillæ (fig. 376, 2), or of hairs (figs. 379, 3, 410 s), and at the period of




Fig. 374.—Transverse section of the style of *Fritillaria imperialis*, or Crown Imperial. The style is composed of three united together. *v v v*, Three vascular bundles, each corresponding to one of the three styles. *p*, Papillæ or cellular bodies projecting into the cavity of the canal.

Fig. 375.—Structure of the canal in the centre of the style of a *Campanula*. *c c*, Cellular tissue forming its parietes traversed by trachæ, *v*. *p p*, Various formed cells, displaced as it were, and along with other elongated and filamentous ones, *f f*, obstructing the canal.

Fig. 376.—1. Stigma, *s*, of *Daphne Laureola*, terminating the style, *t*. *o*, Summit of the ovary. 2. A small portion of the surface of the stigma, much magnified to show its papillæ.

Fig. 377.—Unilateral stigma, *s*, of *Asimina triloba*. *t*, Style.

Fig. 378.—Bilateral stigma, *s s*, of *Plantago saxatilis*. *o*, Ovary. *t*, Style.

fertilization exuding a viscous fluid, which retains the grains of pollen, and causes the protrusion of tubes.

434. A pistil is usually formed by more than one carpel. The carpels may be arranged like leaves, either at the same or nearly the same height in a verticil (figs. 380, 381), or at different heights in a spiral cycle (fig 306 c). When they remain separate and distinct, thus showing at once the composition of the pistil, as in *Caltha*, *Ranunculus*, *Hellebore*, and *Butomus* (fig. 381), the term *apocarpus* ($\alpha\pi\acute{o}\varsigma$, separate, and $\kappa\alpha\rho\acute{o}\varsigma$, fruit) is applied. * Thus, in *Crassula rubens* (fig. 258), the pistil consists of five verticillate carpels, *o*, alternating with the stamens, *e*; and the same arrangement is seen in *Zanthoxylon fraxineum* (fig. 380). In the Tulip-tree (fig. 306), the separate carpels, *c c*, are numerous, and arranged in a spiral cycle, upon an elongated axis or receptacle. In the Raspberry, the carpels are on a conical receptacle; in the Strawberry, on a swollen succulent one; and in the Rose (fig. 270 *o o*), on a concave one, *r r*, covered by the tube of the calyx, *c t*.



435. When the fruit consists of several rows of carpels on a flat receptacle, the innermost have their margins directed to the centre, while those of the outer rows are arranged on the back of the inner ones; if the receptacle is convex, the outer carpels are lowest, as in the Strawberry; if concave, the outer ones are uppermost, as in the Rose. At other times the carpels are united, as in the Pear, *Arbutus*, and Chickweed, so that the pistil becomes *syncarpous* ($\sigma\upsilon\nu$, together or

Fig. 379.—1. Summit of the style, *t*, of *Hibiscus palustris*, dividing into five branches, which are each terminated by a stigma, *s s*. 2. One of these branches highly magnified. 3. Portion of the surface of the stigma still more magnified to show its papillæ, which are elongated like hairs.

Fig. 380.—Pistil of *Zanthoxylon fraxineum*, consisting of five distinct carpels, supported on a gynophore, *g*. Each of the ovaries, *o*, bears a terminal style dilated at its extremity into a stigma, *s*. The five stigmata remain for a long time adherent by their sides.

Fig. 381.—1. Carpels of *Butomus umbellatus*, consisting of folded leaves arranged in different verticils. 2. Section of the same, showing the alternation of the parts of the flower. Three outer leaves of the perianth, *o'*, alternating with three inner ones, *p i*, three rows of stamens, *e o* and *e t*, and the carpels, *c e* and *c t*.

united). In *Dictamnus Fraxinella* (fig. 382), five carpels unite to form a compound pistil. In *Scilla italica* (fig. 259), the three carpels form only one apparently; but on examination it will be found that the pistil consists of three carpels alternating with the three inner stamens. The union, however, is not always complete; it may take place by the ovaries alone, while the styles and stigmata remain free, the pistil being then *gamogastrous* ($\gamma\acute{\alpha}\mu\omicron\varsigma$, union, and $\gamma\alpha\sigma\tau\eta\rho$, ovary); and in this case, when the ovaries form apparently a single body, this organ receives the name of *compound ovary*; or the union may take place by the ovaries and styles, while the stigmata are disunited; or by the stigmata and the summit of the style only (fig. 380). Various intermediate states exist, such as partial union of the ovaries, as in the Rue, where they coalesce at their base; and partial union of the styles, as in *Malvaceæ* (fig. 383). The union is usually most complete at the base; but in *Labiata* the styles are united throughout their length, and in *Apocynaceæ* and *Asclepiadaceæ*, the stigmata only.

436. When the union is incomplete, the number of the parts of a compound pistil may be determined by the number of styles and stigmata (fig. 383 *s*); when complete, the external venation, the grooves on the surface, and the internal divisions of the ovary, indicate the number. When the grooves between the carpels are deep, the ovary is denominated lobed, being one, two, three, four, or five-lobed, according to circumstances. In fig. 383, the nine carpels forming the ovary, *o*, are divided by grooves; and in fig. 384, a transverse section of the ovary of *Fuchsia coccinea*, shows the four carpels which form it. The changes which take place in the pistil by adhesion, degeneration, and abortion, are frequently so great as to obscure its composition, and to lead to anomalies in the




Fig. 382.—Portion of the pistil of *Dictamnus Fraxinella*. Two of the five carpels have been removed in order to show how the styles, *s*, produced on the inner side of the carpels, and at first distinct, approximate and become united into one. *o*, Ovaries, two of which in front show their dorsal surface, *d*, and their lateral surface, *l*. At the base of the gynophore, *g*, are seen the cicatrices, *c*, marking the insertion of the calyx, the petals, and the stamens.

Fig. 383.—Pistil of *Malva Alcea*. *o*, Nine ovaries, united so as to form one. *t*, Column formed by nine styles united to near their summit, where they diverge and separate. Each of the divisions of the style is terminated by a stigma, *s*.

Fig. 384.—Horizontal section of the four-celled (*quadrilocular* or *tetrathecal*) ovary of *Fuchsia coccinea*. *c c c c*, Wall of the ovary, which is formed by four carpellary leaves. *a*, Quadrangular axis to which the carpels are united. *o*, Ovules attached to the inner margin of the carpels.

alternation of parts. The pistil is more liable to changes of this kind than any other part of the flower.



437. The carpels are usually sessile leaves, but sometimes they are petiolate, and then are elevated above the external whorls. This elevation of the pistil may in general, however, be traced to an elongation of the axis itself, in such a way that the carpels, in place of being dispersed over it, arise only from its summit. A monstrosity often occurs in the Rose (fig. 385), by which the axis is prolonged, and bears the carpels, *f*, in the form of alternate leaves. Thus, by the union of the petioles of the carpels, or by lengthening of the axis, the pistil becomes *stipitate* (*stipes*, a trunk), or supported, as in the Passion-flower, on a stalk (figs. 380, 382 *g*), called a *gynophore* (γυνή, pistil, and φορέω, I bear), or *thecaphore* (θήκη, a case). Sometimes the axis is produced beyond the ovaries, and the styles become united to it, as in Geraniaceæ and Umbelliferæ. In this case the prolongation is called a *carpophore* (καρπός, fruit, and φορέω, I bear).


Placenta
438. The ovules are developed on the inner side of the carpel where the two edges of the carpellary leaves unite, and they are connected to it by vascular bundles which proceed from below upwards, traverse the carpel, and send a branch to each of the ovules. At the same place there is a development of cellular tissue in connection with the conducting tissue of the style and with the stigma. By the union of these tissues is formed the *placenta*, or projection to which the ovules are attached. Some restrict the term placenta to the point of attachment of a single ovule, and call the union of placentas, bearing several ovules, *placentaries* or *pistillary cords*. The part of the carpel where the placenta is formed, is the *inner* or *ventral suture*, corresponding to the margin of the folded carpellary leaf, while the *outer* or *dorsal suture* corresponds to the midrib of the carpellary leaf. The placenta is hence sometimes called *marginal*. The placenta is formed on each margin of the carpel, and hence is essentially double. This is seen in cases where the margins of the carpel do not unite, but remain separate, and consequently two placentas are formed in place of one. In fig. 386, the two carpels are folded, so that their margins meet, and the placenta is apparently single; whereas in fig. 387 the margins of each carpel do not meet, and the placenta of each is double. Again, in fig. 388, the two carpels, after meeting in the centre or axis, *a*, are

Fig. 385.—Section of monstrous Rose, as figured at section 324, the axis of which is prolonged beyond the flower, and the envelopes removed to show the abortive stamens, *r*. The carpels are attached alternately along the axis in the form of leaves. *p*, Abortive floral envelopes. *s*, Stamens in imperfect flower at the apex.


reflected outwards towards the dorsal suture, *s d*, and their margins separate slightly, each being placental and bearing ovules, *o*.



386



387



388

439. When the pistil is formed by one carpel, the inner margins unite in the axis, and form usually a common marginal placenta. This placenta may extend along the whole margin of the ovary as far as the base of the style, or it may be confined to the base or apex only. When the pistil is composed of several separate carpels, or, in other words, is apocarpous, there are generally separate placentas at each of their margins. In a syncarpous pistil, on the other hand, the carpels are so united that the edges of each of the contiguous ones by their union form a *septum* (*septum*, a fence or enclosure), or *dissepiment* (*dissepio*, I separate), and the number of these septa consequently indicates the number of carpels in the compound pistil. It is obvious then that each dissepiment is formed by a double wall or two laminæ; that the presence of a septum implies the presence of more than one carpel; and that, when carpels are placed side by side, true dissepiments must be vertical, and not horizontal.

440. When the dissepiments extend to the centre or axis, the ovary is divided into cavities, *cells* or *loculaments* (*loculus*, a box), and it may be *bilocular*, *trilocular*, *quadrilocular*, *quinquelocular*, or *multilocular*, according as it is formed by two, three, four, five, or many carpels, each corresponding to a single cell or loculament (fig. 381, 2, *c e, c i*). In these cases the marginal placentas meet in the axis, and unite so as to form a single *central* one (fig. 386 *a*). Some call this placentation *axile* (belonging to the *axis*), but this term is perhaps properly restricted to cases where the placenta is an actual prolongation of the axis. The number of loculaments is equal to that of the dissepiments. In fig. 384, there is shown a transverse section of the ovary of *Fuchsia coccinea*, *c c c c* being its parietes formed by the union of four carpellary leaves, *a* the axis united to the parietes by dissepiments, and *o* the ovules attached to the placentas at the margin of each carpel. When the carpels in a syncarpous pistil do not fold inwards completely so as to meet in the centre, but only partially, so that the dissepiments appear as projections on the walls of the ovary, then the ovary is *unilocular* (fig. 387), and the placentas are *parietal* (*paries*, a wall).

Figs. 386, 387, 388.—Horizontal sections of ovaries, composed of two carpellary leaves, the edges of which are folded so as to meet in the axis, *a*, in fig. 387; are reflected inwards into the loculaments after meeting in the axis in fig. 388; and do not reach the axis in fig. 387.

A horizontal section of the ovary of *Erythræa Centaurium* (fig. 389), exhibits a unilocular ovary with parietal placentas, *p*, formed at each of the margins of the carpels which do not meet in the centre. In these instances the placentas may be formed at the margin of the united contiguous leaves, so as to appear single, or the margins may not be united, each developing a placenta. From this it will be seen that dissepiments are opposite to placentas, formed by the union of the margins of two contiguous carpels, but alternate with those formed by the margins of the same carpel.

441. The carpellary leaves may fold inwards very slightly, or they may be applied in a valvate manner, merely touching at their margins, the placentas then being parietal, and appearing as lines or thickenings along the walls. In fig. 390, the pistil of *Viola tricolor* is represented, 1, cut vertically, and, 2, cut transversely, the ovules being attached to the walls of the ovary, and the placentas, *p*, being merely thickened portions of the walls. Cases occur, however, in which the placentas are not connected with the walls of the ovary, and form what is called a *free central placenta*. This is seen in many of the Caryo-



390



391



392

phyllaceæ. Thus, in *Cerastium hirsutum* (figs. 391, 392), the ovary, *o*, is composed of five carpels, indicated by the styles, *s*, but there is only one loculament, the placenta, *p*, being free in the centre, and the ovules, *g*, attached to it.

Fig. 389.—Horizontal section of the ovary of *Erythræa Centaurium*. *c*, Wall or paries of the ovary or carpellary leaf. *p*, The edge on which the placenta is formed, bearing the ovules, *o*. *l*, Cell or loculament.

Fig. 390.—Pistil of *Viola tricolor*, or Pansy, cut vertically to show the ovules, *o*, attached to the parietes. Two rows of ovules are seen, one in front, and the other in profile. *p*, A thickened line on the walls forming the placenta. *c*, Calyx. *d*, Ovary. 1. Hooded stigma terminating the short style. 2. Horizontal section of the same. *p*, Placenta. *o*, Ovules. *s*, Suture.


Fig. 391.—Pistil of *Cerastium hirsutum* cut vertically. *o*, Unilocular or monotheical ovary. *p*, Free central placenta. *g*, Ovules. *s*, Styles.

Fig. 392.—The same cut horizontally, and the halves separated so as to show the interior of the cavity of the ovary, *o*, with the free central placenta, *p*, covered with ovules, *g*.

442. In Caryophyllaceæ, however, while the placenta is free in the centre, there are often traces found at the base of the ovary of the remains of septa, as if rupture had taken place; and, in rare instances, ovules are found on the margins. But examples occur of this kind of placentation, as in Primulaceæ, Myrsinaceæ, Santalaceæ, and Theophrasteæ, in which no vestiges of septa or marginal ovules can be perceived at any period of growth. Duchartre states that the free placenta of Primulaceæ is totally different from that of Caryophyllaceæ. It is always free, and rises in the centre of the ovary, and the part uncovered by ovules gradually extends into the style. It is not first continuous with the style, and then free; neither is it originally marginal, and then free; but it is, according to him, wholly throughout its *organogeny* (ὄργανον, organ, and γένεσις, production or development) separate and axile.

443. This placentation, therefore, has been accounted for in two ways, either by supposing that the placentas in the early state were formed on the margins of carpellary leaves, and that in the progress of development these leaves separated from them, leaving the placentas and ovules free in the centre; or by supposing that the placentas are not *marginal* but *axile* formations, produced by an elongation of the axis, the ovules being lateral buds, and the carpels verticillate leaves, united together around the axis. The latter view has been supported by many botanists, and is confirmed by the fact, that in some cases the placenta is actually prolonged beyond the carpels. The first of these views would apply well to Caryophyllaceæ, the second to Primulaceæ. In the latter case, the only way of explaining the appearance on the marginal hypothesis, will be by considering the placentas as formed from the carpels by a process of chorization (§ 383), and united together in the centre.

444. Some indeed, as Schleiden and Endlicher, consider the axile view of placentation as applicable to all cases, the axis in some cases remaining free and independent, at other times sending prolongations along the margins of the carpellary leaves, and thus forming the marginal placentas. The occurrence of placentas over the whole inner surface of the carpels or of the dissepiments, as in *Nymphæa* and in *Butomus umbellatus* (figs. 393, 394); also, though very rarely, along the dorsal suture, as in *Cabomba*, or on lines within the margin, as in *Orobanche*, has been supposed to confirm this view. Schleiden argues in favour of it, from the case of *Armeria*, where there



Figs. 393, 394.—One of the carpels of *Butomus umbellatus*, or flowering Rush, cut transversely in 393, and longitudinally in 394. *l*, Loculament or cavity of the carpel. *a*, Ovules. *s*, Stigmata.

are five carpels and a single ovule attached to a cord, which arises from the axis, and becomes curved at the apex so as to suspend the ovule; also, from cases, such as *Taxus*, where the ovule appears to be naked and terminates a branch.

445. This theory of placentation, however, cannot be easily applied to all cases; and Gray says that it is disproved in cases of monstrosity, in which the anther is changed into a carpel, or where one part of the anther is thus transformed and bears ovules, while the other, as well as the filament, remain unchanged. In the case of *Luffa fœtida*, the entangled fibres of the carpellary leaves, even in the young state, seem to be connected with perpendicular lines forming the placenta. Brongniart mentions a case where the marginal placenta was entire, while the axis was prolonged separately, and totally unconnected with the placenta; he also notices peculiar monstrosities, which seem to prove that, in some cases at least, marginal placentation must take place.

446. Upon the whole, then, it appears that marginal, or, as it is often called, carpellary placentation, generally prevails; that axile placentation explains easily cases such as *Primulaceæ*, while such instances as *Caryophyllaceæ* are explicable on either view.

447. Occasionally, divisions take place in ovaries which are not formed by the edges of contiguous carpels. These are called *spurious dissepiments*. They are often horizontal, and are then called *phragmata* (φραγμα, a separation), as in *Cathartocarpus Fistula* (fig. 395), where they consist of transverse cellular prolongations from the walls of the ovary, only developed after fertilization, and therefore more properly noticed under fruit. At other times they are vertical, as in *Datura*, where the ovary, in place of being two-celled, is thus rendered four-celled; in *Cruciferae*, where the prolongation of the placentas forms a *replum* (*replum*, leaf of a door) or partition; in *Astragalus* and *Thespesia*, where the dorsal suture is folded inwards; and in *Diplophractum*, where the inner margin of the carpels is reflexed (fig. 388).

In *Cucurbitaceæ*, divisions are formed in the ovary, apparently by peculiar projections sent inwards from curved parietal placentas. In some cases, horizontal dissepiments are supposed to be formed by the union of carpels situated at different heights, so that the base of one becomes united to the apex of another. In such cases, the divisions are true dissepiments formed by carpellary leaves. The anomalous divisions in the ovary of the Pomegranate have been thus explained.

448. The ovary is usually of a more or less spherical or curved form, sometimes smooth and uniform on its surface, at other times hairy



395

and grooved. The grooves, especially when deep, indicate the divisions between the carpels, and correspond to the dissepiments. The dorsal suture may be marked by a slight projection, or by a superficial groove.

449. The ovary is either free in the centre of the flower, or it is united to the surrounding parts, more especially to the calyx. The union may take place completely, so that the calyx is *adherent* through-



out, and becomes *superior*, while the ovary is *inferior*, as in the Melon (fig. 396, *o* being the ovary, *l* the upper part of the adherent calyx); or it may take place partially, as in Saxifragaceæ (figs. 397, 398), where

Fig. 396.—Flower of *Cucumis Melo*, or Melon. *o*, Inferior ovary covered by the adherent calyx. *l*, Limb of the calyx appearing above the ovary. *p*, Corolla.

Fig. 397.—Flower of *Saxifraga Geum*, cut vertically to show the ovary, *o*, adherent for half its height to the calyx. *c*, The calyx, which is called half-superior. *p*, Petals. *e*, Stamens. *s*, Styles and stigmas.

Fig. 398.—Pistil of *Hoteia japonica*, one of the Saxifragaceæ, cut vertically in order to show the interior of its two cavities or loculaments. It is a bilocular or dithecal ovary. *o*, Two ovaries consolidated into one, and adherent for half their height to the calyx, *c*. *t*, Styles. *s*, Stigmas. *p*, Placentas covered with ovules. *p e*, Base of the petals.

Fig. 399.—Flower of *Fuchsia coccinea* divided horizontally into two halves through the middle of the ovary, *o*. The lower half, 2, of the ovary has been left untouched, to show its four cavities or loculi, with the ovules attached to their internal angles. Fig. 384 shows the same section more highly magnified. The upper half, 1, has been cut vertically, to show the ovules, *g*, arranged in a row in each loculament. The calyx, incorporated with the ovary below, is prolonged above it in the form of a tube, *t*, and divides at its summit into four segments, *l l*. *p*, Petals inserted on the tube of the calyx at the place where it divides into segments. *e*, Stamens inserted also on the tube, alternately large and small. The style rising from the summit of the ovary, and terminated by an ovoid stigma, *s*.

the ovary, *o*, becomes *half-inferior*, the calyx being *half-superior*. These adhesions between the calyx and the ovary will be found to be of importance, as determining the *epigynous* and *perigynous* (ἐπί, upon or above, and περί, around, and γυνή, pistil) condition of the stamens. Cases of adhesion between the ovary and the calyx, as occur in the Apple, Pear, Gooseberry, and Fuchsia, (fig. 399), must not be confounded with cases such as the Rose (fig. 270), where the tube of the calyx becomes enlarged and hollowed so as to cover the carpels. In the former case a transverse section (fig. 399) shows one or several closed loculements containing ovules; while in the latter, it exhibits one cavity open at the top, and separate carpels scattered over the surface, each having a style and stigma.

450. Peculiar views have been advocated by Schleiden, who considers the ovary in some cases as not formed by carpels, but by a hollowing out of the axis, at other times by these two modes combined. Thus the superior ovary, according to him, is formed of carpellary leaves, while the inferior ovary of the Apple and Pomegranate is composed of the expanded summit of the axis, bearing the carpels in its interior; that of *Epilobium* is formed from the stem alone, and that of *Saxifrage* partly by the peduncle and partly by carpels.

451. **The Style** proceeds from the summit of the carpel, and may be looked upon as a prolongation of it in an upward direction (fig. 372 *t*). It is hence called *apicalar* (*apex*, top). It consists not merely of the midrib, but of the vascular and cellular tissue of the carpel, along with a continuation of the placenta or conducting tissue, which ends in the stigma. In some cases, the carpellary leaf is folded from above downwards, in a hooded manner, so that its apex (as in reclinate veneration, fig. 205 *a*) approaches more or less to the base. When the folding is slight, the style becomes *lateral* (fig. 382); when to a greater extent, the style appears to arise from near the base, as in the Strawberry (fig. 400), or from the base, as in *Chrysobalanus Icaco* (fig. 401), when it is *basilar*. In all these cases the style



still indicates the organic apex of the ovary, although it may not be the apparent apex.

452. The carpel sometimes becomes imbedded in the torus or thalamus, so as to have a projection of the latter on one side; and then, if the style is basilar or lateral, it may adhere to this portion of the torus, and appear to arise from it. This is seen in *Labiatae* (fig. 402), and *Boraginaceae* (fig. 403), where the four carpels, *o*, are sunk in the

Fig. 400.—Carpel of Strawberry. *o*, Ovary. *t*, Style arising from near the base, and becoming basilar by the mode in which the ovary is developed; the style, however, still indicating the organic apex of the ovary.

Fig. 401.—Carpel of *Chrysobalanus Icaco*. *o*, Ovary. *t*, Basilar style. *s*, Stigma.

torus, *r*, in such a way that the common style, *s*, formed by the union of four basilar styles, seems to be actually a prolongation of the torus. When ovaries are thus attached round a central prolongation of the torus, continuous with a united columnar style, the arrangement is called a *gynobase* (γυνή, pistil, and βᾶσις, base). It is well developed in *Ochnaceæ*. In *Geraniums* there is a carpophore or prolongation of the torus in the form of a long beak, to which the styles are attached.



402



403

453. The form of the style is usually cylindrical, more or less filiform and simple; sometimes it is grooved on one side, at other times it is flat, thick, angular, compressed, and even petaloid, as in *Iris* and *Canna*. In *Goodeniaceæ* it ends in a cup-like expansion enclosing the stigma. It may be smooth and covered with glands and hairs. These hairs occasionally aid in scattering the pollen, and are called *collecting hairs*, as in *Goldfussia* or *Ruellia*. In *Campanula* they appear double and retractile. In *Aster* and other *Compositæ* (fig. 404), there are hairs produced on parts of the style, *p c*, prolonged beyond the stigma, *s*; these hairs, while the part is being developed, come into contact with the pollen and carry it up along with them. In *Vicia* and *Lobelia*, the hairs form often a tuft below the stigma.



404

454. The styles of a syncarpous pistil may be either separate or united; when separate, they alternate with the septa. When united completely, it is usual to call the style *simple* (fig. 399); when the union is partial, then the style is said to be *bifid*, *trifid*, *multifid*, according as it is two-cleft, three-cleft, many-cleft; or, to speak more correctly, according to the mode and extent of the union of two, three, or many styles. The style is said to be *bipartite*, *tripartite*, or *multipartite*, when the union of two, three, or many styles only extends a short way above the apex of the ovary. The style from a single carpel, or from each carpel of a compound pistil, may also be divided. In

Fig. 402.—Pistil of *Lamium album*, shown by a vertical section of part of the flower. Two of the four ovaries have been removed to exhibit the connection of the style with the torus, *r*, by adhesion. *o*, The two remaining ovaries. *d*, Glandular disk placed below the pistil. *c*, Part of calyx. *p*, Corolla.

Fig. 403.—Pistil of *Erithricium Jacquemontianum*, with one of the ovaries removed in front, to show the manner in which the ovaries are inserted obliquely on a pyramidal torus, *r*, whence the style appears to arise, ending in a stigma, *s*.

Fig. 404.—Summit of the style, *t*, of an *Aster*, separating into two branches, *s*, each terminated by an inverted cone of collecting hairs, *p c*. The stigma, *s*, is seen below in the form of a band or line on the inner curvature of the branches.

fig. 314, 2, each division of the tricarpellary ovary of *Jatropha Curcas* has a *bifurcate* or forked style, *s*, and in fig. 405, the ovary of *Emblica officinalis* has three styles, each of which is divided twice in a bifurcate manner, exhibiting thus a dichotomous division.



455. The length of the style is determined by the relation which ought to subsist between the position of the stigma and that of the anthers, so as to allow the proper application of the pollen. In some cases the ovary passes directly into the style, as in *Digitalis*, in other instances there is a marked transition from one to the other. The style may remain *persistent*, or it may fall off after fertilization is accomplished, and thus be *deciduous*.


456. **The Stigma** is the termination of the conducting tissue of the style, and is usually in direct communication with the placenta. It may, therefore, in most instances, be considered as the placentiferous portion of the carpel prolonged upwards. In *Armeria*, and some other plants, this connection with the placenta cannot be traced. Its position may be either terminal or lateral. The latter is seen in some cases, as *Asimina triloba*, where it is *unilateral* (fig. 377), and in *Plantago saxatilis* (fig. 378), where it is *bilateral*. Occasionally, as in *Tasmannia*, it is prolonged along the whole inner surface of the style. In *Iris*, it is situated on a cleft on the back of the petaloid divisions of the style. Some stigmata, as those of the *Mimulæ*, present sensitive flattened hands which close when touched. The stigma consists of loose cellular tissue, and secretes a viscid matter which detains the pollen, and causes it to protrude tubes. This secreting portion is, strictly speaking, the true stigma, but the name is generally applied to all the divisions of the style on which the stigmatic apparatus is situated, as in *Labiatae*. It usually alternates with the dissepiments of a syncarpous pistil, or corresponds with the cells; but in some cases, it would appear that half the stigma of one carpel unites with half that of the contiguous carpel, and thus the stigma is opposite the dissepiments, or alternates with the cells. This appears to be the case in the Poppy, where the stigma of a single carpel is two-lobed, and the lobes are opposite the septa.

457. If the stigma is viewed as essentially a prolongation of the placenta, then there is no necessary alternation between it and the placenta, both being formed by the margins of carpellary leaves, which

Fig. 405.—Female flower of *Emblica officinalis*, one of the Euphorbiaceæ. *c*, Calyx. *p.p.* Petals. *z*, Membranous tube surrounding the ovary. *o*, Ovary, crowned by three styles, *s*, each being twice bifurcate.

in the one case are ovuliferous, in the other stigmatiferous. There is often a notch in one side of a stigma (as in some Rosaceæ), indicating apparently that it is a double organ like the placenta. To the division of a compound stigma the terms *bifid*, *trifid*, &c., are applied, according to the number of the divisions. Thus, in Labiatæ (fig. 299), and in Compositæ (figs. 301, 404 s), the stigma is bifid; in Polemonium, trifid. When the divisions are large, they are called *lobes*, and when flattened like bands, *lamellæ*; so that stigmas may be *bilobate*, *trilobate*, *bilamellar*, *trilamellar*, &c.

458. It has already been stated that the divisions of the stigma mark the number of carpels which are united together. Thus, in Campanula (fig. 405 bis), the quinquefid or five-cleft stigma indicates five carpels, the stigmata of which are separate, although the other parts are united. In Bignoniaceæ (fig. 406), as well as in Scrophulariaceæ and Acanthaceæ, the two-lobed or bilamellar stigma indicates a bilocular ovary. Sometimes, however, as in the case of the styles, the stigma of a single carpel may divide. It is probable that in many instances, what is called bifurcation of the style is only the division of the stigma. In Gramineæ and Compositæ (figs. 301, 404), there is a bifid stigma and only one cavity in the ovary. This, however, may be probably traced to subsequent abortion of the ovary of one of the carpels. The stigma presents various forms. It may be globular, as in Mirabilis



Jalapa (figs. 376, 407); orbicular, as in Arbutus Andrachne (fig. 408); umbrella-like, as in Sarracenia, where, however, the proper stigmatic surface is below the points of the large expansion of the apex of the style; ovoid, as in Fuchsia (fig. 399); hemispherical; polyhedral; radiating, as in the Poppy (fig. 409), where the true stigmatic rays are attached to a sort of *peltate* or shield-like body, which may repre-

Fig. 405, bis.—Stigmas, s, of Campanula rotundifolia. l, Style.

Fig. 406.—Bilamellar stigmas of Bignonia pandorea. The two lamellæ are applied naturally against each other in 1, while in 2 they are artificially separated.

Fig. 407.—Globular stigma of Mirabilis Jalapa. l, Style. s, Stigma.

Fig. 408.—Circular stigma, s, and l, style of Arbutus Andrachne.

Fig. 409.—Pistil of Papaver somniferum, or opium Poppy. o, Ovary. s, Radiating stigmas on its summit.

sent depressed or flattened styles. The lobes of which a stigma consists may be flat or pointed, as in *Mimulus* and *Bignonia* (fig. 406); or fleshy and blunt, smooth, granular, feathery, as in many Grasses (fig. 410). In Orchidaceæ, the stigma is placed on the column formed by the union of the styles and filaments. The situation where it occurs has been called *gynæceus* (§ 433). In Asclepiadaceæ the stigmas are united to the face of the anthers, and along with them form a solid mass (fig. 353).

459. In **Cryptogamic Plants** there exist organs called *pistillidia*, which have been supposed to perform the function of pistils. They consist of hollow cavities, like ovaries, to which the names of *sporangia* (σπορά, a spore or seed, and ἄγγος, a vessel), and *thecæ* (θήκη, a sac), have been given, containing bodies called *spores*, equivalent to ovules. These spores being capable of germination, and being devoid of cotyledons, have been termed leafless phytons. The sporangia, or spore-cases, are sometimes immersed in the substance of the plant, as in *Riccia glauca* (fig. 411, 1); at other times they are supported on stalks, or *setæ* (*seta*, a bristle), as in Mosses. In *Marchantia* poly-



morpha, they consist of distinct and separate expansions, having a bottle-like form (fig. 412), the lower part, *o*, being enlarged, contain-

Fig. 410.—Pistil of *Cynodon Dactylon*, a Grass. *o*, Ovary. *s*, Feathery Stigmas.

Fig. 411.—1. Perpendicular section of the frond, *f*, of *Riccia glauca*, and of the sporangium or spore-case, *o*, which is imbedded in it. *s*, Narrow process or style by which the sporangium communicates with the external surface. *l* Its cavity or loculus. *s*, Young spores still united in sets of four in the parent cells. *r*, Cells elongated like roots. 2. One of the cells more highly magnified, with the four spores which it contains. Three of the spores are seen, the fourth being concealed by them.

Fig. 412.—Sporangium or spore-case of *Marchantia polymorpha*. *o*, Hollow swelling containing spores, and which has been compared to the ovary. *t*, Narrow process prolonged upwards, and resembling a style. *s*, Termination of this cellular process, compared to the stigma. *c*, Cellular covering of the sporangium, or spore-case, surrounding it like a calyx.

ing the spores, and being surrounded by a cellular tube resembling a calyx, *c*. From this ovary-like body there is a prolongation which may be considered as a style, *t*, terminated by a cellular enlargement, *s*, which has been compared to a stigma. The styloid prolongation withers and disappears when the spores are mature. Sometimes the thecæ, as in Lichens, consist of a club-shaped elongated cell or *ascus* (fig. 413, 1), containing nuclei or cells in its interior, which form the spores. Sometimes these are single, at other times united in sets of two (fig. 413, 2), or of four (fig. 411, 2), or of some multiple of four. There are various modifications of sporangia in other Cryptogamic tribes. Thus, in Ferns, they are often surrounded by an annular ring, or by elastic bands, which cause their dehiscence; while in the Chara they are called *nucules*, and present an oval form with a spiral arrangement of tubes.



413

460. The Ovule.—The ovule is the body attached to the placenta, and destined to become the seed. It bears the same relation to the carpel that marginal buds do to leaves, and when produced on a free central placenta, it may be considered as a bud developed on a branch formed by the elongated axis. The single ovule contained in the ovaries of Compositæ and Grasses, may be called a terminal bud surrounded by a whorl of adhering leaves or carpels, in the axil of one of which it is produced. In *Delphinium elatum*, Brongniart noticed carpels bearing ovules, which were sometimes normal, at other times mere lobes of the carpellary leaf; and in *Aquilegia*, Lindley saw ovules transformed into true leaves, produced on either margin of the carpel. Henslow has seen the ovules of *Mignonette* become leaves. In such cases the vascular bundles of the placenta (*pistillary cords*) are formed by the lateral veins of the carpellary leaf. These veins pass into the marginal lobes or leaflets which represent ovules, and seem to prove that the placenta, in such cases, must be truly a carpellary, and not an axile, formation. Godron, from observing monstrosities, says, that in Leguminosæ, the pericarp, or seed-vessel, is formed by the common petiole dilated, the style is probably the terminal leaflet, or tendrill, or *apiculum*, while the ovules represent lateral leaflets of the leaf, and are modifications of it.

461. The ovule is usually contained in an ovary, but in Coniferæ and Cyadaceæ it has no proper ovarian covering, and is called *naked*. In these orders the ovule is produced on the edges, or in the axil of altered leaves, which do not present a trace of style or stigma. The

carpellary leaves are sometimes so folded as to leave the ovules exposed, or *seminude*, as in *Mignonette*. In *Leontice thalictroides* (blue cohosh), the ovules rupture the ovary immediately after flowering, and the seeds are exposed. So also in species of *Ophiopogon*, *Peliosanthes*, and *Stateria*. In the species of *Cuphea*, the placenta ultimately bursts through the ovary and corolla, becoming erect, and bearing the exposed ovules.

462. The ovule is attached to the placenta either directly, when it is called *sessile*, or by means of a prolongation called a *funiculus* (*funis*, a cord), *umbilical cord*, or *podosperm* (ποῦς, a foot, and σπέρμα, a seed). This cord sometimes becomes much elongated after fertilization. The placenta is sometimes called the *trophosperm* (τρέφω, I nourish). The part by which the ovule is attached to the placenta or cord, is its *base* or *hilum*, the opposite extremity being its *apex*. The latter is frequently turned round in such a way as to approach the base. The ovule is sometimes imbedded in the placenta.

463. In its simplest form, as in the *Mistletoe*, the ovule presents itself as a small cellular projection, which enlarges, assumes an ovoid form (fig. 414), and ultimately becomes hollowed at the apex (fig. 415 c). The cavity thus produced is surrounded by a mass of cells called the *nucleus*, *n*, and is destined to contain the embryo plant after the process of fertilization has been completed. In this embryonal cavity the young plant is suspended by a thread-like cellular

process called *suspensor*, attached to the summit of the nucleus. This cavity in some plants is surrounded by the cells of the nucleus; but, in other cases, it becomes lined with a regular *epithelial* (§ 431), or thin cellular covering, and constitutes the *embryo sac*, which is produced before fecundation, and contains *amnios* or mucilaginous matter in which the embryo is formed.

464. The nucleus (fig. 421 *n*) may remain naked, and form the ovule, as in the *Mistletoe*, *Veronica hederifolia*, *Asclepias*, &c.; but in most plants it becomes surrounded by certain coverings during the progress of development. These appear first in the form of one or more cellular rings at the base, which gradually spread over the surface. In some cases only one covering is formed, as in *Compositæ*, *Campanulaceæ*, *Walnut*, &c. Thus, in the latter (fig. 416), the nucleus, *n*, is covered by a single envelope, *t*, which, in the first instance, extends over the base, and then spreads over the whole surface (fig. 417), leaving only



Fig. 414.—Ovule of the *Mistletoe* entire.

Fig. 415.—Ovule of *Mistletoe* cut to show the embryo-sac, *c*, and the whole of the rest of the mass, *n*, composed of uniform tissue, and forming a nucleus without integuments.

an opening at the apex. In other instances (fig. 418), the nucleus, *n*, besides the single covering (fig. 418, 2, *ti*), has another developed subsequently (fig. 418, 3, *te*), which gradually extends over the first, and ultimately covers it completely. There are thus two integuments, an outer and an inner—the latter, according to Schleiden, being first produced. Mirbel considers the outer as the first formed, and hence has called it *primine*, *te*, while the inner is denominated *secundine*, *ti*. The

latter names are in the present day used by botanists as denominating the outer and inner covering, without reference to their order of development. At the apex

of the nucleus these integuments leave an opening or foramen composed of two apertures; that in the primine (fig. 418, 3, *ex*), called *exostome* (ἐξω, outside, and στόμα, mouth), that in the secundine (fig. 418, 3, *ed*), *endostome* (ἐνδον, within). The


foramen of the ovule is also called *micropyle* (μικρός, small, and πύλη, a gate); but this name is often restricted to its appearance in the seed after fecundation. The length of the canal of the foramen depends on the development of the nucleus, as well as the thickness of the integuments. The embryo-sac is sometimes prolonged beyond the apex of the nucleus, as noticed by Meyen in *Phaseolus* and *Alsine media*, and by Griffith in *Santalum album* and *Loranthus*. Some authors, as Mirbel, consider the ovule in reference to the embryo, and speak of five coverings of the latter, viz., 1. primine; 2. secundine; 3. tercine, or the nucleus; 4. quartine, a temporary cellular layer, which is occasionally formed at an after period around 5. quintine, or the embryo-sac. By most botanists the nucleus and sac with its two integuments are mentioned as the ordinary structure of the ovule.

465. All these parts are originally cellular. The nucleus and integuments are united at the base of the ovule by a cellulo-vascular

Fig. 416.—Ovule of *Juglans regia*, the Walnut. *n*, Nucleus, the base of which only is covered with integument at the early period of development.


Fig. 417.—The same ovule more advanced, in which the nucleus is nearly completely covered.

Fig. 418.—Ovule of *Polygonum cymosum* at various ages. *n*, Nucleus. *te*, The outer integument or primine. *ti*, The inner integument or secundine. *ex*, Exostome or opening in the primine. *ed*, Endostome or opening in the secundine. 1. Ovule in the early state, when the nucleus is still naked. 2. Ovule in second stage, when the nucleus is covered at its base by the internal integument or secundine only. 3. Ovule in the third stage, when the two integuments, primine, and secundine, form a double covering, at the apex of which the nucleus still appears



membrane, called the *chalaza* (fig. 421 *ch*). This is often coloured, of a denser texture than the surrounding tissue, and is traversed by fibro-vascular bundles, which come from the placenta, in order to nourish the ovule. The hilum indicates the organic base of the ovule, while the foramen marks its apex. When the ovule is so developed that the union between the primine, secundine, and nucleus with the chalaza, is at the hilum or base (next the placenta), and the foramen is at the opposite extremity (figs. 417, 418), the ovule is *orthotropal*, *orthotropous*, or *atropous* (ὀρθός, straight, and τρόπος, mode, or α, privative, and τρέπω, I turn). This is the state of an ovule when it first makes its appearance, and occasionally, as in Polygonaceæ, it remains permanent. In such ovules, a straight line drawn from the hilum to the foramen passes through their axes.

466. In general, however, changes take place on the ovule, so that it deviates from the straight line. Thus it may be curved upon itself, so that the foramen approaches to the hilum or placenta, and ultimately is placed close to it, while the chalaza is only slightly removed from the hilum. This change depends apparently on the ovule increasing more on one side than on the other, and as it were drawing the chalaza slightly to one side of the hilum opposite to that where the foramen is applied. Such ovules are called *campylotropal* or *campylotropous* (καμπύλος, curved), when the portions on either side of the line of curvature are unequal (fig. 419); or *camptotropal* (καμπτός, curved), when




they are equal (fig. 420). Curved ovules are found in Leguminosæ, Cruciferae, and Caryophyllaceæ. The union between the parts of the curved portion usually becomes complete, but in some cases there is no union, and the ovules are *lecotropal*, or horse-shoe shaped (λίξος, a hollow disk, and τρόπος, mode or form).

467. When in consequence of the increase on one side, the ovule is

Fig. 419.—Campylotropal or Campylotropous ovule of the Stock. 1. Ovule entire. 2. Ovule cut lengthwise. *f*, Funiculus or umbilical cord. *c*, Chalaza. *n*, Nucleus. *te*, Primine or outer covering. *ti*, Secundine or inner covering. *ex*, Exostome. *ed*, Endostome.

Fig. 420.—Carpel of *Menispermum canadense*, with a curved or camptotropous ovule, *o*, *f*, Funiculus. *s*, The base of the style.

so changed that its apex or foramen (fig. 421, 4, *n*), is in close apposition to the hilum (fig. 421, 5, *h*) and the chalaza is also carried round



so as to be at the opposite extremity (fig. 421, 5, *c*), then the ovule becomes inverted, *anatropal* or *anatropous* (ἀνατρέπω, I subvert). In this case (fig. 422), the union of the chalaza, *ch*, with the nucleus, *n*, is removed from the hilum, and the connection between the chalaza and placenta is kept up by a vascular cord, *r*, passing through the funiculus, and called the *raphe* (ῥᾶφι, a line). The raphe often forms a ridge along one side of the ovule, and it is usually on the side of the ovule next the placenta. Some look upon this kind of ovule as formed by an elongated funiculus (fig. 421, 5, *f*) folded along the side of the ovule, and becoming adherent to it completely; and support this view by the case of *semi-anatropal* ovules, where the funiculus is only, as it were, partially attached along one side, becoming free in the middle; and also by cases where an anatropal ovule, by the separation of the funiculus from its side, becomes an orthotropal seed.

468. The anatropous form of the ovule is of very common occurrence, and may probably aid in the process of fertilization. Ovules which are at first orthotropous, as in *Chelidonium majus* (fig. 421, 2), become often anatropous in the progress of development (fig. 421, 4). When the ovule is attached to the placenta, so that the hilum is in the middle, and the foramen and chalaza at opposite ends, it becomes transverse, *amphitropal* or *heterotropal* (ἄμφι, around, ἑτέρος, diverse).

469. The nucleus of the ovule becomes hollowed at a particular part (fig. 415 *c*), so as to form a cavity. Mirbel states that the whole nucleus is transformed into a membrane called the *tercine*, lining the secundine, and that in its interior another covering, the *quartine*, and





Fig. 421.—Ovule of *Chelidonium majus* at different stages of development. *h*, Hilum or umbilicus. *ch*, Chalaza. *f*, Funiculus or umbilical cord. *r*, Raphe. *n*, Nucleus. *ti*, Secundine. *te*, Primine. *ed*, Endostome. *ex*, Exostome. 1. First stage: nucleus still naked. 2. Second stage: nucleus covered at its base by the secundine. 3. Third stage: the primine developed and covering the secundine at its base. 4. Fourth stage: the ovule completely reflected, and its point turned downwards. 5 The same cut longitudinally, to show the relation of its different parts

Fig. 422.—Anatropous ovule of Dandelion, cut vertically. *ch*, Chalaza. *r*, Raphe. *n*, Nucleus.

finally, the embryo-sac, are produced. The view, however, generally adopted, is that the embryo-sac is formed within the nucleus, assuming a greater or less size according to circumstances, and in some instances reducing the nucleus to a mere external sac. In the interior of the embryo-sac, cellular layers are deposited from without inwards, the earlier ones probably forming the fugacious quartine of Mirbel.

470. In the Mistletoe there are two or three embryo-sacs. The neck of the embryo-sac in *Veronica* and *Euphrasia* becomes elongated and swollen, and from it are developed certain cellular or filamentous appendages, which are probably connected with the nutrition of the embryo.

471. The position of the ovule relative to the ovary varies. When there is a single ovule, it may be attached to the placenta at the base of the ovary (*basal placenta*), following a straight direction, and being *erect*, as in *Polygonaceæ* and *Compositæ* (fig. 423); or it may be inserted a little above the base, on a parietal placenta, with the apex upwards (fig. 424), and then is called *ascending*, as in *Parietaria*. It may hang from an apicular placenta at the summit of the ovary, the apex being downwards, and be *inverted* or *pendulous*, as in *Hippuris vulgaris* (fig. 425), or from a parietal placenta near the summit, and be *suspended*, as in *Daphne Mezereum* (fig. 426), *Polygalaceæ*, and



Euphorbiaceæ. Sometimes a long funiculus arises from a basal placenta, reaches the summit of the ovary, and suspends the ovule, as in *Armeria*; at other times the hilum or true organic base appears to be in the middle, and the ovule becomes *horizontal*, *peltate* (*pelta*, a shield), or *peritropous* (*περί*, around, and *τρέπω*, I turn). All these modifications are influenced by the relative position of the hilum and

Fig. 423—426.—Carpels belonging to different flowers, cut vertically to show the different directions of the solitary ovule, *o*, contained in them. *f*, Funiculus. *r*, Raphe. *c*, Chalaza. *s*, Base of the style.

Fig. 423.—Carpel of *Senecio vulgaris*, with a straight or erect anatropous ovule.


Fig. 424.—Carpel of *Parietaria officinalis*, pellitory, with an ascending orthotropous ovule.

Fig. 425.—Carpel of *Hippuris vulgaris*, Mare's-tail, with a reversed or pendulous anatropous ovule.

Fig. 426.—Carpel of *Daphne Mezereum*, with a suspended anatropous ovule.

foramen, the length of the funiculus, and its adhesion, as well as the position of the placenta.

472. When there are two ovules in the same cell, they may be either *collateral*, that is, placed side by side (fig. 427), or the one may be erect and the other inverted, as in some species of *Spiræa* and *Æsculus* (fig. 428), or they may be placed one above another, each following the same direction. Such is also the case with ovaries containing a moderate or definite number of ovules. Thus, in the ovary of Leguminous plants (fig. 429), the ovules, *o*, are attached to the extended marginal placenta, one above the other, forming usually two parallel rows corresponding to each margin of the carpel. When the ovules are *definite* (uniform, and can be counted), it is usual to find their attachment so constant as to afford good characters for natural orders. When the ovules are very numerous or *indefinite*, while at the same time the placenta is not much developed, their position exhibits great variety, some being directed upwards, others downwards,



others transversely (fig. 430), and their form is altered by pressure into various polyhedral shapes. In such cases it frequently happens that some of the ovules are arrested in their development and become abortive. In Cryptogamous plants, in place of ovules there are cellular bodies called *spores*, to which allusion will be made when the seed is considered.

4.—FUNCTIONS OF THE FLORAL ENVELOPES.

473. The bracts and calyx, when of a green colour, perform the same functions as leaves, giving off oxygen under the influence of light, and producing the carbonized substance called chlorophylle. They

Fig. 427.—Carpel of *Nuttallia cerasoides*, with two suspended collateral ovules. *o*, One of the ovules. *f*, Funiculus. *s*, The base of the style.

Fig. 428.—One of the loculaments of the ovary of *Æsculus hybrida*, laid open to show two ovules, *o o*, inserted at the same height, but turned in different directions. *m m*, Micropyle indicating their apex. *s*, Base of the style.

Fig. 429.—Carpel or legume of *Ononis rotundifolia*, with several campylotropous ovules, *o*, placed one above the other. *f*, Funiculus. *s*, Base of the style.

Fig. 430.—Loculament of the ovary of *Peganum Harmala*, with numerous ovules, *o*, attached to a projecting placenta, *p*, and pointing in different directions. *s*, Base of style.

are consequently concerned in the assimilation of matters fitted for the nutrition of the flower, and they aid in protecting the central organs. The corolla, along with the thalamus and disk, is concerned rather with development than with respiration. Hence it does not in general produce chlorophylle, nor does it give off oxygen. The fact noticed by Garreau that the petals are generally covered with a waxy or fatty matter over their whole surface, may account for their incapability of absorbing water, as well as their resistance to atmospheric influence. The corolla protects the essential organs, and eliminates carbonic acid by a process of oxidation. The starch granules contained in it, as well as in the thalamus and disk, are not altered by the respiratory process, so as to become more highly carbonized, but are oxidized, so as to be converted into saccharine matter. When thus changed it is absorbed by the flowers for their nourishment. The quantity of oxygen absorbed was determined by Saussure. He found that double flowers absorbed less in proportion to their volume than single flowers; that the essential organs contained more oxygen than the floral envelopes; and that the greatest absorption took place when the stamens and pistil were mature.

474. The following are some of Saussure's experiments:—

Name.	Duration of Experiment.	Oxygen Consumed—	
		By Flowers entire.	By essential Organs only.
Stock, single	24 hours.	11·5 times their vol.	18· times their vol.
Stock, double	—	7·7 —	— —
Tuberose, single	—	9· —	— —
Tuberose, double	—	7·4 —	— —
Indian Cress, single	—	8·5 —	16·3 —
Indian Cress, double	—	7·25 —	— —
Brugmansia arborea	—	9· —	— —
Passiflora serratifolia	—	18·5 —	— —
Gourd, male flower	10 —	7·6 —	16· —
Gourd, female	24 —	3·5 —	— —
Hibiscus speciosus	12 —	5·4 —	6·3 —
Hypericum calycinum	24 —	7·5 —	8·5 —
Cobæa scandens	—	6·5 —	7·5 —
Arum italicum	—	— —	30· —
Typha latifolia	—	9·8 —	— —
White lily	—	5· —	— —
Castanea vulgaris	—	9·1 —	— —

475. While this oxidation is going on, carbon is given off in the form of carbonic acid, and heat is developed by the combination between the oxygen and carbon. The quantity of carbonic acid given off is in a ratio to the amount of oxygen absorbed, and the degree of heat present proportionate to the activity of the chemical and vital changes accomplishing. Experiments have been performed by Lamarck, Schultz, Huber, Saussure, Brongniart, Vrolik, and De Vriese, as to the amount of heat produced during flowering, especially by species of

Arum, Caladium, and Colocasia. These are plants in which the floral envelopes are nearly absent, while the essential organs, the torus and growing point, attain a high degree of development, forming a spadix enclosed in a large spathe. No heat could be detected when the contact of oxygen was prevented, either by putting the plants into other gases, or by covering the surface of the spadix with oil. The surface of the spadix is the part whence the heat was chiefly evolved. The Arum cordifolium occasionally had a temperature 20° or 30° above that of the surrounding air; Arum maculatum 17° to 20° ; and Arum Dracunculus and other species still higher. The following observations were made by Brongniart on the spadix of Colocasia odora. The spathe opened on the 14th of March; the discharge of pollen commenced on the 16th, and continued till the 18th. The maximum temperature occurred at different hours.

	Maximum.	Temperature above the Air.		Maximum.	Temperature above the Air.
14th March.	3 P.M.	4.5° Cent.	17th	5 P.M.	11.0° Cent.
15th	4 —	10.0°	18th	11 A.M.	8.2°
16th	5 —	10.2°	19th	10 —	2.5°

476. Vrolik and De Vriese made a series of observations on the same plant, and have given the results for every half-hour. The following are some of these results:—

Hour.	Temperature of Plant.	Temperature of Air.	Hour.	Temperature of Plant.	Temperature of Air.
11-30	20.6° Cent.	18.3° Cent.	3	25.0° Cent.	15.6° Cent.
12	21.1 —	18.7 —	3-30	24.4 —	15.0 —
12-30	23.3 —	19.4 —	4	23.3 —	15.0 —
1	24.4 —	19.4 —	5	22.2 —	18.7 —
1-30	24.4 —	18.9 —	6	21.0 —	18.7 —
2	25.6 —	17.2 —	7	20.0 —	18.7 —
2-30	26.5 —	15.6 —			

The greatest amount of heat observed was at 2-30 P.M., when it was 10.9° above the temperature of the air. On the previous day the maximum occurred at 3 P.M., and on the following day at 1, but then it was only 8.2° above that of the air. Decandolle states that at Montpellier, Arum italicum attained the maximum of temperature about 5 P.M. Saussure observed similar phenomena, but to a less extent, in the Gourd, where the temperature varied from 1.8° to 3.6° ; also in Bignonia radicans, from 0.9° to 3° . From all these experiments, it would appear that in the Araceæ and some other plants, especially at the period when the essential organs reach maturity, there is a production of heat, which increases during the performance of their functions, attaining a daily maximum, and ultimately declining.

477. While these changes are taking place, the starch is converted into dextrine, and ultimately into grape-sugar, which, being soluble,

can be immediately applied to the purposes of the plant. The honey-like matter thus formed is stored up frequently at the base of the petals, in little pits or nectaries, as in *Fritillary*, *Asarum*, &c. It is considered by *Vaucher* and others as performing an important office in fertilization, covering the stigma, and aiding in the dispersion and rupture of the pollen-grains. Bees and other insects, in collecting the saccharine matter, also scatter the pollen.

478. Flowering takes place usually at a definite period of the plant's existence. It requires a considerable amount of nutrient matter, and its occurrence is accompanied with a greater or less exhaustion of the assimilated products. Hence, a certain degree of accumulation of sap seems necessary in order that flowering may proceed. Annual plants are so exhausted after flowering as to die; but, by retarding the epoch for two or more years, as by nipping off the flower-buds, time is allowed for accumulating sap; the stems, from being herbaceous, become shrubby, and sometimes, as in the *Tree-Mignonette*, they may be made to live and flower for several years. Perennial plants, by the retardation of flowering, are enabled to accumulate a greater amount of nutritive matter, and thus to withstand the exhaustion. Many cultivated plants which lay up a large store of nutriment in the form of starch, lose it when the plants shoot out a flowering stem. This is seen in the case of Carrots and Turnips, in which the succulent roots become fibrous and unfit for food when the plants are allowed to run to seed. The receptacle of the Artichoke, and many *Compositæ*, which is succulent before the expansion of the flowers, becomes dry as the process of flowering proceeds. The juices of plants, then, when required for the purpose either of food or medicine, ought in general to be collected immediately before flowering.

479. By cutting a ring out of the bark of trees, and thus retarding the descent of the sap, the period of flowering is sometimes hastened. Again, when the period of flowering is long delayed, either naturally, as in *Agave* and several palms, or artificially, the process, when it does begin, proceeds with amazing rapidity and vigour. In such cases this vigorous flowering is often followed by the death of the plant. *Richard* mentions, that a plant of *Agave*, which had not flowered for nearly a century, sent out a flowering stem of $22\frac{1}{2}$ feet in 87 days, increasing at one period at the rate of one foot a-day. Common fruit trees, when they begin to flower, often do so luxuriantly; but if, from the season being bad, there is a deficiency in flowering, it frequently happens that, from the accumulation of sap, the next year's produce is abundant.

480. If plants are allowed to send out their roots very extensively in highly nutritive soil, the tendency is to produce branches and leaves rather than flowers. In such cases, cutting the roots or pruning the young twigs may act beneficially in checking the vegetative functions. In pruning, the young shoot is removed, and the buds connected with

the branch of the previous year are left, which thus receive accumulated nourishment. Grafting, by giving an increase of assimilated matter to the scion or graft (see section on Fruiting), and at the same time checking luxuriant branching, contributes to the hastening of the flowering.

481. The period of flowering of the same plant varies at different seasons, and in different countries. During the winter in temperate climates, and during the dry season in the tropics, the vegetative process is checked, more especially by the diminished supply of moisture, and the arrestment of the circulation of the sap. The assimilated matter remains in a state of repose, ready to be applied to the purposes of the plant when the moisture and heat again stimulate the vegetable functions. This stimulation occurs at different periods of the year, according to the nature of the climate. By observing the mode of flowering of the same species of plant in successive years, conclusions may be drawn as to the nature of the seasons in a country; and by contrasting these periods in different countries, comparisons may be instituted as to the nature of their climate. Thus valuable floral calendars may be constructed.

482. Plants are accommodated to the climate in which they grow, and flower at certain seasons; and even when transferred to other climates where the seasons are reversed, they still have a tendency to flower at their accustomed period of the year. Again, in the same climate, some individuals of a species, from a peculiar idiosyncrasy, regularly flower earlier than others. Decandolle mentions a horse-chestnut at Geneva, which flowered always a month before the rest in the neighbourhood. From such individuals, by propagation, gardeners are able to produce early-flowering varieties.

483. There is a periodicity in the hours of the day at which some species open their flowers. Some expand early, some at mid-day, others in the evening. The flowers of Succory open at 8 A.M., and close at 4 P.M.; those of *Tragopogon porrifolius*, or Salsafy, close about mid-day. Linnæus constructed a floral clock or watch, in which the different hours were marked by the expansion of certain flowers. The periods, however, do not seem to be always so regular as he remarked them at Upsal. The following are a few of these horological flowers, with their hours of opening:—

<i>Ipomœa Nil</i>	3 to 4 A.M.
<i>Tragopogon pratense</i>	4 ... 5 —
<i>Papaver nudicaule</i>	5 —
<i>Hypochaeris maculata</i>	6 —
Various species of <i>Sonchus</i> and <i>Hieracium</i>	6 ... 7 —
<i>Lactuca sativa</i>	7 —
<i>Specularia Speculum</i>	} 7 ... 8 —
<i>Calendula pluvialis</i>	
<i>Anagallis arvensis</i>	8 —

<i>Nolana prostrata</i>	8 to 9 A.M.
<i>Calendula arvensis</i>	9 —
<i>Arenaria rubra</i>	9...10 —
<i>Mesembryanthemum nodiflorum</i>	10...11 —
<i>Ornithogalum umbellatum</i> (<i>Dame d'onze heures</i>).....	11 —
Various Ficoideous plants.....	12 —
<i>Scilla pomeridiana</i>	2 P.M.
<i>Silene noctiflora</i>	5...6 —
<i>Oenothera biennis</i>	6 —
<i>Mirabilis Jalapa</i>	6...7 —
<i>Cereus grandiflorus</i>	7...8 —

484. Plants which expand their flowers in the evening, as some species of *Hesperis*, *Pelargonium*, &c., were called by Linnæus *plantæ tristes* on that account. Several species of *Cooperia*, and of *Cereus*, also *Sceptranthus Drummondii*, are nocturnal flowers. Some flowers open and decay in a day, and are called *ephemeral*, others continue to open and close for several days before withering. The corolla usually begins to fade after fecundation has been effected. Many flowers, or heads of flowers, do not open during cloudy or rainy weather, and have been called *meteoric*. Composite plants frequently exhibit this phenomenon, and it has been remarked in *Anagallis arvensis*, which has hence been denominated the "poor man's weather-glass." The closing of many flowers in such circumstances protects the pollen from the injurious effects of moisture.

485. The expansion and closing of flowers is regulated by light and moisture, and also by a certain law of periodicity. A plant accustomed to flower in day-light at a certain time, will continue to expand its flowers at the wonted period, even when kept in a dark room. Decandolle made a series of experiments on the flowering of plants kept in darkness, and in a cellar lighted by lamps. He found that the law of periodicity continued to operate for a considerable time, and that in artificial light some flowers opened, while others, such as species of *Convolvulus*, still followed the clock hours in their opening and closing.

486. Light has been said also to have an effect on the direction which flowers assume. Some *Compositæ*, as *Hypochæris radicata* and *Apargia autumnalis*, are stated by Henslow to have been seen in meadows, where they abound, inclining their flowers towards the quarter of the heavens in which the sun is shining. A similar statement has been made regarding the Sun-flower, but it has not been confirmed in this country at least. Perhaps in its native clime, where the effect of the sun's rays is greater, the phenomenon alluded to may be observable. Vaucher mentions the effects of light on the direction of the flowers of many plants, as *Narcissuses* and certain species of *Melampyrum*.

487. It is of importance, both as regards meteorology and botanical

geography, that observations should be made carefully on what are called the annual and diurnal periods of plants: the former being the space of time computed between two successive returns of the leaves, the flowers, and the fruit; and the latter, the return of the hour of the day at which certain species of flowers open. The same species should be selected in different localities, and care should be taken that the plants are such as have determinate periods of flowering. Rules as to the mode of observing periodical phenomena in plants have been drawn up by the British Association, and a committee has been appointed to carry this into effect. The committee has published (1.) a list of plants to be observed for the periods of foliation and defoliation: (2.) a list of plants to be noticed for flowering and ripening of the fruit; (3.) a list of plants to be observed at the vernal and autumnal equinoxes, and summer solstice, for the hours of opening and closing their flowers.

5.—FUNCTIONS OF THE STAMENS AND PISTIL—FERTILIZATION OR FECUNDATION.

488. The stamens and pistil are called the *Essential Organs* of flowering plants, inasmuch as without them reproduction cannot be effected. In plants which do not flower, this function is performed by special organs, which have been termed antheridia and archegonia, or results from a process of conjugation on the part of peculiar cells. The stamens, considered as the male organs, prepare the pollen, which is discharged by the dehiscence of the anther. The pistil, or the female organ, is provided with a secreting surface or stigma, to which the pollen is applied in order that the ovules contained in the ovary may be fertilized.

489. The existence of separate sexes in plants appears to have been conjectured in early times, as shown by the means taken for perfecting the fruit of the Date Palm. In this palm, the stamens and pistils are on separate plants; and the Egyptians were in the habit of applying the sterile flowers to those in which the rudiments of the fruit appeared, in order that perfect dates might be produced. This practice appears to have been empirical, and not founded on correct notions as to the parts of the plant concerned in the process. In the case of the Fig, they were in the habit of bringing wild figs in contact with the cultivated ones, on the erroneous supposition that a similar result was produced as in the case of the Date, proving that they were not aware of the fact, that in the Fig there are stamens and pistils present on the same receptacle. The effect produced by the wild figs, or the process of *caprification* (*caprificus*, a wild fig-tree), as it was called, seems to depend on the presence of a species of *Cynips*, which punctures the fruit, and causes an acceleration in ripening. The presence of sexual organs in plants was first shown in 1676, by Grew, who communicated

his observation to Sir Thomas Millington. This latter naturalist entertained at the same time similar opinions, which were confirmed by Malpighi, Ray, and others. Linnæus made it the basis of his artificial system of classification.

490. Numerous proofs have been given of the functions of the stamens and pistils, especially in the case of plants where these organs are in separate flowers, either on the same or on different plants. Thus, a pistilliferous specimen of Palm (*Chamærops humilis*), in the Leyden Botanic Garden, which had long been unproductive, was made to produce fruit by shaking over it the pollen from a staminiferous specimen. The same experiment has on several occasions been performed in the Botanic Garden at Edinburgh, and the fruit thus ripened has furnished seeds which have germinated. Similar results were observed in the case of the Pitcher plant. In Cucumbers, when the staminiferous flowers are removed, no perfect fruit is formed. Removing the stamens in the very early state of the flower, before the pollen is perfectly formed, prevents fertilization. Care must be taken, in all such experiments, that pollen is not wafted to the pistil from other plants in the neighbourhood, and the result must be put to the test by the germination of the seed; in some instances, the fruit enlarges independently of the application of the pollen, without, however, containing perfect seed. Thus, a species of *Carica* was fertilized by the application of pollen, and produced perfect fruit and seed, and it continued for at least one year afterwards to have large and apparently perfect fruit, but the ovules were abortive.

491. Some authors maintain, that in the case of Hemp, *Lychnis dioica*, and a plant called *Cœlebogyne*, perfect seeds have been produced without the influence of any substance equivalent to pollen; but these statements have not as yet been confirmed. Henslow explains such cases by regarding them as being analogous to what occurs in Aphides, where several generations result from the one impregnation. On the contrary, in Phanerogamous or flowering plants, all experiments lead to the conclusion that there are distinct sexual organs.

492. In Cryptogamous or flowerless plants, considerable doubts have been entertained as to the existence of such organs. In the simplest form of Cryptogamic plants, as the unicellular Red Snow, production and nutrition simultaneously progress within the same cell. As the scale of vegetation is ascended, and the plant becomes complex, there seem to be cells of different kinds, which require to be brought into contact in order that spores (which are equivalent to seeds) may be produced. These reproductive cells are of two kinds, and they are situated either together or apart, on the same or on different individuals. One of these is the *Antheridium*, a cellular body containing granular matter, and *Phytozoa* (φυτόν, a plant, and ζῶς, living), or minute bodies which exhibit movements; the other is the *Pistilli-*

dium or Archegonium (ἀρχή, beginning, and γένος, offspring), containing spores which germinate, and which are sometimes provided with cilia (figs. 431—434), so as to become *Zoospores* (ζῶς, living, and σπορά, a seed or spore), or moving spores. These phytozoa have been regarded as exercising a function similar to that of the spermatozoa in animals. Their presence has been detected in the Archegonia, wherein the embryo is developed. A cessation of their active movements has been observed coincident with the earliest formation of the fruit. An analogy between animal and vegetable embryonic life is thus maintained. When the contents of the antheridia and archegonia are brought into contact, a cellular body is produced in the latter. Within this cell a perfect germ exists, which may be either separated immediately, or, in conjunction with the plant from which it has been produced, become further matured. In Algæ and Charæ, antheridia have been found containing phytozoa and zoospores. The same has been observed in Fungi and Lichens. The spores of the latter when set free have been noticed to undergo peculiar motions which continued for some hours. Mosses having antheridia and archegonia on the same stem generally prove fruitful; when these organs are situated on different plants, unless they are in close proximity one to the other, fructification does not result. The contact of the Antheridium and Pistillidium is by many considered as necessary for the fertilization of the spore. In other cases, as in Confervæ and Diatomaceæ, there is a union of the cells of the plant by *conjugation*, so as to produce germinating bodies. In these cases, the contents of one cell pass, by the formation of a tube, into the other. In *Zygnema*, this conjugation gives rise to germinating bodies in the interior of one of the cells of the plant; in Diatomaceæ, on the outside of the cells.

493. The union of two kinds of *endochrome* (ἐνδον, within, and χρῶμα, colour), or of two kinds of coloured particles, appears in these plants to give rise to the sporangium or spore-case, and the spore. Sometimes the two kinds of endochrome are in separate plants, as already noticed, and then conjugation unites them, and causes a mixture of the endochrome; while in *Meloseira*, &c., the different kinds are apparently situated in different parts of the same cell, and movements take place towards the centre, by which their union is effected and a spore produced. In Ferns, Mosses, &c., there have been detected separate



Fig. 431.—434.—Spores of different fresh-water Algæ.
 Fig. 431.—Spores of a Conferva, with two vibratile cilia.
 Fig. 432.—Spore of a Charophora, with four cilia.
 Fig. 433.—Spore of a Prolifera, with a circle of cilia.
 Fig. 434.—Spore of a Vaucheria, covered with cilia.

cellular bodies, the union of which is considered necessary for the perfection of the spores. In Ferns, the spores contained in the archegonium result, without a process of impregnation, in the formation of a prothallus. On the under surface of this prothallus, antheridia and archegonia appear. The former consist of a cellular papilla having a central cavity containing free cellulæ, which cellulæ when escaping from the investing papillary membrane, set free by their rupture ciliated spiral filaments resembling spermatozoids. The archegonia, which also exist on the under surface of the prothallus, consist of cellular papillæ, in the centre of which may be traced a closed canal leading to a deeper cell or ovule, which, imbedded in the substance of the prothallus, contains within it the embryo germ, with the maturation of the antheridia, and on the escape of the spermatozoids from their ruptured cells, the ovule of the archegonia undergoes simultaneous changes, the canal leading to it becomes opened, a free cell is developed. This cell if not penetrated is reached by those active spermatozoids, when, as a consequence, vital movements result. The cell undergoes a process of division, a perfect embryo is formed, from which the future plant becomes matured. In many Cryptogamic plants the formation of new cells by a constant process of fissiparous or merismatic division (§ 24), may be considered as analogous to the formation of buds, depending perhaps on certain changes similar to fecundation going on in the interior.

494. In flowering plants, various provisions are made for insuring the application of the pollen to the stigma. The comparative length of the stamens and pistils, their position, and the dehiscence of the anthers, are all regulated with this view. The existence of spiral cells in the endothecium has reference apparently to the bursting of the anther, and the scattering of the pollen. The number of pollen-grains produced is also very great. Hassall says that a single head of Dandelion produces upwards of 240,000, each stamen of a Pæony 21,000, a Bulrush 144 grains by weight. It has been stated, that a single plant of *Wisteria sinensis* produced, 6,750,000 stamens, and these, if perfect, would have contained 27,000,000,000 pollen-grains.* In the case of Evergreens, such as Firs, the quantity of pollen is enormous, apparently to insure its application notwithstanding the presence of leaves. The pollen from pine forests has been wafted by the winds to a great distance, and is said to have fallen on the ground like a shower of

* The following estimate was made of the amount of flowers, stamens, &c., in a single specimen of *Wisteria sinensis*:—

Number of clusters of Flowers	9,000
——— Individual Flowers	675,000
——— Petals	3,375,000
——— Stamens	6,750,000
——— Ovules	4,050,000

For the purpose of fertilizing these ovules, the anthers, if perfect, would have contained about 27,000,000,000 pollen-grains, or about 7000 grains to each ovule.

sulphur. It is thus that the phenomena of coloured rain, occasionally witnessed, may be accounted for. The pollen powder transmitted to considerable distances remains floating in the air till carried down by some descending shower it encounters in its course.

495. The quantity of pollen required for impregnation varies. Koelreuter says, that from fifty to sixty grains of the pollen of *Hibiscus Trionum* are required to fecundate the fruit completely, containing about thirty ovules. The ovary of *Nicotiana*, *Datura*, *Lychnis*, and *Dianthus*, according to Gærtner, may be completely fertilized by the pollen of a single perfect anther. In *Geum*, from eight to ten anthers, out of eighty-four to ninety-six contained in each flower, are sufficient to fertilize from eighty to one hundred and thirty ovules contained in the ovary.

496. In many trees in which the organs of reproduction are in separate flowers (as *Hazel* and *Willow*), the leaves are not produced until fertilization has been effected. The protection of the pollen from the direct influence of moisture, is effected by the closing of the flowers, by the elasticity of the anther-coat only coming into play in dry weather; and in aquatics, either by a peculiar covering, as in *Zostera*, or by the flowers being developed above water, as in *Nymphæa*, *Lobelia*, *Stratiotes*, and *Hottonia*. In *Vallisneria spiralis*, a plant growing in the mud of ditches, the stamiferous plants are detached from the soil, float on the surface of the water, and produce there flowers and pollen; while the pistilliferous plants send up a long peduncle (fig. 228), which accommodates itself to the depth of the water by being spiral, and bears on its summit the flower with the pistil. By this means the two organs are brought into contact, and fertilization is effected. *Lagarosiphon muscoides*, an aquatic plant from Africa, shows similar phenomena in regard to impregnation as are seen in *Vallisneria*. When continued wet weather comes on after the pollen has been matured, and has begun to be discharged, it often happens that little or no fruit is produced. In flowers where the anthers burst in succession, the injury done by moisture is less likely to extend to all.

497. In some plants the stamens, at a certain period of their development, move towards the pistil, so as to scatter the contents of the anther. In *Parnassia palustris* and *Rue*, they do so in succession. In *Kalmia*, the anthers are contained in little sacs or pouches of the corolla, until the pollen is mature, and when the expansion of the corolla, and the elasticity of the filament, combine to liberate them, they spring towards the pistil with a jerk. In *Parietaria officinalis*, and in the *Nettle*, the spiral filament is kept in a folded state until the sepals expand, and then it rises with elastic force and scatters the pollen. Similar phenomena are observed in the *Cornus canadensis*. In the various species of *Barberry*, the inner and lower part of the filament is irritable, and

when touched it causes the stamen to move towards the pistil. This takes place naturally when the anther is ripe, and the recurved valves covered with pollen are ready to be applied to the stigma. The species of *Stylidium* have their anthers and stigma seated on a column, the base of which is slightly swollen and irritable. When a stimulus is applied, this column passes with considerable force from one side of the flower to the other, rupturing the anther lobes, and thus aiding in fertilization.

498. In certain plants the agency of insects is employed to insure fecundation. In species of *Aristolochia*, the tube of the calyx completely encloses the organs of reproduction, and the stamens are placed below the stigma, so that the pollen can neither be applied directly, nor be carried by the winds. These plants are said to be frequented by insects, which enter the tubes and reach the little chamber at the base, with the view of collecting saccharine matter. The deflexed hairs in the interior of the tube prevent their escape, and in their movements they apply the pollen to the stigma. When this is accomplished, the flower withers, and the insects escape. Orchidaceous plants have remarkable flowers, which resemble bees, flies, spiders, and in general the insects of the country in which they grow. The special odour of these flowers subserve the purposes of their attraction. They also contain a large quantity of honey-like matter connected with the essential organs, on which these insects feed. These insects, in searching for such food, detach the pollen masses, which are easily removed from the clinandrium, or the part of the column on which they are placed, and then naturally fall on the stigmatic surface. Bees and aphides may, in many instances, contribute to the process of fertilization.

499. While the pollen is being elaborated, the stigma is also undergoing changes. It becomes enlarged, and secretes a viscid matter ready to detain the pollen-grains when they are discharged. This secretion was represented by Dr. Aldridge to be, in some cases, acid; but it seems more generally to be of a mucilaginous or saccharine nature. Vaucher thinks that the nectariferous fluid usually found in the flower spreads itself over the stigma, and that it is sometimes instrumental in conveying the pollen-grains. In *Goldfussia* or *Ruellia anisophylla*, and the species of *Campanula*, as *media*, *Rapunculoides*, and *Trachelium*, the style is covered with collecting hairs, which appear to aid in the application of the pollen. In the first-mentioned plant, a remarkable curvation of the style takes place, so as to make the stigma come into contact with the hairs. In *Campanula*, the hairs on the upper part of the style seem to collect the pollen, and allow it to be applied to the revolute branches of the stigma. In this genus the style is at first slightly longer than the stamens, but it soon becomes twice their length, and during its elongation, the hairs upon it brush the pollen-grains

out of the anther cases, and thus raise them into a position where they can be applied ultimately to the stigmatic surface. The stigma consists of two branches, which are at first erect, but afterwards, by changes in the cells, become completely revolute, so as to come into contact with the hairs. After the hairs have performed their office, their fine inner membrane collapses by a process of endosmose, and the stiff outer membrane is drawn inwards, so as to retire within its cell. After the pollen reaches the stigma, changes take place in it, by which the fovilla contained in the intine of the grains is allowed to escape. This fovilla consists of minute molecules exhibiting certain movements, which by some have been considered analogous to those of phytozoa in Cryptogamic plants, or spermatozoa in the animal kingdom. Others look upon these motions as entirely molecular.

500. The length of time during which the pollen retains its vitality, or power of effecting fertilization, varies in different plants. According to Gærtner and others, the pollen of some species of *Nicotiana* retains its vitality only for forty-eight hours; pollen of various species of *Datura*, two days; pollen of *Dianthus Caryophyllus*, three days; pollen of *Lobelia splendens*, eight or nine days; pollen of *Cheiranthus Cheiri*, fourteen days; pollen of *Orchis abortiva*, two months; pollen of *Candollea*, one year; pollen of Date Palm, one year or more. Michaux says, that in some Palms, as Date and *Chamærops humilis*, the pollen may be applied successfully after having been carefully kept for eighteen years. The pollen retains its vitality longer when not removed from the anthers; and the finer it is, the more quickly it loses its fecundating property.

501. **Theories of Embryology.**—So far as the application of the pollen to the stigma is concerned, the process of fertilization can be easily traced, but the changes which are subsequently produced on the pollen-grain and the ovule are very obscure, require minute microscopic research, and have led to numerous conflicting theories of Embryology. It has been already stated that some physiologists, especially Bernhardt, believe, that in the case of Hemp, *Lychnis dioica*, and some other plants, an embryo, or young plant, can be produced without the influence of the pollen. These views have not been confirmed. It has been supposed that in such plants as Hemp, where the stamens and pistil are generally on separate individuals, there may occasionally occur instances in which they are developed on the same plant. It is known that this takes place in other cases. Thus, a specimen of *Chamærops humilis*, or European Fan-palm, in the Botanic Garden of Edinburgh, which had for upwards of twenty years shown pistilliferous flowers only, exhibited, in 1847, both pistilliferous and staminiferous clusters,* and produced perfect fruit

* It has done so also in 1848.

without an artificial application of pollen. Again, in Diœcious plants growing in the open air, the pollen may be carried from other plants by the wind, and thus produce perfect seed. There are thus numerous sources of fallacy in Bernhardt's observations; and Gærtner's recent experiments seem to prove that Hemp is no exception to the ordinary rule. Some, with Henslow, on the supposition of the correctness of Bernhardt's views, have thought that the case might be analogous to that of some Aphides, where one impregnation is sufficient to produce several generations. Mr. Smith has recently stated, that a female plant of *Cœlebogyne*, belonging to the natural order of Euphorbiacæ, produced perfect seeds in the garden at Kew, without any apparent contact of pollen; and Gasparrini maintains, that in the case of the cultivated Fig, the seeds are the product of pistillate flowers only. Such cases, if proved, will modify the views entertained relative to the action of pollen. Can it be that, as in the case of some Cryptogamies, there are in these anomalous cases two kinds of cells present in the same organ, some with fertilizing matter, and others containing the rudiments of ovules, or of the embryo?

502. The subject of Embryology, or the development of the embryo in the seed, has attracted much attention of late, and numerous opinions have been advanced. There are many discrepancies as to the contents of the ovule before impregnation; some maintaining that the cavity of the nucleus, or the embryo-sac alone, is developed before the pollen is applied; others, that besides it there is a utricle, or vesicle, in its interior, which forms the first embryonic cell. In some plants, as in *Lygnema*, the primitive embryo results from a process of conjugation in which the different functions of distant cells are, when brought into contact, conjointly exercised for the production of a spore, the united endochromes resulting in its formation, and the development of its proper membrane. The tubes in contact with the ovule are by some said to be derived from the pollen; by others, from the ovule itself; and by a third set, from the conducting tissue of the style.

503. Some maintain that the pollen-grains burst on the stigma, and scatter the fovilla directly upon it, the *influence* of which is conveyed by the conducting tissue to the ovule. Hartig seems to adopt this view in some cases, as well as Gasparrini, who, from observations on the Orange-tree and *Cytinus*, thinks that, as the result of this influence, a filament, or cellular prolongation, is sent from the lower extremity of the style into the ovule. Brongniart believed, that, from the inferior extremity of the pollen-tube the fovilla burst forth, and was conveyed by the conducting tissue to the ovules. Almost all modern physiologists discard this view, and believe in the formation of pollen-tubes, which were so ably demonstrated by Brown in Orchidacæ and Asclepiadacæ, and have subsequently been shown

by Schleiden, Amici, Brongniart, Meyen, Mohl, Mûlder, Griffith, and others. These tubes, which vary in size, being often about $\frac{1}{1000}$ inch in diameter, may be easily traced in many instances, after the pollen has lain for a few hours on the stigma; for instance, in *Crocus*, *Salvia*, *Colocasia odora*, *Gesnera*, *Oenothera*, and *Antirrhinum* (fig. 434 bis). In some instances the emission of tubes commences almost immediately after the pollen has been applied to the stigma; in others, they may, as in *Mirululis Jalapa*, take from 24 to 36 hours, or, as in the Larch, take as many days for their emergence. When an ovary, style, and stigma are present, the tubes pass into the conducting tissue, while in the case of naked ovules, as in *Coniferæ* and *Cycadaceæ*, the pollen comes into direct contact with the foramen.



434 bis

504. The extent to which the tubes penetrate, and the mode in which they give rise to the embryo, are matters of dispute. It is maintained by some, that the tubes formed by the intine proceed only to a certain extent down the style before rupturing to discharge the fovilla, so that the influence of the latter is subsequently conveyed to the ovule by the conducting tissue, and thus causes the formation of an embryo. Mirbel and Spach, from observations made on *Gramineæ*, as *Zea Mais*, and on the Yew and other *Coniferæ*, have been led to support this view. They believe that the tube does not reach the ovule, that a primary utricle or vesicle (the first part of the embryo) exists in the embryo-sac, or cavity of the nucleus, before fertilization, and that the fovilla is the means of determining the future development of an embryo in it. This utricle is attached to the embryo by a cellular process, or *suspensor*. These observations are opposed to Schleiden's views, who, according to the opinions of their authors, was led into the error of confounding a cell of the ovule with the extremity of the pollen-tube. Giraud entertains the same opinion, founded on an examination of the ovule of *Tropæolum majus*. In

Fig. 434 bis.—Portion of the stigma of *Antirrhinum majus* at the time of fecundation. *ps, ps*, Superficial cells forming the papillæ. *tc, tc*, Deep elongated cylindrical cells forming the conducting tissue. *gp*, Grains of pollen attached to the surface of the stigma, the extine having been ruptured, and the intine protruded in the form of tubes, *tp, tp*, which pierce the interstices between the superficial stigmatic cells.

this plant he traces the formation of the amniotic or embryo-sac, and primary utricle or germinal vesicle of the embryo, before impregnation, the latter being at first distinct from the sac, but subsequently attached to it by a suspensor; the fovilla is brought into contact with the outer surface of the embryo-sac, and the first trace of the embryo appears in the formation of a spherical body at the inferior extremity of the primary utricle,—this spherical body resulting from a peculiar process of nutrition, determined by the dynamic influence of the fovilla. Giraud also observed a lengthening of the primary utricle and of its suspensor, so as to protrude through the apex of the embryo-sac, the nucleus, and the foramen, forming cells which partly communicate with the conducting tissue, and partly passed round the ovule within the carpellary cavity. By slight traction of this cellular process, the suspensor with the embryo may be drawn from the embryo-sac through the exostome.

505. Hartig thinks that in some cases where the pollen-tube cannot be traced directly downwards to the ovule, there is a series of cells, which, from their continuity, might be mistaken for it. In some recent observations on *Campanula rotundifolia*, Wilson appears to think that the pollen-tube is prolonged into the foramen of the ovule. Dickie has noticed, in *Narthecium ossifragum*, and *Euphrasia officinalis* and *Odontites*, a cellular process proceeding upwards from the ovule into the style, which he thinks may unite with the pollen-tube. Through Dr. Dickie's kindness, I have had an opportunity of seeing these ovule tubes, which appear to end in shut extremities, and not to have a direct communication with the pollen-tubes. These tubes (less than $\frac{1}{800}$ of an inch in diameter) have been traced by him from the interior of the embryo-sac. They are probably derived from the embryo itself, which, in its early state, may send out tubular prolongations similar to those of the spores of Cryptogamic plants. Tubular prolongations from the ovule were long ago noticed by Mirbel, and of late years by Griffith and Hartig. These authors, however, seem to differ from Dickie, in supposing that the tubes are derived from the embryo-sac, or some of the coverings of the ovule.

506. Another, or what may be called a third view of impregnation, is, that the pollen-tube does not stop short in the style, but proceeds as far as the foramen of the ovule, enters it, and is applied to the embryo-sac. In the case of the Mistletoe, where there are no coverings of the ovule, and consequently no foramen, the pollen reaches the nucleus directly. Meyen, Amici, Mohl, Mülder, and Hofmeister, are in favour of this theory. Meyen states, that after the pollen-tube becomes united to the ovule in the form of a *cul de sac*, the process of absorption goes on so as to allow the fovilla to reach the embryo-sac. Immediately thereafter, a development of cells takes place in the form of a beaded prolongation or suspensor, at the extremity of which the

embryo, in the form of a globular cell, is developed. The free communication between the pollen-tube and the embryo-sac ceases after a time; a constriction takes place by the formation of a diagonal septum, and the pollen-tube either shrivels or continues for some time adherent to the sac.


507. Amici also believes that the pollen-tube is applied to the sac of the embryo. He observes in the nucleus a large cavity, which he has called the embryonic vesicle (fig. 435 *c*). In *Cucurbita Pepo* and *Orchidaceæ*, he traces the tube to a certain depth into the nucleus, and he believes that the granular contents of the tube, which are accumulated at the extremity, are absorbed by the embryonic vesicle, so as to effect impregnation. Cells are then produced in the vesicle, commencing at the base, *i.e.* opposite to the part where the pollen-tube exerts its influence. The vesicle becomes full of granular matter; it then exhibits a contraction in the middle, the lower part becoming appropriated to the embryo, and forming the true embryo-sac, while the upper part, in such plants as *Orchis Morio* and *mascula*, elongates upwards (fig. 435 *e*), forming a compound filament, composed of cells with fluid contents. This filament traverses in an inverse manner the course followed by the pollen-tube, and passes into the interior of the placenta, being quite distinct from the pollen-tube, and probably connected in some way with the nutrition of the embryo. Sometimes the pollen-tube remains after the embryo has multiplied its cells. Hartig thinks, that, in different instances, the mode of impregnation is different. Thus he admits that the true pollen-tube comes into contact with the ovule in *Coniferæ*, that in some *Cruciferæ* the tubes in connection with the ovule are derived from the conducting tissue of the style, as maintained by Gasparrini, while, in certain *Cupuliferæ*, tubes proceed from some part of the ovule itself.

508. A fourth theory is, that the pollen-tube (fig. 436 *p t*), after reaching the ovule, enters the foramen, *ex* and *en*, and then penetrates the embryo-sac, *es*, or pushes the sac before it, becoming thus enclosed in a reflection of it. This view is supported by Schleiden, Wydler, Tulasne, Gelesnow, and Wilson. Schleiden, who has made a very elaborate series of observations on the embryo, is in favour of this view. According to him, the extremity of the pollen-tube does not enter the embryo-sac, but continues on the outside of it, pressing in the sac



Fig. 435.—Ovule of *Orchis mascula*, illustrating Amici's view of fertilization. *a*, Primine. *b*, Secundine. *c*, Embryo. *e*, Conservoid filament which proceeds from the embryo towards the placenta, and is independent of the pollen-tube.

before it, and thus becoming surrounded by a double layer of it. The end of the pollen-tube (fig. 436 *e*) forms the germinal vesicle, in the interior of which nuclei and cells are produced, which ultimately give origin to the embryo, surrounded by the endosperm or albumen which results from the transformation of the contents of the embryo-sac. All the portion of the pollen-tube within the embryo-sac may be developed as the embryo, or a portion may remain in the form of a suspensor, or suspensory filament, attached to the upper part of the sac.



436

509. Wydler, Gelesnow, and Tulasne, maintain that there is no indentation of the sac, but that the tube enters it directly. Schleiden thinks

that this may be true in certain cases where the embryo-sac becomes elongated upwards, and then the membrane is absorbed so as to allow the pollen-tube to penetrate into the interior. Schleiden thus looks upon the embryo as a foreign body entering from without, and supports his views by cases of polyembryony in *Coniferæ*, *Cycadacæ*, *Mistletoe*, *Onion*, &c., where the plurality of embryos, according to him, depends on more than one pollen-tube having entered the foramen of the ovule. Wilson has adopted Schleiden's views, from observations made on *Campanula rotundifolia*. Tulasne, from examining the embryogeny of *Veronica hederifolia*, *triphyllos*, and *præcox*, concurs in Wydler's views. Endlicher supports similar views with Schleiden, considering the stigma, however, as an organ, the peculiar secretion of which acts on the pollen-grain, so as to render it capable of penetrating to the ovule, and developing an embryo. Unger's opinion nearly corresponds with this. Griffith, from his researches on *Viscum*, *Santalum*, *Osyris*, and *Loranthus*, concludes, that the pollen-tube penetrates the embryo-sac or cavity, and passes through it longitudinally; and he seems to think that, in some cases, the embryo proceeds from cells developed from the end of the tube. Klotzsch states that pollen-tubes may be seen entering the embryo-sac in *Lavatera tremestris*, *Tobacco*, and some *Orchidacæ*, after the pollen has lain from twenty-four to thirty-six hours on the stigma. Hofmeister admits that the pollen-tube, in some cases where the embryo-sac is very delicate, pushes it inwards to a certain extent.

Fig. 436.—Section of ovule to illustrate Schleiden's view of fertilization. *r*, Raphe. *ch*, Chalaza. *p*, Primine. *s*, Secundine. *ex*, Exostome. *en*, Endostome. *n*, Nucleus. *es*, Embryo-sac. *p* *t*, Pollen-tube. *s*, The embryo formed by the extremity of the pollen-tube.

510. Those who object to Schleiden's views, think that he has mistaken the primary utricle, or germinal vesicle, which exists in the embryo-sac before impregnation, for the end of the pollen-tube, and that the cellular filament attaching this to the embryo-sac is totally independent of the pollen-tube. Brown finds that, in the seed of Coniferous plants which have several embryos, there are semicylindrical corpuscles, three to six in number, which are arranged in a circle near the apex. In each of these is a distinct embryonal filament. These filaments frequently ramify, each of the ramifications terminating in an embryo. He believes that these corpuscles are not formed by the pollen-tubes, and that the fact of the ramifications giving rise to rudimentary embryos is opposed to Schleiden's views. The corpuscles lie dormant for at least twelve months before being developed. In a female plant belonging to Cycadaceæ, another polyembryonous order of plants, Brown has noted the formation of corpuscles at a time when no male flowers were known to exist in the country. These corpuscles may probably be considered as analogous to embryo-sacs, or embryonal cavities, such as those in the Mistletoe. Henfrey states that in Conifera and Cycads the pollen-grains are applied immediately to the micropylle of the ovule without the aid of the stigma: that those grains traversing the cell of the nucleus, are brought into contact with the embryo-sac. From the cells within this sac corpuscles become developed consisting of a large central cell, which, surrounded by four others, is accessible through a canal existing between them. The pollen-tube, ere discharging its fovilla, enters this canal, and impregnates the large central cell. This central cell may then be regarded as an ovule, from which suspensors, equal in number to the surrounding cells, become developed. At the extremity of each of these suspensors a rudimentary embryo exists, but one of which becomes matured.

511. Some of the supporters of Schleiden's views institute a fanciful comparison between the spores of Cryptogamous, and the pollen of Phanerogamous plants. In the former, the cellular germinating body or spore is contained in a case or theca, just as the pollen-grains are in the anther of the latter. In the first instance, the body when discharged is at once fit for germination; in the second it requires to be transmitted to an ovary, and then to be matured within an ovule, and supplied with a store of nutritious matter, so as to be fitted for independent existence. These views are theoretical, and do not seem to be borne out by facts.

512. It will thus be seen that physiologists are much divided in their views relative to this obscure subject; and when we consider the minuteness of the observations, and the high microscopic powers required, it is not a matter of surprise that there are numerous sources of fallacy. Nearly all agree in the formation of pollen-tubes; these, according to some, end in the cellular tissue of the style; according

to others, they reach the ovule; the influence of the fovilla is communicated to the ovule either directly or indirectly; the first cells of the embryo, some maintain, are formed by the end of the pollen-tube directly, while others say indirectly; and a third party consider them as the result of changes induced by the action of the fovilla.

513. The opinions which have been recently supported by Amici, Mohl, Karl, Mûlder, and Hofmeister, are those which seem to rest on the best foundation, viz., that at the time of the opening of the flower the embryo-sac exists, and that, at its upper or micropyle end, one or more cells or germinal vesicles are produced from cytoblasts; that the pollen-tubes pass down the style to the ovary into the foramen of the ovule, and come into contact with the embryo-sac, either at its apex or a little below it; then an imbibition of fluids takes place, the embryo-sac begins to increase, and the embryo is produced. The chief point to be determined, is the existence or not of the germinal vesicle before impregnation.

514. The formation of the process called the suspensor is variously accounted for. Schleiden considers it as a part of the pollen-tube; Amici thinks that it is part of the embryo-sac prolonged upwards, forming a confervoid filament (fig. 435 e); Mohl, Mirbel, and Spach, maintain that the suspensor is produced from the germinal vesicle, and therefore intimately united to the embryo, which is developed from the lowest cells of that vesicle; Dickie thinks that the suspensor may be a cord-like process sent out from the cellular embryo, reaching a certain degree of development, and sometimes sending off tubular prolongations or filaments, as in *Euphrasia* and *Orchidaceæ*. The suspensor is usually directed to the apex of the nucleus or the micropyle, and it is sometimes of great length. In *Draba verna* (fig. 485, 2), Dickie says he observed in an embryo, $\frac{1}{4}$ of an inch long, a suspensor three times that length. In *Gnetum*, Griffith mentions a tortuous suspensor $3\frac{1}{2}$ to 5 inches long, the whole seed being only one inch long.

515. Taking a comprehensive view of the whole subject, it may be said that the union of two kinds of cells appears to be necessary for fertilization. In Cryptogamic plants this has been traced, particularly in certain cases of conjugation; where the two cells come into contact, a tube is formed between them, and the contents of the one unites with those of the other, giving rise to a germinating body. In Phanerogamic plants, also, there are two cells with different contents—the pollen-grain with its granular fovilla, and the ovule with its mucilaginous fluid. These are brought into connection by means of the pollen-tube, formed from the intine, which either enters the embryo-sac, or comes into contact with it, the union taking place either directly by its extremity, or indirectly by cellular prolongations from the conducting tissue, or from the ovule. By this means the

formation of the embryo is determined, which commences as a cellular body or germinal vesicle, in the interior of which other cells are subsequently formed in a definite order of succession.*

516. The Production of Hybrids.—If the pollen of one species is employed to fertilize the ovules of another, the seed will often produce plants strictly intermediate between the two parents. These are termed *hybrids*, and are analogous to mules in the animal kingdom. As a general rule, hybrids can only be produced between plants which are very nearly allied, as between different species of the same genus. Thus, different species of Heath, Fuchsia, Cereus, Rhododendron, and Azalea, readily inoculate each other, and produce intermediate forms. It is found, however, that species which seem to be nearly related do not hybridize. Thus, hybrids are not met with between the Apple and Pear, between the Gooseberry and Currant, nor between the Raspberry and Strawberry. The ovules of Fuchsia coccinea, fertilized with the pollen of Fuchsia fulgens, produce plants having intermediate forms between these two species. Some of the seedling plants closely resemble the one parent, and some the other, but they all partake more or less of the characters of each. By the examination of the foliage, conclusions may be drawn as to what will be the character of the flower. Mr. Thwaites mentions a case in which a seed produced two plants extremely different in appearance and character; one partaking rather of the character of Fuchsia fulgens, and the other of Fuchsia coccinea.

517. In the case of hybridization, there appears to be a mixture of matters derived from the pollen-grain and the ovule, just like the mixture of two endochromes in flowerless plants; and the nature of the hybrid depends on the preponderance of the one or other. Some have supposed that the pollen-grains require to be of the same form and dimension in order to admit of artificial union taking place; but this is a mere conjecture. It is, however, requisite that, for successful hybridizing, the pollen must be in a state of full maturity, and the stigma be equally perfected. Hybrids perform the same functions as their parents, but they do not perpetuate themselves by seed. They must be propagated by offsets or cuttings. If not absolutely sterile at first, they usually become so in the course of the second or third generation. Herbert mentions instances of hybrid Narcissuses, from which he attempted in vain to obtain seed. The cause of this sterility

* For opinions as to Embryology, see Schleiden's paper in *Nova acta Academ. Cæsar. Leopold Carol. Naturæ Curios.*; Brown on the Fecundation of Orchidaceæ and Asclepiadaceæ, in the *Transactions of the Linnæan Society* for 1833; Brongniart on the same subject, in *Annales des Sciences Naturelles*, 1st series, tom. xxiv., 1831; Meyen on the Act of Impregnation, translated in *Taylor's Scientific Memoirs*, vol. iii.; Amici sur la Fécondation des Orchidées, in *Annales des Sciences Naturelles*, 3d series, tom. vii.; Wilson on *Tropæolum majus*, and *Hairs of Campanula*, *London Journal of Botany*, vol. ii.; Giraud on Embryology, in the *Transactions of the Edinburgh Botanical Society*, and in the *Annals of Natural History*, vol. v.; and on the Ovule of *Tropæolum*, in *Linnæan Transactions*, vol. xix.; Henfrey and Diskie on Embryology, in the *Annals of Natural History* for 1848; Griffith on the Ovule of *Santalum*, *Osyris*, &c., in *Linnæan Transactions*, vol. xix.; also *Notulæ ad Plantas Asiaticas*, Calcutta, 1847.

has not been determined. Some have referred it to an alteration in the pollen. Hybrids may be fertilized, however, by the pollen taken from one of the parents, and then the offspring assumes the characters of that parent.

518. Hybrids are rarely produced naturally, as the stigma is more likely to be affected by the pollen of its own stamens than by that of other plants. In dioecious plants, however, this is not the case, and hence the reason, probably, of the numerous so-called species of Willows. Hybrids are constantly produced artificially, with the view of obtaining choice flowers and fruits, the plants being propagated afterwards by cuttings. In this way many beautiful Roses, Azaleas, Rhododendrons, Pansies, Cactuses, Pelargoniums, Fuchsias, Calceolarias, Narcissuses, &c., have been obtained. By this process of inoculation, and carefully selecting the parents, gardeners are enabled to increase the size of the flowers, to improve their colour, to render tender plants hardy, and to heighten the flavour of fruits. Herbert thinks, from what he saw in Amaryllides, that in hybrids the flowers and organs of reproduction partake of the characters of the female parent, while the foliage and habit, or the organs of vegetation, resemble the male. Hybrids occur in which the characters of the two parents are so intimately blended, that it is impossible to say to which they belong.

519. This subject is important as connected with the origin and limitation of species. If, as some of the old authors believed, there were only a few species originally formed, and all the rest are the result of hybridization, there appears to be no limit to species, and no permanence in their characters. This, however, is not borne out by facts; the generally received opinion being, that types of all the species now in existence were originally placed on the globe, and that these have given origin to an offspring like themselves, capable of reproducing the species. Hybrids, on the other hand, are rare in a wild state, and they are seldom permanent and fertile. Even where they are so, there seems to be a tendency in the offspring to return to one or other of the original types from which they sprang.


6.—FRUIT, OR THE PISTIL ARRIVED AT MATURITY.

520. After fertilization, various changes take place in the parts of the flower. Those more immediately concerned in the process, the anther and stigma, rapidly wither and decay, while the filaments and style often remain for some time; the floral envelopes also become dry, the petals fall, and the sepals are either deciduous, or remain persistent in an altered form; the ovary becomes enlarged, forming the pericarp (περί, around, and καρπός, fruit); and the ovules are developed as the *seeds* containing the *embryo-plant*. The term *fruit*

is strictly applied to the mature pistil or ovary, with the seeds in its interior. But it often includes other parts of the flower, such as the bracts and floral envelopes. Thus, the fruit of the Hazel and Oak consists of the ovary and bracts and calyx combined; that of the Apple, Pear, and Gooseberry, of the ovary and calyx; and that of the Pine-apple, of the ovaries and floral envelopes of several flowers combined. Fruits formed by the ovaries alone, as the Plum and the Grape, seem to be more liable to drop off and suffer from unfavourable weather, than those which have the calyx entering into their composition, as the Gooseberry, the Melon, and the Apple.

521. In general, the fruit is not ripened unless fertilization has been effected; but cases occur in which the fruit swells, and becomes to all appearance perfect, while no seeds are produced. Thus, there are seedless Oranges, Grapes, and Pine-apples. When the seeds are abortive, it is common to see the fruit wither and not come to maturity; but in the case of Bananas, Plantains, and Bread-fruit, the non-development of seeds seems to lead to a larger growth, and a greater succulence of the fruit.

522. In order to comprehend the structure of the fruit, it is of great importance to study that of the ovary in the young state. It is in this way only that the changes occurring in the progress of growth can be determined. The fruit, like the ovary, may be formed of a single carpel, or of several. It may have one cell or cavity, then being *unilocular* (*unus*, one, and *loculus*, box or cavity); or many, *multilocular* (*multus*, many), &c. The number and nature of the divisions depend on the number of carpels, and the extent to which their edges are folded inwards. The appearances presented by the ovary do not, however, always remain permanent in the fruit. Great changes are observed to take place, not merely as regards the increased size of the ovary, its softening and hardening, but also in its internal structure, owing to the suppression, enlargement, or union of parts. In this way the parts of the fruit often become unsymmetrical, that is, not equal to, or some multiple of, the parts of the flower; and at times they are developed more in one direction than another, so as to assume an irregular appearance. In the Ash (fig. 437), an ovary with two cells, each containing an ovule attached to a central placenta, is changed into a unilocular fruit with one seed; one ovule, *l*, having become abortive, and the other, *g*, gradually extending until the septum is pushed to one side, becoming united to the walls of the cell,



437

Fig. 437.—Samara, or Samaroid fruit of *Fraxinus oxyphylla*. 1. Entire, with its wing, *a*. 2. Lower portion cut transversely, to show that it consists of two locuments; one of which, *l*, is abortive, and is reduced to a very small cavity, while the other is much enlarged, and filled with a seed, *g*.

and the placenta appearing to be parietal. In the Oak and Hazel, an ovary with three cells, and two ovules in each, changes into a one-celled fruit with one seed. Similar changes take place in the Horse-chestnut, in which the remains of the abortive ovules are often seen in the ripe fruit. In the Coco-nut, a trilocular and triovular ovary is changed into a one-celled, one-seeded fruit. This abortion may depend on the pressure caused by the development of certain ovules, or it may proceed from the influence of the pollen not being communicated to all the ovules. Again, by the growth of the placenta, or the folding inwards of parts of the ovary, divisions may take place in the fruit which did not exist in the ovary. In *Pretrea Zanguebarica*, a one-celled ovary is changed into a four-celled fruit by the extension of the placenta. In *Cathartocarpus Fistula* (fig. 395), a one-celled ovary is changed into a fruit having each of its seeds in a separate cell, in consequence of spurious dissepiments being produced in a horizontal manner, from the inner wall of the ovary after fertilization. In *Tribulus terrestris*, each cell of the ovary (fig. 438) has slight projections, *c*, on its walls, interposed between the ovules, *o*, which, when the fruit is ripe, are seen to have formed distinct transverse divisions (fig. 439 *c*), or spurious dissepiments, separating the seeds, *g*. In *Astragalus*, the folding of the dorsal suture inwards converts a one-celled ovary into a two-celled fruit.



523. The development of cellular or pulpy matter frequently makes great changes in the fruit, and renders it difficult to discover its formation. In the Strawberry, the axis becomes succulent, and bears the carpels on its convex surface; in the Rose, there is a fleshy lining of the calyx (sometimes called a disk), which bears the carpels on its concave surface. In the Gooseberry, Grape, Guava, Tomato, and Pomegranate, the seeds nestle in pulp formed apparently by the placentas. In the Orange, the pulpy matter surrounding the seeds is formed by succulent cells, which are produced from the inner lining of the pericarp.

524. The pistil, in its simplest state, consists of a carpel or folded leaf, with ovules at its margin; and the same thing will be found in the fruit, where the pericarp, as in the Bean (fig. 440), represents the carpellary leaf, and the seeds correspond to the ovules. The pericarp consists usually of three layers; the external (fig. 440 *e*), or *epicarp* (*ἐπί*, upon, or on the outside, *καρπός*, fruit), corresponding to the lower epidermis of the leaf; the middle (fig. 440 *m*), or *mesocarp* (*μέσος*, middle), representing the parenchyma of the leaf; and the internal (fig. 440 *n*), or *endocarp* (*ἐνδον*, within), equivalent to the

Fig. 438.—Cell or loclament of the ovary of *Tribulus terrestris*, cut vertically, to show the commencement of the projections, *c*, from the paries, which are interposed between the ovules, *o*.

Fig. 439.—The same in a mature state, showing the transverse partitions, *c*, dividing the fruit into cavities, in one of which a seed, *g*, is left.

upper epidermis of the leaf, or the epithelium of the ovary. In some plants, as *Colutea arborescens*, the pericarp retains its leaf-like appearance, but in most cases it becomes altered both in consistence and in colour. Sometimes the three parts become blended together, as in the Nut; at other times, as in the Peach, they remain separable. In the latter fruit, the epicarp is thickened by the addition of cells, and can be taken off in the form of what is called the skin; the mesocarp becomes much developed, forming the flesh or pulp, and hence has sometimes been called *sarcocarp* (σαρκξ, flesh), while the endocarp becomes hardened by the production of woody cells, and forms the *stone* or *putamen* (*putamen*, a shell), immediately covering the kernel or the seed. The same arrangement is seen in the fruit of the Cherry, Apricot, and Plum. In these cases, the mesocarp is the part of the fruit which is eaten. In the Almond, on the other hand, the seed is used as food, while the shell or endocarp, with its leathery covering or mesocarp, and its greenish epicarp, are rejected. The pulpy matter found in the interior of fruits, such as the Gooseberry, Grape, and *Cathartocarpus Fistula* (fig. 395), is formed from the placentas, and must not be confounded with the sarcocarp.

525. In the Date, the epicarp is the outer brownish skin, the pulpy matter is the mesocarp or sarcocarp, and the thin papery-like lining is the endocarp covering the hard seed. In the Pear and Apple, the outer skin or epicarp is composed of the epidermis of the calyx, combined with the ovary; the fleshy portion is the mesocarp, formed by the cellular portion of the calyx and ovary; while the scaly layer, forming the walls of the seed-bearing cavities in the centre, is the endocarp. In the Medlar (fig. 472), the endocarp becomes of a stony hardness. In the Melon, the epicarp and endocarp are very thin, while the mesocarp forms the bulk of the fruit, varying in its texture and taste in the external and internal part. The rind of the Orange consists of epicarp and mesocarp, while the endocarp forms partitions in the interior, filled with pulpy cells.

526. Thus, while normally the divisions of the fruit ought to indicate the number of the carpels composing it, and these carpels should each have three layers forming the walls, it is found that frequently the divisions of a multilocular fruit are atrophied or absorbed, in whole or in part, and the layers become confounded together, so that they appear to be one. Again, in fruits formed of several carpels, the




Fig. 440.—Lower portion of the carpel or legume of the Bean, *Faba sativa*, cut transversely, to show the structure of the pericarp. *e*, Epicarp, or external epidermis. *m*, Mesocarp. *n*, Endocarp. *s d*, Dorsal suture. *s v*, Ventral suture. *g*, A seed situated at the upper part of the section, and cut also transversely.

endocarp and mesocarp are occasionally so much developed as to leave the epicarp only on the free dorsal face of the fruit, forming a covering which is wholly external, as in the Castor-oil plant, Euphorbia, and Mallow. Occasionally, the endocarp remains attached to the centre, forming cells, in which the seeds are placed, while the outer layer separates from it at certain points, and leaves a row of cavities in the substance of the pericarp itself.

527. While in many fruits the calyx becomes incorporated with the pericarp, there are others in which it is closely applied to the ovary, but still separable from it. Thus, in the fruit of *Mirabilis Jalapa* (fig. 441, 1), when a section is made longitudinally (fig. 441, 2), the hardened calyx, *c c*, is observed distinct from the fruit, *f*, which is in this instance incorporated with the seed, but at once distinguished by its style, *s*. The same thing occurs in *Spinacia* or Spinach. Again, in *Hippophæ rhamnoides*, and in the Yew (fig. 442), there is an external succulent covering, *i c*, formed by modified bracts, which here occupy the place of a pericarp, and display the seed, *g*, which is naked, because not contained in a true ovary with a style and stigma.



528. The part of the pericarp attached to the peduncle is called its *base*, and the part where the style or stigma existed is the *apex*. This latter is not always the mathematical apex. In *Alchemilla*, *Labiatae*, and *Boraginaceæ*, it is at the base or side (figs. 400, 401, 402). The style sometimes remains in a hardened form, rendering the fruit *apiculate*; at other times it falls off, leaving only traces of its existence. The presence of the style or stigma serves to distinguish certain single-seeded pericarps from seeds.

529. As in the case of the carpel, so in the mature ovary formed of it, the edges unite towards the axis, and constitute the ventral suture (fig. 443 *s v*), while the back, corresponding with the midrib, is the

Fig. 441.—Fruit of *Mirabilis Jalapa*. 1. Entire. 2. Cut longitudinally, to show its composition. *c c*, Lower part of calyx hardened, and forming an outer envelope. *f*, The true fruit, covered by the calyx. The integuments of the fruit are incorporated with those of the seed, which has been also cut. The fruit is distinguished by the remains of the style, *s*, at the apiculus or summit.

Fig. 442.—Fruit of *Taxus baccata*, the Yew. *b*, Imbricated bracts at its base. *i c*, Fleshy envelope taking the place of the pericarp. This envelope covers the seed, *g*, partially, leaving its apex naked.

dorsal suture (fig. 443 *s d*). The inner suture in some fruits formed of a single carpel, as the Apricot and Bladder senna, is marked by a distinct furrow or depression, consequent on the folding inwards of the carpellary edges; and occasionally the outer or dorsal suture is also thus rendered distinctly visible. When the fruit consists of several mature carpels, all meeting in the centre, and united together, then the dorsal suture is also visible externally; but in cases where the placentation is parietal or free central, then the edges of the separate carpels, being near the surface, may present also externally the marks of the ventral sutures.



443

530. Where the sutures are formed, there are usually two bundles of fibro-vascular tissue (fig. 443), one on each edge. The edges of the sutures are often so intimately united, as not to give way when the fruit is ripe. In this case it is called *indehiscent* (*in*, used in the sense of not, and *dehisco*, I open), as in the Acorn and Nut; at other times the fruit opens between the two vascular bundles, either at the ventral or dorsal suture, so as to allow the seeds to escape, and then it is *dehiscent* (*dehisco*, I open). By this *dehiscence* the pericarp becomes divided into different pieces, which are denominated *valves*, the fruit being *univalvular*, *bivalvular*, or *multivalvular*, &c., according as there are one, two, or many valves. These valves separate either completely or partially. In the latter case, the divisions may open in the form of teeth at the apex of the fruit, the dehiscence being *apicular*, as in Caryophyllaceæ (fig. 444 *v*), or as partial slits of the ventral suture, when the carpels are only free at the apex, as in Saxifragæ.



444


531. **Indehiscent Fruits** are either dry, as the Nut, or fleshy, as the Cherry and Apple. They may be formed of one or several carpels; and in the former case they usually contain only a single seed, which may become so incorporated with the pericarp as to appear to be naked. Such fruits are called *pseudo-spermous* ($\psi\epsilon\upsilon\delta\omicron\varsigma$, false, and $\sigma\pi\epsilon\rho\mu\alpha$, seed), or false-seeded, and are well seen in the grain of Wheat. In such cases the presence of the style or stigma determines their true nature.

532. **Dehiscent Fruits**, when composed of single carpels, may open by the ventral suture only, as in the follicles of Pæony; by the dorsal suture only; or by both together, as in the legume of the Pea and

Fig. 443.—A single carpel of *Helleborus foetidus* after dehiscence. *s d*, Dorsal suture. *s v*, Ventral suture. The carpel, when mature, opens on the ventral suture, and forms the fruit denominated a follicle.

Fig. 444.—Capsule or dry seed-vessel of *Cerastium viscosum* after dehiscence, *c*, Persistent calyx. *p*, Pericarp dividing at the apex, *v*, into ten teeth, which indicate the summits of as many valves united below.

Bean; in which cases the dehiscence is called *sutural*. When composed of several united carpels, the valves may separate through the dissepiments, so that the fruit will be resolved into its original carpels, as in *Rhododendron*, *Colchicum*, &c. This dehiscence, in consequence of taking place through the lamellæ of the septum, is called *septicidal* (*septum* and *cado*, I cut) (figs. 445, 446). The valves may separate



from their commissure, or central line of union, carrying the placentas with them, or they may leave the latter in the centre, so as to form

Fig. 445.—Capsule of *Digitalis purpurea* at the moment of dehiscence, when the two cavities *c c*, separate by division of the septum, *d d*, so as to have the appearance of distinct carpels. At the apex are seen the seeds, *g*.

Fig. 446.—Inferior portion of the same capsule cut transversely, to show the formation of the septum, *d*, formed by the two inner faces of the carpels, *c c*. *p p*, Placentaries reflected and projecting into the anterior of the cavities. *g*, Seeds.

Fig. 447.—Capsule (*triccoccus regina*) of *Ricinus communis*, Castor-oil plant, at the moment of dehiscence. The three carpels or cocci, *c c c*, are separated from the axis, *a*, by which they were at first united (see fig. 453), and which remains in a columnar form. These cocci begin to open by their dorsal suture, *s d*.


Fig. 448.—Capsule of *Iris* opening by loculicidal dehiscence.

Fig. 449.—Capsule of *Hibiscus esculentus*, showing also loculicidal dehiscence. *v v v*, Valves of the seed-vessel. *c*, Septum or partition. *g*, Seeds.

Fig. 450.—Capsule of *Cedrela angustifolia*, the valves of which, *v v v*, separate from the septa, *c c*, by septicidal dehiscence. The separation takes place from above downwards, in such a manner that the axis, *a*, remains in the centre, with five projecting angles, corresponding to the septa. *g*, The seeds contained in the locuments.

with the axis a column of a cylindrical, conical, or prismatic shape, which has received the designation of *columella* (fig. 447 c). The union between the edges of the carpels may be persistent, and they may dehisce by the dorsal suture, or through the back of the loculaments, as in the Lily and Iris (fig. 448). In this case the valves are formed by the halves of the cells, and the septa either remain united to the axis, or they separate from it, carrying the placentas with them (fig. 449), or leaving them in the centre. This dehiscence is *loculicidal* (*loculus*, cell, and *cædo*, I cut). Sometimes the fruit opens by the dorsal suture, and at the same time the valves or walls of the ovaries separate from the septa (fig. 450), leaving them attached to the centre, as in *Datura*. This is called *septifragal* dehiscence (*septum* and *frango*, I break), and may be looked upon as a modification of the loculicidal. The separation of the valves takes place either from above downwards (fig. 450), or from below upwards (fig. 451).

533. Sometimes the axis is prolonged as far as the base of the styles, as in the Mallow (fig. 452), and Castor-oil plant (fig. 453), the carpels being united to it by their faces, and separating from it



without opening. In the Umbelliferæ (fig. 454), the two carpels separate from the lower part of the axis, and remain attached to a prolongation of it, called a *carpopophore* (*καρπός*, fruit, and *φέρω*, I bear), or *podocarp* (*πούς*, foot, and *καρπός*, fruit), which splits into two (fig. 454 a), and suspends them. Hence the name *cremocarp* (*κρεμῶω*,

Fig. 451.—Capsule of *Swietenia Mahagoni*, opening by valves from below upwards. The letters have the same signification as in fig. 450.


Fig. 452.—Fruit of *Malva rotundifolia*, with half the carpels comprising it removed, to show the axis, *a*, to which they are attached. This axis ends at the point where the style, *s*, is produced. *c c*, The carpels which are left attached to the axis, around which they are arranged in a verticillate manner. The lateral surface of the two carpels in front, *g*, is exposed.

Fig. 453.—Trilocular capsule of *Ricinus communis*, Castor-oil plant, cut vertically, to show the axis, *a*, prolonged between the carpels, and terminating by small cords or funiculi, *f*, which project into the loculaments, and are attached to seeds. *g g*, Seeds exposed, each surmounted by a fleshy caruncula, *c*. *p p*, Pericarp.

I suspend), applied to this fruit. In Geraniaceæ, the axis is prolonged beyond the carpels, forming a carpophore, to which the styles are attached, and the pericarps separate from below upwards, before dehiscent by their dorsal suture (fig. 455). Carpels of this kind are called *cocci* (κόκκος, seed, berry), and the fruit is said to be *tricocous*, &c., according to the number of separate carpels. In the case of many Euphorbiaceæ, as *Hura crepitans*, the cocci separate with great force and elasticity, the cells being called *dissilient* (*disilio*, I burst).



534. In the Siliqua, or fruit of Cruciferæ, as Wallflower, (fig. 456), the valves separate from the base of the fruit, leaving a central *replum*, or frame, *r*. The replum is considered as being formed by parietal placentas, which remain attached to the fibro-vascular line of the suture, the valves giving way on either side of the suture. In Orchidaceæ (fig. 457), the pericarp, when ripe, separates into three valves, by giving way only on the margins within the sutures, where the placentas are united; and when the valves fall off, the placentas



are left in the form of three arched repla, or frames, to which the seeds are attached. In the case of a free central placenta, when the valves separate, it is sometimes difficult to tell whether the dehiscence

Fig. 454.—Fruit or cremocarp of *Prangos uloptera*, an umbelliferous plant. The carpels, mericarps, or achenia, *cc*, separate from the axis, *a*, and are each suspended by a carpophore. *ss*, Persistent styles with swollen bases, forming an epigynous disk.

Fig. 455.—Fruit or mature carpel of *Geranium sanguineum*. *c*, Persistent calyx. *a*, Axis prolonged as a beak. *st*, The styles, at first united to the beak, and afterwards separating from below upwards, along with the carpels, *cc*, which dehisce by their dorsal suture. *s*, Stigmas. The fruit is sometimes called gynobasic.

Fig. 456.—Siliqua of *Cheiranthus Cheiri*, Wallflower, dehiscent by two valves, *vv*, which separate from a frame or replum, *r*. *g*, Seeds arranged on either margin. *s*, Two-lobed stigma.

Fig. 457.—Capsule of *Orchis maculata* at the period of dehiscence. *c*, Remains of the limb of the calyx crowning the fruit. *vv*, Segments of the pericarp which are detached in the form of valves. *pp*, Arched repla or placentas which remain persistent, and bear the seeds.

is septicial or loculicidal, inasmuch as there are no dissepiments, and the placentas and seeds form a column in the axis. Their number, as well as their alternation or opposition, as compared with the sepals, will aid in determining whether the valves are the entire carpellary leaves, as in septicial dehiscence, or only halves united, as in loculicidal. In some instances, as in *Linum catharticum*, the fruit opens first by loculicidal dehiscence, and afterwards the carpels separate in a septicial manner.

535. Another mode in which fruits open is transversely, the dehiscence in this case being called *circumscissile* (*circum*, around, and *scindo*, I cut). In such cases, the fruit or seed-vessel may be supposed to be formed by a number of articulated leaves like those of the Orange, the division taking place where the laminæ join the petioles.

In this dehiscence, the upper part of the united valves falls off in the form of a lid or operculum, as in *Anagallis* (fig. 458), and in *Hyoscyamus* (fig. 459), and hence the fruit is often denominated *operculate* (*operculum*, a lid). In some instances the axis seems to be prolonged in the form of a hollow cup, and the valves appear as leaves united to it by articulation, similar to what occurs in the calyx of *Eschscholtzia*. In *Lecythis*, or the Monkey-pot, and in *Couratari*, the calyx is adherent to the seed-vessel, and the lid is formed at the place where the tube of the calyx ceases to be adherent.

536. Transverse divisions take place occasionally in fruits formed by a single carpel, as in the pods of some leguminous plants. Examples are met with in *Ornithopus*, *Hedysarum*, (fig. 460), *Coronilla*, &c., in which each seed is contained in a separate division, the partitions being formed by the folding in of the sides of the pericarp, and distinct separations taking place at these partitions, by what has been termed *solubility*. Some look upon these pods as formed by pinnate leaves folded, and the divisions as indicating the points where




Fig. 458.—Pyxidium or capsule of *Anagallis arvensis*, opening by circumscissile dehiscence. *c*, Persistent calyx. *p*, Pericarp divided into two, the upper part, *o*, separating in the form of a lid or operculum. On the capsule are seen three lines passing from the base to the apex, and marking the true valves. *g*, Seeds forming a globular mass round a central placenta.

Fig. 459.—Operculate capsule or Pyxidium of *Hyoscyamus niger*, Henbane. *o*, Operculum or lid separating and allowing the seeds to appear.

Fig. 460.—Lomentaceous legume or lomentum of *Hedysarum coronarium*. 1. Entire, the upper division being nearly detached from the rest. 2. Two of the joints cut longitudinally to show the spurious loculaments, each containing a seed. This seed-vessel divides into separate single-seeded portions by solubility.

the different pairs of pinnæ are united. Others do not admit this explanation, but regard the legume or pod as formed by the expanded midrib or petiole, and the pinnæ as represented by the seeds. Dehiscence may also be effected by partial openings in the pericarp, called *pores*, which are situated either at the apex, base, or side. In



461



462

the Poppy (fig. 409), the opening takes place by numerous pores under the peltate processes bearing the stigmas. In Campanulas, there are irregular openings towards the middle or base (fig. 461 *t*), which pierce both the pericarp and the adherent calyx. In Frogmouth (fig. 462) or Snapdragon, the pericarp gives way at certain fixed points, forming two or three orifices, one of which corresponds to the upper carpel, and the other to the lower. These orifices have a ragged

appearance at the margins, which has given rise to the name *rupturing*, as applied to this mode of dehiscence.

537. **Carpology.**—Much has been done of late in the study of *carpology* (καρπός, fruit, and λόγος, discourse), or the formation of the fruit; but much still remains to be done ere the terminology of this department is complete. Many classifications of fruits have been given, but they are confessedly imperfect, and unfortunately much confusion has arisen in consequence of the same names having been applied to different kinds of fruit. In many cases, therefore, it is necessary to give a description of a fruit in place of using any single term. There are, however, some names in general use, and others which have been carefully defined, to which it is necessary to direct attention.

538. Fruits may be formed by one flower, or they may be the product of several flowers combined. In the former case, they are either *apocarpous* (από, separate, and καρπός, fruit), or *dialycarpous* (διαλύω, I dissolve, or separate), that is, composed of one mature carpel, or of several separate free carpels; or *syncarpous* (σύν, together), that is, composed of several carpels, more or less completely united. These different kinds of fruits may be *indehiscent* (not opening), or *dehiscent* (opening). When the fruit is composed of the ovaries of

Fig. 461.—Capsule of *Campanula persicifolia*, opening by holes or pores, *t, t*, above the middle. *c*, Persistent calyx, incorporated below with the pericarp, *p*, and separating above into five acute segments, in the midst of which is seen the withered and plaited corolla in the form of involucre. *n*, The holes perforate the walls of the pericarp and the calyx.

Fig. 462.—Capsule of *Antirrhinum majus*, Frogmouth, after dehiscence. *c, c*, Persistent calyx. *p*, Pericarp perforated near the summit by three holes, *t, t*, two of which correspond to one of the locuments, and one to the other. The apex of the capsule is acuminate by the remains of the persistent style, *s*.

several flowers united, it is usual to find the bracts and floral envelopes also joined with them, so as to form one mass; hence such fruits are called *multiple* or *anthocarpous* (ἄνθος, flower, and καρπός, fruit). The term *simple* is perhaps properly applied to fruits, which, when mature, appear to be formed of one carpel only; but it has been also given to those which, when mature, are formed by several separate carpels; while the term *compound* is applied to cases where several carpels are combined. The name *aggregate* is by some made synonymous with *anthocarpous*, while *multiple* is applied to apocarpous fruits formed by several free carpels.

Fruits which are the Produce of a Single Flower.

539. Apocarpous Fruits.—These fruits are formed out of one or several free carpels. They are either *dry* or *succulent*; the pericarp, in the former instance, remaining more or less foliaceous in its structure, and sometimes becoming incorporated with the seed; in the latter, becoming thick and fleshy, or pulpy. Some of these do not open when ripe, but fall entire, the pericarp either decaying, and thus allowing the seeds ultimately to escape, as is common in fleshy fruits, or remaining united to the seed, and being ruptured irregularly when the young plant begins to grow; such fruits are *indehiscent*. Other apocarpous fruits, when mature, open spontaneously to discharge the seeds, and are *dehiscent*.

540. Indehiscent Apocarpous Fruits, when formed of a single mature carpel, frequently contain only one seed, or are *monospermous* (μόνος, one, and σπέρμα, seed). In some instances there may have been only one ovule originally, in others two, one of which has become abortive.

541. The Achænium (α, privative, and χαίνω, I open), is a dry monospermous fruit, the pericarp of which is closely applied to the seed, but separable from it (fig. 463). It may be *solitary*, forming a single fruit, as in the Cashew (fig. 227 a), where it is supported on a fleshy peduncle, *p*; or *aggregate*, as in *Ranunculus* (fig. 464), where several achænia are placed on a common elevated receptacle. In the Strawberry, the achænia are placed on a convex succulent receptacle. In the Rose, they are supported on a concave receptacle, covered by the calycine tube (fig.




Fig. 463.—Achænium or indehiscent monospermous carpel from the pistil of a *Ranunculus*.

Fig. 464.—1. Similar achænium, with rough points on the pericarp, from the pistil of *Ranunculus muricatus*. 2. Achænium cut transversely to show the seed, *g*, not adherent to the parietes.

270), and in the Fig, they are placed inside the hollow peduncle or receptacle (fig. 246), which ultimately forms what is commonly called the fruit. In the Rose, the aggregate achænia, with their general covering, are sometimes collectively called *Cynarrhodum* (κυνᾶν, a dog, and ῥόδον, a rose, seen in the dog-rose). It will thus be remarked, that what in common language are called the seeds of the Strawberry, Rose, and Fig, are in reality carpels, which are distinguished from seeds by the presence of styles and stigmas. The styles occasionally remain attached to the achænia, in the form of feathery appendages, as in Clematis, where they are called *caudate* (*cauda*, a tail).

542. In Compositæ, the fruit which is sometimes called *Cypsela* (κυψέλη, a box), when ripe, is an achænium united with the tube of the calyx (fig. 279 *t*). The limb of the calyx in the Compositæ sometimes becomes pappose, and remains attached to the fruit, as in Dandelion, Thistles, &c. (fig. 279 *l*). When the pericarp is thin, and appears like a bladder surrounding the seed, the achænium becomes a *Utricle*, as in Amaranthaceæ. This name is often given to fruits which differ from the achænium, in being composed of more than one carpel. When the pericarp is extended in the form of a winged appendage, a *Samara* (*samera*, seed of Elm) or *samaroid achænium* is produced, as in the Ash (fig. 437), common Sycamore (fig. 465), and Hiræa (fig. 466). In these cases, there are usually two achænia united,



one of which, however, as in *Fraxinus oxyphylla* (fig. 437), may be

Fig. 465.—Seed-vessel of *Acer Pseudo-platanus*, composed of two samaras or winged mono-sperous carpels united. *a*, Upper part forming a dorsal wing. *l*, Lower portion corresponding to the loculament.

Fig. 466.—Samara taken from the fruit of *Hiræa*. *s*, Persistent style. *l*, Part corresponding to the loculament. *a*, Marginal wing or ala.

Fig. 467.—Caryopsis of *Secale cereale*, Rye. 1. Entire. 2. Cut transversely to show the seed adherent to the parietes of the pericarp.

abortive. The Wing (fig. 465 *a*) is formed by the carpel, and is either *dorsal*, i.e. a prolongation from the median vein (fig. 465 *a*), or *marginal*, that is, formed by the lateral veins (fig. 466 *a*). It surrounds the fruit longitudinally in the Elm. When the pericarp becomes so incorporated with the seed as to be inseparable from it, as in Grains of Wheat, Maize, Rye (fig. 467), and other grasses, then the name *Caryopsis* (καρύα, a nut, and ὄψις, appearance) is given.

543. There are some fruits which consist of two or more achænia, at first united together, but which separate when ripe. Of this nature is the fruit of the Tropæolum or Indian Cress, also that of Labiatae and Boraginaceæ, which is formed of four achænia attached to the axis (fig. 402), whence the common style appears to proceed. Some of these are occasionally abortive. In the ripe state the pericarp separates from the seed in these cases; and thus there is a transition from indehiscent achænia to single-seeded dehiscent pericarps. So also the *Cremocarp* (κρεμάω, to suspend), or the fruit of Umbelliferæ (fig. 454), which is composed of two achænia united by a commissure to a common axis or *carpophore* (καρπός, fruit, and φέρω, I bear), from which they are suspended at maturity. It is sometimes denominated *diachænium* (δις, twice), from the union of two achænia, which in this instance receive the name of *mericarps* (μέρος, part), or *hemicarps* (ἡμισυς, half, and καρπός, fruit).

544. *The Nut or Glans*. This is a one-celled fruit with a hardened pericarp, surrounded by bracts at the base, and, when mature, containing only one seed. In the young state, the ovary contains two or more ovules, but only one comes to maturity. It is illustrated by the fruit of the Hazel and Chestnut, which are covered by leafy appendages, in the form of a *husk*, and by the Acorn, in which the leaves or bracts are united so as to form a *cupula* or *cup* (fig. 257). The parts of the pericarp of the Nut are united so as to appear one. In Sagus, or the Sago Palm, it is covered by peculiar closely applied scales, giving the appearance of a cone.

545. *The Drupe* (*drupæ*, unripe olives). This is a succulent fruit covered by a pericarp, consisting of epicarp, mesocarp, and endocarp; and when mature, containing a single seed. This term is applied to such fruits as the Cherry, Peach, Plum, Apricot, Mango, Walnut, Nutmeg, and Date. The endocarp is usually hard, forming the stone of the fruit, which encloses the kernel or seed. The mesocarp is generally pulpy and succulent, so as to be truly a *sarcocarp* (Peach), but it is sometimes of a tough texture, as in the Almond, and at other times more or less fibrous. There is thus a transition from the Drupe to the Nut. Moreover, in the Almond, there are often two ovules formed, only one of which comes to perfection. In the Walnut, the endocarp, which is easily separable into two, forms prolongations which enter into the interior, and cause a remarkable division in the seed.

It has been sometimes called *Tryma*. In the Raspberry and Bramble, several small drupes or *drupels* are aggregated so as to constitute an *Etaerio* (ἑταῖρος, a companion).

546. **Dehiscent Apocarpous Fruits.**—These open in various ways, and usually contain more than one seed, being either few-seeded, *oligospermous* (ὀλίγος, few, and σπέρμα, a seed), or many-seeded, *poly-spermous* (πολύς, many).




468

547. **Follicle** (*folliculus*, a little bag). This is a mature carpel, containing several seeds, and opening by the ventral suture (figs. 443, 468). It is rare to meet with a solitary follicle forming the fruit. There are usually several aggregated together, either in a circular manner on a shortened receptacle, as in Hellebore, Aconite, Delphinium, and Asclepiadaceæ; or in a spiral manner on an elongated receptacle, as in Magnolias, Banksias, and Liriodendron (fig. 306). Occasionally in Magnolia grandiflora, some of the follicles open by the dorsal suture.

548. **The Legume or Pod** (*legumen*, pulse) is a solitary, simple, mature carpel, dehiscing

by the ventral and dorsal suture, and bearing seeds on the former. It characterizes leguminous plants, and is seen in the Bean and Pea (fig. 469). In the Bladder-senna (fig. 470) it retains its leaf-like appearance, and forms an inflated legume. In some Leguminosæ, as *Arachis*, the fruit must be considered a legume, although it does not dehisce. In place of opening at the sutures, some legumes are contracted at intervals, so



469



470

Fig. 468.—Follicle or dehiscent many-seeded carpel of *Aquilegia vulgaris*, Columbine. The follicle dehisces by the ventral suture only.

Fig. 469.—Legume of *Pisum sativum*, common Pea, opened. It is formed by a single carpel, and dehisces by the ventral and dorsal suture. *v v*, Valves formed by the two parts of the pericarp. *p*, The epicarp or external layer of the pericarp. *p'*, Endocarp or internal layer. Between these the mesocarp is situated. *g*, Seeds placed one over the other, attached to the placenta by short funiculi or cords, *ff*. The placenta forms a narrow line along the ventral suture, *s v*. *s d*, The dorsal suture corresponding to the midrib of the carpellary leaf.

Fig. 470.—Legume of Bladder-senna (*Colutea arborescens*), showing an inflated, foliaceous pericarp.

as to include each seed in a separate cell, and when ripe, the different divisions of the pod separate from each other. This constitutes the *Lomentum* (*lomentum*, bean-meal), or *lomentaceous legume* of *Hedysarum coronarium* (fig. 460), *Coronillas*, *Ornithopus*, &c. In *Medicago*, the legume is twisted like a snail (fig. 471), and in *Cæsalpinia coriaria*, or *Divi-divi*, it is vermiform or curved like a worm; in *Carmichaelia*, the valves give way close to the suture, and separate from it, leaving a division.


549. **Syncarpous Fruits** are formed by several carpels, which are so united together as to appear one in their mature state. These fruits are either dry or succulent; in the former case, being usually dehiscent, in the latter, indehiscent.

550. **Indehiscent Syncarpous Fruits.**—*The Berry (bacca)* is a succulent fruit, in which the seeds are immersed in a pulpy mass, formed by the placentas. The name is usually given to such fruits as the Gooseberry and Currant, in which the calyx is adherent to the ovary, and the placentas are parietal, the seeds being ultimately detached from the placenta, and lying loose in the pulp. Others have applied it also to those in which the ovary is free, as in the Grape, Potato, and *Ardisia*, and the placentas central or free central. The latter might be separated under the name *Uva* (grape). In general, the name of *baccate* or *berried* is applied to all pulpy fruits. In the Pomegranate there is a peculiar baccate many-celled fruit, having a tough rind formed by the calyx, enclosing two rows of carpels placed above each other. The seeds are immersed in pulp, and are attached irregularly to the parietes, base, and centre. The fruit has been called *Balausta* (*balaustium*, flower of pomegranate), and the tough rind is called *malicorium* (a name applied to it by Pliny).

551. The *Pepo* or *Peponida* (πέπων, a pumpkin), is illustrated by the fruit of the Gourd, Melon, and other Cucurbitaceæ, where the calyx is adherent, the rind is thick and fleshy, and there are three or more seed-bearing parietal placentas, either surrounding a central cavity, or sending prolongations inwards. The fruit of the Papaw resembles the Pepo, but the ovary is not adherent to the calyx.


552. The *Hesperidium* (golden fruit in the garden of Hesperides) is the name given to the fruit of the Orange, &c., in which the epicarp and mesocarp form a separable rind, and the endocarp sends prolongations inwards, forming triangular divisions, in which pulpy cells are developed so as to surround the seeds which are attached to the inner angle. Both Pepo and *Hesperidium* may be considered as modifications of the Berry.

553. The *Pome* (*pomum*, an apple) seen in the Apple, Pear, Quince,



471

&c., is a fleshy fruit with the calyx adherent, and forming along with the epicarp and mesocarp a thick cellular mass, which is eatable, while the endocarp is scaly or horny, and forms separate cells enclosing the seeds. The covering of the cells is sometimes stony, as in the Medlar (fig. 472), and the Holly, forming what has been called a *Nuculanium* (*nucula*, a nut). In the Medlar, the stony endocarps are called *pyrenæ* (πυρήν, the stone of the fruit). In *Cornus mas* (fig. 473), there are



two stony cells, *n*, surrounded by the fleshy epicarp and mesocarp, and as they are close together, and one is often abortive (fig. 473 2 *l*), there is a direct transition to the Drupe.

554. **Dehiscent Syncarpous Fruits.**—The *Capsule* (*capsula*, a little chest). This name is applied generally to all dry syncarpous fruits, which open by valves or pores. The valvular capsule is observed in *Digitalis* (fig. 445), *Hibiscus esculentus* (fig. 449), *Cedrela angustifolia* (fig. 450), *Mahogany* (fig. 451), and *Cerastium viscosum* (fig. 444). The porous capsule is seen in the Poppy, *Antirrhinum majus* (fig. 462), and *Campanula persicifolia* (fig. 461). Sometimes the capsule opens by a lid, or by circumscissile dehiscence, and it is then called a *Pyxidium* (*pyxis*, a box), as in *Anagallis arvensis* (fig. 458), *Henbane* (fig. 459), and *Lecythis*. The capsule assumes a spiral form in the *Helicteres*, and a star-like or stellate form in *Illicium anisatum*. In certain instances, the cells of the capsule separate from each other, and open with elasticity to scatter the seeds. This kind of capsule is met with in *Hura crepitans*, and other *Euphorbiaceæ*, where the cocci, containing each a single seed, burst asunder with force (fig. 453); and in *Geraniaceæ*, where the cocci containing more than one seed,* separate from the carpophore, and become curved upwards by their adherent

Fig. 472.—Fruit of common Medlar (*Mespilus germanica*). Transverse section showing, *e*, epicarp. *s*, Sarcocarp. *n*, Endocarp forming stony coverings of the seeds. The fruit has been called *nuculanium*, and the hard central cells *pyrenæ*.

Fig. 473.—Fruit of *Cornus mas*, common Cornel. 1. Transverse section detaching the upper half of the fleshy portion, *s*, so as to show the central kernel, *n*. 2. Transverse section of the fruit through the central portion, *n*, showing that it consisted of two locuments. *l*, One of the locuments empty, the other containing a seed, *g*.

* The individual cocci of *Geraniaceæ* contain only one seed each.

styles (fig. 455). In the former case, the fruit collectively has been called *Regma* (ῥήγμα, a rupture).

555. The *Siliqua* (*siliqua*, a husk or pod) (fig. 456), may be considered as a variety of the capsule, opening by two valves; these are detached from below upwards, close to the sutures, bearing thin parietal placentas, which are united together by a prolongation called a *replum*, or spurious dissepiment, dividing the fruit into two. The seeds are attached on either side of the replum, either in one row or in two. When the fruit is long and narrow, it is called *Siliqua*; when broad and short, it is called *Silicula*. It occurs in cruciferous plants, as Wallflower, Cabbages, Cresses, &c. The siliqua may be considered as formed of two carpels, and two parietal placentas united together so as to form a two-celled seed-vessel. Some say that in its normal state it consists of four carpels, and that two of these are abortive. There are four bundles of vessels in it, one corresponding to each valve, which may be called *valvular* or *pericarpial*, and others running along the edge called *placental*. The replum consists of two lamellæ. It sometimes exhibits perforations, becoming *fenestrate* (*fenestra*, a window). At other times its central portion is absorbed, so that the fruit becomes one-celled.

Fruits which are the produce of several Flowers united.

556. It sometimes happens that the ovaries of two flowers unite so as to form a double fruit. This may be seen in many species of Honeysuckle. But the fruits which are now to be considered, consist usually of the floral envelopes, as well as the ovaries of several flowers united into one, and are called *Multiple* or *Anthocarpous*.

557. The *Sorosis* (σωρός, a congeries or cluster) is a multiple fruit formed by a united spike of flowers, which becomes succulent. The fruit of the Pine-apple (fig. 474) is composed of numerous ovaries, floral envelopes, and bracts combined so as to form a succulent mass. The scales outside, *c c*, are the modified bracts and floral leaves, which, when the development of the fruit-bearing spike terminates, appear in the form of ordinary leaves, and constitute the crown, *f*. Other instances of a sorosis are the Bread-fruit and Jack-fruit. Sometimes a fruit of this kind resembles that formed by a single flower, and a superficial observer might have some difficulty in marking the difference. Thus, the Strawberry, Mulberry, and



Fig. 474.—Anthocarpous fruit of *Ananassa sativa*, Pine-apple. Axis bearing numerous flowers, the ovaries of which are combined with the bracts, *c c*, to form the fruit. *f*, Crown of the Pine-apple, consisting of empty bracts or floral leaves.

Raspberry appear to be very like each other, but they differ totally in their structure. The Strawberry and Raspberry are each the produce of a single flower, the former being a succulent edible receptacle bearing achenia on its convex surface; the latter being a collection of



475

drupes placed on a conical unpalatable receptacle; while the Mulberry (fig. 475) is a sorosis formed by numerous flowers united together, the calyces becoming succulent, and investing the pericarps.

558. *Syconus* (σῦκον, a fig,) is an anthocarpous fruit, in which the axis, or the extremity of the peduncle, is hollowed, so as to bear numerous flowers, all of which are united in one mass to form the fruit. The Fig (fig. 246) is of this nature, and what are called its seeds are the achenia or monospermal seed-vessels of the numerous flowers scattered through the pulpy hollowed axis. In *Dorstenia* (fig. 245), the axis is less deeply hollowed, and of a harder texture, the fruit exhibiting often very anomalous forms.

559. *Strobilus* (στέβιλος, fir-cone,) is a fruit-bearing spike more or less elongated, covered with scales, each of which represents separate flowers, and has two seeds at its base (fig. 476). The scales may be considered as bracts, or as flattened carpellary leaves, and the seeds are naked, as there is no true ovary present with its style or stigma. This fruit is seen in the cones of Firs, Spruces, Larches, Cedars, &c., which have received the name of Coniferae, or cone-bearing, on this account. The scales of the strobilus are sometimes membranous and thin,



476



477



478

as in the Hop; at other times they are thick and closely united, so as to form a more or less angular and rounded mass, as in the Cypress (fig. 477); while in the Juniper they become fleshy, and are so incorporated as to form a globular fruit like a berry (fig. 478), which has received the name of *Galbulus* (*galbulus*, nut of the cypress).

Fig. 475.—Anthocarpous fruit of the Mulberry, formed by the union of several flowers.

Fig. 476.—Cone of *Pinus sylvestris*, Scotch Fir, consisting of numerous bracts or floral leaves, each of which covers two winged seeds. These seeds are called naked, in consequence of not being contained in an ovary, with a style or stigma.

Fig. 477.—Cone of *Cupressus sempervirens*, Cypress; one of the Gymnospermous or naked-seeded plants, like the Pine.

Fig. 478.—Succulent cone or Galbulus of *Juniperus macrocarpa*. e e e e, The different scales or bracts united so as to enclose the seeds.

560.—TABULAR ARRANGEMENT OF FRUITS.

A. Fruits formed from a single flower, and consisting of one or more Carpels either separate or combined; thus including Apocarpous, Aggregate, and Syncarpous Fruits.

I. Indehiscent Pericarps.

1. Usually containing a single seed:

Covered by a dry simple Pericarp.	Separable from the seed.....	{ Achaenium (<i>Lithospermum</i>). <i>Mericarp</i> and <i>Cremocarp</i> in Umbelliferae, and <i>Cypsela</i> in Compositae.
	Achaenia enclosed in fleshy tube of Calyx, <i>Cynarrhodum</i> (Rose).	
	Inseparable from the seed.....	Caryopsis (Grasses).
	Inflated	Utricle (<i>Chenopodium</i>).
	Having a cupulate involucre.....	Glans (Acorn).
Covered by a Pericarp, consisting of Epicarp, Sarcocarp, and Endocarp.....	Having winged appendages.....	Samara (<i>Sycamore</i>).
	Drupe, with a two-valved Endocarp, having divisions extending from its inner surface, <i>Tryma</i> (Walnut).	Drupa (Cherry).
	Aggregate Drupes, <i>Etaerio</i> (Raspberry).	

2. Containing two or more seeds:

Enclosed in a fleshy or pulpy mass.	{ Ovary adherent to Calyx, Placenta parietal, attachment of seeds lost when ripe.....	Bacca (Gooseberry).
		attachment permanent, rind thick and hard.....
	{ Peculiar berried many-celled fruit, with two or more rows of Carpels.....	Pepo (Gourd).
		Balausta (Pomegranate).
	{ Ovary not adherent to Calyx, Placenta central.....	Uva (Grape).
		Placenta parietal.....
	{ Having a spongy separable rind, and separable pulpy cells.....	Papaw fruit.
		Hesperidium (Orange).
	{ Walls of cells or Endocarp horny, covered by a fleshy Mesocarp and Epicarp.....	Pomum (Apple).
		Nuculanium (Medlar).

II. Dehiscent Pericarps.

Seeds in a dry Pericarp.	Opening by Ventral Suture only		Follicle (<i>Pæony</i>).
	Opening by Ventral and Dorsal Suture		Legume (<i>Pea</i>).
	<i>Lomentum</i> , a Legume separating into distinct pieces, each containing a seed (<i>Ornithopus</i>).		
	{ Opening by two valves which separate from a Central Replum or Frame		Siliqua (<i>Cabbage</i>).
		Silicula	(<i>Capsella</i>).
	Opening by Transverse or Circumscissile Dehiscence.....		Pyxidium (<i>Henbane</i>).
	{ Opening by several valves or pores, without Ventral or Dorsal Suture or Replum.....		Capsule (<i>Poppy</i>).
	Capsule adherent to Calyx.....		Diplogegia (<i>Campanula</i>).
	A long pod-like Capsule		Ceratium (<i>Glaucium</i>).
	Opening by separation of elastic Cocci.....		Regma (<i>Hura</i>).

B. Fruits formed by the union of several Flowers, and consisting of Floral Envelopes, as well as Ovaries; these are Multiple or Anthocarpous.

Hollow Anthocarpous Fruit.—*Syconus* (Fig).

Convex Anthocarpous Fruit.	{	formed by Indurated Catkin.— <i>Strobilus</i> (Fir Cone).
		formed by Succulent Spike.— <i>Sorosis</i> (Bread-fruit).

7.—MATURATION OF THE PERICARP.

561. After fertilization, the parts of the ovary begin to swell, the foramen of the ovule is more or less closed, the stigma becomes dry, and the style either withers and falls off, or remains attached as a hardened process or apiculum; while the embryo plant is developed in the ovule. It has been stated that fruits, such as Oranges and Grapes, are sometimes produced without seeds. It does not appear, therefore, necessary for the production of fruit in all cases, that the process of fertilization should be complete. In speaking of seedless Oranges, Dr. Bullar states that the thinness of the rind of a St. Michael Orange, and its freedom from pips, depend on the age of the tree. The young trees, when in full vigour, bear fruit with a thick pulpy rind and abundance of seeds; but as the vigour of the plant declines, the peel becomes thinner, and the seeds gradually diminish in number, till they disappear altogether.

562. While the fruit enlarges, the sap is drawn towards it, and a great exhaustion of the juices of the plant takes place. In Annuals this exhaustion is such as to destroy the plants; but if they are prevented from bearing fruit, they may be made to live for two or more years. Perennials, by acquiring increased vigour, are able better to bear the demand made upon them during fruiting. If large and highly-flavoured fruit is desired, it is of importance to allow an accumulation of sap to take place before the plant flowers. When a very young plant is permitted to do so, it seldom brings fruit to perfection. When a plant produces fruit in very large quantities, gardeners are in the habit of thinning it early, in order that there may be an increased supply of sap to that which remains. In this way, Peaches, Nectarines, Apricots, &c., are rendered larger and better flavoured. When the fruiting is checked for one season, there is an accumulation of nutritive matter, which has a beneficial effect on the subsequent crop.

563. The pericarp is at first of a green colour, and performs the same functions as the other green parts of plants, decomposing carbonic acid under the agency of light, and liberating oxygen. Saussure asserts that all fruits in a green state are adequate to, and perform this process of deoxidation. As the pericarp advances to maturity, it either becomes dry or succulent. In the former case, it changes into a brown or a white colour, and has a quantity of ligneous matter deposited in

its substance, so as to acquire sometimes great hardness, when it is incapable of performing any active process of vegetable life; in the latter, it becomes fleshy in its texture, and assumes various bright tints, as red, yellow, &c. In fleshy fruits, however, there is frequently a deposition of ligneous cells in the endocarp, forming the stone of the fruit; and even in the substance of the pulpy matter or sarcocarp, there are found isolated cells of a similar nature, as in some varieties of Pear, where they cause a peculiar grittiness. The contents of the cells near the circumference of succulent fruits are thickened by exhalation, and a process of endosmose goes on, by which the thinner contents of the inner cells pass outwards and thus cause swelling of the fruit. As the fruit advances to maturity, however, this exhalation diminishes, the water becoming free, and entering into new combinations. In all pulpy fruits which are not green there are changes going on by which carbon is separated in combination with oxygen.

564. Dry fruits may remain attached to the tree for some time before they are fully ripe, and ultimately separate by disarticulation. Occasionally, when the pericarp is thick, it separates in layers like the bark. Succulent fruits contain a large quantity of water, along with cellulose or lignine, sugar, gummy matter or dextrine, albumen, colouring matter, various organic acids, as citric, malic, and tartaric, combined with lime and alkaline substances, besides a pulpy gelatinous matter, which is converted by acids into pectine or pectose, whence pectic acid is formed by the action of albumen. Pectine is soluble in water, and exists in the pulp of fruits, as Apples, Pears, Gooseberries, Currants, Raspberries, Strawberries, &c. Pectic acid is said to consist of $C^{14} H^3 O^{12} + HO$. It absorbs water, and is changed into a jelly-like matter; hence its use in making preserves. Each kind of fruit is flavoured with a peculiar aromatic substance. Starch is rarely present in the pericarp of the fruit, although it occurs commonly in the seed. In Plantains, Bananas, and Bread-fruit, however, especially when seedless, there is a considerable quantity of starchy matter, giving rise to mealiness when these fruits are prepared as fritters. Oily matters are also found in the cellular tissue of many fruits. Thus, a fixed oil occurs in the Olive, and essential oils in the Orange, Lemon, Lime, Rue, Dictamnus, &c.

565. During ripening, much of the water disappears, while the cellulose or lignine, and the dextrine, are converted into sugar. Berard is of opinion that the changes in fruits are caused by the action of the oxygen of the air. Freney found that fruits covered with varnish did not ripen. As the process of ripening becomes perfected the acids combine with alkalies, and thus the acidity of the fruit diminishes, while its sweetness increases. The formation of sugar is by some attributed to the action of organic acids on the vegetable constituents, gum, dextrine, and starch; others think that the cellulose and lignine

are similarly changed by the action of acids. The formation of sugar is said to be prevented by watering the tree with alkaline solutions. In the Grape, when young, there is abundance of tartaric acid; but as the fruit advances to maturity, this combines with potash, so as to diminish the acidity. Certain fruits owe their astringent qualities to the saline matter which they contain. In seasons when there is little sun, and a great abundance of moisture, succulent fruits become watery, and lose their flavour. The same thing frequently takes place in young trees with abundance of sap, and in cases where a large supply of water has been given artificially.

566. The following analysis of the Cherry in its unripe and ripe state, as given by Berard, exhibits generally the chemical composition of succulent fruits:—

	Unripe.		Ripe.
Chlorophylle.....	0·05	—
Sugar.....	1·12	18·12
Gum or dextrine.....	6·01	3·23
Cellulose.....	2·44	1·12
Albumen.....	0·21	0·57
Malic Acid.....	1·75	2·01
Lime.....	0·14	0·10
Water.....	88·28	74·85
	<hr/> 100·00		<hr/> 100·00

The following table shows the changes produced on the water, sugar, and cellulose, in 100 parts of unripe and ripe fruits:—

	Water.			Sugar.			Cellulose.	
	Unripe.	Ripe.		Unripe.	Ripe.		Unripe.	Ripe.
Apricot.....	89·39	74·87	6·64	16·48	3·61	1·86
Peach.....	90·31	80·24	0·63	11·61	3·01	1·21
Cherries.....	88·28	74·85	1·12	18·12	2·44	1·12
Plums.....	74·87	71·10	17·71	24·81	1·26	1·11
Pears.....	86·28	83·88	6·45	11·52	3·80	2·19

567. It is not easy in all cases to determine the exact time when the fruit is ripe. In dry fruits, the period immediately before dehiscence is considered as that of maturation; but, in pulpy fruits, there is much uncertainty. It is usual to say that edible fruits are ripe, when their ingredients are in such a state of combination as to give the most agreeable flavour. This occurs at different periods in different fruits. After succulent fruits are ripe in the ordinary sense, so as to be capable of being used for food, they undergo further changes, by the oxidation of their tissues, even after being separated from the plant. In some cases, these changes improve the quality of the fruit, as in the case of the Medlar, the austerity of which is thus still further diminished. In the Pear, this process, called by Lindley *bletting* (from the French, *blessi*), renders it soft, but still fit for food; while in the

Apple, it causes a decay which acts injuriously on its qualities. By this process of oxidation, the whole fruit is ultimately reduced to a putrefactive mass, which probably acts beneficially in promoting the germination of the seeds when the fruit drops on the ground.

568. The period of time required for ripening the fruit, varies in different plants. Most plants ripen their fruit within a year from the time of the expansion of the flower. Some come to maturity in a few days, others require some months. Certain plants, as some Coniferæ, require more than a year, and in the *Metrosideros*, the fruit remains attached to the branch for several years. The following is a general statement of the usual time required for the maturation of different kinds of fruit:—

Grasses	13 to 45 days.
Raspberry, Strawberry, Cherry.....	2 months.
Bird-cherry, Lime-tree.....	3 —
Roses, White-thorn, Horse-chestnut,.....	4 —
Vine, Pear, Apple, Walnut, Beech, Plum, Nut, Almond, 5 to 6	—
Olive, Savin.....	7 —
Colchicum, Mistletoe.....	8 to 9 —
Many Coniferæ.....	10 to 12 —
Some Coniferæ, certain species of Oak, <i>Metrosideros</i> , above 12	—

The ripening of fruits may be accelerated by the application of heat, by placing dark-coloured bricks below it, and by removing a ring of bark so as to lead to an accumulation of sap. It has been observed that plants subjected to a high temperature not unfrequently prove abortive, which seems to result from the over stimulation causing the production of unisexual flowers alone. Trees are sometimes made to produce fruit, by checking their roots when too luxuriant, and by preventing the excessive development of branches.

569. **Grafting.**—A very important benefit is produced, both as regards the period of fruiting and the quality of the fruit, by the process of grafting. This is accomplished by taking a young twig or scion, called a *graft*, and causing it to unite to a vigorous stem or *stock*, thus enabling it to derive a larger supply of nutritive matter than it could otherwise obtain, and checking its vegetative powers. In place of a slip or cutting, a bud is sometimes taken. In order that grafting may be successfully performed, there must be an affinity between the graft and the stock as regards their sap, &c. It has often been supposed that any kinds of plants may be grafted together, and instances are mentioned by Virgil and Pliny, where different fruits are said to have been borne on the same stock. This was probably produced by what the French call *Greffe des charlatans*,—cutting down a tree within a short distance of the ground, and then hollowing out the stump, and planting within it several young trees of different species; in a few years they grow up together so as to fill up the cavity, and appear to

be one. The deception is kept up better, if some buds of the parent stock have been kept alive.

570. The object which gardeners wish to secure by grafting, is the improvement of the kinds of fruit, the perpetuation of good varieties, which could not be procured from seed, and the hastening of the period of the fruit-bearing. Grafting a young twig on an older stock, has the effect of making it flower earlier than it would otherwise do. The accumulation of sap in the old stock is made beneficial to the twig, and a check is given at the same time to its tendency to produce leaves. Plants that are grafted the one on the other must, if not botanically allied, have a close similarity in the composition of their sap. This general law appears to be departed from in the case of some parasites. The Mistletoe has been found on plants belonging to the Rosaceæ, Leguminosæ, Coniferæ, and other families. In such instances, however, the seed derives its life from the plant on which it grows, and was not perfected in its vitality before being brought to its position.

571. Mr. Knight did much to improve fruits by grafting. He believed, however, that a graft would not live longer than the natural limit of life allowed to the tree from which it has been taken. In this way he endeavoured to account for the supposed extinction of some valuable varieties of fruits, such as the Golden pippin, and many cider apples of the seventeenth century.* He conceived that the only natural method of propagating plants was by seed. His views have not been confirmed by physiologists. Many plants are undoubtedly propagated naturally by shoots, buds, tubers, &c., as well as by seed; and it is certain that the life of slips may be prolonged by various means, much beyond the usual limit of the life of the parent stock. The Sugar-cane is propagated naturally by the stem, the Strawberry by runners, the Couch-grass by creeping stems, Potatoes and Jerusalem Artichokes by tubers, the Tiger-lily by bulblets, and Achimenes by scaly bodies, like tubers. The fruits, moreover, which Mr. Knight thought had disappeared, such as Red streak, Golden pippin, and Golden Harvey, still exist, and any feebleness that they exhibit does not appear to proceed from old age, but seems to be owing to other causes, such as the nature of the soil, cold, violence, and mutilation. Vines have been transmitted by perpetual division from the time of the Romans. A slip taken from a Willow in Mr. Knight's garden, pronounced by him as dying from old age, was planted in the Edinburgh Botanic Garden many years ago, and is now a vigorous tree, although the original stock has long since undergone decay. It is true, however, that a cutting taken from a specimen already exhausted by excessive development of its parts, will partake of the impaired vigour of its parent, and will possess less constitutional energy than that taken from a vigorous stock.

572. In grafting, various methods have been adopted. One of these

* See Knight's Horticultural Papers, 8vo, London, 1841, p. 81.

is *grafting by approach*, or *inarching*, when two growing plants are united together, and after adhesion one is severed from its own stock, and left to grow on the other. This kind of adhesion sometimes takes place naturally in trees growing close together. It is well seen in a fir-tree in the burying-ground at Killin. The branch of the same tree may also be bent, so as to become united to the stem at two points. This is often seen in the Ivy. The roots of contiguous trees occasionally unite by a process of grafting, and to this is attributed the continued vigour of the stump of Spruce-trees cut down on the Swiss mountains. This natural grafting of roots has been observed in the White Pine (*Abies pectinata*), and sometimes in the Red Pine (*Abies excelsa*), as well as in the Scotch Fir (*Pinus sylvestris*).

573. The usual method of grafting is by *scions* or *slips*, which are applied to the stock by a sloping surface, or are inserted into slits in it by *cleft-grafting*, or into perforations by *wimble* or *peg-grafting*. Sometimes several slips are placed in a circular manner round the inside of the bark of the stock, by *crown-grafting*; or the bark of a portion of the stock is removed, and that of the scion is hollowed out, so as to be applied over it like the parts of a flute, hence called *flute-grafting*. *Budding* is practised by the removal of a bud from one plant, along with the portion of the bark and new wood, and applying it to another plant, in which a similar wound has been made. Grafting is usually performed between the woody parts of the plants, but herbaceous parts may also be united in this way. It is requisite that the growing parts be brought into apposition — the two albumens and the two libers. Union will only take place where the active process of life are freely exercised. The graft and stock are secured together by means of clay, or a mixture of bees'-wax and tallow, or by bits of Indian rubber.

574. By grafting, all our good varieties of apples have been produced from the Crab Apple. The seeds of the cultivated apples, when sown, produce plants which have a tendency to revert to the original sour Crab. Grafted varieties can only be propagated by cuttings. The influence exercised by the stock is very marked, and it is of great importance to select good stocks on which to graft slips. In this way the fruit is often much improved by a process of *ennobling*, as it is called. The scion also seems in some cases to exercise a remarkable effect on the stock. Slips taken from varieties with variegated leaves, grafted on non-variegated, have caused the leaves of the latter to assume variegation, and the effect, when once established, has continued even after the slip was removed. The effects of grafting are well seen in the case of the Red Laburnum, when united to the Yellow species. The Red Laburnum is a hybrid between the common Yellow Laburnum and *Cytisus purpureus*, or the Purple Laburnum. The branches below the graft produce the ordinary yellow laburnum

flowers of large size; those above exhibit often the small purple laburnum flowers, as well as reddish flowers intermediate between the two in size and colour. Occasionally, the same cluster has some flowers yellow and some purplish.

8.—SEED OR FERTILIZED OVULE ARRIVED AT MATURITY.

575. While the pistil undergoes changes consequent on the discharge of the pollen on the stigma, and ultimately becomes the fruit, the ovule also is transformed, and, when fully developed, constitutes the *seed*. After fertilization, the foramen of the ovule contracts, the embryo or young plant gradually increases in its interior, by the absorption of the fluid matter contained in the sac of the amnios, solid nutritive matter is deposited, and a greater or less degree of hardness is acquired. The seed then is the fecundated mature ovule containing the embryo, with certain nutritive and protective appendages. When ripe, the seed contains usually a quantity of starchy and ligneous matter, various azotised compounds, as caseine, vegetable albumen, oily and saline matters. It sometimes acquires a stony hardness, as in the case of vegetable ivory, the seed of *Phytolophos macrocarpa*. Care must be taken not to confound it with single-seeded pericarps, such as the *Achæmium* and *Caryopsis*, in which a style and stigma are present; nor with bulbils or bulblets, as in *Lilium bulbiferum*, and *Dentaria bulbifera*, which are germs or separable buds developed without fecundation.

576. Seeds are usually enclosed in a seed-vessel or pericarp, and hence the great mass of flowering plants are called *angiospermous* (ἄγγος, or ἀγγεῖον, a vessel, and σπέρμα, a seed). In *Coniferæ* and *Cycadaceæ*, however, the seeds have no true pericarpial covering, and fertilization




479

takes place by the direct application of the pollen to the seed, without the intervention of stigma or style. Hence the seeds, although sometimes protected by scales, are truly naked, and the plants are called *gymnospermous* (γυμνός, naked, and σπέρμα, a seed). Occasionally, by the early rupture of the pericarp, seeds originally covered become exposed. This is seen in *Leontice*, *Cuphea*, &c. In *Mignonette*, the seed-vessel (fig. 479), opens early, so as to expose the seeds, which are called *seminude*.

577. Besides being contained in a pericarp, the seed has its own peculiar coverings. Like the ovule, it consists of a *nucleus* or *kernel*, and *integuments*. In some instances, although rarely, all the parts of the ovule are visible in the seed, viz., the embryo-sac, or quintine, the quartine, the tercine formed from the nucleus, the secundine, and

the primine. In fig. 480, there is a representation of the seed of *Nymphæa alba*, in which *se* indicates the embryo-sac, containing the embryo, *e*; *n*, the cellular farinaceous covering (quartine), formed round the embryo-sac; *m*, *t*, membrane formed from the nucleus (tercine); *m i*, the secundine; *t*, the primine. In general, however, great changes take place by the development of the embryo; the embryo-sac is often absorbed, or becomes incorporated with the cellular tissue of the nucleus; the same thing occasionally takes place in the secundine, so that in the ripe seed, all that can be detected is the embryo and two coverings. The general covering of the seed is called *spermoderm* (σπέρμα, seed, and δέρμα, covering). In order to correspond with the name applied to the covering of the fruit, it ought more properly to be denominated *perisperm* (περί, around, and σπέρμα, seed). This latter term, however, has been appropriated to a certain portion of the seed, to be afterwards noticed under the name of *albumen*.



480

578. **The Spermoderm** usually consists of two parts, an *external membrane*, called the *episperm* or *testa* (ἐπί, upon, or on the outside, and σπέρμα, a seed, or *testa*, a shell), and an *internal membrane*, called *endopleura* (ἐνδον, within, and πλευρά, side). The former may consist of a union of the primine and secundine, or of the primine only, when as occasionally happens, the secundine is absorbed; the latter, of a combination between the membrane of the nucleus and the embryo-sac, or of one of these parts alone. Sometimes the secundine remains distinct in the seed, forming what has been called a *mesosperm* (μέσος, middle); and when it assumes a fleshy character, it has received the name of *sarcosperm* or *sarcoderm* (σάρξ, flesh).

579. **The Episperm** consists of cellular tissue, which often assumes various colours, and becomes more or less hardened by depositions in its interior. In *Abrus precatorius*, and *Adenantha pavonina*, it is of a bright red colour; in French beans, it is beautifully mottled; in the Almond, it is veined; in the Tulip and Primrose, it is rough; in the Snapdragon, it is marked with depressions; in Cotton and Ascle-

Fig. 480.—Young seed of *Nymphæa alba* cut vertically. *f*, Funiculus or umbilical cord. *a*, Arillus derived from the placenta. *r*, Raphe. *c*, Chalaza or cotyledonary end of the seed. *h*, Hilum or base of the seed. *m*, Micropyle or foramen. *t*, Testa or Primine. *m i*, Secundine. *n*, Tercine or membrane of the nucleus. *n*, Farinaceous external perisperm or albumen formed by the nucleus, and probably constituting the quartine of Mirbel. *se*, *se*, Internal perisperm or endosperm formed by the embryo-sac. *e*, The embryo.

pias, it has hairs attached to it; and in Mahogany and Bignonia, it is expanded in the form of wing-like appendages. In *Salvia*, *Collomia*, *Acanthodium*, and other seeds, it contains spiral cells, from which, when moistened with water, the fibres uncoil in a beautiful manner, having a membranous covering. In the epispERM of the seed of *Ulmus campestris*, the cells are compressed, and their sinuous boundaries are traced out by minute rectangular crystals adhering to their walls.

580. **The Endopleura** is also cellular. It is often thin and transparent, but it sometimes becomes thickened. It is applied more or less closely to the embryo, and sometimes follows a sinuous course, forming folds on its internal surface, and separating from the epispERM.

When the embryo-sac remains distinct from the nucleus in the seeds, as in *Nymphæa*, *Zingiber*, *Piper*, &c., it forms a covering to which the name of *vitellus* (*vitellus*, yolk of an egg), was given by Gærtner.

581. **Arillus.**—Sometimes there is an additional covering to the seed, derived from an expansion of the funiculus or placenta after fertilization, to which the name *arillus* has been given. This is seen in the Passion-flower, where the covering commences at the base, and proceeds towards the apex, leaving the foramen uncovered. In the Nutmeg and Spindle-tree, this additional coat is said to commence at the side of the exostome, and to proceed from above downwards, constituting, in the former case, the substance called *mace*; and in the



481

latter, the bright scarlet covering of the seeds (figs. 481, 482). In such instances, it has been called by some an *arillode*. This arillode, after growing downwards, may be reflected upwards, so as to cover the foramen.

582. On the testa, at various points, there are produced at times cellular bodies, which are not dependent on fertilization, to which the name of *strophioles* (*strophiolium*, a little garland), or *caruncles* (*carun-*

Fig. 481.—1, 2, 3, and 4, Various states of the arillus of *Euonymus*, the Spindle-tree. The figures show the mode in which it is developed from the edges of the foramen. *a a a*, Arillode. *f f f f*, Foramen or exostome.

cula, a little piece of flesh), has been given, the seeds being strophio-late or carunculate. These tumours may occur near the base or apex



of the seed, they may be swellings of the exostome, as in *Ricinus* (fig. 483 *c*), or they may occur in the course of the raphe.


583. Seeds are attached to the placenta by means of a funiculus or umbilical cord, which varies much in length. In *Magnolias* it attains a great length, and when the seed is ripe it appears like a cord suspending it from the follicle. The point of the seed by which it is united to the cord or the scar left on its separation, is called the *hilum* or *umbilicus*, and represents its base. It frequently exhibits marked colours, being black in the Bean, white in many species of *Phaseolus*, &c. It may occupy a small or large surface, according to the nature of the attachment. What constitutes the foramen of the ovule, becomes the *micropyle* (*μικρός*, small, and *πύλη*, gate) of the seed, with its exostome and endostome. This may be recognizable by the naked eye, as in the Pea and Bean tribe, Iris, &c., or it may be very minute, and microscopic. It indicates the true apex of the seed, and is important as marking the part to which the root of the embryo is directed. At the micropyle in the Bean, is observed a small process of integument, which, when the young plant sprouts, is pushed up like a lid, and is called *embryotega* (*tego*, I cover.) The fibro-vascular bundles, from the placenta pass through the funiculus and reach the seed, either entering it directly at a point called the *omphalode* (*ὀμφαλός*, navel,) which forms part of the hilum, or being prolonged between the



Fig. 482.—Development of the same arillus, *a*, around the ovule, *o*, exhibited in a different position. 1, 2, 3, 4, are four successive stages of development. In fig. 4, the arillus has been cut vertically, to show its relation to the ovule, which it surrounds completely.

Fig. 483.—Vertical section of a carpel of *Ricinus communis*, and of the seed which it contains. *a*, Pericarp. *l*, Loculament. *f*, Funiculus or umbilical cord. *t*, Integuments of the seed, having at their apex a caruncula, *c*, which is traversed by the small canal of the exostome. The exostome does not correspond exactly with the endostome, which is immediately above the radicle. *r*, Raphe. *ch*, Chalaza. *p*, Perisperm or albumen, the upper portion of which only is seen. *e*, Embryo, with its radicle, *er*, and its cotyledons, *ec*.

outer and inner integument in the form of a *raphe*, and reaching the *chalaza* or organic base of the nucleus, where a swelling or peculiar expansion may often be detected, as in *Crocus*. In fig. 480, the spiral vessels, *r*, are seen entering the cord, *f*, passing through the hilum, *h*, forming the raphe, *r*, between the testa, *t*, and endopleura, *m* i, and ending in the chalazal expansion, *c*. So also in fig. 484, where *f* is the funiculus, *r* the raphe united to the hilum and chalaza, *c*, whence vessels, *v*, penetrate the seed. In some seeds, as *Narthecium ossifragum*, the vessels are said not to appear till after fertilization, and in *Habenaria viridis*, none have been detected. The chalaza is often of a different colour from the rest of the integuments. In the Orange, it is of a reddish-brown colour. Sometimes, however, its structure can only be recognized by careful dissection. It indicates the cotyledonary extremity of the embryo. The hilum and chalaza may correspond, or they may be separated from each other and united by the raphe (fig. 484). The raphe is generally on the side of the seed next the ventral suture.




584. The positions of the hilum, micropyle, and chalaza, are of importance in determining the nature of the seed. The hilum is the base of the seed, and the micropyle its apex, while the chalaza is the organic base of the nucleus. The hilum and chalaza may correspond, the micropyle being at the opposite extremity, and then the seed is *orthotropal* (ὀρθός, straight). The seed may be curved so that the micropyle is close to the hilum, and the chalaza, by the growth of the seed on one side, may be slightly removed from the hilum, then the seed is *campylotropal* (καμπύλος, curved). The micropyle may be close to the hilum, and the chalaza in the progress of development may be removed to the opposite end, then the seed is *anatropal* (ἀνατρέπω, I reverse).*

585. The position of the seed as regards the pericarp, resembles that of the ovule in the ovary, and the same terms are applied—erect, ascending, pendulous, suspended, curved, &c. (figs. 423, 424, 425, 426, 420.) These terms have no reference to the mode in which the fruit is attached to the axis. Thus the seed may be erect while the fruit itself is pendent, in the ordinary meaning of that term. The part of the seed next the axis or the ventral suture is its *face*, the opposite side being the *back*. Seeds exhibit great varieties of forms. They may be flattened laterally, *compressed*; or from above downwards, *depressed*. They may be round, oval, triangular, polygonal, rolled up like a snail, as in *Physostemon*; or coiled up like a snake, as in *Ophiocaryon paradoxum*.

* See ¶ 467, where these terms are more fully explained when treating of the ovule.

Fig. 484.—Seed of the Hazel. *f*, Funiculus. *r*, Raphe. *c*, Chalaza. *n*, Veins spreading in a radiating manner over the integuments of the seed.

586. The great object of fertilization is the formation of the embryo in the interior of the seed. In general, one embryo is produced, constituting what is denominated *monembryony* ($\mu\acute{o}\nu\omicron\varsigma$, one); but in Coniferæ, Cycadaceæ, Mistletoe, &c., there are frequently several embryos, giving rise to what is called *polyembryony* ($\pi\omicron\lambda\acute{\upsilon}\varsigma$, many). Sometimes two embryos become united together in the same seed. In the coniferous seeds, numerous corpuscles are seen whence the embryos proceed. The process of fertilization has already been traced until the embryo appears as a rounded cellular body, enclosed in the embryo-sac, and attached to a suspensor. In fig. 480, *e* is the embryo, and *s e* the embryo-sac. In this sac there is at first a mucilaginous fluid, the *amnios*, in which cells are speedily developed, commencing on its inner surface, and extending towards the interior. The embryonic-cell (fig. 485 *v*), still attached to the sac by its suspensor, *s*, contains in the early state semi-fluid granular matter, which becomes organized, producing distinct nucleated cells (fig. 485, 2, *e*). These gradually multiply, and form at length a cellular mass, at first undivided (fig. 485, 3, *e*), but afterwards showing a separation of parts, so that the



485

487

490

491

axis and lateral projections or rudiments of leaves can be distinguished. In figs. 486 to 491, all the stages of the formation of embryo can be traced; appearing first as a simple cell (figs. 486, 488), forming others in its interior (figs. 489, 490); and finally, the parts of the embryo becoming visible, as in fig. 491, where *g r* is the axis representing the

Fig. 485.—First development of the embryo of *Draba verna*. *o*, Suspensor, which in this plant is very long. *v*, Embryonic or germinal vesicle. *e*, Embryo. 1. First stage, in which the embryonic vesicle only is seen. 2. Second stage, showing several cells formed in the embryonic vesicle. 3. Third stage, in which the embryo becomes more conspicuous in consequence of the formation of numerous small cells.

Fig. 486.—Monocotyledonous embryo of *Potamogeton perfoliatus* in its early stage, appearing as a vesicle or simple cell.

Fig. 487.—The same further advanced, showing radicle, *r*, gemmule or plumule, *g*, and the cotyledon, *c*.

Fig. 488.—Dicotyledonous embryo of *Oenothera crassipes* in its early stage, appearing as a vesicle or cell.

Fig. 489.—The same further advanced, showing three united utricle or cells.

Fig. 490.—The same more developed, showing numerous cells

Fig. 491.—The same in a more developed state, showing radicle, *r*, gemmule, *g*, and cotyledons, *c*.

stem and roots, and *c c* are the lateral projections, which are developed as leaf-like bodies, called *cotyledons* (κοτυληδών, the name of a plant, having leaves like seed-lobes.)

587. **Perisperm or Albumen.**—As the embryo increases in size, it gradually causes absorption of the cellular tissue in the embryo-sac, and it is sometimes developed to such a degree as to reduce the nucleus and embryo-sac to a thin integument. In such a case the seed consists of integuments and embryo alone. In *Santalum*, *Osyris*, and *Loranthus*, Griffith says the ovule is sometimes reduced entirely to a sort of embryonary sac. In *Avicennia*, the embryo, at its maturity, is on the outside of the nucleus and body of the ovule. In other cases it enlarges to a certain extent, filling the embryo-sac completely or partially, and only encroaching slightly on the cells of the nucleus. The cells surrounding the embryo then become filled with a deposit of solid matter called *albumen*, consisting of starchy, oily matter, and nitrogenous compounds. To this matter some have applied the term *perisperm* (περί, around, and σπέρμα, seed); others, that of *endosperm* (ἐνδον, within). The name *perispermic albumen*, or *perisperm*, is often restricted to that found in the cells of the nucleus alone (fig. 480 *n*); *endospermic albumen*, or *endosperm*, to that found within the embryo-sac alone (fig. 480 *s e*), as in *Chelidonium majus*, *Ranunculaceæ*, *Umbelliferae*, &c. Sometimes both kinds of albumen occur in the same seed, as in *Nymphaeaceæ*



and *Piperaceæ*. Schleiden states, that in some instances the albumen is produced in the region of the chalaza. He also remarks, that endospermic albumen is common in *Endogens*. In some *Scrophularias*, the embryo-sac forms little cavities or bags, which in the ripe seed remain as appendages to the albumen. Seeds in which the embryo occupies

Fig. 492.—Anatropal mature seed of *Helleborus niger* cut vertically. The embryo, *e*, is small, as compared with the perisperm or albumen, *p*. *t*, Spermoderm or coverings of the seed. *f*, Funiculus. *h*, Hilum. *c*, Chalaza.

Fig. 493.—Mature seed of *Diphyllela peltata*, showing an embryo, *e*, which occupies a larger portion of the seed than in fig. 494. Letters indicate the same parts as in the previous figure.

Fig. 494.—Ripe seed of *Berberis vulgaris*, exhibiting a larger embryo, *e*, as compared with the perisperm, *p*. Letters as in figs. 492 and 493.

the entire seed, are called *exalbuminous* (*ex*, without), as *Compositæ*, *Cruciferae*, and most *Leguminosæ*, while others having separate albumen are *albuminous*. The larger the quantity of albumen in a seed, the smaller the embryo. In figs. 492 to 494, the relative proportion which the embryo bears to the albumen or perisperm in different seeds is shown; *e* being the embryo with its cotyledons and young root, *p* the perisperm, *t* the coverings of the seed, *f* the funiculus or cord, *h* the hilum, and *c* the chalaza. In fig. 492, the embryo is minute, and occupies only a small part of the apex of the albumen; in fig. 493, it is larger, and has encroached on the perisperm; while in fig. 494, it is still more developed, much of the albumen having been absorbed.

588. The albumen varies much in its nature and consistence, and furnishes important characters. It may be *farinaceous* or *mealy*, consisting chiefly of cells filled with starch (fig. 495), as in Cereal grains, where it is abundant; *fleshy* or *cartilaginous*, consisting of thicker cells which are still soft, as in the Coco-nut, and which sometimes contain oil, as in the oily albumen of Croton (fig. 496), Ricinus, and Poppy; *horny*, when the matter in the cells is of a hard consistence, and often arranged in a concentric manner, so as nearly to fill the entire cavity, as in Date, Ivory-Palm, and Coffee. The albumen may be uniform throughout, or it may present a mottled appearance, as in the Nutmeg, the seeds of *Anonaceæ*, and some Palms (fig. 497), where it is called



495



496



497

ruminated. This mottled appearance depends on the endopleura or inner integument forming folds on which the albumen is deposited, and when the seed is ripe, these foldings of the membrane divide the albumen in a sinuous or convoluted manner.

589. The albumen is a store of matter laid up for the nourishment of the embryo. In the Coco-nut and double Coco-nut, it forms the great bulk of the seed, weighing many ounces, while the embryo is

Fig. 495.—Section of a small portion of the farinaceous perisperm or albumen of *Zea Mais*, Indian corn. *c c c*, Cells. *f f f*, Grains of starch in the cells.

Fig. 496.—Section of a small portion of the oily perisperm or albumen of *Croton Tigllum*. *c c c*, Cells. *h h h*, Drops of oil contained in the cells.

Fig. 497.—Vertical section of the fruit of *Areca Catechu*. *c*, Perianth. *f*, Pericarp. *p*, Ruminated perisperm or albumen. *e*, Embryo.

minute, weighing a few grains, and lies in a cavity at one extremity. In Coffee, the albumen is the horny portion, the infusion of which is used for a beverage. In *Phytelephas* it is called vegetable ivory from its hardness, and is used for the same purposes as ivory. In the horny albumen of this Palm, as well as in that of the *Attalea funifera*, the Date and the Doom Palm, the concentric deposition of secondary layers, leaving a small cavity in the centre of the cells, and radiating spaces uncovered with thickening matter, is well seen under the microscope.

590. The embryo consists of *cotyledons* or rudimentary leaves, the *plumule* (*plumula*, a little feather), or *gemma* (*gemma*, a bud), representing the ascending axis, the *radicle* (*radix*, root), or the descending axis, and their point of union the *collum*, collar or neck; that part of the axis which intervenes between the collar and cotyledons being the *caulicule* (*cauliculus*, a little stalk), or *tigelle* (*tigellus*, a little stalk). The embryo varies in its structure in the different divisions of the vegetable kingdom. In acrogenous and thallogamous plants, it continues as a cell or spore, with granular matter in its interior (fig. 498), without any separation of parts or



498



499

500

501

the production of cotyledons. Hence these plants are called *acotyledonous* (α privative $\kappa\omicron\tau\upsilon\lambda\eta\delta\acute{\omega}\nu$). Endogenous and Exogenous plants, on the other hand, exhibit a marked separation of parts in their embryo, the former having one cotyledon, and hence being *monocotyledonous* ($\mu\acute{\iota}\nu\omicron\varsigma$, one); the latter two, and hence *dicotyledonous* ($\delta\acute{\iota}\varsigma$, twice). Thus, the whole vegetable kingdom is divided into three

Fig. 498.—Acotyledonous embryo of *Marchantia polymorpha*. Such embryos bear the name of spores.

Fig. 499.—Monocotyledonous embryo of *Potamogeton perfoliatus* nearly mature. *r*, Radicle. *t*, Caulicule or tigellus. *c*, Cotyledon. *g*, Gemmule or plumule.

Fig. 500.—Mature dicotyledonous embryo of the common Almond. *r*, Radicle or young root.

Fig. 501.—The same, with one of the cotyledons removed. *r*, Radicle. *t*, Tigelle or caulicule. *c*, One of the cotyledons left. *ie*, Cicatrix left at the place where the other cotyledon was attached. *g*, Gemmule composed of several small leaves.

grand classes by the nature of the embryo, the first of which classes correspond with the Cryptogamic division of plants, the second with the endogenous division of phanerogamous or flowering plants, the third with the exogenous division of the same. Fig. 499 represents a monocotyledonous embryo, with its cotyledon, *c*; while figs. 500 and 501, exhibit a dicotyledonous embryo, with its cotyledons, *c c*.

591. **The Spore** of acotyledonous plants (fig. 498) is a cellular body, from which a new plant is produced. Germination takes place in any part of its surface, and not from fixed points. Some consider it as produced independent of any process of fertilization, others consider the union of two kinds of cells as necessary for its formation. When formed, it sometimes presents filaments or vibratile cilia on its surface (figs. 431-434), by means of which it moves about in fluids like some of the Infusoria. When it germinates, these cilia disappear. Sometimes spores are united in definite numbers, as in fours, surrounded by a cellular covering, or *perispore* (περί, around, and σπορά, a spore), or *sporidium*, and thus forming the reproductive body called a *tetraspore* (τετρας, four), which is common in Algæ.

592. **Embryo.**—In the *embryo* or *corculum* (*corculum*, a little heart), the first part formed is the *axis*, having one of its extremities turned towards the suspensor, and the other in the opposite direction; the former indicating the point whence the young root or *radicle* is to proceed, and the latter that whence the leafy *stem* is to arise. As the first leaves produced are the cotyledons, this stem is called the *cotyledonary* extremity of the embryo, while the other is the *radicular*. The radicular is thus continuous with the suspensor, and consequently points towards the micropyle (fig. 494 *h*), or the summit of the nucleus, an important fact in practical botany; while the cotyledonary, being opposite, is pointed towards the base of the nucleus or the chalaza (fig. 494 *c*). Hence, by ascertaining the position of the micropyle and chalaza, the two extremities of the embryo can in general be discovered. In some rare instances, in consequence of a thickening taking place in the coats of the seed, as in *Ricinus* (fig. 483), and some Euphorbiaceæ, there is an alteration in the micropyle, so that the radicle does not point directly to it.

593. The part of the axis which unites the radicle and the cotyledon or cotyledons, is denominated *caulicule* or *tigelle* (figs. 499 *t*, 501 *t*). This is sometimes very short. From the point where the cotyledons are united to the axis, a bud is developed (in the same way as from the axil of leaves); this bud contains the rudiments of the true or *primordial* (*primus*, first, and *ordo*, rank) leaves of the plant, and has been called *plumule* or *gemmule*. This bud may be seen usually lying within the cotyledons. Thus, in fig. 501, the embryo of the Almond exhibits the *gemmule*, *g*, lying on one of the cotyledons, the other having been removed and leaving a cicatrix, *i c*; while in fig. 499,

the gemmule, *g*, of *Potamogeton perfoliatus*, is covered by the single cotyledon, *c*.

594. The gemmule as well as the cotyledon are sometimes obscurely seen. Thus, in *Cuscuta* (fig. 502), the embryo appears as an elongated axis without divisions; and in *Pekea butyroza* (fig. 503),



the mass of the embryo is made up by the radicular extremity and tigelle, *t*, in a groove of which, *s*, the cotyledonary extremity lies embedded, which when separated, as in the figure, shows only very small cotyledons. In some monocotyledonous embryos, as

Orchidaceæ, it requires a microscopic examination to detect the cotyledonary leaf.

595. **Monocotyledonous Embryo.**—In this embryo, the single cotyledon in general encloses the gemmule at its lower portion, and exhibits on one side a small slit (fig. 504 *f*), which indicates the edges of the vaginal, or sheathing portion of the cotyledonary leaf. The embryo presents commonly a cylindrical form, rounded at the extremities, or a more or less elongated ovoid (fig. 504). At first sight there seems to be no distinction of parts; but on careful examination, by moistening the embryo, and making a vertical section, there will be detected, at a variable height, a small projecting mammilla, buried a little below the surface. This is the gemmule which marks the termination of the axis. From the lower extremity proceeds the radicular portion (figs. 499 *t r*, 504 *r*), which may be said to represent both the tigelle and radicle. The upper portion or chalazal end of the embryo, is the cotyledon (figs. 499, 504 *c*), which is sheathing at its base,

so as to enclose the gemmule. The length of the radicular portion, or that below the gemmule, varies. It is usually shorter than the cotyledon, and is denser in structure; but in some instances it becomes much larger, giving rise to what has been called a *macrapodous* embryo (*μακρός*, long, and *πούς*, a foot). Thus, in fig. 505, *t* represents the long radicular portion in the young state, whence ultimately the root, *r*, proceeds. Occasionally, the radicular portion becomes very thick and large, so as to form a considerable portion of the embryo; and in all monocotyledons, it may be considered as an enlarged mammillary projection, whence the rootlets proceed by bursting through it, and carrying with them a covering or sheath (¶ 127, fig. 124).

Fig. 502.—Spiral embryo of *Cuscuta* or Dodder.

Fig. 503.—Embryo of *Pekea butyroza*. *t*, Thick tigelle or caulicule, forming nearly the whole mass, becoming narrowed and curved at its extremity, and applied to the groove, *g*. In the figure this narrowed portion is slightly separated from the groove. *c*, Two rudimentary cotyledons.

Fig. 504.—Embryo of *Triglochin Barrelieri*. *r*, Radicle. *f*, Slit corresponding to the gemmule. *c*, Cotyledon.

596. When considering endogenous or monocotyledonous stems, it was shown that the leaves are produced singly and alternately, in a sheathing manner, each embracing the subsequently developed bud. So it is in the monocotyledonous embryo. There is a single leaf or cotyledon produced, and if in any instance there is more than one, it is alternate with the first formed. The cotyledon (fig. 504 *c*) is folded either partially, as in *Dioscorea*, or completely. Its sheathing portion (vagina) embraces the bud or gemmule, which appears as a mammillary projection; its position being indicated by a cleft or slit (fig. 504 *f*), where the edges of the sheath unite. All the portion of the embryo above the gemmule, is the cotyledon; all below, the radicle.

597. **Dicotyledonous Embryo.**—The form of this embryo varies much; and although sometimes resembling in its general aspect that of monocotyledons, yet it is always distinguished by a division taking place at the cotyledonary extremity, by which it is separated into two, more or less evident, lobes. The parts of this embryo are easily traced in the Bean, Pea, Acorn, and Almond. In the latter (fig. 500), the embryo has an oval form, consisting of two thick cotyledons, *c c*, and a radicle, *r*. When one of the cotyledons is removed (fig. 501), leaving scars, *i c*, the gemmule or plumule, *g*, is seen included between them, with its caulicle or tigelle, *t*.

598. The cotyledons are not always, however, of the same size. Thus, in a species of *Hiræa* (fig. 506), one of them, *c'*, is smaller than the other; and in *Carapa guianensis* (fig. 507), there appears to be only one, in consequence of the intimate union which takes place between the two as indicated by the dotted line, *c*. The union between the cotyledonary leaves may continue after the young plant begins to germinate. Such embryos have been called





Fig. 505.—Monocotyledonous embryo of *Zannichellia palustris* germinating. *m*, Collum or neck, the point intermediate between the stem or tigelle, *t*, and the radicle or root, *r*. *c*, Cotyledon. *g*, Gemmule or plumule.

pseudo-monocotyledonous (ψευδής, false). When there are two cotyledons, they are opposite to each other. In some cases there are more than two present, and then they become verticillate. This occurs in Coniferæ, especially in the Fir (fig. 508), Spruce, and Larch, in which



six, nine, twelve, and even fifteen have been observed. In such instances the cotyledons have been split by collateral choris, and thus become divided into several. They are linear, and resemble in their form and mode of development the clustered or fasciculated leaves of the Larch. Plants having numerous cotyledons are occasionally denominated *polycotyledonous*. Duchartre thinks that the multiple cotyledons of the Firs are not verticillate, but occur in two opposite groups, placed like two ordinary cotyledons. Hence he considers the plants to be truly dicotyledonous, with the cotyledons deeply divided into a number of segments. Between the two cotyledons there is a slit which is well seen in *Pinus Pinaster* and *excelsa*. Thus, the arrangement of the cotyledons follows the same law as that of the leaves in dicotyledonous or exogenous plants, being opposite or verticillate according to the mode of formation of the axis.

599. The texture of the cotyledons varies. They may be thick, as in the Bean, exhibiting only slight traces of venation, with their flat internal surfaces in contact, and their backs more or less convex; or they may be in the form of thin and delicate laminæ, flattened on both sides, and having distinct venation, as in *Ricinus* (fig. 509), *Jatropha*, *Euonymus*, &c. In the former case they are called *fleshy*, or seminal lobes; in the latter, *foliaceous*, or seminal leaves.

600. Cotyledons are usually entire and sessile. But they occasionally become lobed, as in the Walnut and the Lime (fig. 510), where

Fig. 506.—Embryo of *Hlræa Salzmanniana*, cut vertically, to show the inequality of the two cotyledons, one of which, *c*, forms almost the whole mass of the embryo. *c'*, The small cotyledon. *g*, Gemmule or plumule. *r*, Radicle.

Fig. 507.—Embryo of *Carapa guianensis*, cut vertically, to show the union of the cotyledons, the distinction between which is only indicated by a faint line, *c*. *r*, Radicle. *g*, Gemmule.

Fig. 508.—Embryo of Fir. 1. Taken from the seed. 2. Beginning to germinate. *r*, Radicle. *c*, Cotyledons, which are numerous; the plant being polycotyledonous.

the cotyledon, *c*, has five lobes; or petiolate, as in *Geranium molle* (fig. 511 *p*); or auriculate, as in the Ash (fig. 512 *o*). Like leaves in the bud (see Vernalion, ¶ 184), cotyledons may be either applied directly to each other (fig. 509), or may be folded in various ways. In the Almond (fig. 500) they lie in the direction of the axis. In other cases they are folded laterally, *conduplicate* (fig. 513); or from



apex to base, *reclinate* (fig. 205 *a*); or rolled up laterally, so as partially to embrace each other, *convolute* (fig. 514); or rolled up like the young fronds of ferns, *circinate* (fig. 515.) In these cases, both cotyledons follow the same direction in their foldings or convolutions; but, in other instances, they are folded in opposite directions, resembling the equitant (fig. 205 *m*) and semi-equant (fig. 205 *n*) vernalion.

601. The radicle may be either straight or curved, and, in particular instances, it gives a marked character to the seed. Thus, the divisions of the order Cruciferæ are founded on the relative position and folding of the radicle and cotyledons. In the division *Pleurorrhizæ*

Fig. 509.—Embryo of *Recinus communis* taken out of the seed (see fig. 483), and cut transversely. The two halves are separated so as to show the two cotyledons, *c*, applied to each other. *r*, Radicle.

Fig. 510.—Embryo of the Lime. *r*, Radicle. *c*, One of the divided or palmate cotyledons.

Fig. 511.—Embryo of *Geranium molle*. *r*, Radicle. *c*, Cotyledons attached to the collar by a stalk or petiole, *p*.


Fig. 512.—Embryo of the Ash. *r*, Radicle. *c*, One of the cotyledons. *o o*, Auricular appendages to the cotyledon.

Fig. 513.—Embryo of *Brassica oleracea*, Cabbage. *r*, Radicle. *c*, Cotyledon. 1. Entire embryo. 2. Embryo cut transversely, showing the cotyledons folded on the radicle or conduplicate. The radicle is dorsal, or on the back of the cotyledons.

Fig. 514.—Embryo of *Punica Granatum*, Pomegranate, cut into two halves. The upper half removed to show the convolute cotyledons. *r*, Radicle.

Fig. 515.—Circinate embryo of *Bunias orientalis*.

(πλευρά, side, and ρίζα, root), the cotyledons are applied by their faces, and the radicle (figs. 516, 517 *r*) is folded on their edges, so as to be *lateral*, while the cotyledons, *c*, are *accumbent* (*accumbo*, I lie at the side). In *Notorhizææ* (νωτον, back), the cotyledons (fig. 518 *c*)



are applied to each other by their faces, and the radicle, *r*, is folded on their back, so as to be *dorsal*, and the cotyledons are *incumbent* (*incumbo*, I lie upon or on the back). In *Orthoploceæ* (ορθός, straight, and πλόκος, a plait), the cotyledons are *conduplicate* (fig. 513, 1, 2, *c*), while the radicle, *r*, is dorsal, and enclosed between their folds. In other divisions, the radicle is folded in a spiral manner (fig. 515), and the cotyledons follow the same course. In the Dodder (fig. 502), the embryo appears as an axis without divisions, having several turns of the spiral on different planes.


602. The seed sometimes is composed of the embryo and integuments alone, the former being either straight or folded in various ways, as already shown. In other cases there is an addition of perisperm or nutritive matter, in greater or less quantity, according to the state of development which the embryo attains (figs. 492, 493, 494). When the embryo is surrounded by the perisperm on all sides except its radicular extremity (fig. 494,) it becomes *internal* or *intrarius* (*intra*, within); when lying outside the perisperm, and only coming into contact with it at certain points, it is *external* or *extrarius* (*extra*, without). When the embryo follows the direction of the axis of the seed, it is *axile* or *axial*, and it may be either *external*, so as to come into contact with the perisperm only by its cotyledonary apex (fig. 519), or *internal* (figs. 492, 493, 494). In the latter case, the radicular extremity may, as in some Coniferæ, become incorporated with the perisperm apparently by means of a thickened suspensor. When the embryo is not in the direction of the axis, it becomes *abaxile* or

Fig. 516.—Embryo of a Pea, cut transversely. Upper half separated to show the fleshy accumbent cotyledons, *c*. *r*, Radicle applied laterally.

Fig. 517.—Embryo of *Isatis tinctoria*. *c*, Accumbent cotyledons. *r*, Radicle. 1. Embryo entire. 2. Transverse section of the embryo.

Fig. 518.—Embryo of *Cheiranthus Cheiri*, Wallflower. *c*, Incumbent cotyledons. *r*, Radicle. 1. Embryo entire. 2. Transverse section of the embryo.

abaxial (fig. 520 *e*); and in this case it may be either straight or curved (fig. 521), internal or external. In the straight seed of Grasses, the perisperm is abundant, and the embryo lies at a point



on its surface, immediately below the integuments, being straight and external. In Campylotropous ovules, the embryo is curved, and in place of being imbedded in perisperm, is frequently external to it, following the concavity of the seed (fig. 522), and becoming *peripheral* (περιφέρω, I carry round), with the chalaza situated in the curvature of the embryo.

603. It has been already stated, that the radicle of the embryo is directed to the micropyle, and the cotyledons to the chalaza. In some cases, by the growth of the integuments, the former is turned round so as not to correspond with the apex of the nucleus, and then the embryo has the radicle directed to one side, and is called *excentric*, as is seen in Primulaceæ, Plantaginaceæ, and many Palms, especially the Date (fig. 520). The position of the embryo in different kinds of seeds varies. In all cases the radicle or base of the embryo points more or less directly to the micropyle, while the cotyledonary extremity is directed towards the chalaza. In an orthotropal seed, then, the embryo is inverted or *antitropal* (αντι, opposite, τρέπω, I turn), the radicle pointing to the apex of the seed, or to the part opposite the hilum (fig. 521). Thus, fig. 523 represents an orthotropal seed of Sterculia Balanphas, attached to the pericarp, *p c*, by the funiculus, *f*. The chalaza and hilum are confounded together at *ch*, the micropyle being at the opposite end. The integuments of the seed, *t*, cover the embryo with its perisperm, *p s*; the cotyledons, *c*, point to the hilum and chalaza; while the radicle, *r*, points to the


Fig. 519.—Grain of *Carex depauperata*, cut vertically. *t*, Integuments. *p*, Perisperm. *c*, Embryo.

Fig. 520.—Seed or kernel of the Date. *p*, Perisperm or horny albumen. *e*, Embryo. 1. Entire seed. 2. Seed cut transversely at the point where the embryo, *e*, is situated.


Fig. 521.—Winged fruit of *Rumex*, cut vertically, to show the abaxile or abaxial slightly curved embryo.

Fig. 522.—Carpel of *Mirabilis Jalapa*, cut vertically, with the seed which it contains. *a*, Pericarp crowned with the remains of the style, *s*. *t*, Integuments of the seed or spermoderm. *c*, Peripheral embryo with its radicle, *r*, and its cotyledons, *c*. *p*, Perisperm or albumen surrounded by the embryo.

micropyle, and the embryo is thus reversed or inverted. Again, in an anatropal seed (figs. 493, 494), where the micropyle is close to the




523



524

hilum, and the chalaza at the opposite extremity, the embryo is *erect* or *homotropal* (*ὁμοιοτροπός*, like, and *τερέπω*, I turn), the radicle or base of the embryo being directed to the base of the seed. In some anatropal ovules, as in Castor oil (fig. 483), the exostome is thickened or carunculate, *c*, and the endostome does not correspond exactly to it, so that the radicle, *er*, of the embryo is directed to a point a little removed from the exostome. In curved or campylotropal seeds, (fig. 419), the embryo is folded so that its radicular and cotyledonary extremities are approximated, and it becomes *amphitropal* (*ἀμφιτροπός*, around, and *τερέπω*, I turn). In this instance the seed may be exalbuminous, and the embryo may be folded on itself (fig. 524), or albuminous, the embryo surrounding more or less completely the perisperm, and being peripheral (fig. 522). In fig. 524, the seed of *Erysimum cheiranthoides* is shown, with the chalaza, *ch*, and the hilum, *h*, nearly confounded together, the micropyle, *m*, the embryo occupying the entire seed, with the radicle, *r*, folded on the cotyledons, *c*, which enclose the plumule, *g*. Thus, by determining the position of the hilum, chalaza, and micropyle, the direction of the embryo may be known.



525

604. According to the mode in which the seed is attached to the pericarp, the radicle may be directed upwards or downwards, or laterally, as regards the ovary. In an orthotropal ovule, attached to the base of the pericarp, it is superior (fig. 521.) So also in a suspended anatropal ovule, as in fig. 483. In other anatropal ovules, as in figs. 492, 504, 525, the radicle is inferior. When the ovule is horizontal as regards the pericarp (fig. 523), the radicle, *r*, is either centrifugal, when it

Fig. 523.—Orthotropal seed of *Sterculia Balanngas*, cut longitudinally, with the portion of the pericarp, *pc*, to which it is attached. *f*, Funiculus. *ch*, Chalaza and hilum confounded together. *t*, Integuments of the seed, or spermoderm. *ps*, Perisperm, the summit of which only is seen. *c*, One of the cotyledons. The other cotyledon has been removed to show the gemmule, *g*. *r*, Radicle which is directed to the foramen at the apex of the seed. The embryo is antitropal or inverted.

Fig. 524.—Campylotropal seed of *Erysimum cheiranthoides*, cut longitudinally. *m*, Micropyle. *ch*, Chalaza not far removed from the hilum, *h*. *t*, Testa or epispERM. *mi*, Inner covering of the seed or endopleura. *r*, Radicle. *c*, Cotyledons. *g*, Gemmule. The embryo is curved or amphitropal.

Fig. 525.—Vertical section of the carpel of *Triglochin Barrelieri*. *p*, Pericarp crowned by the sessile stigma, *s*. *g*, Seed. *f*, Funiculus. *r*, Raphe. *c*, Chalaza.

points to the outer wall of the ovary; or centripetal, when it points to the axis or inner wall of the ovary.

9.—FUNCTIONS OF THE SEED.

605. The seed contains the embryo or germ, which, when placed in favourable circumstances, is developed as a new plant. The embryo is usually of a whitish or pale colour, resembling the perisperm when present, and sometimes scarcely distinguishable from it at first sight. Occasionally, however, it is of a greenish or yellow hue. Instances of this occur in the *perispermic* or albuminous seed of *Euonymus*, and the *aperispermic* or exalbuminous seeds of most *Cruciferae*. The changes which take place in the composition of the seed, and in its coats, are with the view of protecting the embryo from vicissitudes of temperature, moisture, &c., and of laying up a store of nourishment for its after growth. The coats become thickened and hardened by the deposition of lignine; and in its interior, starch, nitrogenous compounds, phosphates, and sulphates, besides oily and fatty matters, various organic acids, tannin, and resins, are found. The specific gravity of the seed is much increased, so that it usually sinks in water, and it becomes more capable of resisting decomposition, and preserving the vitality of the embryo. In some instances where air is contained in their envelopes seeds float in water.

606. When seeds are matured, they are detached from the plant in various ways. They separate from the funiculus at the hilum, and remain free in the cavity of the pericarp, which either falls along with them, or opens in various ways so as to scatter them. The elasticity with which some seed-vessels open during the process of desiccation is very great. It may be seen in *Hura crepitans*, Common Broom, and *Cardamine*. In the *Geranium* (fig. 455), the seed-vessels are coiled upwards on the elongated beak, and in this way the seeds are dropped. In the *Cyclamen* the peduncle curves towards the earth so as to place the seed-vessels in a position suitable for germination. In the succulent fruit of *Momordica Elaterium*, or squirting Cucumber, the cells vary in their size and contents in different parts; some containing thick matter become distended at the expense of others with thinner contents, and the force of endosmose ultimately causes rupture of the valves at their weakest point, viz., where they are united to the peduncle. When this takes place, the elasticity of the valves sends out the seeds and fluid contents with great force through the opening left by the separation of the peduncle. In the *Impatiens* or *Balsam*, the seed-vessel opens with force by a similar process, the five valves curving inwards in a spiral manner, in consequence of the distension of the outer large cells. The seeds are discharged before they are dry. In the case of *Mignonette* (fig. 479), the seed-vessel opens

early, so as to expose the seeds; and in *Cuphea*, the placenta pierces the ovary and floral coverings early, so as to render the seeds naked. Fleshy fruits, which fall from the tree when ripe, supply by their succulent portion the most suitable nutriment for the young embryo in its earliest stages of growth.

607. Wind, water, animals, and man, are instrumental in the dissemination of seeds. Some seeds, as those of Mahogany, Bignonia, Tecoma, Pine, Asclepias, Epilobium, and the Cotton plant, have winged or feathery appendages, by means of which they are wafted to a distance. The same thing occurs in some indehiscent seed-vessels, as the samara of the Sycamore and Ash, and the achænia of Dandelion, Thistles, &c. Moisture, as well as dryness, operates in the bursting of seed-vessels. The pod of the *Anastatica*, or Rose of Jericho, and the capsule of some *Mesembryanthemums*, exhibit the effects of moisture in a remarkable degree. Animals, by feeding on fleshy fruits, the kernels of which resist the action of the juice of the stomach, disseminate seeds; and man has been the means of transporting seeds from one country to another. In some cases, the pericarps ripen their seeds under ground, and are called *hypocarpogean* (ὑπὸ, under, καρπός, fruit, and γῆα, earth). This is seen in the *Arachis hypogæa*, or Ground nut. Others, as *Vicia amphicarpos*, have both aerial and subterranean fruit. Many seeds are used for food by animals, and a great destruction of them takes place from decay: but this is compensated for by the vast number produced, so as to secure the continuance of the species. The quantity of seeds produced by many plants is very great. In single capsules of Poppy and Tobacco, upwards of 40,000 have been counted.

608. **Germination.**—The act by which the embryo of a seed leaves its state of torpidity, and becomes developed as a new plant, is called *germination* (*germinatio*, springing). In order that this process may go on, a certain combination of circumstances is necessary. The chief requisites are moisture, air, and a certain temperature. Exclusion from light is also beneficial. Generally speaking seeds do not begin to sprout until they are placed in such a position as presents those circumstances favourable to their growth. In some instances, however, more particularly in the case of plants whose seeds contain much albumen, germination commences when the period suitable for their sowing arrives.

609. *Moisture* is necessary in order that the nutritive matters may be taken up in a state of solution, and that certain changes may take place in the seed. Dry seeds will not germinate. Until water be absorbed no circulation of fluids in the seed can take place. The quantity of water absorbed by seeds is often very large. Decandolle found that a French bean, weighing 544 miligrammes, absorbed 756 of water. The swelling of Peas by absorption of water is familiar to

all. The kernels or seeds by this means are enabled to burst their stony coverings.

610. *The temperature* required for germination varies in different seeds. Some demand a tropical heat, others are satisfied with the warmth of our spring. In general, the requisite temperature may be said to vary from 60° to 80° F. Some seeds can bear a temperature which would kill others. Some have been known to germinate after exposure for a short time to the heat of boiling syrup; others after exposure to a cold of —39° F. Many plants grow in the immediate vicinity of very hot springs, others in cold regions. Edwards and Colin, from their experiments, were led to fix 95° F. as the highest limit of prolonged temperature which cereal grains can bear in water; and 113° F. as the highest they can bear in sand or earth. Vegetable life has been observed progressing under much higher temperatures. In the Manilla Islands, a hot spring which raised the thermometer to 187° had plants flourishing in it and on its borders. A species of *Chara* has been observed growing in the hot springs of Iceland, as well as in the boiling springs of Arabia and of the Cape of Good Hope. Wheat, Oats, and Barley, are said to thrive in any country where the mean temperature exceeds 65° F. The spores of certain cryptogamic plants are especially fitted for cold countries. Edwards and Colin found that seeds in a dry air bore a higher temperature than in water or steam.

611. *Air*, or rather *oxygen*, was shown by Scheele to be necessary for germination. Seeds deeply buried in the soil, and excluded from air, do not spring. The depth at which seeds should be sown, varies from half an inch to two inches, according to the nature of the soil. The following experiments were made by Petri:—

Seed sown to the depth of	Came above ground in	No. of plants that came up.
$\frac{1}{8}$ inch	11 days	7-8ths.
1 —	12 —	all.
2 —	18 —	7-8ths.
3 —	20 —	6-8ths.
4 —	21 —	4-8ths.
5 —	22 —	3-8ths.
6 —	23 —	1-8th.

Shallow sowing is thus proved to be the best.

612. Seeds, when buried deep in the soil sometimes lie dormant for a long time, and only germinate when the air is admitted by the process of subsoil ploughing, or other agricultural operations. When ground is turned up for the first time, it is common to see a crop of white clover and other plants spring up, which had not been previously seen in the locality. After the great fire in London, plants sprung up, the seeds of which must have long lain dormant; and the same thing is observed after the burning of forests, and the draining

of marshes. Gardner says that the name *capoeira* is given in Brazil to the trees which spring up after the burning of the virgin forests (*matos virgens* and *capoes*), and that they are always very distinct from those which constituted the original vegetation. Mr. Vernon Harcourt mentions a case where turnip seeds lay in a dormant state for seven or eight years, in consequence of being carried down to a great depth in the soil. On the Calton Hill, at Edinburgh, when new soil was turned up some years ago for building, a large crop of *Fumaria micrantha* sprung up; and seeds gathered from under six feet of peat-moss in Stirlingshire have been known to germinate. Mr. Kemp mentions the germination of seeds found at the bottom of a sand-pit 25 feet deep, which he concludes, from various circumstances, to have been deposited more than 2000 years ago. The seeds were farinaceous, belonging to the natural order Polygonaceæ. A weak solution of chlorine is said to accelerate germination, probably by the decomposition of water, and the liberation of oxygen.

613. *Darkness* is favourable to germination. Seeds germinate best when excluded from light. M. Boitard showed this by experiments on *Auricula* seeds, some of which were covered by a transparent bell-jar, others by a jar of ground glass, and a third set by a jar enveloped in black cloth. The last germinated most rapidly. Senebier concluded that the height and size of a plant was proportionate to the intensity of the illumination, its verdure dependent on the quality of the rays. Mr. Hunt says that the luminous or light-giving rays, and those nearest the yellow, have a marked effect in impeding germination; the red or heat-giving rays are favourable to the process, if abundance of water is present; while the blue rays, or those concerned in chemical action or *actinism*, accelerate the process and cause rapid growth. His experiments were performed by making the sun's rays pass through different kinds of coloured glass. He believes that the scorching effect of the sun on leaves may be prevented by the use of blue glass, and that a high temperature might be obtained by red glass. He has suggested a pale-green glass made with oxide of copper, as that best fitted for conservatories. By this means he expects that the scorching rays of light will be excluded, while no hindrance is given to the passage of the others; the green colour being a compound of yellow or luminous, and of blue or chemical rays. A delicate emerald-green glass has been employed lately at his suggestion, in glazing the large Palm-house at Kew.

614. Some have said that *electricity* prevents germination, but facts are wanting to confirm this. The experiments of Dr. Fife,* Mechi, Coventry, and others, have shown that the statements made in regard to the efficacy of electro-culture are erroneous.

615. In order that plants may germinate vigorously, moisture,

* See Trans. of Soc. of Arts, vol. iii. part II. p. 109.

heat, and air must be supplied in due proportion. If any of them are deficient, or in excess, injury may be done. It is of great importance, therefore, in agricultural operations, that the ground should be well pulverized, the seeds regularly sown at a proper and equal depth, and the soil drained. Pulverized soil, when examined, is found to consist of small particles having cavities in their interior, and separated from each other by interstitial spaces. In a very dry soil, all these cavities are full of air; in a very wet undrained soil, they are full of moisture; while in a perfectly drained soil, the interstices are full of air, while the particles themselves are moist. The seed in such a soil is under the influence of heat, air, and moisture, and is excluded from light. Hence it is in very favourable circumstances for germination. Frost has an important effect in pulverizing the soil, by the expansion of the water contained in the particles, when it is converted into ice. Snow again, acts in giving a covering to the young plant, protecting it from intense frost and sudden alternations of temperature, and by its slow melting allows the plant to accommodate itself to the mild atmosphere. Snow contains often much oxygen.

616. If a field is not equally planted, the seeds will sink to different depths, and will spring up very irregularly. The seeds should be placed at a depth not greater than two inches. *Draining* acts not merely in removing superfluous moisture, but in allowing a constant renewal of nutritive matter, more especially of ammonia and carbonic acid from the atmosphere, in giving a supply of air, and in keeping up a proper temperature in the soil. In an undrained soil the water is stagnant, and there is little supply of fresh nutriment, and much cold is produced. Of late there has been a discussion as to whether shallow or deep draining is the best. Much depends on the nature of the soil, and it is impossible to lay down any fixed rule applicable to all cases. Mr. Smith says that drains in very stiff soils should be fifteen feet apart, and in very light soils thirty or forty; the depth being from thirty to thirty-six inches, and the main drains six inches deeper than the parallel ones. In extremely stiff clays, he makes drains two and a-half feet deep. He was the first to advocate the system of parallel drains, or what is called *thorough-draining*.

617. **Vitality of Seeds.**—Some seeds lose their vitality soon, others retain it for a long time. Coffee seeds, in order to grow, require to be sown immediately after ripening. On the other hand, Melon seeds have been known to retain their vitality for upwards of forty years, and those of the Sensitive plant for more than sixty years. Oily seeds in general lose their vitality quickly, probably from their power of absorbing oxygen, and the chemical changes thus induced. Considerable discussions have taken place as to the length of time during which seeds will retain their germinating powers. Lindley mentions a case in which young plants were raised from seeds found in an ancient

barrow in Devonshire, along with some coins of the Emperor Hadrian; and M. des Moulins relates an instance of seeds capable of germinating, which were discovered in a Roman tomb, supposed to be fifteen or sixteen centuries old. In these instances, it is to be remarked, that the seeds were protected from the influences required for growth, and were preserved in circumstances which cannot be easily imitated. There seems to be great doubts as to the seeds found in the catacombs of Egypt, and in mummy cases, having actually produced living plants. The statements relative to Mummy Wheat are not fully confirmed, and there are many sources of fallacy.

618. With the view of preserving seeds, it is of importance that they should be thoroughly ripened, kept in a uniform temperature, and in a dry state, and not directly exposed to the oxygen of the air. They are often best kept in their seed-vessels. The hard coverings of many foreign legumes, and of the cones of Firs, &c., seem to be of importance in preserving the germinating power of seeds. Seeds not fully ripened are very apt to decay, and are easily affected by moisture. Seeds, although fit for food, may have lost their germinating power. Corn, pulse, and farinaceous seeds generally will live for a long time if gathered ripe, and preserved quite dry. In sending seeds from foreign countries, they should be put into dry papers, and exposed to free ventilation in a cool place; as, for instance, in a coarse bag suspended in a cabin. Oily seeds and those containing much tannin, as beech-mast, acorns, and nuts, must not only be ripe and dry, but also must be excluded from the air. When transported, they are often put into dry earth and sand, and pressed hard, or preserved in charcoal powder, the whole being covered with tin, and put into a stout box. Some have suggested their preservation in hermetically-sealed bottles full of carbonic acid gas. Seeds enveloped in wax sent from India germinated well. They had been kept for three months, and were quite firm and fresh. They did not sprout for a month, but afterwards grew strong and healthy. Seeds sent in cotton and brown paper had grown considerably in their transit, and, when potted, grew fast, but soon displayed symptoms of debility. Spanish chestnuts and filberts have been sent enveloped in wax to the Himalayahs, and are now growing there. Cuttings of fruit trees, with their ends enveloped in wax, were also sent, and arrived in a living state. In this way also, apples, pears, and plums have been sent.

619. M. Alphonse Decandolle made experiments on the vitality of seeds. He took 368 species of seed, fifteen years old, collected in the same garden, and sowed them at the same time, and in the same circumstances as nearly as possible. Of the 368, only seventeen germinated, and comparatively few of the species came up. The following are the results:—

				Per cent.
Malvaceæ.....	5	came up out of.....	10 species.....	0.50
Leguminosæ.....	9	— — — — —	45 — — — — —	0.20
Labiata.....	1	— — — — —	30 — — — — —	0.03
Scrophulariaceæ.....	0	— — — — —	10 — — — — —	0.00
Umbelliferae.....	0	— — — — —	10 — — — — —	0.00
Caryophyllaceæ.....	0	— — — — —	16 — — — — —	0.00
Gramineæ.....	0	— — — — —	32 — — — — —	0.00
Cruciferae.....	0	— — — — —	34 — — — — —	0.00
Compositæ.....	0	— — — — —	45 — — — — —	0.00

In 357 species, of which the duration of life was known, the results were:—

				Per cent.
Annuals.....	9	came up out of.....	180 species.....	5.0
Biennials.....	0	— — — — —	28 — — — — —	0.0
Perennials.....	4	— — — — —	105 — — — — —	3.8
Ligneous.....	3	— — — — —	44 — — — — —	6.7
	16		357	4.4

or it may be thus given—

				Per cent.
Monocarpic.....	9	came up out of.....	208 species.....	4.3
Polycarpic*.....	7	— — — — —	149 — — — — —	4.7
	16		357	4.4

620. Woody species thus seem to preserve the power of germinating longer than others, while biennials are at the opposite end of the scale; perennials would appear to lose their vitality sooner than annuals. Large seeds were found to retain the germinating power longer than small ones, and the presence or absence of separate albumen or perisperm did not seem to make any difference. Compositæ and Umbelliferae lost their germinating power very early. From these experiments, Decandolle concludes that the duration of vitality is frequently in an inverse proportion to the rapidity of the germination. This subject has been investigated by a committee of the British Association, under the direction of Professor Daubeny.

621. **Chemical changes during Germination.**—During the process of germination, certain changes take place in the contents of the seed, by which they are rendered fit for the nourishment of the embryo. In exalbuminous or aperispermic seeds, where the embryo alone occupies the interior, these changes are effected principally in the matters stored up in the cotyledons. In albuminous or perispermic seeds, on the other hand, the changes occur in the substance of the perisperm. One of the most remarkable of these changes is the conversion of starch into dextrine and grape sugar by a process of oxidation, the object being the conversion of an insoluble into a soluble substance. A nitrogenous compound, called Diastase (§ 310), is developed during


* For an explanation of these terms, see § 631.

germination, and is said to act on the starch. This diastase may be probably a portion of gluten passing into a state of decomposition, and acting as a ferment. The change of starch into dextrine and sugar is referred by chemists to catalytic action, or the action of contact, and to the influence exercised by diastase and other matters in making a new arrangement of the molecules. While this conversion of starch into sugar proceeds, oxygen is absorbed, carbonic acid is given off, and heat is produced. It is probable that at this period there is a certain amount of electric disturbance. Carpenter states that the conversion of the starch of the seed into sugar involves the liberation of carbonic acid, with a small quantity of acetic acid, all acids being negative, and like electricities repelling each other. It is probable when such a process is being accomplished that the seed is at the time in an electro-negative condition. These phenomena are well seen in the malting of barley. The changes produced in the air by germinating seeds have been investigated by Saussure, who showed that in all cases carbonic acid was evolved at the expense of the carbon of the seed.

622. When all the requisites for germination are supplied, the seed, by the absorption of moisture, becomes softened and swollen. When albumen or the perisperm is present, it undergoes certain chemical changes by the action of the air and water, so as to be rendered fit for the nutrition of the embryo. These changes consist partly in the conversion of starch into sugar, and are accompanied with the evolution of carbonic acid, and the production of heat. As the fluid matters are absorbed by the cells of the embryo, the latter continues to increase until it fills the cavity of the seed, and ultimately bursts through the softened integuments. In cases where there is no perisperm, the exalbuminous embryo occupies the entire seed, and the process of germination goes on with greater rapidity. The embryo speedily swells, ruptures the integument, and is nourished at the expense of the cotyledons, which are often fleshy, containing much starchy matter, as in the Bean and Pea, along with oily matter, as in the Nut and Rape seed. There are thus two stages of germination—that in which the embryo undergoes certain changes within the seed itself, and that in which it protrudes through the integuments and becomes an independent plant.

623. The embryo, then, nourished at the expense of its perisperm and cotyledons, continues to grow, and usually protrudes its radicular extremity (fig. 526, 1) in the first instance, which is nearest the surface, and next the micropyle. This, which in the embryo is very short, and confounded with the cauliculus so as to form the first internode, becomes thickened by addition to its extremity (fig. 526, 2), and the division between the ascending and descending axis becomes more marked. The caulicule or axis also elongates, bearing at its

summit the plumule, which now appears outside the integuments (fig. 526, 3, *g*), forming the second internode, either accompanied by the cotyledons, or leaving them still within the seed coats. In the latter case, the cotyledons are usually fleshy and of a pale colour, and become gradually absorbed like the perisperm. In the former they assume a more or less leafy aspect, exercising the functions of leaves for a certain period, and ultimately decaying. While the radicle descends towards the centre of the earth, producing roots of a pale colour, the plumule has a tendency to ascend, forming the leafy axis, and assuming a green colour under the influence of light and air.



624. **Direction of Plumule and Radicle.**—Various attempts have been made to explain the ascent of the plumule, and the descent of the radicle, but none of them are satisfactory. Physiologists have not been able to detect any law to which they can refer the phenomena, although certain agencies are obviously concerned in the effects. Some have said that the root is especially influenced by the attraction of the earth, while the stem is influenced by light. Experiments have shown that the direction of the root is not owing to the moisture of the soil, and that the ascent of the stem is not due to the action of light and air; for roots descend, and stems ascend, even when the latter are placed in contact with the earth, and the former submitted to the action of light. Knight thinks that the direction of stem and roots may be traced to gravitation, and the state of the tissues. When a branch is horizontal, the fluids gravitate towards the lower side; a vigorous growth takes place there; the tissues enlarge, and, by increasing more than those on the upper side, an incurvation is produced, the convexity of which looks downwards, and thus the extremity of the branch is directed upwards. Again, in the root the increase takes place by the extremity, and the fluids by their gravity cause this to retain always a descending direction. A similar explanation is given by Dodart. Dutrochet refers the phenomena to endosmose, which varies in its effects according to the comparative size of the cells in the centre and circumference of an axis. In young stems with large pith, the central cells are larger, and they diminish towards the circumference; whereas in roots, according to him, the diminution takes place in the reverse manner. Large cells distend more rapidly

Fig. 526.—Germination of the dicotyledonous aperispermic seed of *Acacia Julibrissin*. *e*, Spermoderm or testa. *r*, Radicle of the embryo. *t*, Tigellus or cauliculus. *c*, Cotyledons. *g*, Gemmule or plumule. 1. First stage: in which the radicle ruptures the envelope or spermoderm, and appears externally at the micropyle. 2. Second stage: where the parts of the embryo are further disengaged from the covering, the summit of the cotyledons only being retained by the spermoderm. 3. Third stage: where the embryo is entirely disengaged from the envelope or spermoderm, and the cotyledons, *c c*, are separated so as to exhibit the plumule, *g*.

than small ones; and, according to their position in the axis, will thus cause curvature outwards or inwards, the largest occupying the convexity of the arch, the smallest the concavity. When a branch or root is laid horizontally, the force of endosmose is weakened on the lower side, and, consequently, will cease to neutralize the tendency to incurvation on the upper side, which will therefore be directed either upwards or downwards, according to the position of its layers of small cells,—in the case of a branch with large central cells, curving upwards; and in the case of a root with larger hemispherical cells, downwards.


625. These explanations do not appear, however, to be altogether satisfactory. It is known that the stem is directed upwards, the root downwards, but, as yet, physiologists have not been able to ascertain the laws which regulate them. The tendencies of the root and stem are not easily counteracted. When a seed is planted in moist earth, and suspended in the air, the root will, in the progress of growth, leave the earth and descend into the air in a perpendicular direction, while the stem will pass through a quantity of moist earth in an upward direction. If their positions are reversed they will become twisted, so as to recover their natural positions.

626. The effect of light on the stem may be illustrated by the growth of plants in circumstances where a pencil of light only is admitted on one side. Dr. Paggioli of Bologna was the first who observed the influence exercised by the rays of the spectrum in causing flexion of plants. Experiments on this subject have been made by Payen, Dutrochet, and Gardner. They consider the blue rays as those which have the greatest effect on the plumule. Hunter observed, that if a barrel filled with earth, in the centre of which are some beans, be rotated for several days horizontally, the roots pointed in a direction parallel to the axis of rotation. Knight* put Mustard seeds and French beans on the circumference of two wheels, which were put in rapid motion, the one in a horizontal, and the other in a vertical manner; and he found that, in the former, the roots took a direction intermediate between that impressed by gravitation and by the centrifugal force, viz., downwards and outwards; while the stems were inclined upwards and inwards. In the latter, where the force of gravitation was neutralized by the constant change of position, the centrifugal force acted alone, by which the roots were directed outwards, at the same time that the stem grew inwards. To explain these results, there must be allowed—1. A more or less liquid condition of the new parts of the young plant. 2. A different density in the different parts of the latter. 3. A tendency of the denser parts of new plants, during germination, towards the root. On the vertical wheel, the parts of the young plants submitted to the centrifugal force only, had their roots or densest parts at the circumference. On the horizontal, the effect

* See Knight's Horticultural Papers, London, 1841, p. 124.

was intermediate between centrifugal force and gravity. The upper side of leaves is under the influence of light in a marked degree, for, when placed in the reverse position by the turning of a branch, they twist round so as to resume their natural posture. During darkness, on the contrary, many leaves fold in such a way that their lower surface is exposed. Some plants grow indifferently in all directions at the period of germination. The Mistletoe and other parasites direct their radicles towards the centre of the tree or plants to which they are attached, while the plumule grows perpendicularly to the surface.

627. Monocotyledonous Germination.—In Monocotyledons, there is generally a perisperm present, often in large quantity, and in them the cotyledon remains more or less within the seed at the period of germination. The intra-seminal portion of the cotyledons, as in *Canna*, and especially in the Coco-nut, becomes developed as a pale cellular mass, which increases much, and absorbs the nutriment required for the embryo. In some Monocotyledons the perisperm disappears entirely; in others, as in the Phytelphas or Ivory Palm, while certain soluble matters are removed, the perisperm still retains its original form. The intra-seminal part may be said to correspond to the limb or lamina of the cotyledonary leaf. The extra-seminal portion, corresponding to the petiole, becomes often much elongated, as in the double Coco-nut, and ends in a sheath which envelopes the axis or cauliculus, and the plumule. Sometimes, however, there is no marked elongation of the cotyledon, the sheath being at once formed on the outside of the seed, so that the plumule and radicle are, as it were,




sessile on its surface. These phenomena are well seen in *Canna indica* (fig. 527), where *e* is the envelope of the seed; *p*, the perisperm or

Fig. 527.—Germination of the monocotyledonous perispermic seed of *Canna indica*. The seed is cut to show the relation between the perisperm and the embryo at different stages, the former diminishing, while the latter increases. *e*, Envelope or spermoderm. *o*, Its upper part, which is separated like a lid or operculum, to allow the passage of the radicle. *p*, Perisperm or albumen. *c*, Cotyledon. *r*, Radicle or young root. *r' r'*, Secondary radicles. *co*, Coleorhiza or sheath of the roots. *f*, Slit indicating the position of the gemmule; at this slit an elongated sheath, *v*, is protruded. *p c*, Narrow portion of the cotyledon (corresponding to the petiolar portion), intermediate between its enlarged portion, *c*, (corresponding to the lamina or limb of the leaf), and its sheathing or vaginal portion, *v*. *t*, Tigellus or cauliculus. *g*, Gemmule or plumule. 1. First stage, in which the radicle, *r*, begins to appear through the integuments or spermoderm. 2. Second stage, where the slit, *f*, is seen also on the outer surface, indicating the situation of the gemmule. The true radicle, *r*, has pierced the envelope of the seed, and at its base shows a small sheath or coleorhiza. One of the small radicles, *r'*, is also seen with a coleorhiza. 3. Third stage, when all the parts are more developed, and the gemmule, *g*, appears on the outside of the slit, *f*, the edges of which are prolonged in the form of a sheath or vagina, *v*.

albumen; *c*, the intra seminal portion of the cotyledon, which absorbs the nourishment; *p c*, the petiolar or extra-seminal portion of the cotyledon, which varies in length, and may be wanting; *v*, the sheathing portion of the cotyledon, from a slit in which, *f*, the plumule, *g*, protrudes, supported on the axis or cauliculus, *t*; while the radicles, *r* and *r'*, pierce the integument at the base, and are each covered with a separate sheath, *c o*, called coleorhiza (fig. 124). In aperispermic Monocotyledons, as Alismaceæ and Potameæ (fig. 505), the cotyledon does not remain within the seed, but is raised above the ground, *c*, giving origin to the plumule, *g*, which is at first enclosed in its sheath.

628. Thus the cotyledon follows the development of leaves. Its limb is first produced, and is either pushed above ground, or is confined within the seed. In the latter case it is arrested in its progress;



subsequently, a sheath is formed which may either be a direct continuation of the limb, or may be separated from it by a petiolar

Fig. 528.—Germination of the dicotyledonous embryo of *Acer Negundo*. *m*, Collum, collar or neck. *r*, Root. *t*, Cauliculus or stem. *c c*, Cotyledons. *g*, Gemmule or plumule.

Fig. 529.—Upper part of the same embryo more developed. *c c*, Cotyledons. *g*, Gemmule, the first leaves of which are already expanded. *t*, Cauliculus or stem.

Fig. 530.—Acotyledonous embryos or spores of *Marchantia polymorpha*, germinating. 1. Spore in the early stage of germination. 2. In a more advanced stage. The spores are simple cells, which elongate during germination at some point of their surface. They are heterorhizal. They may be compared to naked embryos rather than to seeds.

portion. When the limb is confined in the seed, and ceases to be developed, the sheath often continues to grow, forming a marked covering of the axis. The roots in Monocotyledons during germination (fig. 124 *r r*), pierce the radicular extremity of the embryo, and become covered with sheaths or coleorhizas, *c c*, formed by a superficial layer of cellular tissue. As the radicular extremity thus remains within the embryo, and sends out radicles from its surface, the plants are said to be *endorhizal* (ἐνδον, within, and ῥίζα, a root). See ¶ 127.

629. Dicotyledonous Germination.—In Dicotyledons, the cotyledons generally separate from the integuments, and either appear above ground in the form of temporary leaves (figs. 528, 529 *c c*), which differ in form from the permanent leaves of the plant (fig. 529 *g*), or remain below as fleshy lobes. In the former case they are *epigeal* (ἐπι, upon or above, and γῆα, the earth); in the latter case (as in Beans, Arachis, &c.), they are *hypogeal* (ὑπὸ, under). The cotyledons usually separate, but sometimes they are united, and appear as one. In all cases, the plumule (figs. 528, 529 *g*) proceeds from between the two cotyledons, and does not pierce through a sheath as in monocotyledons. The root (fig. 528 *r*) is a direct prolongation of the axis, *t*, in a downward direction, separating from it at the collar, *m*, and the embryo is here *exorhizal* (ἐξω, outwards). See ¶ 126.

In Acotyledons, the spore (fig. 530) has no separate embryo in its interior, but germinates from any part of the surface; hence it is called *heterorhizal* (ἑτερος, diverse). See ¶ 128. The spore may be considered as a cellular embryo rather than a seed.

630. Some seeds commence the process of germination before being detached from the plant. This occurs in a remarkable degree in the Mangrove trees, or Rhizophoras, which grow at the muddy mouths of rivers in warm climates. Coco-nuts often begin to germinate during a voyage from the tropics to Britain, and germinating seeds have been found in the interior of Gourds, as well as the fruit of *Carica Papaya*, the Papaw.

631. Proliferous Plants.—In place of seeds, some plants produce buds which can be detached and produce separate individuals. Flowers which are thus changed into separable buds, are called *proliferous* (*proles*, offspring, and *fero*, I bear), or *viviparous* (*vivus*, alive, and *pario*, I produce). They are met with in many alpine grasses, as *Festuca ovina*, *var. vivipara*, *Aira cæspitosa*, *var. alpina*, *Poa alpina*, &c., as well as in *Alliums*, *Trifoliums*, &c. Buds of a similar kind may be produced on the edges, or in the axil of leaves, as in *Byrophyllum calycinum*, *Malaxis paludosa*, many species of *Gesnera*, *Gloxinia*, and *Achimenes*; and the bulbils of *Lilium*, *Ixia*, and *Dentaria*, seem to be peculiar forms of buds, capable of being detached, and of assuming independent growth. Buds, however, differ from embryos of seeds in the direction of the roots being towards the axis of the plant.

632. The length of time required for the protrusion of the radicle varies in different plants. Some seeds, as garden cresses, germinate in the course of twenty-four hours, others require many days or many months. Seeds with hard coverings, or a stony perisperm, may lie dormant in the soil for a year or more. The following experiments were made in the Geneva garden, on seeds similarly watered, and exposed to a medium temperature of 53° F. It was ascertained that one-half of the species of the following families germinated after the lapse of the number of days here mentioned:—

Amaranthaceæ,.....	9 days.
Cruciferae,.....	10 —
Boraginaceæ, Caryophyllaceæ, Chenopodiaceæ, Malvaceæ,.....	11 —
Compositæ, Convolvulaceæ, Plantaginaceæ,.....	12 —
Polygonaceæ,.....	13 —
Campanulaceæ, Leguminosæ, Valerianaceæ,.....	14 —
Gramineæ, Labiatae, Solanaceæ,.....	15 —
Rosaceæ,.....	17 —
Ranunculaceæ,.....	20 —
Antirrhinums, Onagrariaceæ,.....	22 —
Umbelliferae,.....	23 —

Temperature has a great effect in accelerating germination. Thus, *Erigeron caucasicum*, at a temperature varying from 49° to 53°, germinated in ten days; at a temperature from 66° to 72°, in two days; *Dolichos abyssinicus*, at the former temperature, in ten days, at the latter, in three; *Zinnia coccinea*, in twenty-two, and five days respectively.

633. **Duration of the Life of Plants.**—Plants, according to the duration of their existence, have been divided into *annual*, *biennial*, and *perennial*. The first of these terms imports that the seed germinates, and that the plant produces leaves and flowers, ripens its seed, and perishes within twelve months; the second, that a plant germinates and produces leaves the first year, but does not produce a flowering stem, nor ripen its seed, till the second, after which it perishes: while the third intimates, that the process of flowering and fruiting may be postponed till the third year, or any indefinite period. The first two exercise the function of flowering in general only once, while the last may do so several times before dying. Under different climates, however, and under different modes of management, the same species may be annual, biennial, or even perennial. Thus, Wheat in this country is annual if sown early in spring, but biennial if sown in autumn; in hot climates, *Lolium perenne* proves annual; the Castor-oil plant in this country is annual, while in Italy it is a shrub of several years' duration; the annual Mignonette, by removing its flower-buds the first year, and keeping it in a proper temperature during the winter, may be rendered perennial and shrubby. Many flowering garden plants, as Neapolitan Violet and Lily of the Valley, may be brought

into flower at a late period of the year, by pinching off the blossoms in the early part of the season.

634. Plants as regards their flowering and fruiting, have also been divided into *monocarpic* (μόνος, one, and καρπός, fruit), or those which flower once only and then die; and *polycarpic* (πολύς, many), or those which flower and fruit several times before the entire plant dies. Thus, annuals and biennials, which flower the first or second year and die, as well as the Agave, and some Palms which flower only once in forty or fifty years, and perish, are monocarpic; while perennials are polycarpic. Some perennial woody plants live to a great age. Some specimens of *Adansonia digitata*, the Baobab of Senegal, are said to be more than 5000 years old. The Yew, the Oak, the Lime, the Cypress, the Olive, the Orange, Banyan, and Chestnut, often attain great longevity.

635. The following is a notice of the size and age of some trees:—

Height to which forest trees grow in France,.....	120 to 130 feet.
Height to which forest trees grow in America,.....	150 —
Trunks of some Baobabs have a girth of.....	90 —
Trunk of <i>Dracæna</i> of the Canaries has a girth of.....	45 —
That of an <i>Acer</i> in South Carolina has a girth of.....	62 —
In France trees have often a girth of.....	25 to 30 —
Oaks in Britain planted before the Conquest, more than.....	800 years old.
Yew at Fountain's Abbey, Ripon.....	1200 —
Yews in churchyard of Crowhurst, Surrey.	1450 —
Yew at Fortingal, Perthshire.....	2500 to 2600 —
Yew at Brabourn churchyard, Kent.....	3000 —
Yew at Hedsor, Bucks, 27 feet diameter.....	3200 —

A specimen of *Ficus indica*, or the Banyan, on an island in the river Nerbudda, is believed to be identical with one that existed in the time of Alexander the Great, and which, according to Nearchus, was then capable of overshadowing 10,000 men. Parts of it have been carried away by floods, but it can shade 7000 men, and its circumference, measuring its principal trunk only, is 2000 feet. The chief trunks of this tree greatly exceed our English Oaks and Elms in thickness, and are above 350 in number. The smaller stems are more than 3000 in number.

636. The Maronites believe that some Cedars near the village of Eden in Lebanon, are the remains of the forest which furnished Solomon with timber for the temple, full 3000 years ago. These Cedars were visited by Belonius in 1550, who found them twenty-eight in number; Rawolf, in 1575, makes them twenty-four; Dandini, in 1660, and Thevenot, about fifty years after, make them twenty-three; Maundrell, in 1696, found them reduced to sixteen; Pococke, in 1736, found fifteen standing; in 1810, Burckhardt counted eleven or twelve; and Dr. Richardson, in 1818, states them to be no more than seven. They must be of great antiquity, seeing they were counted

old 300 years ago. Maundrell mentions the size of some of the Cedars. The largest he measured was 36 feet 6 inches in circumference, and 117 feet in the spread of its boughs.

637. Decandolle gives a list of the ascertained ages of certain trees :—

Elm.....	335 years.
Cypress, about.....	350 —
Cheirostemon (Hand-tree), about.....	400 —
Ivy.....	450 —
Larch	576 —
Sweet Chestnut, about.....	600 —
Orange.....	630 —
Olive.....	700 —
Platanus Orientalis.....	720 —
Cedar.....	800 —
Many tropical trees, according to Humboldt, about	1000 —
Lime.....	1076, 1147 —
Oak.....	810, 1080, 1500 —
Yew	1214, 1458, 2588, 2820 —
Taxodium, upwards of.....	4000 —
Adansonia.....	5000 —

10.—GENERAL OBSERVATIONS ON THE ORGANS OF PLANTS, AND ON THE MODE IN WHICH THEY ARE ARRANGED.

638. Before concluding the consideration of the elementary and compound organs of plants, it is proposed to make some general observations on their arrangement and development. The following is a tabular view of the various organs to which attention has been directed :—

GENERAL ARRANGEMENT OF THE ORGANS OF PHANEROGAMOUS PLANTS.

I.—Elementary Organs.

Membrane.....	{ Vesicles or Cellules..... Cellular Tissue. Spiral Vessels or Tracheae. } Ducts or Tubes..... } Vascular Tissue.
Fibre.....		

II.—Compound Organs.

1. General Integument.

Cuticle or Pellicle....	{Epidermis.
Stomata.....	

Hairs, Prickles or Aculei, Stings, Glands.

2. Nutritive Compound Organs.

Spongioles....	{ Roots.
Fibrils.....	

Pith.....	} Stem and Branches.	Rhizome.
Medullary Sheath.....		Tuber.
Heartwood or Duramen.....		Runner.
Sapwood or Alburnum.....		Sucker.
Medullary Rays or Plates.....		Corm.
Liber or Endophlœum.....		Bulb.
Cortical Layers or Epi- and Mesophlœum.....		Thorn.
Petiole.....	} Leaf.	Phyllodia.
Limb or Lamina....		Tendrils.
Stipules.		Ascidia.

3. Reproductive Compound Organs.

Bract.			Involucre.	
		Sepals—Calyx,	} Perianth or Perigone.	
		Petals—Corolla,		
Fovilla } Pollen }	} Anther.....	} Stamens	} Flower.	
granules.... } grains, }				
	Filament... }			
Carpels.... } Ovary, }	} Pistil.....			
Ovules.... } Style, }				
	{ Stigma, }			

4. Composition of Ripe Fruit.

Radicle.....	} Embryo.....	} Pericarp. }	} Fruit.
Cotyledon.....			
Plumule.....			
	Spermoderm.....	} Seed..... }	
	Albumen or Perisperm.....		



639. Plants may be said to be composed of numerous individuals, each having a sort of independent existence, and all contributing to the general growth of the compound individual formed by their union. In the case of a tree there are a vast number of buds, each of which is capable of being removed, and made to grow on another tree by grafting; and although each has thus a vitality of its own, it is nevertheless dependent on the general vitality of the tree, so long as it is attached to it. The same thing is seen in Sertularian Zoophytes. Each of the individuals forming a compound plant is called by Gaudichaud a *phyton* (*φυτόν*, a plant), and in it he recognizes three parts or *merithalli* (*μέρος*, a part, and *θαλλός*, a frond), the *radicular merithal* corresponding to the root, the *cauline* to the stem, and the *foliar* to the leaf.

640. In the Acotyledonous plants, the embryo or spore consists of cells united together, and it is only during germination that it exhibits these different parts. In Monocotyledons, the embryo consists of a single phyton, with a radicular merithal or radicle, a cauline or tigellus, and a foliar or cotyledon. In Dicotyledons, the embryo consists of two or more pytons united, with their foliar merithals (cotyledons) distinct, while their cauline and radicular merithals form each a single organ.

641. In tracing the various parts of plants, it has been shown that all may be referred to the leaf as a type. This morphological law was propounded by Linnæus and Wolff, but it is to Goethe we owe the full enunciation of it. Vegetable morphology, the study of forms, or the reference of the forms of the parts of plants to the leaf, is now the basis of organography; and it will be observed, that in considering the various organs, this has been kept constantly in view. The calyx, corolla, stamens, and pistil, are only modifications of the leaf adapted for peculiar functions. It is not meant that they were originally leaves, and were afterwards transformed; but that they are formed of the same elements, and arranged upon the same *plan*, and that in the changes which they undergo, and the relation which they bear to each other, they follow the same laws as leaves do. The different parts of the flower may be changed into each other, as into true leaves; or, in other words, the cellular papillæ from which they are formed are capable of being developed in different ways, according to laws which are still unknown. These changes may take place from without inwards, by an *ascending* or *direct metamorphosis*, as in the case of petals becoming stamens; or from within outwards, by *descending* or *retrograde metamorphosis*, as when stamens become petals.

642. Bracts are very evidently allied to leaves, both in their colour and form. Like leaves, too, they produce buds in their axil. The monstrosity called Hen and Chicken Daisy, depends on the development of buds in the axil of the leaves of the involucre. The sepals frequently present the appearance of true leaves, as in the Rose. The petals sometimes become green like leaves, as in a variety of *Ranunculus Philonotis*, mentioned by Decandolle, and in a variety of *Campanula rapunculoides*, noticed by Dumas. At other times they are changed into stamens. Decandolle mentions a variety of *Capsella Bursa-pastoris*, in which there were ten stamens produced in consequence of a transformation of petals. The stamens in double flowers are changed into petals, and in *Nymphæa alba* there is a gradual transition from the one to the other. Sometimes the stamens are changed into carpels, and bear ovules. This has been seen in Wall-flower, some Willows, Poppy, &c. Petit-Thouars noticed a plant of House-leek, in which the one-half of the anthers bore ovules, and the other half pollen. The carpels, as in the double Cherry, may be seen in the form of folded leaves; in double flowers they are transformed into petals, and in other cases they are developed as stamens. It is said that increase of temperature, and luxuriance of growth, sometimes make flowers produce stamens only. In plants having unisexual flowers, this is more liable to take place, as in Melon, Cucumber, &c. Increased vigour seems to be required for the development of stamens, for some fir trees in their young state bear cones, and produce male flowers only when they reach the prime of life.

643. **Symmetry of Organs.**—In the progress of growth, the plants belonging to the different divisions of the vegetable kingdom follow certain organogenic laws ($\delta\epsilon\gamma\alpha\nu\nu\sigma$, an organ, and $\gamma\epsilon\nu\nu\acute{\alpha}\epsilon\iota\nu$, to produce), the operation of which is seen in the definite arrangement of their organs. The flower consists sometimes of three, at other times of four or five equal sets of organs, similarly and regularly disposed. Thus, the Iris has three straight parts of its perianth, and three reflexed ones alternately disposed, while the Fuchsia has four parts of the calyx alternating with four petals, and the Rose has five alternating portions. This orderly and similar distribution of a certain number of parts is called *symmetry*, and flowers are thus said to be symmetrical with various numbers of members. When the number of parts is two, the



flower is *dimerous* ($\delta\iota\varsigma$, twice, and $\mu\acute{\epsilon}\rho\omicron\varsigma$, a part) (fig. 531), and the symmetry two-membered. When the number of parts is three, the flower is *trimerous* ($\tau\epsilon\acute{\iota}\varsigma$, three), and when the parts are arranged in an alternating manner (fig. 532), the symmetry is *trigonal* or *triangular* ($\tau\epsilon\acute{\iota}\varsigma$, three), and *γωνία*, an angle), as in the Lily. When there are four parts, the flower is *tetramerous* ($\tau\epsilon\tau\alpha\acute{\rho}\varsigma$, four), and the symmetry is *tetragonal* or *square* (figs. 533, 534), as in Galium and Paris.

When there are five parts, the

flower is *pentamerous* ($\pi\acute{\epsilon}\nu\tau\epsilon$, five), and the symmetry *pentagonal* (fig. 535), as in Ranunculus. The number of parts in the flower is indicated by the following symbols:—Dimerous $\sqrt{2}$ Trimerous $\sqrt[3]{}$ Tetramerous $\sqrt[4]{}$ Pentamerous $\sqrt[5]{}$.

644. There are also other kinds of arrangements in flowers, which may be referred to certain modifications in the organogenic law. Thus, what is called *oblong* or *two and two-membered* symmetry, occurs in cases where the opposite ends are similar, and the opposite sides as in the arrangement of the stamens of Cruciferæ. Again, *simple* symmetry is that in which the two sides of the object are exactly alike, without

Fig. 531.—Diagram of the dimerous flower of *Circea Lutetiana*, Enchanter's Nightshade. There are two carpels, two stamens, two divisions of the corolla, and two of the calyx. The flower is Isostemonous.

Fig. 532.—Diagram of the trimerous Isostemonous flower of *Cneorum tricoccon*. The floral envelopes are arranged in sets of three, and so are the essential organs.

Fig. 533.—Diagram of the tetramerous Isostemonous flower of *Zieria*. The organs are arranged in verticils of four parts each.

Fig. 534.—Diagram of the tetramerous Diplostemonous flower of *Ruta graveolens*. There are four carpels, eight stamens, or four in each verticil, four foliols of the calyx, and four petals.

any further repetition, as in papilionaceous, personate, and labiate flowers, as well as in most leaves. The term symmetry, however, is properly confined to cases where the parts are arranged alternately, and are either equal or some multiple of each other, and has no reference to the forms of the different parts. In the very young state, the parts of the flower appear as a shallow rim, from which the petals and sepals arise as mammillæ, in a symmetrical manner. In the case of irregular corollas, the parts at first appear regular, as shown by Barneoud.* In speaking of flowers, it is usual to call them symmetrical when the sepals, petals, and stamens follow the law mentioned, even although the pistil may be abnormal. Thus, many Solanaceæ are pentamerous, and have a dimerous ovary, yet they are called symmetrical. In Cruciferae, the flowers are, properly speaking, unsymmetrical, for while there are four sepals and four petals, there are six stamens in place of four. This depends apparently on the long stamens being in reality composed only of two, the filament of each of which is split by a process of chorization (§ 383), and each division forms for itself by multiplication a perfect anther. In Papilionaceous flowers, the parts are usually symmetrical, there being five divisions of the calyx, five petals, and ten stamens in two rows.



645. It will be seen that flowers constituting trigonal or pentagonal symmetry, may present what has been called simple symmetry, when one of the petals or sepals becomes more developed than the others. In Dicotyledonous plants, it is common to meet with pentagonal (figs. 535, 536, 537) and tetragonal (figs. 533, 534), symmetry, the parts being arranged in fives and fours, or in multiples of these numbers. It is common to find the stamens more numerous than the petals, and in that case they are arranged in different verticils, each alternating

* Annales des Sciences Naturelles, November, 1846.

Fig. 535.—Diagram of the pentamerous Isostemonous flower of *Crassula rubens*. c c c c c, Parts of the calyx. p p p p p, Petals alternating with the leaves of the calyx. e e e e e, Stamens alternating with the petals. a, Accessory bodies in the form of scales, or a disk alternating with the stamens. These scales are often an abortive row of stamens. o, Carpels alternating with the stamens, and opposite to the scales.

Fig. 536.—Diagram of the pentamerous flower of *Sedum Telephium*. The stamens are ten, arranged in two alternating verticils. The flower is Diplostemonous.

Fig. 537.—Diagram of the pentamerous Diplostemonous flower of *Coriaria myrtifolia*; the parts of the four whorls alternating, the verticil of stamens being double.


Fig. 538.—Diagram of the trimerous Diplostemonous flower of *Ornithogalum pyrenaicum*. Stamens six in two alternating verticils.

with that next it. Thus if there are five sepals, five petals, and twenty stamens, the latter are considered as forming four verticils. No doubt the verticils are often traced with difficulty, more especially when adhesions take place. In Monocotyledons (fig. 538), the parts are usually in sets of three, or in some multiple of that number, exhibiting trigonal symmetry. In Acotyledons, when any definite number can be traced, it is found to be two, or some multiple of two. The teeth of Mosses are in sets of four, or some multiple of four. The spores of many Acotyledons are also arranged in fours.

646. **Teratology.**—There has thus been traced a tendency to symmetrical arrangement in plants. But the parts of plants are often modified by natural causes which cannot be explained. It is assumed that each of the similar members of a flower have the same organization, and a similar power of development; and hence, if among these similar parts some are less developed than others, they are considered as *abortive*, and these abnormal states are traced to changes which take place in the earlier stages of growth. Such changes often interfere with the symmetry of the flower. Alteration in the symmetrical arrangement, as well as in the forms of the different parts of plants, have been traced to *suppression* or the *non-development* of organs, *degeneration* or imperfect formation, *adhesion* or union of one part to another, *multiplication* of parts, and *unlining* or *chorization*. The study of *Teratology* (τέρας, a monstrosity, and λόγος, treatise), or of the monstrosities occurring in plants, has led to many important conclusions relative to the development of organs, and it is only by tracing the parts of plants through all their stages and transformations, that correct ideas can be formed as to their relations and forms.

647. By *suppression* is meant the non-appearance of an organ at the place where it ought to appear if the structure was normal; the organ being wanting to complete the symmetry. This suppression is liable to occur in all the parts of plants, and gives rise to various abnormalities. Suppression may consist in the non-appearance of one or more parts of certain verticils, or of one or more entire verticils. In the flowers of *Staphylea* (fig. 539), there are five parts of the calyx, five petals, five stamens, and only two carpels; in many *Caryophyllaceæ*, as *Polycarpon* and *Holosteum* (fig. 540), while the calyx and corolla are pentamerous, there are only three or four stamens and three carpels; in *Impatiens noli-me-tangere* (fig. 541), the calyx is composed of three parts, while the other verticils have five; in *Labiæ* flowers, there are five parts of the calyx and corolla, and only four stamens; and in *Tropæolum pentaphyllum* (fig. 542), there are five sepals, two petals, eight stamens, and three carpels. In all these cases, the want of symmetry is traced to the suppression of certain parts. In the last mentioned plant, the normal number is five; hence it is said that there are three petals suppressed, as shown by the position of the two

- remaining ones (fig. 542); there are two rows of stamens, in each of which one is awanting, and there are two carpels suppressed. In many



instances the parts which are afterwards suppressed can be seen in the early stages of growth, and occasionally some vestiges of them remain in the fully developed flower. Sometimes the whorl of the petals is awanting, the flowers being *apetalous* (α , privative, and $\pi\acute{\epsilon}\tau\alpha\lambda\omicron\nu$, a leaf) (fig. 543), and in such cases it is common to see the stamens opposite to the segments of the calyx, as in *Chenopodiaceæ* (fig. 544). That this suppression of the petals takes place is shown in the case of certain allied plants, as in the natural orders *Caryophyllaceæ* and *Paronychiaceæ*, where some species have petals and others want them.



648. By the suppression of the verticil of the stamens or of the carpels, flowers become *unisexual* (*unus*, one, and *sexus*, sex), or *diclinous* ($\delta\iota\varsigma$, twice, and $\kappa\lambda\acute{\iota}\nu\eta$, a bed), and are marked thus, ♂ ♀; the first of these symbols indicating the male, and the second the female flower. Thus, in *Jatropha Curcas* (fig. 314), the flowers have five segments of the calyx, and five petals, while in some (fig. 314, 1) the pistil is awanting; in others (fig. 314, 2), the stamens. In the genus

Fig. 539.—Diagram of the flower of *Staphylea pinnata*. The parts of the calyx, corolla, and stamens are pentamerous, while the pistil, in consequence of the suppression of three carpels, is dimerous.

Fig. 540.—Diagram of the flower of *Holosteum umbellatum*. There are five calycine divisions, and five petals; but the stamens, by the suppression of one, are only four in number; while the carpels are, by suppression, reduced to three. Thus the flower is unsymmetrical.

Fig. 541.—Diagram of the flower of *Impatiens parviflora*, with one of the calycine leaves spurred. There are five carpels, five stamens, five petals, one of which is larger than the rest, but only three parts of the calyx, in consequence of suppression.


Fig. 542.—Diagram of the flower of *Tropæolum pentaphyllum*, with a spurred or calcarate calycine leaf. The petals, by suppression, are reduced to two; the stamens are eight in place of ten, and the carpels three in place of five.

Fig. 543.—Diagram of the flower of *Glaux maritima*, showing the suppression of the verticil of the corolla. There are five divisions of the calyx, five stamens alternating with them, and five divisions of the ovary, with a central placentation.

Fig. 544.—Diagram of the flower of *Chenopodium album*, showing the suppression of the verticil of the corolla. The five stamens, in this case, are opposite to the divisions of the calyx, thus exhibiting the arrangement which might be expected from a non-development of the corolla. The divisions of the ovary are not easily seen, the placentation being central.

Lychnis, there are usually stamens and pistil present, or the flower is hermaphrodite, or *monoclinous* (μόνος, one, and κλίνη, a bed); but in Lychnis dioica, some flowers have stamens only; others pistils only. Thus it is that *monoecious* and *dioecious* (μόνος, one, δις, twice, and οίκιον, a habitation) plants are produced by the suppression of the essential organs of the flowers, either of the same or of different individuals of the same species; while *polygamous* (πολύς, many, and γάμος, marriage) plants are those in which, besides unisexual, there are also hermaphrodite or perfect flowers.

649. Some parts of the pistil are generally suppressed in the progress of growth, and hence it is rare to find it symmetrical with the other whorls. When the fruit was treated of (§ 522), it was shown that carpels and ovules often become abortive by pressure and absorption, so that the pericarp and seeds differ in their divisions and numbers from the ovary and ovules. If the whorls of the calyx and corolla are awanting, the flower becomes *naked* or *achlamydeous* (§ 351). It may still, however, be fitted for the functions of producing seed; but if the essential organs, viz., the verticils of stamens and pistils, are suppressed, then the flower, however showy as regards its envelopes, is unfit for its functions, and is called *neuter*. Flowers having stamens only, are *staminiferous*, *staminal*, *sterile*; or those having pistils only, are *pistilliferous*, *pistillate*, or *fertile*. The suppression of



various verticils, and parts of them, is well seen in the family of the Euphorbiaceæ (figs. 545—550). Thus, in fig. 545 is delineated an apetalous trimerous staminal flower; in fig. 546 one of the stamens is suppressed, and in fig. 547 two of them are awanting. Again, in figs. 548, 549, 550, the calyx is suppressed, and its place occupied by one, two, or three bracts (so that the flower is, properly speaking, *achlamydeous*), and only one or two stamens produced. In fig. 550, 1, there is a sterile flower, consisting of a single stamen with a bract; and in fig. 552, 2, a fertile flower, consisting of a single carpel with a

Figs. 545—550.—Diagrams of flowers of Euphorbiaceæ plants, becoming more and more simple. (1.) The calyx is the only envelope, and consists of three parts in figs. 545, 546, and 547. It is completely suppressed in figs. 548, 549, and 550, and its place is occupied by a bract, in the axil of which the flower is produced; this bract being accompanied in figs. 548 and 549 with two small bractlets. (2.) The male flowers in fig. 545 have three stamens, in figs. 546 and 548 they have two, in figs. 547 and 549 one stamen only is developed, and in fig. 550, 1, the solitary stamen has only one anther-lobe. (3.) The female flower in fig. 550, 2, is reduced to a single carpel, with a bract in the axil of which it is produced.

Fig. 545.—Diagram of a staminiferous flower of *Tragia cannabina*.

Fig. 546.—Diagram of a staminiferous flower of *Tragia volubilis*.

Fig. 547.—Diagram of a staminiferous flower of *Anthostema senegalense*.

Fig. 548.—Diagram of a staminiferous flower of *Adenopeltis colliguaya*.

Fig. 549.—Diagram of a staminiferous flower of a *Euphorbia*.

Fig. 550.—1. Diagram of a staminiferous flower of *Najas minor*. 2. Of a pistilliferous flower of *Najas major*.

bract. There is thus traced a *degradation*, as it is called, from a flower with three stamens and three divisions of the calyx, to one with a single bract and a single stamen or carpel.

650. It is common to find some of the buds of a plant suppressed, thus altering the spiral arrangement. Such buds, however, are often capable of being developed, if any accident occurs, or if the plant is pruned. Deficiency of light and of air, and want of proper nourishment, are capable of producing abortions of various kinds. The non-development of a branch gives rise to *clustered* or *fascicled* (*fascis*, a bundle of twigs) leaves, as in the Larch, and to fascicled twigs, as in a common bird-nest-like monstrosity of the Birch. When the true leaves of a plant are suppressed, their place may be occupied by a tendril, as in *Lathyrus Aphaca*, in which the stipules perform the functions of leaves (§ 201); or the petiole may be developed in a peculiar way, as in the phyllodia (§ 157) of some *Acacias*.


651. *Degeneration*, or the transformation of parts, often give rise either to an apparent want of symmetry, or to irregularity in form. Branches, when not properly developed, may assume the form of thorns or spines (§ 200), as in the Hawthorn and Wild-plum; and by culture these spines may be converted into leaf-bearing branches. Leaves often become mere scales, as in *Lathræa*, *Orobanche*, and in *Bulbs*. The limb of the calyx may appear as a rim, as in some *Umbelliferæ*; or as pappus, in *Compositæ* and *Valeriana*. In *Scrophularia*, the fifth stamen appears as a scale-like body, called *staminodium* (fig. 346); in many other plants belonging to the *Scrophulariaceæ*, it assumes the form of a filament, with hairs at its apex in place of an anther. In unisexual flowers, it is not uncommon to find vestiges of the undeveloped stamens in the form of filiform bodies or scales. To many of these staminal degenerations, Linnæus gave the name of *nectaries*. In double flowers, transformations of the stamens and pistils take place, so that they appear as petals. In *Cannæ*, what are called petals are in reality metamorphosed stamens. Allusion has already been made to the various changes which the different parts of the flower thus undergo. The object of the florist is to produce such monstrosities; and flowers, which by him are considered perfect, are looked upon by the botanist as imperfect, from the want of the essential organs.

652. *Adhesion*, or the growing together of parts, is a very common cause of changes both as regards form and symmetry. The union of stems gives rise occasionally to anomalies, as in the *fasciated* stalk of *Cockscomb* (fig. 230), and the flattened stems of some *Coniferæ* (§ 197), and probably also the peculiar stems of certain *Sapindaceæ* and *Menispermaceæ* of Brazil (§ 90). Some of these, however, may perhaps be traced not to adhesion, but to an abnormal development of buds, producing wood only in one direction in place of all round.

Natural grafts occasionally occur from one branch of a tree uniting to another. Roots also sometimes become grafted, and to this has been attributed the vitality occasionally preserved by the stumps of Spruce-firs which have been felled on the Swiss Alps. The union of two leaves by their base, forms a connate leaf, and the adhesion of the lobes of a single leaf on the opposite side of the stalk, gives rise to perfoliate leaves (fig. 156). The union of the edges of a folded leaf forms *Ascidia*, or pitchers (figs. 184, 187). The different parts of the same verticil of the flower unite often more or less completely, giving rise to a monophyllous or gamophyllous involucre (§ 347); a monosepalous or gamosepalous calyx (fig. 273); a monopetalous or gamopetalous corolla (figs. 293, 294, &c.); monadelphous (figs. 307, 314, 1), diadelphous (§ 399), and polyadelphous (figs. 315, 551) stamens; syngenesious anthers (§ 417); a gynandrous column (§ 400); and a syncarpous ovary (fig. 383). The different verticils of the flower are frequently adherent. The calyx is often



551



552


united to the corolla or to the stamens, or both (fig. 308); the stamens may adhere to the corolla (fig. 552); or there may be a union of the four verticils of the flower, so that the calyx becomes superior (fig. 309). In some instances, when the axis is elongated, adhesions take place between it and certain whorls of the flower. Thus, in some Caryophyllaceæ (fig. 553), the calyx, *c*, bearing the stamens, *e*, and petals, *p*, becomes united to the axis, *g*, which supports the ovary, *o*. In Capparidaceæ (fig. 554), the calyx, *c*, and petals, *p*, occupy their usual position, but the axis is prolonged in the form of a gynophore, *ag*, to which the stamens, *e*, are united. Occasionally, contiguous flowers may unite, giving rise to double fruits, as is sometimes seen in Apples, Grapes, and Cucumbers.

653. *Multiplication*, or an increase of the number of parts, gives rise

Fig. 551.—One of the five bundles of stamens taken from the polyadelphous flower of *Malva miniata*. Stamens are united by their filaments.

Fig. 552.—Portion of the gamopetalous or monopetalous corolla, *p*, of a *Collomia*, showing part of the tube, *t*, terminated by two lobes of the limb, *l*, and having the stamen, *e*, inserted into it, and united to it, so that the upper part of the filament, *f*, only is free.

to changes in plants. It is often found, that in plants belonging to the same natural order, the number of stamens in one is greater than that



553



554

in another, either in consequence of additional stamens being developed in the verticil, or on account of the production of additional verticils. The same thing is met with in the case of the other whorls, and is well illustrated in the formation of the disk (§ 428). Multiplication causes a repetition of successive whorls, which still follow the law of alternation.

654. Parts of the flower are often increased by a process of *deduplication*, *unlining*, *dilamination*, or *chorization*, i.e. the separation of a lamina from organs already formed (§ 383). This is believed to take place in a remarkable degree in the case of appendages to petals. Thus, in *Ranunculus*, the petal (fig. 555) has a scale at its base, *a*, which is looked upon as a mere fold of it. This fold may in some cases be more highly developed, as in *Caryophyllaceæ*, and in *Crasula rubens* (fig. 258 *a*), and it may even assume the characters of a stamen, which will therefore be opposite the petal, as in *Primulaceæ*. Some do not consider the production of scales or stamens opposite to the petals, as the result of chorization. Lindley argues against it from what is observed in *Camellia japonica*, in which the petals are usually alternate; but, by cultivation, the law of alternation is interfered with, and the parts are so developed that the petals are

Fig. 553.—Flower of *Lychnis viscaria*, one of the *Caryophyllaceæ*, cut lengthwise, to show the relation of its different parts. *c*, Gamosepalous calyx. *p*, Petals with their elongated unguis or claw, *u* *u*, their limb, *l* *l*, and the appendages, *a* *a*, in the form of dilaminated scales of the petals. *e* *e*, Stamens. Pistil consists of the ovary, *o*, and five styles, *s*. *g*, Prolongation of the axis in the form of a gynophore, or anthophore bearing the petals, the stamens, and the pistil.

Fig. 554.—Flower of *Gynandropsis palmipes*, one of the *Capparidaceæ*. *c*, Calyx. *p*, Petals. *e* *e*, Stamens. *a* *g'*, Gynophore or elongated internode or axis bearing the stamens. *a* *g''*, Gynophore or elongated internode bearing the pistil. *o* *f*, Pistil composed of an ovary, *o*, a style and a stigma, *f*.

opposite, and run in several regular lines, from the centre to the circumference. Again, by this process of chorization, one stamen may give rise to several. Thus, in *Luhea paniculata* (fig. 316, 1), in place of five stamens there are five bundles (fig. 316, 2), composed partly of sterile filaments, *f s*, and partly of filaments bearing anthers, *f a*; and each of these bundles is traced to a deduplication of a single stamen, inasmuch as they arise from one point, and do not follow the law of alternation. Thus, dilamination repeats the single organs, and causes opposition of parts. In the case of the four long stamens of *Cruciferae* (§ 644), chorization is said to take place by a splitting of the filaments of two stamens; and thus the two stamens on each side are, by *geminatio* (*gemini*, twins), normally one. This view is confirmed by cases in which the filaments of the long stamens are more or less united; also, by cases in which the shorter filaments exhibit tooth-like processes on either side, while the longer ones have them only on the outer side. In such cases, the two long filaments, if united, would present the same appearance as the shorter ones, and occupy their usual position of alternation with the petals. In some instances, by *pelorization* (*πελώριος*, monstrous), it is found that tetradynamous plants become tetrandrous with equal stamens alternating with the petals.

655. *Cultivation* has a great effect in causing changes in the various parts of plants. Many alterations in form, size, number, and adhesion of parts, are due to the art of the horticulturist. The development of cellular tissue and of starchy matter is often thus much increased, as may be seen in the case of Turnips, Carrots, and Potato. The succulence of the leaves of the Cabbage and Lettuce, and the formation of a *heart*, as it is called, is due to cultivation; so also the curled leaves of Savoys, Cress, Endive, &c. The changes in the colour and forms of flowers thus produced are endless. In the *Dahlia*, the florets are rendered quilled, and are made to assume many glowing colours. In *Pelargonium*, the flowers have been rendered larger and more showy; and such is also the case with the *Ranunculus*, the *Auricula*, and the *Carnation*. Some flowers, with spurred petals in their usual state, as *Columbine*, are changed so that the spurs disappear; and others, as *Linaria*, in which one petal only is usually spurred, are altered so as to have all the petals spurred.




Fig. 555.—Petal of *Ranunculus Ficaria*, viewed on the inside. *l*, The limb. *a*, Small scaly appendage at its base formed by chorization or dilamination.

SECTION IV.—SOME GENERAL PHENOMENA CONNECTED WITH VEGETATION.

1.—VEGETABLE IRRITABILITY.

656. Under this head are included certain sensible movements of living plants, exercised without any direct application of mechanical force, and not referable to mere elasticity, or the hygroscopic nature of the tissues. These motions are influenced chiefly by light and heat, and, like many phenomena occurring in organized beings, they cannot at present be explained by chemical or mere mechanical laws. They may, however, be excited by a stimuli of a chemical or mechanical nature. Although the cause of them is obscure, still, in some instances, their use is obvious.

657. Among the lowest classes of plants there are some peculiar movements of this kind. The simplest members of the Sea-weed tribe occasionally move throughout their whole substance. Oscillatorias, which are filaments composed of cells placed end to end, containing fluid and granular matter, have an undulating movement, by means of which they advance. When placed in fluids under the field of the microscope, some of them may thus be seen to pass from one side to the other. The filaments sometimes twist up in a spiral manner, and then project themselves forward by straightening again. The motions are influenced by temperature and light, and by some are considered as being connected with the production of new cells. The spores of many Cryptogamic plants, especially species of *Vaucheria*, *Conferva*, and *Prolifera*, exhibit motions which, according to Thuret and Decaisne, depend on the presence of cellular hair-like processes, called *cilia* or *tentacula*. These motive organs are in a state of constant agitation, which lasts for some hours, becoming slower, and finally ceasing after germination has commenced. In the spores of *Conferva glomerata* and *rivularis* (fig. 431), there are two of these cilia or filiform tentacula, which project from a colourless rostrum. In *Chætophora elegans*, *var. fusiformis*, four have been seen (fig. 432); in *Prolifera* (fig. 433), there is a circle of cilia, and in *Vaucheria* (fig. 434), the spore is entirely covered with very short cilia, the vibration of which determines their forward movement. These spores, from their movements, have received the name of Zoospores (§ 492). Mr. Thwaites accounts for the rhythmical movements of cilia by electrical currents. In certain cells of Cryptogamic plants, especially in what are called Antheridia, bodies are met with called Phytozoa (§ 492), which also exhibit movements during a part of their existence.

658. Remarkable movements have also been observed in the higher

classes of plants. The fovilla contained in the pollen-grain in a young state, when moistened with water, exhibits movements when viewed under the microscope. These movements have by some been referred to irritability, but by Brown and other accurate observers, they are considered as merely molecular, and similar to what takes place between the minute particles of inorganic matter—as, for instance, finely powdered Gamboge suspended in water. These fovilla movements are easily seen in the very young pollen of *Antirrhinum majus*. Certain movements also take place in the floral envelopes. Thus many flowers open and close at particular periods (§ 483—485); these phenomena depending on light, temperature, and moisture. Leaves also, especially those which are compound, are folded at certain periods in a distinct and uniform manner. What was called by Linnæus *the sleep of plants*, is the change produced on leaves by the absence of light. It is by no means analogous to the sleep of animals. During darkness some are slightly twisted and hang down; others, such as pinnate and ternate leaves, have the leaflets folded together, and frequently the common petiole depressed. The youngest leaflets first exhibit these changes; and when the plants become old, and their tissues hardened, the irritability is often much diminished, as is seen in Oxalises. The folding of the leaflets of compound leaves usually takes place from below upwards, but sometimes in the reverse manner, as in *Tephrosia Caribæa*; so also with the common petiole, which is directed upwards during sleep in the *Cassias*, and downwards in *Amorpha*. When besides the common petiole there are partial petioles, as in the Sensitive plant, they may be bent inwards towards each other, while the former is bent downwards.

659. *Mimosa sensitiva* and *pudica*, commonly called *sensitive plants*, display these movements of their leaves in a remarkable degree, not only under the influence of light and darkness, but also under mechanical and other stimuli. They have bipinnate leaves with four partial petioles proceeding from a common rachis, and each of the petioles is furnished with numerous pairs (about twenty) of leaflets, which are expanded horizontally during the day. During darkness, or when touched or irritated in any way, each leaflet moves upwards towards its fellow of the opposite side, which in its turn rises up, so that their upper surfaces come into contact. When the movement commences at the apex of the leaf, it usually proceeds downwards to the base, and thence may be communicated to the leaflets of the next partial petiole, and ultimately to the common petiole, which falls down towards the stem. The partial petioles then converge towards each other, and have a tendency to become parallel to the common petiole, at the extremity of which they are suspended. When the plant is shaken as by the wind, all the leaflets close simultaneously, and the petioles drop together. If, however, the agitation is long continued,

the plant seems as it were to become accustomed to the shock, and the leaflets will expand again. The stem itself is not concerned in the movements. It may be cut and wounded cautiously without causing any change in the leaves, and a portion of it may be removed with a leaf attached, and still remaining expanded. If, however, a mineral acid is applied to the stem, after some time the petioles will fall and the leaflets collapse—the leaves perishing with the stem which has been moistened. The chemical action of the acid and absorption cause these phenomena. When a sensitive plant is exposed to artificial light during the night, Decandolle found that its leaves expanded, and that they closed when put into a dark room during the day, showing the influence which light has on these phenomena. It is to be remarked, however, that if the plant is kept for a long period of time in darkness, it will ultimately expand its leaves, and the phenomena of folding and opening will go on, although at very irregular intervals.

660. The ternate leaves of many species of *Oxalis* fold not merely during darkness, but also when agitated or struck lightly and repeatedly. Each of the leaflets folds upon itself, and then bends downwards upon the common petiole. The plant called *Desmodium gyrans*, the moving plant of India, has compound leaves consisting of a large terminal leaflet, and usually two smaller lateral ones. The latter are in constant movement, being elevated by a succession of little jerks, until they come into contact, and sometimes even slightly cross each other; after remaining in this position for a short time, they separate from each other, and move downwards by rapid jerks on opposite sides of the petiole. This process is constantly repeated, and goes on in a greater or less degree, both during day and night, but is most vigorous during warm moist weather. The large terminal leaflet undergoes movements also, oscillating very gradually from one side to the other, and becoming horizontal or depressed. By the lateral oscillatory movement, the leaf becomes inclined in various ways, often assuming a remarkable oblique direction. The upward and downward movements seem to depend on the influence of light and darkness. During the day the leaf becomes more or less horizontal, while during darkness it hangs down. Similar movements are seen in other species of *Desmodium*, as *D. gyroides* and *vespertilionis*.

661. The movements in these cases have, by Martius and Meyen, been referred to the presence of some structure analogous to the nervous system in animals. There is, however, no evidence of such a structure in plants, and these authors have not pretended to prove its existence. It is to be remarked, that the movements differ in many respects from animal contractility. They are usually *hinge* movements, and may be referred to certain changes in the organs, causing distention or contraction of these tissues. Dutrochet and Morren refer them to alterations in the circulation of fluids and air in the vessels

and cells. In plants with irritable leaves, there are frequently swellings where the leaflets join the stalk, as well as where the stalk joins the stem. These swellings contain cells which differ in their dimensions and their contents, and the movements are considered as being produced by changes in the contents of the cells, some of which become more distended than others, and thus cause incurvation or folding. In these swellings the vascular bundles are disposed in a circle near the periphery, and may be concerned in the movements. Mechanical and chemical stimuli are supposed to act by inducing alterations in the contents of the vessels and cells.

662. In the case of the sensitive plant, if the swelling at the base of the common petiole is touched even slightly on its lower side, it is followed by instant depression of the whole leaf, but no such effect is produced if the upper portion of the swelling is lightly touched. Again, touching the little swelling at the base of each leaflet on its upper side, causes the upward movement of the leaflet, but no such effect follows cautious touching of the lower part of the swelling only. If a pair of leaflets is touched at the extremity of a petiole, the irritation is usually continued downwards from apex to base; but if a pair at the base are touched, the progress of folding is reversed. Clear warm weather, with a certain degree of moisture, seem to be the conditions most favourable for these movements. They are seen best in young plants. The leaves of the sensitive plant contract under the action of electricity and galvanism. Some suppose, that in the sensitive plant there are two kinds of cells connected with the upper and lower sides of the leaves and petioles; the one set being contractile, and causing the closing of the leaflet and the fall of the petiole, the other being acted on chiefly through the circulation. In the case of the petiole, it is conceived that the tissue on the lower side of the swellings is contractile, while that in the upper is distensible. The turgescence of the latter, which is kept up by light, counteracts the contractility of the former, and maintains an equilibrium, so as to keep the petiole erect; but when acted on by cold, mechanical irritation, &c., the equilibrium is disturbed, and the contractility operates in depressing the petiole. A careful microscopic dissection of the swellings, shows peculiar cells in some parts, which seem to differ in their contents from others in their vicinity.

663. In the sensitive species of the *Desmodium* and *Oxalis*, the movements are not so evidently influenced by mechanical irritation. In the former, the little leaflets are supported on swollen petiolules, and it is to the curvation and twisting of these in different directions, that the movement seemed to be owing. The leaflets remain flat and do not fold on themselves. It is said that by arresting the vital actions going on in the leaflets, by giving them a coating of gum, and thus preventing transpiration and respiration, the movements are

stopped, and that they recommence when the gum is removed by water. Cutting a leaflet across, and only leaving a small portion of its lamina attached to the petiole, does not immediately stop the movement of gyration. In such a case, however, the motion ultimately ceases, while it continues in the uncut leaflet. So also, if a leaflet is divided longitudinally into two parts, each of them continues to move for a time, but the motions cease as the process of desiccation goes on.

664. There are occasionally cellular prolongations from the leaf, which, when touched, cause folding. Thus, in *Dionæa muscipula*, or Venus's fly trap, the lamina is articulated to the petiole, and consists of two free portions which are united together by a joint along the midrib. On the upper side of each part of the lamina are situated three hairs with swellings at the base, and when these hairs are touched, the halves approach each other from below upwards, so as to enclose any object, as a fly, which may happen to light on them. Similar movements, but in a much less obvious manner, are said to take place in the leaves of *Drosera* or Sun-dews. The movements are attributed to the same causes as those already mentioned, but the ultimate object is not known.

665. Movements take place in some parts of the flower, occasionally with the view of scattering the pollen on the stigma. The stamens of various species of *Berberis* and *Mahonia* are articulated to the torus or thalamus, and when touched at their inner and lower part, move towards the pistil. In *Parnassia palustris*, the stamens move towards the pistil in succession to discharge their contents. The *Helianthemum vulgare* or common Rock-rose, exhibits staminal movements also connected with the bursting of the anthers. Morren has noticed sensitiveness in the andrœcium of *Sparmannia africana* and *Cereus grandiflorus*. In the Nettle and Pellitory, the filaments are confined in a peculiar way by the perianth, and at a certain period of growth they are released so as to allow their elasticity to come into play, by means of which the pollen is forcibly scattered (§ 497). In *Goldfussia* or *Ruellia anisophylla*, the style has a curved stigmatic apex, which gradually becomes straightened, so as to come into contact with the hairs of the corolla, upon which the pollen has been scattered; and in *Mimulus* and *Bignonia* (fig. 406), the stigma has two expanded lobes which close when touched, a movement apparently in some way connected with fertilization. In the Passion-flower, and some Cacti, the styles move towards the stamens. The species of *Stylidium* have the filaments and styles united in a common column, at the upper part of which the anther-lobes and stigma are placed. The column often projects beyond the flower, and is jointed. At the articulation an irritable swelling occurs, which when touched or acted upon by heat and light, causes a sudden incurvation by which the column is thrown to the opposite side of the flower, bursting the anthers and scattering

the pollen on the central stigma. After a time the column recovers its position. These movements take place in the flower for some time after it has been removed from the plant and kept in water (§ 497). Certain petals in some flowers, as in Orchidaceæ, are said to move. Morren notices this in the case of species of *Megaclinium* and *Pterostylis*. *Gentiana sedifolia* closes its petals when touched. *Drakæa elastica*, a Swan River terrestrial Orchid, is remarkable for the irritability of the stalk of the labellum. This stalk exhibits a moveable joint like an elbow.

666. Chemical agents have an effect on the movements of plants. Some act by causing irritation, others by destroying irritability. Narcotic poisons, as opium, belladonna, and hydrocyanic acid, either taken up by the roots or applied externally, destroy the irritability of plants. They cause closure of the leaves of the sensitive plant, and render it insensible to the action of stimuli. Their prolonged action causes death, but if they are applied in moderate quantity, the plant may recover, and again unfold its leaflets. It frequently happens, however, that the irritability continues for some time much impaired; so that mechanical stimuli do not act in the same rapid and energetic manner as at first. Similar effects are produced by ether and chloroform when sensitive plants are introduced into an atmosphere through which these substances are diffused. The effects may be produced locally by applying the vapour only to certain parts of the plant. Experiments on the action of poisonous agents, both in the fluid and gaseous state, have been performed by Marcet, Christison, Turner, and others (§ 292—296).*

2.—TEMPERATURE OF PLANTS.

667. The heat developed during the expansion of flowers and the preparation of the pollen, especially in the case of Aroideæ, and also at the period of germination, has been already considered (§ 475, 476, &c.) These phenomena appear to be strictly of a chemical nature, and may be traced to the absorption of oxygen, and its combination with the carbon of the starch, the latter being converted into dextrine and grape sugar. It is now proposed to consider the observations which have been made relative to the general temperature of plants.

668. Great differences of opinion have prevailed as to the existence of a *proper* heat in plants. Hunter examined the heat of the internal parts of the trunks of trees by boring holes of different depths in them, and inserting thermometers; and similar experiments were made by Schubler at Tubingen. The result of these experiments were,

* For further observations on Irritability, see Dutrochet, *Recherches Anat. et Physiol. sur la Structure intime des Animaux et Vegetaux*, 1844. Morren on the movements of the leaves of *Oxalis*; on the movement of the style of *Goldfussia anisophylla*, and of the column of *Stylidium*, in *Bulletin. Acad. Roy. de Bruxelles*, tom. iv. v. vi. and xii. Thuret and Decaisne on the movements of spores, in *Annales des Sciences Naturelles*.

1st. That the temperature of trees is higher than that of the air in winter, and lower in summer; 2d. That the temperature corresponds to the depth in the soil to which the roots penetrate; and, 3d. That it depends on the temperature of the fluid matters taken up by the roots, as well as the bad conducting power of the wood of the trees. Dr. Hooker believes that the temperature of the juices of a plant is chiefly dependent on that of the soil at its spongeoles. Dutochet instituted a series of experiments to determine the temperature of the growing parts of plants. He found, by means of a thermo-electric apparatus, that this varied from two or three-tenths of a degree, to one degree above that of the air. This generation of heat only takes place when the plant is active and vigorous, and seems to be connected with processes going on in the interior of the cells. When the cells cease to be actively engaged in the processes of vegetable life, they cease to manifest this vital heat. It reaches a daily maximum, the period of which varies in different plants, according to their vigour. Rameaux has confirmed Dutochet's observations. He observed that during the day, the temperature of the various layers diminished from the surface to the centre, while during the night the inverse took place, and that both of these processes were materially influenced by the nature of the surrounding temperature and the diameter of the tree. There appear, therefore, to be two sources of heat in plants, one depending on organic actions carried on in the growing parts, and the other on meteorological influences, which either act directly through the air, or indirectly through the fluid matters brought up from a certain depth in the earth.*

3.—LUMINOSITY OF PLANTS.

669. Luminous appearances have been observed in certain plants. These have been long noticed in the lower classes of plants, such as Fungi. Decaying wood, in which Fungi are developed, is sometimes luminous. Mr. James Drummond describes some species of Agaric, near the Swan River, growing on the trunks of Banksias and other trees, which emitted at night a phosphorescent light sufficient to enable him to read. A phosphorescent Agaric, with the upper surface of the pileus black, while the centre and gills were white, was noticed by him on the trunk of a dead *Eucalyptus occidentalis*. The *Agaricus Gardneri*, found in Brazil, gives out a light of a pale greenish hue, similar to that of fire-flies. It is found growing on a Palm, and is called *Flor de Coco*. Dedile found luminosity in the *Agaricus olearius*, near Montpellier. In the coal mines of Dresden, certain

* For further remarks on the subject of Vegetable Heat, see Rameaux's papers in the *Annales des Sciences Naturelles*, January, 1843. Schubler's experiments in *Poggendorff Annalen* x. 592; translated in Thomson's *Chemistry of Vegetables*, p. 959. Dutochet, *Ann. des Sc. Nat.* (n. s.) tom. xii. Gardner in *Linn. Trans.* for Dec. 1841, and in *Philosoph. Mag.* for July, 1842.

Rhizomorphous fungi have long been celebrated for the light which they emit. The spawn of the Truffle (*Tuber cibarium*) is said to present similar appearances. Some have said that the luminosity of these fungi, as well as of decaying wood, is increased by exposure to oxygen gas. Some consider it as connected with the absorption of oxygen, being in reality a slow spontaneous combustion; while, according to others, it is referable to the liberation of phosphorus from some of its combinations in the plant.

670. These luminous appearances are said not to be confined to fungi. The younger Linnæus states, that the flowers of *Nasturtium*, Orange Lily, and African Marigold, at the end of a hot summer day, give out intermittent light. Mr. Dowden and Mr. James confirmed this by observations on the common Marigold and *Papaver pilosum*; while other observers have noticed the phenomena in the Sun-flower, French Marigold, species of *Oenothera*, and *Arum*. It is to be remarked, that the flowers said to be thus luminous, are all of a more or less orange colour, and that the phenomenon takes place in still warm summer evenings, towards twilight. Hence, Professor Allman is disposed to attribute them to optical illusions, depending on a peculiar intermittent effect on the retina. Some authors mention the occurrence of luminous sap in plants with milky juices, as the *Euphorbia phosphorea* of Brazil. A rhizome of an endogenous plant from India, is said, when moistened, to acquire a phosphorescent appearance, and to lose this property when dry.

4.—COLOURS OF PLANTS.

671. Colour is not of much importance in Botany as regards classification and arrangement. It is chiefly in the case of Fungi that it is employed as a means of diagnosis. Perhaps the want of an accurate nomenclature of colours in Botany may have in part led to this. Mirbel and Henslow have proposed a nomenclature, which consists in referring all natural colours to certain absolute tints and shades, determined according to fixed laws. Thus, the latter assumes three primaries, as red, blue, and yellow, which together give white light, and derives all others from admixtures of these in definite proportions. On this principle he has constructed a chromatometer (*χρῶμα*, colour, and *μέτρον*, a measure), or measure of colour, the employment of which would lead to an accurate nomenclature.

672. It has already been remarked, that the green colour of the leaves, young bark, calyx, and carpels, depends on the presence of chlorophylle (§ 19). This waxy substance is contained in the deep cells or mesophyllum of leaves, and depends on the action of light for its elaboration. When leaves are grown in darkness, they become colourless from the absence of chlorophylle. Light acts by the fixation

of carbon. The different rays of the spectrum appear to vary in their power of developing the green colour. Senebier performed experiments on the subject, by making the light pass through coloured media, and he was led to the conclusion, that while the yellow rays had the greatest effect on the growth of the plant, the blue and chemical rays were those chiefly concerned in the production of the green colour. Hunt seems to agree with Senebier. Other experimenters, however, as Morren, Daubeny, Draper, and Gardner, think the yellow rays are the most active in producing the green colour. The following table shows the result of some of Gardner's experiments. The rays are denominated *active* or *inactive* in relation to their power of producing a green colour, and the figures under each of them show their power in this respect, 1 being the highest value. The sign — indicates that the effect was not satisfactorily tested:—

Ex. Plants.	Hours of sunshine.	Total time.	Red.	Active Rays.			Inactive.					
				Or.	Yel.	Gr.	Bl.	In.	Vio.			
1. Turnips.....	22	...	109	...	4	2	1	3	...	0	0	0
2. Beans.....	14	...	95	...	—	2	1	3	...	0	—	—
3. Turnips.....	8	...	69	...	4	2	1	3	...	—	—	—
4. Turnips.....	23	...	101	...	—	—	—	1	...	0	0	0
5. Turnips.....	17·5	...	52	...	—	2	1	3	...	4	0	0
6. Turnips.....	5·5	...	6	...	4	2	1	3	...	0	0	0

673. The ray producing the green colour is found to be that which acts most efficiently on the decomposition of carbonic acid, as shown by the following table:—

Places of spectrum examined.	Production of chlorophylle.	Decomp. of CO ₂	Illuminating power.
Extreme Red.....	0·000	0·0000
Commencement of Orange.....	—	·5500
Centre of Orange.....	·777	—
Centre of Yellow.....	1·000	1·0000
Centre of Green.....	·583	—
Centre of Blue.....	·100	—

674. The green colour becomes lighter or deeper according to the quantity of chlorophylle and the aggregation of the cells. It is usually paler on the lower sides of leaves. The dark shades of green in the Yew, Bay, and Holly, are the effect of an immense crowding together of green cells.

675. As light decreases in autumn, the chlorophylle, in many cases, diminishes, and is probably altered by the loss of a portion of carbon. Thus, Evergreen leaves become of a paler colour, and deciduous leaves assume various hues, commonly called autumnal tints. The leaves of the Poplar, Ash, and Beech, before falling, become yellow; those of some species of Rhus, bright red; those of Cornus sanguinea, dull red; those of the Vine, yellow and purple. Berzelius states, that the leaves become red in plants having red fruits. Robinet and Guibourt maintain that the Vines which produce bluish grapes, have red leaves

in autumn, while such as produce white grapes have yellow leaves. These yellow and red colours by some are said to depend on changes in the state of oxidation of the chlorophylle, and have been traced by others to the production of peculiar waxy substances, one red, called *erythrophylle*, the other yellow, *xanthophylle*. Marquart believes that the action of water on chlorophylle, in different proportions, gives rise to yellow and blue matters. Ellis supposed the change of hues to be due to the prevalence of acid and alkaline matters.

676. Dr. Hope endeavoured to show that there is in plants a colourable principle, *chromogen* (χρῶμα, colour, and γεννάω, I generate,) consisting of two separate principles, one of which forms a red compound with acids, while the other forms a yellow with alkalies, and he attributes the green colour produced by the latter to the mixture of the yellow matter with the blue infusion. The two principles, according to him, may exist together, or separate, in different parts of the same plant. In some very fleshy leaves, as Agave, the central cells are pale, while those of the cuticle are coloured and much thickened. Although leaves are usually of a green colour, still they frequently assume various tints. In certain varieties of Beech and Beet, they become of a uniform red or copper colour. In some cases, only one of the surfaces of the leaf is coloured, as in many species of Begonia, Saxifraga, Cyclamen, and Tradescantia, in which they are green above and red or brown below; while in others there is a variation of colour, giving rise to variegation, as in *Acuba japonica*, *Carduus marianus*, and *Calathea zebrina*, where there are yellowish spots, or in many Arums, where they are of a red colour. The whitish or brown spots which occur on leaves, are often produced by thickened cells containing peculiar colouring matter, underlying the chlorophylle cells. In such cases, variegation might be traced to an alteration in the epidermal cells, and the same is true of certain bright colours assumed by the surfaces of some leaves. The juices of many plants are colourless when contained in the vessels, but become milky or coloured by exposure to the air. Thus, the sap of *Ceanothe crocata* becomes yellow, that of *Chelidonium* orange, that of *Madder* changes from yellow to red, and that of some *Boletuses* becomes blue or bluish-green. In some instances, the changes have been prevented by keeping the cut or broken surfaces in nitrogen or hydrogen, or carbonic acid, and thus preventing their exposure to oxygen. It is said, however, that the change of colour in the *Madder* does not take place in pure oxygen.

677. The bark, at first green, becomes often of a brown colour from the thickening of the cell-walls, as well as the deposition of brown matter. Similar changes take place in the woody fibres, giving rise to the coloured duramen of many trees, as the *Laburnum*, *Guaiac*, *Ebony*, &c. Such changes, however, depend on chemical actions going

on in the interior of stems, and are not due to the direct influence of the air. The colour of wood, however, is generally deepened when exposed to the atmosphere.

678. The red, blue, and yellow colours of flowers depend on fluid or semifluid matters contained in superficial cells, which can be detached with the cuticle. This coloured cellular tissue is called by Nourse the *Rete*, and lies immediately below the epidermis. In this respect these colours differ from the green colour in the leaf which is confined to the central cells, and which, as already stated, owes its origin to a granular matter of a peculiar nature. In petals, different cells frequently contain different kinds of colouring matter, thus giving rise to variegation. By the juxtaposition and mechanical mixture of various cells, different tints are produced; and the colours are also modified by the nature of the cuticle through which they are seen. In the interior of petals, the colour is generally more or less yellow, but it is modified when seen through superficial cells. Along with the colouring matter, there is a colourless substance present, the relative quantity of which varies, and hence the colour may be deeper or fainter. In flowers as well as in leaves, the colours appear to depend on the action of light. It has been said, however, that the powerful action of solar light, in some cases, tends to decolorize flowers. Hence, tulips are screened by floriculturists from the direct rays of the sun. The leaves of herbaceous plants also, when exposed to the direct rays of the sun, do not acquire so deep a green as when they are subjected merely to a bright daylight.

679. The colours of flowers have been arranged in two series:— 1st. The *xanthic* (ξανθός, yellow) or yellow; and 2d. The *cyanic* (κύανος, blue) or blue; and it has been shown that plants in general may be referred to one or other of these series, while red is common to both series, and green, as composed of blue and yellow, is intermediate between them. White is considered as depending on absence or extreme dilution of the colouring principles, while brown or black depends on their accumulation or concentration. Even in white flowers there will be seen a slight admixture of a yellowish or bluish tint.

Green.

Cyanic series.	{	Greenish-blue.		Yellowish-green.	}	Xanthic series.
		Blue.		Yellow.		
		Violet-blue.		Orange-Yellow.		
		Violet.		Orange.		
		Violet-red.		Orange-Red.		
		Red.		Red.		

680. Some starting from greenness, as a state of equilibrium between the two series, pass through the blue and violet to red, by a process of oxidation, while the transition from red to orange and yellow has been traced to deoxidation. As illustrations of the cyanic series may be

mentioned, all, or nearly all, the species of *Campanula*, *Phlox*, *Epilobium*, *Hyacinth*, *Geranium*, *Anagallis*; of the xanthic series, *Cactus*, *Aloe*, *Cytisus*, *Oxalis*, *Rose*, *Verbascum*, *Potentilla*, *Oenothera*, *Ranunculus*, *Adonis*, *Tulip*, *Dahlia*.

681. Plants belonging to either series, vary in colour usually by rising or falling in the series to which they belong, and not by passing from one to the other. Thus, a plant belonging to the blue series does not usually become yellow, nor does one in the yellow series change into a pure blue. This remark will not apply in all cases, although it is generally true. It cannot be said to hold good in regard to genera, as at present determined; thus, in the genus *Gentian*, there are blue and yellow species. It seems, however, to be applicable to individual species; thus, the *Dahlia* belonging to the yellow series has been made to pass to all varieties of that series, but has never been produced of a blue colour; so also with the *Tulip*, the *Rose*, &c. Even in the case of species, however, there are anomalies. Thus, the rule does not apply to such plants as *Myosotis versicolor* and *Dendrobium sanguinolentum*, where there are different yellow and blue colours on the corolla. Notwithstanding, however, all the exceptions, the general law already mentioned as to the variation of colour in flowers, seems to be founded on correct observations.

682. Changes are produced in the colour of flowers, by bruising and injuring the petals. The pure white flowers of *Camellia* easily become brown, while those of *Calanthe veratrifolia* and *Bletia Tankervilleæ* assume a deep blue. By drying, many flowers become of a brown or black colour: this is particularly the case with *Orchidaceæ*, *Melampyrum*, and *Orobis niger*. It would appear to depend on the combination between the colouring principle and the oxygen of the air, and may in some cases be traced to the existence of tannin, gallic acid, and iron. Blue flowers, under the process of desiccation, are often whitened. *Ipomœa Learii*, in drying, changes from blue to red.

683. Remarkable changes take place in the colour of some flowers during the course of the day. The flowers of the common pink *Phlox*, early in the morning, have a lightish blue colour, which alters as the sun advances, and becomes bright pink. The *Oenothera tetraflora* has white flowers which change to red. *Hibiscus variabilis* has its flowers white in the morning, pink at noon, and bright red at sunset. The colour of many flowers of *Boraginaceæ*, before expansion, are red; after expansion, blue. The bracts of *Hakea Victoria* are yellowish-white in the centre the first year; the second year, what was white becomes a rich golden yellow; the third year, the yellow becomes rich orange; the fourth year, the colour becomes blood-red; the green portion of the bracts becomes annually darker. It has been stated that soils have an effect on the colour of flowers. The flower of the common *Hydrangea hortensis* may be changed from pink and rose-coloured to

blue, by growing the plant in certain kinds of loam and peat earth. Alum in the soil is said to produce a similar effect.

684. Köhler and Schubler have endeavoured to determine the relative proportions between the different colours met with in flowers. They examined upwards of 4000 species, belonging to twenty-seven natural orders, of which twenty were dicotyledonous, and seven monocotyledonous. The following are some of their conclusions:—

1. White.....	1193	6. Green.....	153
2. Yellow.....	951	7. Orange.....	50
3. Red.....	923	8. Brown.....	18
4. Blue.....	594	9. Nearly Black.....	8
5. Violet.....	307		

685. The proportion of white, cyanic, and xanthic flowers varies in different quarters of the globe, and at different elevations. The following are the proportions of colour in different natural orders, deduced from the examination of about 120 species of each:—

	Red.	Violet.	Blue.	Green.	Yel.	Orange.	White.
Nymphæaceæ.....	11	—	14	—	28	—	46
Rosaceæ.....	32	1	—	—	52	—	40
Primulaceæ.....	41	7	6	2	15	10	27
Boraginaceæ.....	10	9	28	3	13	1	35
Convolvulaceæ.....	39	10	12	—	7	2	27
Ranunculaceæ.....	16	4	15	2	42	1	19
Papaveraceæ.....	38	9	—	—	36	7	7
Campanulaceæ.....	5	21	58	—	3	1	10

Thus, Nymphæaceæ and Rosaceæ, according to Schubler and Köhler's observations, contain a large number of white flowering species; Primulaceæ and Convolvulaceæ, red; Campanulaceæ, blue; Ranunculaceæ, yellow.

686. In arranging flowers in a garden, it is of importance to place the complementary colours together, in order to produce the best effect. The complementary colour of red, or that which is required to make white light, is green; of orange, blue; of yellow, violet; consequently blue and orange coloured flowers, yellow and violet, may be placed together; while red and rose-coloured flowers harmonize well with their own green leaves. When the colours do not agree, the interposition of white often restores harmony.

5.—ODOURS OF FLOWERS.

687. The peculiar odours of plants depend on various secreted volatile matters, which are often so subtle as to be incapable of detection by ordinary chemical means. Nothing is known of the causes which render one flower odoriferous and another scentless. In some cases the odours of plants remain after being dried, but in general they

disappear. Some leaves, as of the Woodruff, become scented only after drying; and certain woods, as Teneriffe rosewood, give out their odour only when heated by friction. Meteorological causes have a great influence on the odours of living plants. Dew, or gentle rain with intervals of sunshine, seem to be the circumstances best fitted for eliciting vegetable perfumes. Light has a powerful effect on the odour as well as the colour of flowers. Plants, when etiolated by being kept in darkness, generally lose their odour. In certain cases, the perfumes of flowers are developed in the evening. Some of these plants were called *tristes* by Linnæus, as *Hesperis tristis*, or night-scented stock. Many orchidaceous plants are fragrant at night only, as some *Catasetums* and *Cymbidiums*. *Cestrum nocturnum* and the white flowers of *Lychnis vespertina* are also night-scented. The odours of some plants are peculiarly offensive. This is the case with *Phallus impudicus*, and with the flowers of many *Stapelias*.

688. Schubler and Köhler, whose investigations in regard to colour have been noticed, have also made observations on the odours of plants in the same monocotyledonous and dicotyledonous orders. The following tables show some of their results:—

Colour.	No. of species.	Odoriferous.		Agreeable.		Disagreeable.	
White.....	1193	187	175	12
Yellow.....	951	75	61	14
Red.....	923	85	76	9
Blue.....	594	31	23	7
Violet.....	307	23	17	6
Green.....	153	12	10	2
Orange.....	50	3	1	2
Brown.....	18	1	0	1

Thus, of the plants examined, those having white flowers presented the larger proportion of odoriferous species. The orange and brown coloured flowers often gave a disagreeable odour. In examining numerous species from various natural orders, they found that out of 100 species of—

Nymphæacæ.....	22	were odoriferous.
Rosacæ.....	13	—
Primulacæ.....	12	—
Boraginacæ.....	6	—
Convolvulacæ.....	4	—
Ranunculacæ.....	4	—
Papaveracæ.....	2	—
Campanulacæ.....	1	—

6.—DISEASES OF PLANTS.

689. Great obscurity attends this department of botany, and much remains to be done ere a system of vegetable nosology (*νόσος*, disease)

can be completed. It is, however, of great importance, whether we regard its bearing on the productions of the garden or the field. Some have divided the diseases of plants into *general*, or those affecting the whole plant, and *local*, or those affecting a part only. A better arrangement seems to be founded on their apparent causes, and in this way they have been divided by Lankester into four groups. 1. Diseases produced by changes in the external conditions of life; as by redundancy or deficiency of the ingredients of the soil, of light, heat, air, and moisture. 2. Diseases produced by poisonous agents, as by injurious gases, or miasmata in the atmosphere, or poisonous matter in the soil. 3. Diseases arising from the growth of parasitic plants, as Fungi, Dodder, &c. 4. Diseases arising from mechanical injuries, as wounds and attacks of insects.

690 Plants are often rendered liable to the attacks of disease by the state of their growth. Thus, cultivated plants, especially such as become succulent by the increase of cellular tissue, appear to be more predisposed to certain diseases than others. Concerning the first two causes of disease very little is known. Absence of light causes *blanching*, which may be looked upon as a diseased state of the tissues. Excess of light may cause disease in plants whose natural habitat is shady places. Excess of heat is sometimes the occasion of a barren or diseased state of some of the organs of the flowers, and frost acts prejudicially on the leaves, stem, and flowers. By excess of moisture, a dropsical state of the tissue is induced.

691. Concerning the influence of atmospheric changes on plants, very little has been determined. Many extensive epidemics seem to depend on this cause. Thus, the late potato disease must be traced, apparently, to some unknown miasma conveyed by the air, and operating over large tracts of country; the disease probably affecting some plants more than others, according to their state of predisposition, and in its progress leading to disorganization of the textures, alteration in the contents of the cells and vessels, and the production of Fungi, &c. In the early stage of the disease, as witnessed in 1845, Harting observed that a brown granular matter was deposited in the interior of the cells, beginning with those near the surface. For some time the cell-walls and starch-grains remained uninjured, but were ultimately attacked, the former losing their transparency, and the latter becoming agglomerated in masses. Subsequently to this, parasitic organisms of various kinds made their appearance, cavities were formed, and rapid decay took place. Among the vegetable parasites, were detected species of *Fusisporium*, *Oidium*, *Botrytis*, *Capillaria*, *Polyactis*, &c. Solly objects to the Fungus theory of Potato disease, and believes the development of parasites a secondary step in the morbid process. The prevalence of hot or cold weather, the amount of light and moisture, changes in the atmosphere, and electrical conditions of

the air and earth, are in all probability connected with epidemic diseases. Some, with Liebig, attribute the late potato disease to suppressed evaporation and transpiration, depending on the hygrometric state of the atmosphere. The vessels and cells are said to become charged with fluids, stagnation of the circulation takes place, and thus disease and death are induced.* The general conclusions to be drawn from the various researches that have been instituted and opinions which have been advanced respecting the cause and progress of this epidemic may be thus stated. Coincident with, if not consequent on certain meteorological and epidemic influences, alterations are accomplished in the cellulose and contents of the cells. Decay follows on those changes. In the decayed and decaying matter, parasitic Fungi find a nidus for growth and progress equally with the disease of which they are primarily a consequence, and to the extension of which they in their turn contribute.

692. *Gangrene* in plants is caused by alterations in the contents of the cells, leading to death of a part. In succulent plants, as Cactuses, this disease is apt to occur. It is capable of extension by contact of the diseased cells. Sometimes excision of the diseased part checks the progress of the gangrene. *Canker*, which attacks Apple and Pear trees, is a kind of gangrene. Some of the most important diseases of corn and other agricultural crops, are owing to the production of Fungi. These have been divided into 1. Those attacking the grain, as *Uredo fœtida*, or pepper-brand. 2. Those attacking the flower, as *Uredo segetum*, or smut. 3. Those attacking the leaves and chaff, as *Uredo Rubigo*, or rust. 4. Those attacking the straw, as *Puccinia graminis*, or corn mildew.

693. *Smut-balls*, *pepper-brand*, or *blight*, is a powdery matter, occupying the interior of the grain of wheat, &c. When examined under the microscope, it consists of minute balls, four millions of which may exist in a single grain, and each of these contains numerous excessively minute sporules. It is caused by the attack of *Uredo Caries*, or *fœtida*. In this disease the seed retains its form and appearance, and the parasitic fungus has a peculiarly fetid odour, hence called *stinking rust*.

694. *Smut* or *dust-brand* is a sooty powder, having no odour, found in Oats and Barley, and produced by *Uredo segetum*. The disease shows itself conspicuously before the ripening of the crop. Bauer says that in $\frac{1}{10000}$ part of a square inch he counted 49 spores of the *uredo*.

695. *Rust* is an orange powder, exuding from the inner chaff scales, and forming yellow or brown spots and blotches in various parts of corn plants. It owes its presence to the attack of *Uredo Rubigo*. It

* See remarks on this subject by Klotzsch, translated by Gregory, in the Appendix to Liebig's work on the Motion of the Juices. London, 1848.

is sometimes called *red gum*, *red robin*, *red rust*, and *red rag*. Some consider Mildew (*Uredo linearis*) as another state of the same disease.

696. Those Fungi which are developed in the interior of plants, and appear afterwards on the surface, are called *entophytic* (ἐντός, within, and φυτόν, a plant). Their minute sporules are either directly applied to the plants entering by their stomata, or they are taken up from the soil. Many other Fungi grow parasitically on plants, and either give rise to disease, or modify it in a peculiar way. Among them may be mentioned species of *Botrytis*, *Fusisporium*, *Depazia*, *Sclerotium*, *Fusarium*, and *Erysiphe*. *Fusisporium solani* is considered by Martius as the cause of a certain disease in the Potato. In the recent potato disease, the *Botrytis infestans*, a species of *Fusarium*, and other Fungi, committed great ravages, spreading their mycelium or spawn through the cells of the leaves and the tuber, and thus accelerating their destruction. Berkeley, Morren, and Townley, consider the *Botrytis* as the cause of the disease. Various species of *Botrytis* also attack the Tomato, Beet, Turnip, and Carrot. A species of *Depazia* sometimes causes disease in the knots of Wheat. A diseased state of Rye and other grasses, called *ergot*, owes its production to the presence of a species of *Spermœdia*. By the action of the fungus, the ovary becomes diseased and altered in its appearance, so as to be dark-coloured, and project from the chaff in the form of a spur. Hence the name *spurred rye*. The nutritious part of the grain is destroyed, and it acquires certain qualities of an injurious nature. Spontaneous gangrene is the consequence of living for some time on diseased rye. Ergot has been seen in *Lolium perenne* and *arvense*, *Festuca pratensis*, *Phleum pratense*, *Dactylis glomerata*, *Anthoxanthum odoratum*, *Phalaris arundinacea*, &c.

697. Fruits when over-ripe are liable to attacks of Fungi, which cause rapid decay; Wood also, especially Alburnum or sap-wood, is injured by the production of Fungi. Dry rot is the result of the attack of *Merulius lacrymans*, which in the progress of growth destroys its texture, and makes it crumble to pieces. Some kinds of wood are much more liable to decay than others.

698. The diseases caused by attacks of Fungi may be propagated by direct contact, or by the diffusion of the minute spores through the atmosphere. When we reflect on the smallness of the spores, the millions produced by a single plant, and the facility with which they are wafted by the wind in the form of the most impalpable powder, we can easily understand that they may be universally diffused and ready to be developed in any place where a nidus is afforded. Perhaps some of the diseases affecting man and animals may be traced to such a source. Quekett found that he could propagate the ergot by mixing the sporules with water, and applying this to the roots.

699. In order to prevent these diseases, it has been proposed to steep the grains in various solutions previously to being sown. For this purpose, alkaline matters and sulphate of copper have been used. In all cases, the seed should be thoroughly cleansed. Smut and pepper-brand have been averted by these means. In the case of the latter, diseased grains are easily removed by being allowed to float in water, and the grains that remain are washed with a solution of lime, common potash, or substances containing ammonia, which form a soapy matter with the oil in the fungus. A weak solution of sulphate of copper acts by destroying the fungus. To prevent wood from dry rot, the processes of kyanizing and burnetizing have been adopted: the former consists in making a solution of corrosive sublimate enter into the cells and vessels; the latter, in impregnating the wood with a solution of chloride of zinc. Creosote has also been used to preserve wood. Boucherie proposed that a solution of pyrolignite of iron should be introduced into trees before being felled, by making perforations at the base of the trunk, and allowing the absorbing power of the cells and vessels to operate. This plan does not appear to have been successful, although reported favourably to the French Academy, and also recommended by Mr. Hyett.

700. Other diseases in plants owe their origin to *insects*. *Earcokkles*, *purples*, or *pepper-corn*, is a disease affecting especially the grains of wheat. The infected grains become first of a dark green, and ultimately of a black colour. They become rounded like a small pepper-corn, but with one or more deep furrows on their surface. The glumes spread open, and the awns become twisted. The blighted grains are full of a moist white cottony matter, which, when moistened and put under the microscope, is seen to consist of a multitude of minute individuals of the *Vibrio tritici*, or eel of the wheat. The animalcules deposit their eggs in the ovary, and their young are hatched in eight or ten days. Henslow calculates that 50,000 of the young might be packed in a moderately sized grain of wheat. The *Vibrio* retains its vitality long. It will remain in a dry state for six or seven years, and when moistened with water will revive. The wheat-fly, or *Cecidomyia tritici*, is another destructive insect. It deposits its eggs by means of a very long retractile ovipositor, and is seen abundantly in warm evenings. The *Cecidomyia* destructor, or Hessian fly, also causes injury, and is said to be very destructive to wheat in America. These insects are destroyed in numbers by the *Ichneumon*s, which deposit their ova in their bodies. The Apple-tree mussel, or dry-scale, *Aspidotus conchiformis*, attacks the bark of Apples, Pears, Plums, Apricots, and Peaches. Many of the *Coccus* tribe are highly injurious to plants. One of this tribe, in 1843, destroyed the whole orange trees in the island of Fayal, one of the Azores. Many insects cause the rolling up of leaves. *Tortricida viridana* acts thus on the leaves of the Oak, and

various species of *Losotænia* do so with other trees. *Sacchiphantes abietis* is the aphid which causes the leaves of the Spruce-fir to be united together, so as to have the appearance of a cone.

701. Many insects, called miners, make their way into the interior of leaves, and hollow out tortuous galleries, sometimes causing an alteration in the colour of the leaves. Galls are caused by the attacks of species of *Cynips*, which are provided with ovipositors, by means of which they pierce the bark or leaves with the view of having a nidus for their ova. These galls are very common on the Oak, and are called *oak-apples*. Sometimes they have one cavity, at other times they are divided into numerous chambers, each containing a grub, pupo, or perfect fly, according to the season. Galls are produced on the twigs, catkins, and leaves of the Oak. The artichoke gall of the Oak depends on an irregular development of a bud, caused by the attack of insects, and consists of a number of leafy imbricated scales resembling a young cone. On examining the galls of commerce, the produce of the *Quercus infectoria*, some are of a blue colour, containing the larva of the insect; others are pale, and are marked with a perforation by which the insect has escaped. Extensive ravages are committed in Elms and other trees by the attacks of *Scolyti*. The presence of much moisture, such as the rapid flow of sap, destroys them. Mr. Robert found that the flow might be promoted by taking off the suberous layer of the bark, and he proposes this as a method of getting rid of the insects. Some galls are formed in the substance of leaves, and burst through the cuticle in the form of ovate bodies, with crenate borders and opercula, which are perforated in the centre. These galls resemble parasitic fungi. *Oak-spangles* are galls of this nature. They are attached by a central point to the outer surface of the leaf, the inner side being smooth—the outer red, hairy, and fringed. Each contains a single insect, which retains its habitation till March, long after the leaves have fallen to the ground.

702. It is impossible in this place to enumerate all the insects which attack plants. Almost every species has certain insects peculiar to it, which feed on its leaves, juices, &c., and often cause great injury. Those which are common to hot-houses and green-houses, have called for the special attention of horticulturists, and various means have been suggested for their removal or prevention. Among them may be enumerated, vapour of tobacco and ammoniacal liquor of gas-works, to kill aphides; vapour of sulphur, for the red spider; vapour of turpentine, for the wasp; vapour of crushed laurel leaves, for the white bug; coal tar, for the wire-worm, &c.*

* For further remarks on the Diseases of Plants, see Henslow's paper in the Journal of the Royal Agricultural Society of England.

PART II.

SYSTEMATIC BOTANY, TAXONOMY, OR THE CLASSIFICATION OF PLANTS.

CHAPTER I.

SYSTEMS OF CLASSIFICATION.

703. THIS department of Botany may be considered as a combination of all the observations made on the structure and physiology of plants, with the view of forming a scientific arrangement. It can only, therefore, be prosecuted successfully after the student has acquired a complete knowledge of Organography. In every branch of science, arrangement is necessary in order that the facts may be rendered available, and this is more especially the case when a knowledge of species is to be acquired. When it is considered that there are upwards of 120,000 known species of plants, it is obvious that there must be a definite nomenclature and classification, were it only to facilitate reference and communication. Taxonomy has sometimes been pursued with no higher aim than that of knowing the names of plants. When prosecuted in such a spirit, it does not lead to an enlarged and philosophical view of the vegetable kingdom. In all truly scientific systems, regard is paid, not merely to the determination of the names of the species, but to their relations and affinities, so as to give some conception of the order which has been followed in the plan of creation.

704. In Classifactory Sciences, the arrangements are founded upon an *idea of likeness*—an idea, however, which is applied in a more exact and rigorous manner than in its common and popular employment. The resemblances of the objects must rest, not on vague generalities, but upon an accurate scientific basis. In order that an arrangement may be constructed on philosophical principles, and that it may be rendered useful for the purpose of science, the following steps are required:—1. A Technical (τεχνικὸς, artificial or conventional) language,

rigorously defined, or what is termed Glossology (γλῶσσα, a tongue or language, and λόγος, a discourse), and Termonology (τέρμων, -ονος, a termination). The meaning of the terms in this descriptive language must not depend on fancied resemblances, but must have a precise definition, and be constant. In acquiring a knowledge of the conventional terms, or of the vocabulary of the science, the student at the same time fixes in his mind the perceptions and notions which these terms convey, and thus, in reality, becomes acquainted with important elementary facts. 2. A Plan of the system, or the principles on which the divisions and subdivisions of the system are made, Diataxis (διάταξις, orderly arrangement), or what is properly called Taxonomy (τάξις, order, and νόμος, law). There have been two great plans proposed in Botany, one denominated *artificial*, the other *natural*. The first is founded on characters taken from certain parts of plants only, without reference to others; while the second takes into account all the parts of plants, and involves the idea of *affinity* in essential organs. 3. There must be also the means of detecting the position of a plant in a system by short diagnostic marks. In doing so, a few essential characters are selected in accordance with natural affinities. The division into genera is a most valuable help in determining plants. Linnæus did great service to science by his generic divisions, and by adopting a *binominal* (*bis*, twice, and *nomen*, a name) system of nomenclature, in which the genus and species are included in the name of the plant.

705. **Species.**—No classification can be made unless the meaning of the term *species* is defined. By species, then, are meant so many individuals as are presumed to have been formed at the creation of the world, and to have been perpetuated ever since. A species embraces individuals which resemble each other more closely than they do any other plant, so that they are considered as originating from a common parent; and their seeds produce similar individuals. There may be differences in size, colour, and other unimportant respects; and thus *varieties* may exist, exhibiting minor differences, which are not, however, incompatible with a common origin. Varieties owe their origin to soil, exposure, and other causes, and have a constant tendency to return to their original type. They are rarely propagated by seed, but can be perpetuated by cuttings and grafts. By cultivation, permanent varieties of *races* have been produced, the seeds of which give rise to individuals varying much from the original *specific* type. Such races are kept up entirely by the art of the gardener, and may be illustrated in the case of the Cereal grains, and of culinary vegetables, such as Cabbages, Cauliflower, Brocoli, Turnips, Radishes, Peas. It is only after a series of years that these permanent varieties have been established, and there is still a tendency in their seeds, when sown in poor soil and neglected, to produce the original wild form. Permanent varieties in the animal kingdom may be illustrated by the

different races of mankind. By scattering the pollen of one plant on the pistil of an allied species, seeds are formed, which, when sown, produce intermediate forms or hybrids (§ 516). Hybrids, however, are rarely perpetuated by seed.

706. Many species vary in a remarkable manner, without any external influences to account for it. Thus, a plant of *Fuchsia* has produced, in successive years, flowers differing so much in form and shape, that, if they had not been known to be produced by the same plant, they would have been considered as belonging to distinct species. Such is also the case with *Calceolarias*, some species of *Amaryllis*, and many *Orchids*. Hence there is sometimes considerable difficulty in determining what are true species, and what are only varieties, more especially when these varieties are permanent and reproduce themselves. To this must in part be attributed the disputes which have arisen among botanists as to the species of many British genera, such as *Roses*, *Rubi*, *Salices*, and *Hieracia*. The following table shows the number of British species in some of the British genera, as given by different authors, and exhibits the uncertainty which still exists as to the limits of species:—

	<i>Salix.</i>	<i>Hieracium.</i>	<i>Mentha.</i>	<i>Rosa.</i>	<i>Rubus.</i>	<i>Saxifraga.</i>
Hudson (1798).....	16	8	10	5	5	9
Smith (1824-28).....	64	18	13	22	14	25
Lindley (1835).....	29	19	9	17	21	24
Hooker (1842).....	70	13	13	19	14	16
Babington (1843).....	57	19	8	19	24	20
Watson (1844).....	38	17	8	7	34	16
Babington (1847).....	58	21	9	19	36	20

707. It is only after a careful study of such forms during a series of years, that any conclusion can be drawn in regard to them. It is important to record all the varieties which occur, but great care is necessary not to raise to the rank of species what are mere accidental aberrant forms. Some have of late years advocated the doctrine of the transmutation of species or the conversion of one species into another. It has been said, that Oats may be changed into Rye, by being constantly cut down for a series of years before flowering. Such statements are not founded on correct data, and have led to very erroneous views and doctrines, which have been recently promulgated with much apparent plausibility. All that has been observed in the vegetable kingdom leads to the conclusion, that there are distinct species which continue to be perpetuated by seed, and that, although these may vary within certain limits, there is always a typical form to which the varieties have a tendency to revert. By grafting and other horticultural operations, changes of a marked kind may be produced in fruits; but the seeds of such fruit, when sown, give rise to individuals resembling

the original stock—they perpetuate the *typical* form, not the artificially-produced variety.

708. **Genera.**—Certain species are more nearly allied than others, and are conveniently grouped together so as to form a distinct *kind* or *genus*. A genus, then, is an assemblage of nearly related species, agreeing with one another in general structure and appearance more closely than they accord with any other species. Thus, the various species of *Roses* compose one genus, which is distinguished by marked characters. Occasionally a *subgenus* is formed by grouping certain species, which agree more nearly with each other in some important particulars than the other species of the genus. The characters of the genera are taken exclusively from the parts of fructification, while all parts of the plant furnish specific characters. In the name of a plant, the genus is given as well as that of the species. The latter was called by Linnæus the *trivial* name. Thus, a particular species of Rose is called *Rosa spinosissima*; the first being the genus, and the second the specific or trivial name. As regards the definition of genera and species, and nomenclature in general, no one has conferred so much benefit on science as the great Linnæus. This may be considered as among his highest titles to fame.

709. **Orders.**—Several genera agreeing in more general characters, although differing in their special conformation, are grouped together so as to form an *order* or *family*. As genera include allied species, so orders embrace allied genera. Subdivisions are also made to facilitate reference, so that *suborders* and *tribes* are formed, consisting of certain genera, more nearly related in particular characters than others. Thus, the order *Rosacæ*, or the Rose family, includes the genera *Rosa*, *Rubus*, *Potentilla*, *Fragaria*, *Prunus*, &c., which all agree in certain general characters; and the order is divided into various suborders, such as the true *Roses*, the *Amygdalæ*, comprehending the Plum, Almond, Peach, &c.; the *Potentillæ*, embracing the Cinquefoil, Strawberry, Raspberry, &c. (§ 854).

710. **Classes.**—Orders having some general characters in common, are united together in *classes*, and *subclasses* are formed in the same way as *suborders*. This is the general plan upon which botanical classification proceeds. With the exception of the individual species, all the divisions are more or less arbitrary. In making them, however, the object of the enlightened botanist is to follow what he considers to be the natural affinities, and thus to trace, as far as possible, the order which pervades the vegetable creation.

711. **Essential Characters.**—Each of the divisions of a system is accurately defined, the characters being as short as is consistent with precise diagnosis. Such characters are called *essential*, and they embrace only those points by which the group is distinguished from the others in the same section. The complete description of an individual

species, from the root to the flower and fruit, is called the *natural* character, and embraces many particulars which are not requisite for the purpose of diagnosis. The essential characters of genera, when in Latin, are put in the nominative case, while those of species are in the ablative.

712. Nomenclature.—The names of genera are variously derived, from the structure or qualities of the group, from the name of some eminent botanist, &c. ; while specific names have reference also to the country where the plant is found, the locality in which it grows, the form of the leaves, root, stem, or the colour of the flowers, &c. When a species is named in honour of its discoverer or describer, his name is put in the genitive, as *Carex Vahlîi*, or the *Carex* detected by Vahl ; but if it is merely in compliment to a botanist his name is added in an adjective form, as *Jungermannia Doniana*, or a *Jungermannia* named in honour of Don, as a botanist. Sometimes two nouns are united in a specific name, as *Dictamnus Fraxinella*. In such cases, the specific name is often an old generic one, has a capital letter prefixed, and does not necessarily agree in gender with the name of the genus. The name of the orders in what is called the natural system, are derived from one of the typical genera included under them.

713. Abbreviations and Symbols.—It is of great importance that correct descriptions should be given of species, for without them it is impossible to form the groups accurately. The difficulties of the taxonomist are often greatly increased by imperfect and careless descriptions. Valuable directions are given in Lindley's Introduction to Botany, as to the proper method of describing plants. There are certain abbreviations in constant use among botanists, which it may be of importance to notice here. The authorities for genera and species are given by adding the abbreviated name of the botanist who described them. Thus *Veronica L.*, is the genus *Veronica* as defined by Linnæus ; *Veronica arvensis L.*, is a certain species of *Veronica*, defined by the same author ; *Oxytropis DC.*, is the genus as defined by De Candolle. It is usual in descriptive works to give a list of the authors, and the symbols for their names. The abbreviation *v. s. sp.*, means *vidi siccam spontaneam*, or that the author has seen a dried wild specimen of the plant ; *v. s. c.*, means *vidi siccam cultam*, or that he has seen a dried cultivated specimen ; *v. v. s.*, means *vidi vivam spontaneam*, or that he has seen a living wild specimen ; while *v. v. c.*, means *vidi vivam cultam*, or that the author has seen a living cultivated specimen. The asterisk prefixed to a name (**L.*), indicates that there is a good description at the reference given to the work ; while the dagger (†*L.*), implies some doubt or uncertainty. The point of admiration (!*DC.*), marks that an authentic specimen has been seen, from the author named ; and the point of interrogation (?) indicates doubts as to the correctness of genus, species, &c., according as it is placed

after the name of the one or other. $\odot, \circ, \textcircled{1}$, or Δ , annual; $\textcircled{2}$, or $\textcircled{3}$, or B, biennial; H , Δ , or P, perennial; h , $\bar{\text{S}}$, or Sh, shrub; (, twining to the right;), twining to the left; H , hermaphrodite; ♂ , male; ♀ , female; $\text{♂} - \text{♀}$, monœcious, or the male and female on one plant; $\text{♂} : \text{♀}$, diœcious, or the male and female on different plants; 00 or ∞ , means indefinite in number. After the description of a plant, its *habitat*, or the country and locality in which it grows, is given. If the plant has been described by others, reference is given to the work in which the description may be found. If it has received different names, the *synonymes* must be carefully detailed, and ought to be arranged in chronological order.

714. **Systems.**—Various attempts have been made at different times to classify plants. One of the earliest methodical arrangements was that of Cæsalpinus, in 1583. It was entirely artificial, and the same thing may be affirmed of those of Gesner, Morison, Rivinus, and Tournefort. The system propounded by Tournefort was for a long time adopted by the French school, but was ultimately displaced by that of Linnæus, who must be looked upon as the great promulgator of the artificial method. In 1682, Ray published a system which laid the foundation of the natural method of classification. It was long neglected, and did not receive the attention it deserved, until Jussieu entered the field, and developed his views. Since that time, the natural method has been advanced by the labours of De Candolle, Brown, Endlicher, Lindley, and many others.

715. **Linnæan System.**—Although the Linnæan system is not in conformity with natural affinities, and does not tend to comprehensive views of structure, still it is useful to the student as an *index*. Linnæus himself did not consider it as occupying a higher position, and he stated distinctly that a natural method was the great object of scientific inquiry. When not elevated to a rank which its author never meant it to occupy, this system may, with all its imperfections, be employed as a useful artificial key, and as such may be combined with the natural system. In many works of the present day, as in Babington's *Manual of British Botany*, the Linnæan system is used as an index to the genera. In the Linnæan or sexual system, twenty-three classes are founded on the number, position, relative lengths, and connection of the stamens; while the orders in these classes depend on the number of the styles, the nature of the fruit, the number of stamens in the classes where this character is not used for distinguishing them, and the perfection of the flowers. The twenty-fourth class includes plants having inconspicuous flowers, and in it the orders are formed according to natural affinities. Under these classes and orders, all the known genera and species were arranged. It is in the higher divisions that the system is artificial, for, as regards genera, the Linnæan rules are followed even in the natural systems of the present day.

716. TABULAR VIEW OF THE CLASSES OF THE LINNÆAN SYSTEM.

A. Flowers present (Phanerogamia).

I. Stamens and Pistil in every flower.

1. Stamens Free.

a. Stamens of equal length, or not differing in certain proportions; ἀνῆς, male or stamen.

in number 1.....	Class I. Monandria.....	μείνος, one.
— 2.....	II. Diandria.....	δύο, two.
— 3.....	III. Triandria.....	τρῆς, three.
— 4.....	IV. Tetrandria.....	τετράς, four.
— 5.....	V. Pentandria.....	πέντε, five.
— 6.....	VI. Hexandria.....	ἕξ, six.
— 7.....	VII. Heptandria.....	ἑπτὰ, seven.
— 8.....	VIII. Octandria.....	ὀκτώ, eight.
— 9.....	IX. Enneandria.....	ἐννέα, nine.
— 10.....	X. Decandria.....	δέκα, ten.
— 12-19.....	XI. Dodecandria.....	δώδεκα, twelve.
— 20 } Inserted on Calyx	XII. Icosandria.....	ἰκσοσι, twenty
or more } — on Receptacle	XIII. Polyandria.....	πολὺς, many.

b. Stamens of different lengths;

two long and two short..... XIV. Didynamia...
four long and two short.. ... XV. Tetrodynamia } δύναμις, power.

2. Stamens united;

by Filaments in one bundle..... XVI. Monadelphia..
— in two bundles..... XVII. Diadelphia...
— in more than two bundles. XVIII. Polyadelphia.. } ἀδελφός, brother.

by Anthers (Compound flowers) XIX. Syngenesia... { σὺν, together.
γένεσις, origin.

with Pistil on a Column..... XX. Gynandria..... γυνή, female.

II. Stamens and Pistil in different

flowers; on the same Plant..... XXI. Monœcia.....
on different Plants..... XXII. Diœcia..... } οἶκος, house.

III. Stamens and Pistil in the same

or in different flowers on the } XXIII. Polygamia..... γάμος, marriage.
same or on different plants.... }

B. Flowers Absent..... XXIV. Cryptogamia..... κρυπτός, concealed.

717. TABULAR VIEW OF THE ORDERS OF THE LINNÆAN SYSTEM.

γυνή, female or pistil.

Class I. } Monogynia.....	1	Free Style	μείνος, one.
II. } Digynia.....	2	Free Styles	δύο, two.
III. } Trigynia.....	3	—	τρῆς, three.
IV. } Tetragynia.....	4	—	τετράς, four.
V. } Pentagynia.....	5	—	πέντε, five.
VI. } Hexagynia.....	6	—	ἕξ, six.
VII. } Heptagynia.....	7	—	ἑπτὰ, seven.
VIII. } Octogynia.....	8	—	ὀκτώ, eight.
IX. } Enneagynia.....	9	—	ἐννέα, nine.
X. } Decagynia.....	10	—	δέκα, ten.
XI. } Dodecagynia.....	12-19	—	δώδεκα, twelve.
XII. } Polygynia.....	20 and upwards	πολὺς, many.
XIII. }				

XIV.	{	Gymnospermia.....	Fruit formed by four Achænia	γυμνός, naked.
		Angiospermia	Fruit, a two-celled Capsule... with many seeds.....	ἄγγος, a vessel. σπίγμα, a seed.
XV.	{	Siliculosa.....	Fruit, a Silicula.	
		Siliquosa	Fruit, a Siliqua.	
XVI.				
XVII.		Triandria, Decandria, &c. (number of Stamens), as in the Classes.		
XVIII.	{	Polygamia Æqualis.....	Florets all hermaphrodite.	
		———— Superflua, ...	Florets of the disk hermaphrodite, those of the ray pistilliferous and fertile.	
		———— Frustranea...	Florets of the disk hermaphrodite, those of the ray neuter.	
		———— Necessaria...	Florets of the disk staminiferous, those of the ray pistilliferous.	
		———— Segregata....	Each floret having a separate involucre.	
		Monogamia	Anthers united, flowers not compound.	
XX.				
XXI.		Monandria, Diandria, &c. (number of Stamens), as in the Classes.		
XXII.	{	Monœcia	Hermaphrodite, staminiferous, and pistilliferous flowers on the same plant.	
		Diœcia	on two plants.	
		Triœcia.....	on three plants.	
XXIII.	{	Filices	Ferns.	
		Musci.....	Mosses.	
		Hepaticæ.....	Liverworts.	
		Lichenes.....	Lichens.	
		Algæ.....	Sea-weeds.	
		Fungi.....	Mushrooms.	
XXIV.				

718. Even as an artificial method, this system has many imperfections. If plants are not in full flower, with all the stamens and styles perfect, it is impossible to determine their class and order. In many instances, the different flowers on the same plant vary as regards the number of the stamens. Again, if carried out rigidly, it would separate in many instances the species of the same genus; but as Linnæus did not wish to break up his genera, which were founded on natural affinities, he adopted an artifice by which he kept all the species of a genus together. Thus, if in a genus nearly all the species had both stamens and pistils in every flower, while one or two were monœcious or diœcious, he put the name of the latter in italics, in the classes and orders to which they belonged according to his method, and referred the student to the proper genus for the description,

719. **Natural System.**—It has been already stated, that a natural system endeavours to bring together plants which are allied in all essential points of structure. It purposes to ascertain the system of nature, and the affinities of plants; and, in doing so, it takes into account all their organs. Every natural method, however, is, to a certain extent, artificial, and is likely to be so. It is impossible to show the affinities of plants in a *lineal* series; many orders pass insen-

sibly into others, so that their limits cannot be accurately defined; and no perfect system can be constituted until all the plants of the globe are known. Moreover, many artificial means are avowedly used in all natural systems to aid the student.

720. The early natural systems were very imperfect, being founded on comparatively vague views of structure and affinity. Such were the systems of Magnolius and Adanson. The sketch of a natural system by Linnæus was very incomplete, and even that of the celebrated Ray was imperfect. It was not until the knowledge of structural botany had advanced, that the affinities of plants were ascertained, and the relative importance of the different characters discovered. The natural systems of the present day recognize a certain subordination of characters, founded on the fact that some organs are of more importance to the life of plants than others. The relative values of these characters are determined by the study of organization, and are not fixed by arbitrary rules. The following table will illustrate this subordination of character:—

721. SUBORDINATION IN VALUE OF THE ORGANS OF THE SAME CLASS.

Relative Values.	Elementary.	Nutritive.	Reproductive.
1.	Cellular Tissue.		
	{ Vascular Tissue.....	{ Embryo.	—
	a. Spiral Vessels..	a. Cotyledon.	—
2.	b. Porous Vessels.	b. Radicle.	—
	c. Scleriform	c. Plumule.	—
	Vessels.....	Spore.	—
	{ Stomata.....	Prothallus.....	
3.	—	{ Root, Stem, Leaf, Frond, Thallus.	{ 1. Stamens and Pistil. 2. Antheridia and Arche- gonia. 3. Fruit. Pericarp. Theca.
4.	—	—	{ Perianth. a. Corolla. b. Calyx.
5.	—	—	{ Inflorescence. Torus, Nectary. Bract, Involucre.

722. Thus, cellular tissue occupies the highest place, as being most universally diffused, and capable of carrying on all the functions; next comes vascular tissue. By the consideration of these, the two great divisions of cellular and vascular plants are determined. There is nothing in the nutritive and reproductive systems of the same value as cellular tissue. The embryo and its parts are reckoned as occupying the highest place in the nutritive system, and as corresponding in value with the vascular among the elementary tissues. In the same way the other values are determined. In examining organs, it is essential

to compare those which belong to the same series; for an organ which occupies the highest place in one series, may be inferior in value to a second-rate organ in another. The comparative importance of the different series must be taken into account also. Thus, the nutritive may be considered as of more importance than the reproductive function, as being more essential for the life of the individual; and an organ of first-rate value in the one, will therefore assume a higher function than one of the same value in the other. The changes which take place in any one set of organs are often accompanied with changes in others; and thus it is found that natural divisions may be arrived at by different routes—for instance, by the elementary, nutritive, and reproductive functions. This gives the true notion of affinity; and classifications formed on such principles will obviously be more valuable, in a practical and physiological point of view, than those which adopt characters in an arbitrary manner.

723. Primary Divisions of the Vegetable Kingdom.—In taking a survey of the Vegetable Kingdom, some plants are found to be composed of cells only, and are called *Cellular* (§ 8); while others consist of cells and vessels, especially spiral vessels, and are denominated *Vascular* (§ 28). If the embryo is examined, it is found in some cases to have cotyledons or seed-lobes, in other cases to want them; and thus some plants are *cotyledonous*, others *acotyledonous* (§ 590); the former being divisible into *monocotyledonous*, having one cotyledon; and *dicotyledonous*, having two cotyledons. The radicle, or young root of acotyledons, is *heterorhizal* (§ 629), that of monocotyledons is *endorhizal* (§ 628), that of dicotyledons, *exorhizal* (§ 629). When the stems are taken into consideration, it is seen that marked differences occur here also, acotyledons being *acrogenous*, monocotyledons *endogenous*, and dicotyledons *exogenous* (§ 107). The venation of the leaves, whether parallel or reticulated (§ 143), establishes the same great natural divisions; and similar results are obtained from a consideration of the flowers, monocotyledons and dicotyledons being *phanerogamous*, and acotyledons *cryptogamous* (§ 323).

724. Thus, the following grand natural divisions are arrived at:—

- | | | | | | |
|----------------|---------------------|---------------|-------------|------------------|---|
| 1. Cellular... | Acotyledonous. | Heterorhizal. | Acrogenous. | Cryptogamous. | ~ |
| 2. Vascular.. | { Monocotyledonous. | Endorhizal. | Endogenous. | { Phanerogamous. | |
| | { Dicotyledonous. | Exorhizal. | Exogenous. | | |

These larger groups are, on similar principles, subdivided, until at length genera and species are reached by a process of analysis. Similar results will be obtained by a synthetical process, conducted on the same principles, and proceeding from species upwards.

725. Henslow illustrates the divisions and subdivisions of a natural system by reference to *Anthyllis Vulneraria*, thus:—

B. CELLULARES OR ACOTYLEDONEÆ.

Class III. Acotyledones.

- Subclass 1. Foliosæ.....Having leaves.
 ——— 2. Aphillæ.....Leafless.

729. By some recent authors, this system has been modified, so as to include, under Corollifloræ, all Dicotyledons with united petals, whether hypogynous or not, and to exclude from Class II. all plants without flowers. It is then presented in the following form:—

MODIFICATION OF DECANDOLLE'S SYSTEM.

Class I. Dicotyledones or Exogenæ.

- | | | | |
|--|---|--------------------------------|--------------------------------------|
| Dichlamydeæ,
having calyx
and corolla.
Having a single
perianth. | { | Subclass 1. Thalamifloræ,..... | Petals distinct, stamens hypogynous. |
| | | ———— 2. Calycifloræ,..... | Petals distinct, stamens perigynous. |
| | | ———— 3. Corollifloræ,..... | Petals united, bearing the stamens. |
| | | ———— 4. Monochlamydeæ,..... | A calyx only, or none. |

Class II. Monocotyledones or Endogenæ.

- Subclass 1. Petaloideæ or Floridæ,.....Floral envelopes verticillate.
 ——— 2. Glumaceæ,.....Floral envelopes imbricated.

Class III. Acotyledones or Acrogenæ.

- Subclass 1. Ætheogamæ,.....Having vascular tissue.
 ——— 2. Amphigamæ or Cellulares,.....Entirely cellular.

730. SYSTEM OF ENDLICHER.

REGION I.—THALLOPHYTA (θαλλίτης, frond, φυτόν, a plant). No opposition of stem and root. No spiral vessels, and no sexual organs. Propagated by spores.

SECTION I. PROTOPHYTA (πρόφυτος, first or originating). Developed without soil; deriving nourishment all around; fructification indefinite.

SECTION II. HYSTEROPHYTA (ὕστερος, posterior or derivative). Developed on decaying organisms; nourished internally from a matrix; all the organs appearing at once, and perishing in a definite manner.

REGION II.—CORMOPHYTA (κορμός, a stalk or trunk). Opposition of stem and root. Spiral vessels and sexual organs distinct in the more perfect.

SECTION III. ACROBRYA (ἄκρον, summit, and βεῖον, to germinate). Stem increasing by the apex, the lower part being unchanged, and only conveying fluids.

Cohort 1. Anophyta (ἄνω, above). No spiral vessels. Both sexes present. Spores free within spore-cases.

Cohort 2. Protophyta. Bundles of vessels more or less perfect. No male organs. Spores free within one- or many-celled spore-cases.

Cohort 3. Hysterophyta. Both sexes perfect. Seeds without an embryo, consisting of many spores. Parasitic.

SECTION IV. AMPHIBRYA (ἄμφω, around). Stem increasing at the circumference. Vegetation peripheral.

SECTION V. ACRAMPHIBRYA (ἀκράμβρυα, ἀκράμβρι, and βερία). Stem increasing both by apex and circumference. Vegetation peripherico-terminal.

Cohort 1. Gymnospermæ (γυμνός, naked, and σπείριμα, seed). Ovules naked, receiving the fecundating matter directly at the micropyle.

Cohort 2. Apetalæ (α, privative or without, and πέταλον, a petal). Perigone either wanting or rudimentary or simple, calycine or coloured, free or adherent to the ovary.

Cohort 3. Gamopetalæ (γάμος, union). Perigone double; outer calycine, inner corolline; gamopetalous, rarely wanting by abortion.

Cohort 4. Dialypetalæ (διαλίω, I separate). Perigone double; outer calycine, parts distinct or united, free or attached to the ovary; inner corolline, parts distinct or very rarely cohering by means of the base of the stamens; insertion hypogynous, perigynous, or epigynous; sometimes abortive.

Under these sections, Endlicher enumerates 279 natural orders, which are grouped under 61 classes.

731. DIVISION OF THE VEGETABLE KINGDOM BY LINDLEY. 1839.

Exogens.....	{ Cyclogens. (Wood in circles), Wood in wedges,	{ Class I. Exogens (proper). II. Gymnogens (naked seeds). III. Homogens.
Endogens	{ Spermogens, (Bearing seeds), (Bearing spores,	{ IV. Dictyogens (leaves reticulated). V. Endogens (proper). VI. Sporogens or Rhizanth.
Acrogens	{ Distinct Stem, Only a Thallus.	VII. Cormogens. VIII. Thallogens.

In the Exogens and Endogens, the following subordinate series of sub-classes are formed:—

1. Consolidated. Floral envelopes are united both with each other and the stamens, and with the ovary.
2. Separated. Floral envelopes and stamens are united to each other, but the ovary is consolidated and free.
3. Adherent. Petals and sepals adhere to each other and the stamens, and ovary, but have their parts disunited.
4. Disunited. Petals and sepals adhere to each other and the stamens, but have their parts disunited, and do not adhere to the consolidated ovary.
5. Dissolved. Petals and sepals are distinct from the stamens, and also from the ovary, whose carpels are disunited, either wholly or by the styles.

In each of these subdivisions, the orders are arranged in two series, the one Albuminous, the other Exalbuminous.

732. LINDLEY'S DIVISION OF THE VEGETABLE KINGDOM. 1846.

ASEXUAL OR FLOWERLESS PLANTS.

Stem and leaves undistinguishable.....	Class I. Thallogens.
Stem and leaves distinguishable.....	II. Acrogens.

SEXUAL OR FLOWERING PLANTS.

Fructification springing from a thallus.....	III. Rhizogens.
Fructification springing from a stem,	
Wood of stem youngest in the centre, cotyledon single.	
Leaves parallel-veined, permanent, wood of stem always confused.....	IV. Endogens.
Leaves net-veined, deciduous, wood of stem, when perennial, arranged in a circle with a central pith.....	V. Dictyogens.
Wood of stem youngest at the circumference, always concentric, cotyledons two or more.	
Seeds quite naked.....	VI. Gymnogens.
Seeds enclosed in seed-vessels.....	VII. Exogens.

The following are the subclasses of Endogens and Exogens adopted by Lindley :—

ENDOGENS.

- Subclass 1. Glumaceous.—Floral envelopes imbricated.
 ——— 2. Petaloid.—Floral envelopes verticillate.
 a. Unisexual, often achlamydeous.
 b. Hermaphrodite, ovary adherent.
 c. Hermaphrodite, ovary free.

EXOGENS.

- Subclass 1. Diclinous.—Flowers unisexual.
 ——— 2. Hypogynous.—Flowers usually hermaphrodite, stamens completely hypogynous, free from the calyx or corolla.
 ——— 3. Perigynous.—Flowers usually hermaphrodite, stamens growing to the side of either the calyx or corolla; ovary superior, or nearly so.
 ——— 4. Epigynous.—Flowers usually hermaphrodite, stamens growing to the side of either the calyx or corolla; ovary inferior, or nearly so.

Under the classes, Lindley enumerates 303 natural orders, which are grouped together under 56 alliances. In this system of Lindley, the divisions of Asexual and Sexual plants correspond to Endlicher's 2 Regions; the 7 classes represent Endlicher's 5 sections; and the 56 alliances are equivalent to the 61 classes in Endlicher's system.

733. This division may be presented thus :—

Classes.	Wood.	Leaves.	Floral Envelopes.	Sexes.	Embryo.
1. Exogenæ.....	Exogenous..	Netted.....	Quinary.	Perfect ...	Dicotyledonous.
2. Gymnospermæ.....	Exogenous..	Parallel or forked	None....	Imperfect.	Dicotyledonous.
3. Endogenæ.....	Endogenous	Parallel.....	Ternary..	Perfect...	Monocotyledonous.
4. Dictyogenæ....	Endogenous	Netted.....	Ternary..	Perfect...	Monocotyledonous.
5. Rhizanthæ.....	None.....	None.....	Variable.	Imperfect.	A cotyledonous ?
6. Acrogenæ.....	Acrogenous..	Forked or	None....	None.....	A cotyledonous.
7. Thallogenæ....	None.....	None.....	None....	None.....	A cotyledonous.

734. Henslow has given a comparative view of all these systems, pointing out, in a tabular form, the corresponding divisions in each of them :—

Henslow's approximate Comparison of the Systematic Views of Jussieu, De Candolle, Endlicher, and Lindley.

JUSSIEU, 1789.	DE CANDOLLE, 1812.	ENDLICHER, 1836.	LINDLEY, 1846.
I. ACOTYLEDONES 1.	* Cellulares v. Acotyledonæ. <div> aphyllæ <div> { algæ, 2. lichenes, fungi, 1. } </div> </div> <div> ACOTYLEDONES.....IV. <div> foliosæ <div> ** V asculares v. Cotyledonæ. Monocotyledones v. Endogenæ. </div> </div> </div>	* ThallopHYTA. I. PROTOPHYTA. II. HYSTEROPHYTA. ** Cormophyta. <div> 1. Anophyta. 2. Protophyta. 3. Hysterophyta. </div>	* Flowerless (alliances). I. THALLOGENS... <div> 1. algaes. 2. lichinales. 3. fungales. </div> <div> 4. muscales. </div> <div> II. ACROGENS. <div> 5. lycopodales. 6. filicales. </div> </div> <div> ** Flowering. </div> <div> III. RHIZOGENS. </div> <div> IV. ENDOGENS and V. DICTYOGENS. <div> (11 alliances). </div> </div>
II. MONOCOTYLEDONES. <div> Dielines 4 irregulares 1 Apetalæ 2 Monopetalæ 3 Polypetalæ </div> <div> { 2 4 3 } </div> <div> { 15 5 6 7 8 9 10 11 12 13 } </div> <div> { { hypogynæ. epigynæ. perigynæ. } { (-gymnospermæ.) (-angiospermæ.) } { (-stamineæ.) } { (-corollæ.) } { (-petalæ.) } </div> <div> { epi- peri- hypo- hypo- peri- epi- } { corisan- theræ. synan- theræ. } </div>	MON. PHANEROGAMÆ.....II. <div> † † Dicotyledones v. Exogenæ. </div> <div> 4. Monochlamydeæ.... 3. Corollifloræ.. 2. Calycifloræ... 1. Thalamifloræ. </div> <div> DICY-LEDONESI. </div>	IV. AMPHIBRYA. 1. Gymnospermæ. 2. Apetalæ. V. ACRAMPHIBRYA. 3. Gamopetalæ. 4. Dialypetalæ.	VI. GYMNOGENS. <div> 1. Diclinous (8 alliances). 2. Hypogynous (14 alliances). 3. Perigynous 10 alliances). 4. Epigynous (17 alliances). </div>
III. DICOTYLEDONES.			

735. In the succeeding pages the natural orders will be grouped under the following divisions:—

A. PHANEROGAMOUS, OR VASCULAR FLOWERING PLANTS.

Class I. Dicotyledones or Exogenæ.

Dichlamydeæ having calyx and corolla.	{	Subclass 1. Thalamifloræ.....Petals distinct, stamens hypogynous.....	{	Polypetalæ of Jus- sieu.
		2. Calycifloræ.....Petals distinct, stamens perigynous.....		
		3. Corollifloræ.....Petals united, bearing the stamens.....		Monopetalæ of Jussieu.
Having a sin- gle perianth.	{	4. Monochlamydeæ...A calyx only, or none.....	{	Apetalæ and partly Diclines of Jussieu.
		a. Angiospermæ, seeds in an ovary.		
		b. Gymnospermæ, seeds naked.		

Class II. Monocotyledones or Endogenæ.

- Subclass 1. Dictyogenæ.....Floral envelopes verticillate, leaves reticulate.
 ——— 2. Petaloideæ or Floridæ, Floral envelopes verticillate, leaves parallel-veined.
 a. Hermaphrodite, ovary adherent.
 b. Hermaphrodite, ovary free.
 c. Unisexual, often achlamydeous.
 ——— 3. Glumacæ.....Floral envelopes imbricated, leaves parallel-veined.

B. CRYPTOGRAMOUS, OR CELLULAR FLOWERLESS PLANTS.

Class III. Acotyledones or Acrogenæ.

- Subclass 1. Ætheogamæ or Cormogenæ,.....Having vascular tissue.
 ——— 2. Amphigamæ, Thallogenæ, or Cellulares.....Entirely cellular.

CHAPTER II.

ARRANGEMENT AND CHARACTERS OF THE CLASSES AND NATURAL ORDERS.

SECTION I.—PHANEROGAMOUS PLANTS.

CLASS I.—DICOTYLEDONES AND EXOGENÆ, *Juss.* and *DC.* ACRAMPHIBRYA, *Endl.*


736. This is the largest class in the vegetable kingdom. The plants included under it have a *cellular* and *vascular* system, the latter consisting partly of elastic spiral vessels (fig. 49). The *stem* is more or less conical, and exhibits wood and true bark. The *wood* is exogenous, i.e. increases by additions at the periphery, the hardest part being internal (§ 72, &c.) It is arranged in concentric circles. Pith exists in the centre, and from it diverge medullary rays. The *bark* is separable, and increases by additions on the inside. The *epidermis* is fur-

nished with stomata (§ 50). *The leaves* are reticulated (§ 143,), usually articulated to the stem. *The flowers* are formed upon a quinary or quaternary type, and have stamens and pistils. *The ovules* are either enclosed in a pericarp, and fertilized by the application of the pollen to the stigma, or they are naked and fertilized by the direct action of the pollen. *The embryo* has two or more opposite cotyledons, and is exorhizal in germination (§ 629).

Subclass 1.—THALAMIFLORÆ.*

737. Calyx and corolla present; petals distinct,[†] inserted into the thalamus or receptacle; stamens hypogynous. This includes the hypogynous polypetalous orders of Jussieu, and a diclinous order (Menispermaceæ).

738. Order 1.—**Ranunculaceæ**, the Crowfoot Family. (*Polypetalæ* *Hypogynæ*.) Sepals 3-6, frequently 5, deciduous (fig. 556 *c*). Petals 5-15 (fig. 556 *p e*), rarely abortive, sometimes anomalous in form, (figs. 284 *p*, 285), occasionally with scales at the base (fig. 555 *a*). Stamens usually indefinite, hypogynous (fig. 556 *e*); anthers adnate (figs. 558, 559); carpels numerous, 1-celled (fig. 556 *p i*), distinct, or united into a single many-celled pistil; ovary containing one anatropal ovule (figs. 492, 560 *g*), or several united to the inner edge. Fruit various, either dry



achænia (figs. 463, 561), or baccate or follicular (figs. 443, 468).

* *Thalamus*, receptacle, and *flos*, flower.

† Sometimes the petals are abortive, and it is then difficult to determine whether the plant belongs to this subclass or to Monochlamydeae.

Figs. 556-561.—Exhibit the organs of fructification of *Ranunculus acris*, to illustrate the natural order Ranunculaceæ.

Fig. 556.—Flower cut vertically. *c*, Calyx. *pe*, Petals. *e*, Stamens. *pi*, Pistil composed of several carpels on an elongated receptacle or axis.

Fig. 557.—Diagram of the flower, showing 5 imbricated sepals, 5 petals alternating with the sepals, indefinite stamens in several whorls, multiples of the petals, and numerous carpels or achænia in the centre.

Fig. 558.—Adnate anther seen on the outer side. The anther is in this instance extrorse. In *Pæonia* and other *Ranunculaceæ* it is introrse.

Fig. 559.—Adnate anther viewed on the inside.

Fig. 560.—Vertical section of the ovary, *o*, showing the ovule, *g*. *s*, Stigma.

Fig. 561.—Fruit or achene cut vertically. *f*, Pericarp. *z*, Spermoderm or integument of the anatropal seed. *p*, Perisperm or albumen, between fleshy and horny. *e*, Minute embryo.

Seeds albuminous, erect, or pendulous; albumen horny (fig. 561 *p*); embryo minute (fig. 561 *e*).—Herbaceous, suffruticose, or rarely shrubby plants, having alternate or opposite, simple, much-divided leaves, with dilated sheathing petioles (fig. 233). Juice watery. Hairs, if present, simple.

739. The plants of the order are found in cold damp climates, and in the elevated regions of warm countries. Europe contains one-fifth of the order, and North America about one-seventh. The order is divided into five suborders:—1. Clematideæ; 2. Anemoneæ (fig. 247); 3. Ranunculeæ (fig. 233); 4. Helleboreæ (fig. 443); 5. Actææ, or Pæoniæ (fig. 370), according to the æstivation of the calyx, the nature of the fruit, &c. Henslow gives the following analysis of these suborders, with the number of British species in each:—

	Spec. Brit.	Anther.	Carpel.	Seed.	Æstiv.
1. Clematideæ.....	1	} extrorse	} mono-sperm.	•	valvate.
2. Anemoneæ.....	9			} pendulous erect.	imbricate.
3. Ranunculeæ.....	20				•
4. Helleboreæ.....	9		polysperm.	•	•
5. Pæoniæ.....	2	introrse	•	•	•

Lindley enumerates 41 known genera, comprising 1000 species. *Examples of the genera*—Clematis, Anemone, Ranunculus, Helleborus, Delphinium, Aconitum, Actæa, Pæonia, Podophyllum.

740. The order has narcotico-acrid properties, and the plants are usually more or less poisonous. The acidity is frequently volatile, and disappears when the plants are dried or heated. It varies in different parts of the plants, and at different seasons. *Ranunculus* (the genus whence the order is named) contains many acrid species, such as *R. sceleratus*, *alpestris*, *bulbosus*, *gramineus*, *acris*, and *Flammula*; while others, such as *R. repens*, *aquatilis*, *Lingua*, and *Ficaria*, are bland. The acidity is entirely lost by drying, and it disappears in the pericarps as the seeds, which are themselves bland, ripen. The leaves of *Aconitum Napellus*, Monkshood, contain a narcotic alkaloid, called aconita or aconitina. They are used as an anodyne in neuralgic affections, in the form of extract and tincture. The root of *Aconitum ferox* furnishes the powerful East Indian poison, called Bikh or Nabee. The leaves of *Clematis recta* and *Flammula* have been used as vesicants. The seeds of *Delphinium Staphisagria*, Stavesacre, are irritant and narcotic, and are used for destroying vermin. They owe their activity to an alkaloid principle, called delphinia. The Hellebores have been long noted for their irritant qualities. *Helleborus officinalis*, *niger*, *fetidus*, and *viridis*, act as drastic purgatives; hence the use of some of them in ancient times in cases of mania. The rhizome of *Podophyllum peltatum*, May-apple, is employed in America as a purgative. Some of the Ranunculaceæ are chiefly marked by bitter tonic properties. This order, in the position, number, and structure of its parts

of fructification generally, presents a close resemblance to others widely differing in their nature. It may, however, be distinguished from each by many particulars, as from Dilleniaceæ, in the want of an aril, a deciduous calyx, and whole habit; from Magnoliaceæ, in want of true stipules; from Papavaraceæ and Nymphæaceæ, in the distinct, not concrete carpels, watery, not milky fluids, and acrid, not narcotic properties. It closely approaches the Berberidaceæ, but differs in its stamens not bursting by recurved valves. In its numerous carpels, floral divisions, and indefinite stamens, it agrees with the Rosewort tribe, but differs in its stamens being hypogynous instead of perigynous, in the albumen surrounding the minute ovary, in want of true stipules, and acrid properties. Crowfoots and Umbellifers agree in many peculiarities, the latter have, however, their calyx concrete with the ovary, and are invariably definite in their stamens.

741. Order 2.—**Dilleniaceæ**, the Dillenia Family. (*Polypet. Hypog.*) Sepals 5, persistent. Petals 5, deciduous, in a single row. Stamens indefinite, hypogynous, either distinct or combined into bundles; filaments dilated at the base or apex; anthers adnate, introrse, with longitudinal dehiscence. Ovaries definite, more or less distinct, with a terminal style and simple stigma; ovules ascending. Fruit of 2-5 capsular or baccate unilocular carpels, which are either distinct or coherent. Seeds universally arillate, several in each carpel, or only two, or one by abortion; testa (spermoderm) hard; embryo straight, minute, at the base of fleshy albumen.—The plants of the order are trees, shrubs, or under-shrubs, having alternate, exstipulate, coriaceous, or rough leaves. They are found chiefly in Australasia, Asia, and the warm parts of America. The Indian species are remarkable for their beauty, the grandeur of their foliage, and the magnificence of their flowers. They have astringent properties, and some of the species afford excellent timber. Lindley enumerates 26 genera, including 200 species. *Examples*—Dillenia, Delima.

742. Order 3.—**Magnoliaceæ**, the Magnolia Family. (*Polypet. Hypog.*) Sepals 2-6, usually deciduous. Petals 2-30, hypogynous, often in several rows. Stamens indefinite, distinct, hypogynous; anthers adnate, long, dehiscing longitudinally. Carpels numerous, 1-celled, arranged upon a more or less elevated receptacle; ovules anatropal, suspended or ascending; styles short. Fruit consisting of numerous distinct or partially coherent carpels, which are either dehiscent or indehiscent, sometimes samaroid. Seeds, when ripe, often hang suspended from the carpels by a long slender cord; embryo minute, at the base of a fleshy perisperm.—Trees and shrubs, with alternate coriaceous leaves, and deciduous convolute stipules. They abound in North America, and some species occur in South America, China, Japan, New Holland, and New Zealand. The order has been divided into two suborders:—1. Winteræ; aromatic plants, in which the

leaves are dotted, the carpels are in a single verticil, and the wood often consists of punctated tissue (fig. 47). 2. *Magnoliæ*; bitter plants with fragrant flowers, in which the carpels are arranged in several rows on an elevated receptacle (fig. 306), and the leaves are not dotted. Lindley mentions 11 known genera, comprising 65 species. *Examples*—*Illicium*, *Drimys*, *Magnolia*, *Liriodendron*.

743. The properties of the order are bitter, tonic, and often aromatic. *Illicium anisatum*, Star-anise, is so called from its carpels being arranged in a star-like manner, and having the taste and odour of anise. Its fruit is employed as a carminative. *Drymis Winteri* or *aromatica*, brought by Captain Winter from the straits of Magellan in 1579, yields Winter's bark, which has been employed medicinally as an aromatic stimulant. It somewhat resembles *Canella* bark. *Magnolias* are remarkable for their large odoriferous flowers, and their tonic aromatic qualities. The bark of *Magnolia glauca*, Swamp Sassafras or Beaver-tree, is used as a substitute for Peruvian bark. *Liriodendron tulipifera*, the Tulip-tree, marked by its truncate leaves, has similar properties.

744. Order 4.—*Anonaceæ*, the Custard-apple Family. (*Polypet. Hypog.*) Sepals 3-4, persistent, often partially cohering. Petals 6, hypogynous, in two rows, coriaceous, with a valvate æstivation. Stamens indefinite (very rarely definite); anthers adnate, extrorse, with a large 4-cornered connective. Carpels usually numerous, separate or cohering slightly, rarely definite; ovules anatropal, solitary or several, erect or ascending. Fruit succulent or dry, the carpels being one or many-seeded, and either distinct or united into a fleshy mass; spermoderm brittle; embryo minute, at the base of a ruminated perisperm, which forms the great feature of the order, there being no exception and very few parallels. Trees or shrubs, with alternate, simple, exstipulate leaves, found usually in tropical countries. Lindley enumerates 20 genera, including 300 species. *Examples*—*Anona*, *Uvaria*, *Guatteria*.

745. Their properties are generally aromatic and fragrant. Some of the plants are bitter and tonic, others yield edible fruits. The custard-apples, Sweetshops, and Soursops of the East and West Indies, are furnished by various species of *Anona*, such as *A. squamosa*, *reticulata*, and *Muricata*. *Anona cherimolia* furnishes the Cherimoyer, a well-known Peruvian fruit. The fruit of *Xylopia aromatica* is commonly called Ethiopian Pepper, from being used as pepper in Africa. *Xylopia glabra* is called Bitter-wood in the West Indies. The Lancewood of coachmakers appears to be furnished by a plant belonging to this order, called by Schomburgk *Duguetia quitarensis*.

746. Order 5.—*Menispermaceæ*, the Moon-seed Family. (*Polypet. Hypog.*) Flowers usually unisexual (often diœcious), generally of a pale-greenish hue. Sepals and petals similar in appearance, in one or

several rows, 3 or 4 in each row, hypogynous, deciduous. Stamens monadelphous, or occasionally free; anthers adnate, extrorse. Carpels solitary or numerous, distinct or partially coherent, unilocular; ovule solitary, curved (fig. 420). Fruit a succulent 1-seeded oblique or lunate drupe. Embryo curved or peripheral; radicle superior; albumen fleshy, sometimes wanting.—The plants of this order are sarmentaceous or twining shrubs, with alternate leaves, and very small flowers. The wood is frequently arranged in wedges, and hence the order was at one time put under the division called Homogens by Lindley (§ 90 and 731). The order is common in the tropical parts of Asia and America. There are 23 known genera, including 202 species. *Examples*—*Menispermum*, *Cissampelos*, *Cocculus*, *Lardizabala*, *Schizandra*.

747. The species are bitter and narcotic. Some are employed as tonics, others have poisonous properties. The root of *Cocculus palmatus*, a plant found in the eastern part of Africa, is known as Calumbaroot, and is used as a pure bitter tonic in cases of dyspepsia, in the form of infusion or tincture. It contains a bitter crystallizable principle called Calumbin. *Cocculus indicus* is the fruit of *Anamirta Cocculus*. It is extremely bitter, and contains a crystalline poisonous narcotic principle, Picrotoxin, which is its active ingredient. It has been used externally in some cutaneous affections. At one time it was employed, most prejudicially, to give bitterness to porter. *Cissampelos Pareira*, Wild-vine or Velvet-leaf, furnishes Pareira-bravaroot, which is employed as a tonic and diuretic, and has been recommended in chronic inflammation of the bladder. Several plants of this order are characterized by their remarkable tenacity of life. It is stated that if a large branch be broken at a considerable distance from the ground, the upper portion immediately throws out a slender filiform shoot, which establishes a connection with the soil and preserves the plant. These shoots have been seen as long as eight feet, and filiform in their appearance.

748. Order 6.—**Berberidaceæ**, the Berberry Family. (*Polypet. Hypog.*) Sepals 3-4-6, deciduous, in a double row. Petals hypogynous, equal in number to the sepals, and opposite to them, or twice as many, often having an appendage at the base on the inside. Stamens equal in number to the petals, and opposite to them; anthers adnate, bilocular (dithecal), each of the loculi opening by a valve from the bottom to the top. Carpel solitary, unilocular, containing 2-12 anatropal ovules; style sometimes lateral; stigma orbicular. Fruit baccate or capsular, indehiscent. Albumen fleshy or horny; embryo straight, sometimes large (fig. 494).—Shrubs or herbaceous perennial plants with alternate, compound, exstipulate leaves. The true leaves are often changed into spines by the absorption of their parenchyma and induration of their ribs (fig. 231 f). Found chiefly in the mountainous parts of the temperate regions of the northern hemisphere.

The plants of the order have bitter and acid properties. The bark and stem of *Berberis vulgaris*, common Berberry, are astringent, and yield a yellow dye; the fruit contains oxalic acid, and is used as a preserve. Lindley enumerates 12 genera, including 100 species. *Examples*—*Berberis*, *Epimedium*, *Leontice*.

749. Order 7.—**Cabombaceæ**, the Watershield Family. (*Polypet. Hypog.*) Sepals 3-4. Petals 3-4, alternate with the sepals. Stamens hypogynous, arising from an inconspicuous torus, two or three times the number of the petals; anthers linear, introrse, continuous with the filament. Carpels 2 or more; stigmas simple; ovules orthotropical. Fruit indehiscent, tipped with the indurated styles, containing one or two pendulous seeds. Embryo minute, enclosed in a vitellus (the sac of the amnios), and placed at the base of a fleshy perisperm.—American aquatic plants, with floating peltate leaves, said to possess nutritious but slightly astringent properties, which have been found useful as a remedy for phthisis and dysentery. Lindley mentions 2 genera, including 3 species. *Examples*—*Hydropeltis*, *Cabomba*.

750. Order 8.—**Nymphæaceæ**, the Water-lily Family (fig. 562). (*Polypet. Hypog.*) Sepals usually 4, sometimes confounded with the petals. Petals numerous, often passing gradually into stamens (fig. 310, 2). Stamens indefinite, inserted above the petals into the torus (fig. 562 c); filaments petaloid; anthers adnate, introrse, opening by two longitudinal clefts. Torus large, fleshy, surrounding the ovary more or less (fig. 562 t). Ovary multilocular, many-seeded, with radiating stigmas (fig. 562 s); numerous anatropal ovules. Fruit many-celled, indehiscent. Seeds very numerous, attached to spongy dissepiments; albumen farinaceous; embryo small, enclosed in a fleshy vitellus, and situated at the base of the perisperm (fig. 480).—Aquatic plants, with peltate or cordate fleshy leaves, and a rootstock or stem which extends itself into the mud at the bottom of the water. Lindley enumerates 5 genera, comprehending 50 species. *Examples*—*Nymphaea*, *Nuphar*, *Victoria*, *Euryale*.

751. The plants of this order are found throughout the northern hemisphere, and are generally rare in the southern hemisphere. Little is known in regard to their properties. Some of them are astringent and bitter, while others are said to be sedative. They have showy flowers, and their petioles and




Fig. 562.—Section of a flower of *Nymphaea alba*, white Water-lily, showing the pistils and the receptacle or torus bearing the stamens and petals. *p*, Peduncle or flower-stalk. *t*, Elevated torus or receptacle. *s*, Radiating stigmas. *a*, Sepal. *b*, Petal. *c*, Stamens.

peduncles contain numerous air-tubes. *Victoria regia* is one of the largest known aquatics. It is found in the waters of South America, and is said to range over 35 degrees of longitude. The flowers have a fine odour. When expanded they are a foot in diameter. The leaves are from four to six and a-half feet in diameter. The seeds and rootstocks of many plants of this order contain much starch, and are used for food. It has been said that the rhizomes of *Nymphæa alba* are better than Oak-galls for dyeing gray; that they have been long employed advantageously for tanning leather. The leaves of *Nuphar luteum* are reported to be styptic.

752. Order 9.—**Nelumbiaceæ**, the Water-bean Family. (*Polypet. Hypog.*) Sepals 4-5. Petals numerous, in many rows. Stamens indefinite, in several rows; filaments petaloid; anthers adnate, introrse, opening by a double longitudinal cleft. Torus large, fleshy, elevated, enclosing in hollows of its surface numerous carpels. Nuts numerous, inserted, but loose, into the depressions of the torus. Seeds 1-2; perisperm none; embryo enclosed in a vitellus, large, with two fleshy cotyledons.—Aquatic herbs, with showy flowers, peltate floating leaves, and prostrate rootstocks, found in the temperate and tropical regions of the old and new world. Lindley enumerates 1 genus, including 3 species. *Example*—*Nelumbium*.

753. The flower of *Nelumbium speciosum* is supposed to be the Lotus figured on Egyptian and Indian monuments, and the fruit is said to be the Pythagorean Bean. The plant is said to have dissappeared from the Nile, where it used to abound. The petioles and peduncles contain numerous spiral vessels, which have been used for wicks of candles. Dr. Wight states that those wicks "on great and solemn occasions are burnt in the lamps of the Hindoos placed before the shrines of their gods."

754. Order 10.—**Sarraceniaceæ**, the Sidesaddle-flower Family. (*Polypet. Hypog.*) Sepals 5, persistent, imbricated in æstivation, often with coherent bracts outside. Petals 5, hypogynous, concave; occasionally the corolla is absent, and the calyx consists of 4-6 segments. Stamens 00; anthers adnate, dithecal, introrse, with longitudinal dehiscence. Ovary free, tri-quinquelocular; style single; stigma persistent, either a truncated point, or large and peltate with 5 angles; ovules anatropal. Capsule 3-5-celled, with loculicidal dehiscence. Seeds very numerous, small, attached to large placentas which project from the axis into the cavity of the cells; albumen copious; embryo cylindrical, lying at the base of the seed; radicle pointing to the hilum.—Herbaceous plants, found in boggy places, having radicle leaves, the petioles of which are folded, and cohere so as to form ascidia or hollow tubes, which being lined by hairs of a singular nature, appear to act as secreting organs (fig. 187). Scapes one or more flowered. The plants are found in North America and

Guiana. Their properties are not known. Lindley enumerates 2 genera, including 7 species. *Examples*—*Sarracenia*, *Heliamphora*.

755. Order 11.—**Papaveraceæ**, the Poppy Family. (*Polypet. Hypog.*) Sepals 2, deciduous. Petals hypogynous, usually 4, cruciate, sometimes a multiple of 4, regular, rarely wanting. Stamens hypogynous, usually 00, sometimes a multiple of 4; anthers dithecal, innate. Ovary solitary; style short or none; stigmas 2, or many and radiating (fig. 409); ovules 00, anatropal (fig. 421). Fruit unilocular, either siliquæform with two, or capsular with several parietal placentas. Seeds numerous; albumen between fleshy and oily; embryo minute, at the base of the albumen, with plano-convex cotyledons.—Herbs or shrubs, usually with milky or coloured juice, having alternate exstipulate leaves, and long one-flowered peduncles. The plants belonging to this order are chiefly European. The species, however, are found scattered over tropical America, Asia, China, New Holland, Cape of Good Hope, &c. Lindley mentions 18 known genera, and 130 species. *Examples*—*Papaver*, *Meconopsis*, *Eschscholtzia*, *Glaucium*, *Chelidonium*.

756. The order possesses well-marked narcotic properties. Opium is the concrete milky juice procured from the unripe capsules of *Papaver somniferum*, and its varieties. The plant is a native of Western Asia, and probably also of the south of Europe; but it has been distributed over various countries. There are five kinds of opium known, viz. Turkey, Egyptian, East Indian, European, and Persian; of which the first is the kind chiefly used in Britain. The most important active principle in opium is the alkaloid called morphia. There are other crystalline principles found in it, such as codeia and narcotine; and an acid called meconic acid, which constitutes with sulphuric acid the solvent of the active principles. According to various circumstances, opium may act as a stimulant, a narcotic, anodyne, or diaphoretic. The seeds of the Opium Poppy yield a bland, wholesome oil. *Chelidonium majus*, Celandine, yields an orange-coloured juice, which is said to have acrid properties. In this plant, observations were made by Schultz on Cyclosis (§ 261, and fig. 223). *Eschscholtzia* is remarkable for the dilated apex of the peduncle, from which the calyx separates in the form of a calyptra, resembling an extinguisher of a candle. *Sanguinaria canadensis*, Blood-root, or Puccoon, has emetic and purgative properties.

757. Order 12.—**Fumariaceæ**, the Fumitory Family. (*Polypet. Hypog.*) Sepals 2, deciduous. Petals 4, cruciate; one or both of the two outer gibbous at the base, the two inner cohering at the apex. Stamens hypogynous, usually 6, diadelphous; anther of middle stamen of each parcel bilocular, outer ones unilocular. Ovary free, 1-celled; style filiform; stigma with 2 or more points; ovules amphitropal. Fruit either an achæmium, or a 2-valved 2-seeded capsule, or a many-


Fumariaceæ two Caducous Sepals.

Impatiens *Conium* *Petals* *6* *diadelphous* *the*

Papaveraceæ 2. Caducous Sepals. Crucial Petals.

seeded siliqua. Seeds crested; albumen fleshy; embryo minute, excentric.—Herbaceous plants, with a watery juice, and alternate multifid leaves. Although at first sight very unlike the Poppy family, the Fumitories resemble this order in their deciduous sepals, in their seeds, and, in many cases, in their fruit. The two outer unilocular stamens of each parcel may be considered as forming one perfect stamen, thus making the whole number four. They are found chiefly in northern temperate latitudes. Two are found at the Cape of Good Hope. They are usually scentless, and are said to be bitter and diaphoretic in their properties. The tuber of *Corydalis bulbosa* has been used as a substitute for Birthworts in expelling intestinal worms, and as an emmenagogue. Lindley notices 15 genera, including 110 species. *Examples*—*Fumaria*, *Corydalis*, *Diclytra*.

758. Order 13.—**Cruciferae**, the Cruciferous or Cresswort Family. Brassicaceæ of Lindley. (*Polypet. Hypog.*) Sepals 4, deciduous, the two lateral ones gibbous at the base. Petals 4, hypogynous, alternating with the sepals, deciduous, cruciate (fig. 291). Stamens 6, tetradynamous (figs. 345, 565), two shorter solitary (fig. 565 *e'*), opposite the lateral sepals, occasionally toothed; four longer (fig. 565 *e''*), opposite the anterior and posterior sepals, generally free, sometimes partially united and furnished with a tooth on the inside; anthers bilocular, introrse (fig. 564). Torus with green glands between the petals and stamens and ovary (fig. 565 *g*). Ovary superior, with parietal placentas, which meet in the middle, forming a spurious dissepiment or replum (fig. 566 *c*); stigmas 2, opposite the placentas, or anterior and posterior (fig. 456 *s*). Fruit a siliqua (figs. 567, 568), or a silicula, rarely 1-celled and indehiscent, usually spuriously



Figs. 563-570.—Organs of fructification of *Erysimum lanceolatum*, one of the Cruciferae.

Fig. 563.—Diagram of the flower, showing the arrangement of four sepals, four petals alternating with them, six tetradynamous stamens, and a siliqua with replum.

Fig. 564.—Vertical section of the flower. *c*, Calyx. *p*, Petals. *e*, Stamens. *o*, Ovary laid open. *s*, Stigma.

Fig. 565.—Flower deprived of its envelopes. *c c*, Cicatrices left by the fall of the sepals. *g*, Glands which are situated at the base of the stamens. *e'*, Two short stamens opposite lateral sepals. *e''*, Four long stamens opposite anterior and posterior sepals. *p*, Pistil.

Fig. 566.—Horizontal section of the ovary. *g*, Ovules. *c*, Spurious dissepiment or replum, which divides the ovary into two cavities. This replum is formed by the placentas.

Cruciferae
Cruciate Petals -
Tetradynamous
Stamens -
Siliqua

2-celled and dehiscing by two valves, which separate from the replum (figs. 456, 568), one or many-seeded. Seeds Campylotropous (figs. 419, 524), pendulous, attached in a single row by a funiculus to each side of the placentas (fig. 568); perisperm none; embryo with the radicle folded upon the cotyledons which are next the placenta (figs. 524, 569 *r*).—Herbaceous plants seldom undershrubs, with alternate leaves, and yellow or white, rarely purple, flowers, without bracts. This order is well distinguished by having tetradynamous stamens. Most of the plants belonging to the order are European. The species, however, are found scattered all over the world. De Candolle, in 1821, calculated that the species then known were distributed as follows:—

In the frigid zone of the northern hemisphere.....	205
In all the tropics (chiefly mountainous parts).....	30
In the temperate zone, { of northern hemisphere.....548 of southern hemisphere..... 86 }	634

Lindley enumerates 173 genera, including 1600 species. *Examples*—*Draba*, *Lepidium*, *Isatis*, *Brassica*, *Sinapis*, *Bunias*, *Senebiera*, *Schizopetalon*.

759. The order has been subdivided into sections, according to the mode in which the radicle of the embryo is folded on the cotyledons, as well as according to the nature of the fruit. The suborders founded on the embryo are—1. *Pleurorhizæ* (πλευρῆς, side, and ρίζα, root), 0 = cotyledons accumbent, radicle lateral, *i.e.* applied to their edge (fig. 517). 2. *Notorhizæ* (νότον, back), 0 || cotyledons incumbent, radicle dorsal, *i.e.* applied to their back (fig. 518). 3. *Orthoplocæ* (ὀρθός, straight, and πλόκος, a plait or fold), 077 cotyledons conduplicate (folded), radicle dorsal (figs. 513, 570). 4. *Spirolobæ* (σπείρα, a coil, and λοβός, a lobe), 0 || cotyledons twice folded, radicle dorsal (fig. 515). 5. *Diplecolobæ* (δίς, twice, πλέκω, I fold or plait, and λοβός, a lobe), 0 || || cotyledons three times folded, radicle dorsal.

760. The divisions founded on the seed-vessel, are—1. *Siliquosæ*, a silique, linear or linear-lanceolate, valves opening longitudinally. 2. *Latisepthæ* (*latus*, broad, and *septum*, partition), a silicula, partition in its broadest diameter, oval or oblong, valves flat or convex, opening longitudinally. 3. *Angustiseptæ* (*angustus*, narrow), a silicula, parti-

Fig. 567.—Silique or long pod.

Fig. 568.—Silique with one of its valves removed, in order to show the seeds attached to the replum.

Fig. 569.—Vertical section of the seed. *f*, Funiculus or umbilical cord. *t*, Spermoderm or testa swollen at the chalaza. *c*, *r*, Radicle. *c*, Cotyledons.

Fig. 570.—Horizontal section of the seed. *t*, Spermoderm or testa. *r*, Radicle. *c*, Incumbent cotyledons.

tion in its narrow diameter, linear or lanceolate, valves opening longitudinally, folded and keeled. 4. Nucumentacæ (*nucumentum*, a nut), silicula, valves indistinct or indehiscent, often 1-celled, from the absence of the replum or partition. 5. Septulatæ (*septa*, partitions), valves opening longitudinally, furnished with transverse partitions in their interior. 6. Lomentacæ (*lomentum*, an articulate legume), siliqua or silicula, dividing transversely into single-seeded cells, the true siliqua being often barren and all the seeds placed in the beak.

761. In this order there is a want of symmetry as regards the number of stamens, compared with the floral envelopes. The two long stamens placed close together may, however, be looked upon as one divided by a process of deduplication, so that the actual number will thus be reduced to four. This view is confirmed by the shorter stamens having teeth on each side, while the longer ones are toothed on one side only. By pelorization, too, some Cruciferæ become tetrandrous. While there is a splitting of the filaments, there is also the production of two additional anther-lobes. In regard to the fruit it has been stated, that normally there are four carpidia or carpels, two of which are constantly abortive. In some species of Iberis there have been seen four sepals, four petals, four stamens, and four carpels. Thus the floral type of Cruciferæ is quaternary: calyx having four sepals, corolla four petals, receptacle four staminiferous glands, andrœcium four stamens, gynœcium four phyllidia, fruit four carpidia.

762. There are no truly poisonous plants in the order. In general, it possesses antiscorbutic and stimulant qualities, with a certain degree of acridity. Many of the most common culinary vegetables belong to the order, such as Cabbages, Cauliflower, Turnip, Radish, Cress, Horseradish, &c. They contain much sulphur and nitrogen, and, on this account, when decaying, give off a disagreeable odour. Many garden flowers, such as Wall-flower, Stock, Rocket, and Honesty, are found in this order. *Brassica oleracea* is the original species whence all the varieties of Cabbage, Cauliflower, Brocoli and Savoys have been obtained by the art of the gardener. The part of the Cauliflower used as food is the deformed flower stalks. *Brassica rapa* is the common Turnip, while *Brassica campestris* is the source of the Swedish Turnip. *Brassica chinensis* yields Shanghæ oil. *Lepidium sativum* is the common Cress, and *Raphanus sativus* the Radish. *Crambe maritima* is the Sea-kale. The seeds of *Sinapis nigra* (*Brassica nigra* of some), furnish table mustard. These contain a bland fixed oil, a peculiar bitter principle, myronic acid, and another principle analogous to albumen or emulsin, called myrosine. When water is added, the myronic acid and myrosine, by their combination, form a pungent volatile oil, containing sulphur and nitrogen, which gives to mustard its peculiar properties as a physiological agent. *Sinapis alba* furnishes white mustard, which contains more fixed oil than black mustard. It does not, however,

contain myronic acid, but an analogous principle called sinapin, or sinapisin, which, by combination with another principle, forms an acrid compound, but not a volatile oil. The mustard of Scripture is not a species of *Sinapis*, but *Salvadora persica*, according to Royle, belonging to the natural order *Chenopodiaceæ*. Many other Cruciferous plants yield volatile oils containing sulphur, and the seeds of many yield by expression a bland fixed oil, such as Rape-seed oil. *Cochlearia officinalis*, common Scurvy-grass, is used as a stimulant. *Cochlearia Armoracia*, or *Armoracia rusticana*, the Horse-Radish, has irritant and even vesicant qualities. *Anastatica hierochuntina*, Rose of Jericho, is remarkable for the hygrometric property of the old withered annual stems, which are rolled up like a ball in dry weather, and drifted about by the winds in the deserts of Syria and Egypt. If rain falls, they resume their original direction. They thus continue for many years to curl up and expand, according to the state of the atmosphere. The genus *Schizapetalon* is remarkable on account of its tetracotyledonous (having four cotyledons) embryo. *Isatis tinctoria*, Woad, when treated like Indigo, yields a blue dye. *Isatis indigotica* is the Tein-Ching, or Chinese Indigo.

763. Order 14.—**Capparidaceæ**, the Caper Family. (*Polypet. Hypog.*) Sepals 4, often more or less cohering (fig. 554 c). Petals 4, sometimes 0, cruciate (fig. 554, p), usually unguiculate and unequal. Stamens hypogynous, 4, 6 (fig. 554 e), or 00, but in general some high multiple of four, placed on an elongated hemispherical and often glandular torus (fig. 554 a g'). Ovary usually stalked (fig. 554 o); styles filiform, sometimes 0; ovules curved. Fruit unilocular, siliqueæ-form and dehiscent, or fleshy and indehiscent, rarely monospermous, usually with two polyspermous placentas. Seeds generally reniform and exalbuminous; embryo curved; cotyledons foliaceous, flattish.—Herbs, shrubs, sometimes trees, with alternate, stalked, undivided, or palmate leaves, which are either exstipulate or have spines at their base. Capparids may be distinguished from Crucifers by their stamens being often indefinite, or, if definite, scarcely ever tetradynamous, while their seeds are generally reniform. They are found chiefly in warm countries, and are abundant in Africa. There are 28 genera, and 340 species. The order is divided into two suborders:—1. Cleomeæ, with capsular fruit. 2. Cappareæ, with baccate fruit. *Examples*—Cleome, Capparid.

764. The plants of this order have stimulant qualities. The flower-buds of *Capparis spinosa* furnish capers. The plant is a native of the south of Europe, and, according to Royle, is the Hyssop (היסופ) of Scripture. Some species of *Cleome* and *Polanisia* are very pungent, and are used as substitutes for mustard. The pungency of some is so great that they act as blisters. The root of *Cleome dodecandra* is used as an anthelmintic.

765. Order 15.—**Resedaceæ**, the Mignonette Family. (*Polypet.*

Hypog.) Calyx many-parted. Petals 4-6, unequal, entire, or lacerated, in the latter case consisting of a broad scale-like claw with a much-divided limb. Stamens 10-24, hypogynous, attached to a glandular torus; filaments variously united; anthers bilocular, innate, with longitudinal dehiscence. Ovary sessile, 3-lobed, 1-celled, multiovular, with 3-6 parietal placentas; stigmas 3. Fruit either a unilocular many-seeded capsule, opening at the apex so as to render the seeds seminude (fig. 479), or 3-6 few-seeded folicles. Seeds reniform, usually exalbuminous; embryo curved; radicle superior; cotyledons fleshy.—Herbaceous plants, rarely shrubs, with alternate, entire, or divided leaves, having gland-like stipules. They inhabit chiefly Europe and the adjoining parts of Asia. A few are found in the north of India and south of Africa. The uses of the order are unimportant. *Reseda Luteola*, Weld, yields a yellow dye. *Reseda odorata* is the fragrant Mignonette. The Mignonette is rendered suffruticose, by preventing the development of its blossoms. This is the origin of the tree Mignonette, which is much cultivated in France. There are 6 known genera, and 41 species, according to Lindley. *Example*—*Reseda*.

766. Order 16.—**Flacourtiaceæ**, the Arnotto Family. (*Polypet. Hypog.*) Sepals 4-7, slightly cohering. Petals equal to and alternating with the sepals, or wanting. Stamens hypogynous, equal in number to the petals, or some multiple of them. Ovary roundish, sessile, or slightly stalked; style either none or filiform; stigmas several, more or less distinct; ovules attached to parietal placentas, which sometimes branch all over the inner surface of the valves. Fruit 1-celled, containing a thin pulp, either fleshy and indehiscent, or capsular with 4 or 5 valves. Seeds numerous, enveloped in a covering formed by the withered pulp; albumen fleshy, somewhat oily; embryo axile, straight; radicle turned towards the hilum; cotyledons flat, foliaceous.—Shrubs or small trees, with alternate, simple, usually exstipulate leaves, which are often dotted. The plants are chiefly natives of the warmest parts of the East and West Indies, and of Africa. The order is divided into two sections:—1. Flacourtiæ, having the placentas ramifying over the inner surface of the fruit. 2. Bixæ, placentas narrow, and running in lines along the parietes. Many of the plants yield edible fruits. The pulp is often sweet and wholesome. Some are astringent, others purgative. The reddish pulp surrounding the seeds of *Bixa orellana* supplies the substance called arnotto, which is used for yielding a red colour, and for staining cheese. Martius says that the seeds are cordial, astringent and febrifugal. There are 31 genera, and 85 species, according to Lindley. *Examples*—*Patrisia*, *Flacourtia*, *Prockia*, *Erythrospermum*, *Bixa*.

767. Order 17.—**Cistaceæ**, the Rock-Rose Family. (*Polypet. Hypog.*) Sepals usually 5, persistent, unequal, the three inner with contorted

æstivation. Petals 5, caducous, hypogynous, æstivation corrugated, and twisted in an opposite direction to that of the sepals. Stamens usually 00, free, hypogynous; anthers 2-celled, adnate. Ovary syncarpous, 1- or many-celled; style single; stigma simple. Fruit capsular, 3-5-10-valved, either 1-celled or imperfectly 5-10-celled, with loculicidal dehiscence. Seeds usually indefinite; embryo inverted, either spiral or curved, in the midst of mealy albumen; radicle remote from the hilum.—Shrubs or herbaceous plants with entire, opposite, or alternate, stipulate or exstipulate leaves. They inhabit chiefly the southern regions of Europe, and the north of Africa. Some of the species are remarkable for the irritability of their stamens (§ 665). Many of them yield a resinous balsamic juice, which imparts viscosity to the branches. The resinous matter called ladanum or labdanum, is yielded by *Cistus creticus*. There are 7 known genera, and 185 species, according to Lindley. *Examples*—*Cistus*, *Helianthemum*, *Lechea*.

768. Order 18.—**Violaceæ**, the Violet Family. (*Polypet. Hypog.*) Sepals 5, persistent usually elongated at the base, æstivation imbricated. Petals 5, hypogynous, equal or unequal, generally withering, æstivation obliquely convolute. Stamens 5, alternate with the petals, sometimes opposite to them, inserted on a hypogynous torus; anthers dithecal, introrse, often cohering, with a prolonged connective sometimes spurred (fig. 343); filaments dilated, two of them in the irregular flowers having an appendage at their base. Ovary unilocular, with many (rarely one) anatropal ovules; style single, usually declinate, with an oblique hooded stigma (fig. 390, 1 s). Fruit a 3-valved capsule, dehiscence loculicidal, placentas on the middle of the valves (fig. 390). Seeds 00 or definite; embryo straight, erect, in the axis of a fleshy perisperm.—Herbs or shrubs, with alternate, rarely opposite, leaves, having persistent stipules, and an involute vernation. They are natives of Europe, Asia, and America. The herbaceous species inhabit chiefly the temperate parts of the northern hemisphere, while the shrubby species are found in South America and India. They have been divided into two suborders:—1. *Violeæ*, with irregular flowers. 2. *Alsodeieæ*, with regular flowers. Their distinctive peculiarity may be regarded as resting in their definite stamens, whose anthers turn inwards, and extend their connective into a crest. There are 14 known genera, and 315 species. *Examples*—*Viola*, *Ionidium*, *Alsodeia*.

769. They are distinguished by the emetic properties of their roots, which contain an active principle called violine, similar in its qualities to emetin. The roots of *Viola odorata* have been used medicinally as emetics. Some species of *Ionidium* are used in South America as substitutes for *Ipecacuanha*. The petals of the Sweet or March Violet, the *ῥο* of the Greeks, are laxative, and are used in the form of infu-

Bracææ - involute Stip. to the Leaves, Sep. Pet. Stam. 5 - unilocular loculicidal Fruit

sion mixed with sugar; and a violet or purple colouring matter is procured from them, which is employed as a test for acids and alkalies, being changed into red by the former, and into green by the latter. *Viola tricolor*, Heart's-ease, and other species, have been used as demulcent expectorants. *V. tricolor* is the origin of all the cultivated varieties of pansy.

770. Order 19.—**Droseraceæ**, the Sundew Family. (*Polypet. Hypog*). Sepals 5, persistent, equal; æstivation imbricated. Petals 5, hypogynous. Stamens free, withering, alternate with the petals, or 10 or more; anthers bilocular, with longitudinal dehiscence. Ovary single; styles usually 3-5, sometimes 1 or wanting. Fruit, a unilocular or spuriously trilocular capsule, 3-5-valved, with loculicidal dehiscence, occasionally indehiscent. Seeds numerous, either albuminous or exalbuminous; embryo minute and erect.—Herbaceous plants with alternate leaves, usually inhabiting marshy places. They are found in various parts of the world, in Europe, North and South America, China, East Indies, &c. The order has been divided into two sections:—1. *Drosereæ*, styles 3-5, albumen present, leaves with a circinate veneration, and furnished with glandular hairs; stipules in the form of fringes. 2. *Parnassieæ*, some of the stamens sterile; style 1 or none; no albumen nor glandular hairs. There are 8 known genera, and upwards of 90 species. *Examples*—*Drosera*, *Aldrovanda*, *Dionæa*, *Parnassia*.

771. The *Droseras* have a more or less acid taste, combined with slight acidity. Some of them are said to be poisonous to cattle. Their leaves are furnished with glandular capitate hairs, which are covered with drops of fluid in sunshine; hence the name Sundew or *Ros solis*. An Italian liqueur, called Rossoli, derives its name from a *Drosera* used in its manufacture. Some of the *Droseras* have dyeing properties. *Dioncea muscipula*, Venus's fly-trap, is a North American plant, having the laminæ of the leaves in two halves, each furnished with three irritable hairs, which, on being touched, cause the folding of the divisions in an upward direction (figs. 186, 664). *Aldrovandra vesiculosa*, an aquatic found in the South of Europe, is distinguished by its whorled cellular leaves, or floating bladders. *Parnassia palustris*, Grass of Parnassus, has remarkable gland-like bodies between the stamens (fig. 304). These are probably an abortive form of the staminal organs. Lindley looks upon them as bundles of stamens, and places the genus among Hypericaceæ. The stamens of *Parnassia* are irritable, and move towards the pistil in succession (fig. 665).

772. Order 20.—**Polygalaceæ**, the Milkwort Family. (*Polypet. Hypog.*) Sepals 5, very irregular, distinct; 3 exterior, of which 1 is superior, and 2 inferior; 2 interior, usually petaloid, lateral; æstivation imbricated. Petals hypogynous, unequal, usually 3, of which

Poingalaceae - Spice, flower - 1 lateral sepals - Pitcairnia - retardance -
Mon - divided

1 is anterior, and larger than the rest, and 2 are alternate with the upper and lateral sepals; sometimes there are 5 petals, 2 of them very minute; the anterior petal, called the keel, is often crested. Stamens hypogynous, 8, monadelphous or diadelphous; anthers clavate, usually 1-celled, and having porous dehiscence. Ovary mostly bilocular; ovules solitary, rarely 2; style simple, curved; stigma simple. Fruit dehiscing in a loculicidal manner, or indehiscent. Seeds pendulous, anatropal, strophiolate at the hilum; albumen fleshy; embryo straight.—Shrubs or herbs with alternate or opposite exstipulate leaves. They are found in all quarters of the globe. Lindley mentions 19 genera, including 495 species. *Examples*—*Polygala*, *Securidaca*, *Krameria*.

773. In the appearance of their flowers, the plants of this order have a resemblance to *Papilionaceæ*. They are distinguished, however, by the odd petal being inferior, and the sepal superior. They are generally bitter, and their roots yield a milky juice. *Polygala Senega*, Snake-root, is a North American species, the root of which is used medicinally, in large doses, as emetic and cathartic; and in small doses as a stimulant, sudorific, expectorant, and sialagogue. It contains an acrid principle, called polygalin, senegin, and polygalic acid. The name of Snake-root was given from its supposed use as an antidote to the bite of the rattle-snake. *Krameria triandra*, a Peruvian plant, furnishes Rhatany-root, which is employed as a powerful and pure astringent in cases of hæmorrhage, and chronic mucous discharges. Its infusion is of a blood-red colour, and has been employed to adulterate port wine.

774. Order 21.—*Tremandraceæ*, the Porewort family. (*Polypet. Hypog.*) Sepals 4-5, slightly coherent, deciduous, with a valvate æstivation. Petals 4-5, deciduous, with an involute æstivation. Stamens hypogynous, distinct, 8-10, 2 before each petal; anther di- or tetralthecal, with porous dehiscence (fig. 324). Ovary bilocular, with 1-3 pendulous ovules in each cell; style, 1; stigmas 1-2. Fruit a 2-celled, 2-valved capsule, with loculicidal dehiscence. Seeds anatropal, pendulous, with a caruncula at the apex; embryo cylindrical, straight, in the axis of fleshy albumen.—Heath-like shrubs, with hairs usually glandular, alternate or verticillate exstipulate leaves, and solitary axillary 1-flowered pedicels. They are natives of New Holland. Nothing is known regarding their properties. Lindley mentions 3 genera, including 16 species. *Examples*—*Tetralthea*, *Tremandra*.

775. Order 22.—*Tamaricaceæ*, the Tamarisk Family. (*Polypet. Hypog.*) Calyx 4-5 partite, persistent, with imbricated æstivation. Petals 4-5, hypogynous, or perhaps inserted at the base of the calyx, marcescent, with imbricated æstivation. Stamens hypogynous, free or monadelphous (fig. 311), equal to the petals in number, or twice as


many; anthers dithecal, introrse, with longitudinal dehiscence. Ovary unilocular; styles, 3. Fruit a 3-valved, 1-celled capsule, with loculicidal dehiscence. Seeds numerous, anatropal, erect or ascending, comose; albumen 0; embryo straight, with the radicle next the hilum. —Shrubs or herbs, with alternate scale-like leaves, and racemose or spiked flowers. They abound in the Mediterranean region, and are confined chiefly to the eastern half of the northern hemisphere. Many are found in the vicinity of the sea. They have a bitter astringent bark, and some of them yield a quantity of sulphate of soda when burned. The saccharine substance called Mount Sinai Manna, is yielded by *Tamarix mannifera*. Lindley mentions 3 genera, comprising 43 known species. *Examples*—*Tamarix*, *Myricaria*.

776. Order 23.—**Frankeniaceæ**, the Frankenia Family. (*Polypet. Hypog.*) Sepals 4-5, cohering into a tube, persistent. Petals 4-5, alternate with the sepals, hypogynous. Stamens hypogynous, equal in number to the petals, and alternate with them, sometimes more numerous; anthers bilocular, with longitudinal dehiscence. Ovary unilocular, with parietal placentas; style filiform, often trifid. Fruit a 1-celled, usually 3-valved capsule, with septicidal dehiscence. This latter distinguishes them from the Violet-worts to which they are allied. Seeds very minute, numerous, anatropal; embryo straight, in the axis of fleshy albumen.—Herbs or undershrubs, with opposite exstipulate leaves. They are found chiefly in the southern parts of Europe, and in the north of Africa. Two species are natives of the Cape of Good Hope, one of South America, four of New Holland, and three of temperate Asia. None have been found in tropical India, or North America. They are said to have mucilaginous and slightly aromatic properties. Genera, 4; species, 24. *Example*—Frankenia.

777. Order 24.—**Elatinaceæ**, the Water-pepper Family. (*Polypet. Hypog.*) Sepals 3-5, free, or slightly coherent at the base. Petals alternate with the sepals, hypogynous. Stamens hypogynous, equal to, or twice as many as, the petals. Ovary tri-quinquelocular; styles 3-5; stigmas, capitate. Fruit capsular, 3-5-celled, 3-5-valved, loculicidal; placenta central. Seeds 00, exalbuminous, anatropal; embryo cylindrical and slightly curved.—Annual marsh plants, with hollow creeping stems, and opposite stipulate leaves. They are found in all parts of the globe. Some of them have acidity, and hence the name Water-pepper. Genera 6, and species 22, according to Lindley. The *Elatines* are natives of Europe and Asia, *Bergias* of the Cape of Good Hope and the East Indies, *Merimea* of South America, and *Tetradiclis* of the Syrian region. *Examples*—*Elatine*, *Bergia*, *Merimea*, *Tetradiclis*.

778. Order 25.—**Caryophyllaceæ**, the Chickweed Family. (*Polypet. Hypog.*) Sepals 4-5 (fig. 571), free (fig. 269), or united in a tube (figs. 290 c, 553 c), persistent. Petals 4-5 (fig. 571), hypogynous,

unguiculate (fig. 281), often bifid or bipartite (fig. 283), occasionally 0. Stamens (fig. 572 *e*) usually double the number of the petals, or, if equal, usually alternate with them; filaments subulate, sometimes united; anthers innate, bilocular, dehiscence longitudinal. Ovary



single, often stalked or supported on a gynophore (fig. 553 *g*), composed of 2 to 5 carpels, which are usually united by their edges, but sometimes the edges are turned inwards, so as to form partial dissepiments; stigmas 2-5 (fig. 391 *s*), with papillæ on their inner surface (fig. 572 *s*). Capsule unilocular (figs. 391, 574, 2), or imperfectly bi-quinquelocular (fig. 573), 2-5 valved, opening either by valves, or more commonly by twice as many teeth as stigmas (figs. 444, 574); placenta in the axis of the fruit (figs. 391, 574, 2, *p*). Seeds usually 00, amphitropal with mealy albumen, and a peripheral embryo (fig. 575).—Herbs, sometimes suffruticose plants, with opposite, entire, exstipulate, sometimes connate leaves, and usually cymose inflorescence (figs. 249, 250.)

Fig. 571-575.—Illustrations of the natural order Caryophyllaceæ.

Fig. 571.—Diagram of the flower of *Alsine media*, common Chickweed, belonging to the natural order Caryophyllaceæ, suborder Alsineæ. The flower consists of five imbricate sepals, five alternate petals, five stamens, a unilocular ovary, with a free central placenta, and numerous ovules.

Fig. 572.—Section of the flower of *Dianthus Caryophyllus*, Carnation. *c*, Calyx; *p*, petals, cohering with the stamens at the base; *e*, stamens; *g*, gynophore or thecophore, i.e. the stalk supporting the ovary; *o*, ovary with central placenta and ovules; *s*, two stigmas, which are papillose all along their inner surface.

Fig. 573.—Horizontal section of the ovary in a very young state, showing the partitions, *c*, which divide the ovary into two cavities. These divisions ultimately disappear, leaving the placenta, *p*, bearing the ovules free in the centre.

Fig. 574.—Capsule of *Lychnis Githago* at the period of dehiscence, when the pericarp separates into five valves at the summit. 1. The capsule entire. 2. Capsule cut vertically, to show the seeds, *s*, grouped in a mass, on a free central placenta, *p*.

Fig. 575.—Seeds. 1. Entire seed. 2. Seed cut vertically. *t*, Spermoderm. *c*, Peripheral embryo, surrounding the mealy perisperm, *p*.

They inhabit chiefly temperate and cold regions. According to the calculation of Humboldt, Cloveworts constitute $\frac{1}{2}$ of the flowering plants of France, $\frac{1}{7}$ of Germany, $\frac{1}{7}$ of Lapland, and $\frac{1}{3}$ of North America. Those found within the tropics are usually met with on high elevations and mountainous tracts, many of them exclusively vegetate in regions of the lowest temperature. The order has been divided into three suborders:—1. Alsineæ, sepals distinct (fig. 269), and opposite the stamens, when the flowers are isostemonous. 2. Sileneæ, sepals cohering in a tube (fig. 290 c), opposite the stamens, when the flowers are isostemonous. 3. Mollugineæ, sepals distinct or nearly so, alternate with the stamens, when isostemonous. Lindley mentions 53 genera, and 1055 species. *Examples*—Alsine, Cerastium, Dianthus, Silene, Mollugo.

779. The plants of this order are usually insipid, but according to Malapert and Bonnet, some are poisonous. The poisonous quality is attributed by them to the principle called Saponine, which exists in many of the species of *Saponaria*, *Silene*, *Lychnis*, and *Dianthus*. To Saponine, also, is due the saponaceous or soap-like properties of the plants. *Arenaria peploides* has been used as a pickle. In Iceland it serves as an article of food. The greater part of the plants of the order are weeds, but some are showy garden flowers. To the latter may be referred all the varieties of *Dianthus Caryophyllus*, Clove-pink or Carnation, Picotees, Bizarres, and Flakes, numerous species of Pink, Campion, &c. The varieties of Carnation depend on the mode in which the coloured stripes or dots are arranged on the petals, and the entire or serrated appearance of their edges. The formation of the placenta in the Caryophyllaceæ has given rise to much discussion, some looking upon it as a marginal, others as an axile formation (§ 443).

780. Order 26.—**Vivianiaceæ**, the Viviania Family. (*Polypet. Hypog.*) Sepals 5, united. Petals 5, hypogynous, unguiculate, persistent, with twisted æstivation. Stamens 10, hypogynous; filaments free; anthers bilocular, opening longitudinally. Ovary free, 3-celled; stigmas 3. Capsule 3-celled, 3-valved, loculicidal; seeds, 2 in each cell, with a curved embryo lying among fleshy albumen.—Herbaceous or suffruticose plants, with opposite or verticillate exstipulate leaves. All the members of this order which have yet been discovered inhabit Chili and South Brazil, they have no properties of importance. Genera, 4; species, 15. *Examples*—Viviania, Cæsarea.


781. Order 27.—**Malvaceæ**, the Mallow Family. (*Polypet. Hypog.*) Sepals 5 (fig. 577), rarely 3 or 4, more or less cohering at the base (fig. 274 c), with a valvate æstivation (fig. 263), often bearing an external calyx (*epicalyx*) or involucre (fig. 274 b). Petals equal in number to the sepals; æstivation twisted (fig. 264). Stamens ∞ (fig. 578 a), hypogynous, all perfect; filaments monadelphous (fig. 578 t) or

Picotees
Valvate calyx - twisted
Bilobed calyx - twisted
Monadelphous Stamens - 1 cell
Polypet. Hypog.
Artificial Stipulate Leaves - 1 cell
Monadelphous Stamens - 1 cell
Polypet. Hypog.

polyadelphous (fig. 551); anthers monothecal (fig. 328), reniform (fig. 579), with transverse dehiscence. Ovary formed by the union of several carpels round a common axis (figs. 452, 580), either distinct



or cohering; styles as many as the carpels (fig. 578 s), united or free. Fruit capsular or baccate; carpels one or many-seeded, sometimes closely united, at other times separate or separable (figs. 383, 580); dehiscence loculicidal (fig. 449), or septicidal. Seeds amphitropal or semi-anatropal; albumen 0, or in very small quantity; embryo



Figs. 576-584.—Organs of fructification of *Malva sylvestris*, to illustrate the natural order Malvaceæ.

Fig. 576.—Flower viewed from above, with its five petals, monadelphous stamens, peduncle or flower-stalk, and two stipules, s.

Fig. 577.—Diagram of the flower, showing the different whorls or verticils; five valvate or imbricate sepals, five twisted petals, indefinite monadelphous stamens, and united carpels.

Fig. 578.—Vertical section of the flower. t, Calliculus, epicalyx, or involucre; c, calyx; p, petals; t, tube of monadelphous stamens, forming an arch above the ovary, o, and united at the base to the petals; a, anthers at the summit of the filaments, free; s, styles free at the summit, united below.

Fig. 579.—A reniform monothecal anther, dehiscing transversely, separated with the upper part of the filament.

Fig. 580.—Fruit, surrounded by the persistent calyx, c, consisting of whorled carpels united together by the axis, a.

Fig. 581.—A separate carpel viewed laterally.

Fig. 582.—Exalbuminous amphitropal seed.

Fig. 583.—Curved embryo.

Fig. 584.—Section of the embryo, to show the doubled cotyledons.

curved (fig. 583); cotyledons twisted or doubled (fig. 584).—Herbaceous plants, trees or shrubs, with alternate stipulate leaves (fig. 576 s), more or less divided, and often with stellate hairs (fig. 82). They are found chiefly in tropical countries, and in the warm parts of the temperate zone. This family is closely related to the Sterculiads, including which they have been calculated to form $\frac{1}{80}$ of the flowering plants in Sicily, $\frac{1}{43}$ in France, $\frac{2}{33}$ in Sweden. In Lapland they are unknown. They form $\frac{1}{23}$ of those plants met with in temperate parts of North America, $\frac{1}{47}$ in the equinoctial parts of the same continent; taking into account the vegetation of the valleys, they, according to Humboldt, form $\frac{1}{50}$ of the flowering plants in the tropics, $\frac{2}{100}$ in the temperate zone, and are not found in the frigid zone. Lindley enumerates 37 genera, including 1,000 species. The order has been divided into three tribes:—1. Malvæ, an involucre present; fruit apocarpous or separable. 2. Hibiscæ, an involucre; fruit syncarpous. 3. Sidææ, no involucre; fruit either apocarpous or syncarpous. *Examples*—Lavatera, Malva, Hibiscus, Sida.

782. The plants of the order are all wholesome, and yield mucilage in large quantity. Some furnish materials for cordage, others supply cotton. *Malva sylvestris*, Common Mallow, and *Althæa officinalis*, Marsh Mallow, are employed medicinally, as demulcents and emollients. The latter is the Guimauve of the French. The flowers of *Althæa rosea*, the Hollyhock, are officinal in Greece for similar purposes; the plant also yields fibres and a blue dye. The petals in *Malva Alcea*, and in *Hibiscus Rosa sinensis*, possess astringent properties, of which the Chinese make use to blacken their eyebrows and the leather of their shoes. The flowers of *Abutilon esculentum*, and the fruit of *Abelmoschus esculentus*, called Ochro, are used as articles of food. *Hibiscus cannabinus* is the source whence sun-hemp is procured in India. Other species of *Hibiscus* also yield useful fibres. The bark of *Paritium tiliaceum* furnishes valuable materials for cordage. Cotton is composed of the hairs surrounding the seeds of various species of *Gossypium*. These hairs, when dry, exhibit under the microscope a peculiar twisted appearance. *Gossypium barbadense* seems to be the species which yields the best cotton; the Sea-Island, New Orleans, and Georgian cotton being produced by varieties of it. *Gossypium peruvianum* or *acuminatum*, furnishes the South American cotton; *Gossypium herbaceum*, the common cotton of India. *G. arboreum* is the Indian-tree cotton. The Chinese Nankin cotton is furnished by a variety of *G. herbaceum*. The quality of cotton-wool depends on the length, strength, and firmness of the tissue, or, as it is called, the *staple*. These essential attributes are modified by the cleanliness and the colour. Long-stapled cottons are generally used for the twist or warp, and short-stapled for the weft. The value of cotton varies not only according to the species,

but also according to the nature of the climate in which it grows. The total import of Cotton into Britain in 1845, was upwards of 716 millions of pounds; while in 1846 and 1847, it amounted only to about 480 millions. The seeds of the cotton-plants yield oil which has been used for lamps.

783. Order 28.—*Sterculiaceæ*, the Sterculia and Silk-cotton Family. (*Polypet. Hypog.*) Calyx of 5, more or less united, sepals, often surrounded by an involucre; æstivation usually valvate. Petals 5 or none, hypogynous, æstivation twisted. Stamens usually ∞ ; their filaments variously united; anthers 2-celled, extrorse. Pistil of 5 (rarely 3) carpels, either distinct or cohering; styles equal in number to the carpels, free or cohering; ovules orthotropal (fig. 523) or anatropal. Fruit capsular, usually with 5 cells, or follicular or succulent. Seeds often with a woolly covering; with a fleshy or oily perisperm (rarely 0), and either a straight or a curved embryo; cotyledons leafy or thick, plaited or rolled round the plumule.—Trees or shrubs, with alternate leaves, which are either simple or compound, deciduous stipules, and often a stellate pubescence. They are distinguished from Malvaceæ by their dithecal extrorse anthers. They inhabit warm climates. The order has been divided into the following sections:—1. Bumbaceæ, with hermaphrodite flowers and palmate or digitate leaves, are found most abundant in America. 2. Helicterææ, with hermaphrodite flowers and simple leaves, seem to be unknown in Africa. 3. Sterculiææ, with unisexual flowers, and either simple or palmate leaves, preferring India and Africa. Lindley mentions 34 genera, including 125 species. *Examples*—*Bombax*, *Helicteres*, *Sterculia*.

784. The plants are mucilaginous and demulcent; many are used for food, others supply a material like cotton. The silky hairs surrounding the seeds of *Bombax Ceiba*, the Silk-cotton tree, are used for stuffing cushions and chairs, and for various other domestic purposes. It cannot be manufactured, in consequence of no adhesion existing between the hairs. The trunk of the tree is made into canoes. *Adansonia digitata*, the Baobab tree of Senegal, or monkey-bread, is one of the largest known trees. Its trunk sometimes attains a diameter of thirty feet, while its height is by no means in proportion. The pulp of its fruit (amphisarca) is used as an article of food. It is emollient and mucilaginous in all its parts. The dried leaves when powdered constitute salo, a favourite article with the Africans, which they mix with their food for the purpose of diminishing the excessive perspiration to which in those climates they are subject. It is found by Europeans to be most serviceable in cases of diarrhœa, fevers, and other maladies. *Durio zibethinus* furnishes the fruit called Durian in the Indian Archipelago. The fruit is much prized, although it has a fœtid odour, which has given rise to the name Civet Durian. *Cheiro-*

temon platanoides is called the hand-plant of Mexico, on account of its five peculiarly curved anthers, which resemble a claw. *Helicteres* (from *helix*, a snail) is so named on account of its twisted fruit. The Kola, mentioned by African travellers as being used to sweeten water, is the seed of a species of *Sterculia*.

785. Order 29.—**Byttneriaceæ**, the Byttneria and Chocolate Family. (*Polypet. Hypog.*) Calyx 4-5 lobed, valvate in æstivation (fig. 261 c). Petals 4-5 or 0, often elongated at the apex, with a twisted or induplicate æstivation (fig. 261 p). Stamens hypogynous, either equal in number to the petals, or some multiple of them, more or less monadelphous, some of them sterile; anthers bilocular, introrse. Ovary free, composed usually of 4-10 carpels arranged round a central column; styles terminal, as many as the carpels, free or united; ovules 2 in each loculament. Fruit capsular, either with loculicidal dehiscence, or the carpels separating from each other. Seeds anatropal, often winged; embryo straight or curved, lying usually in fleshy albumen; cotyledons either plaited or rolled up spirally.—Trees, shrubs or undershrubs, with alternate leaves, having either deciduous stipules or 0, and stellate or forked hairs. They abound in tropical climates. Lindley enumerates 45 genera, embracing 400 species. Byttneriads are often united with Sterculiads, from which they are distinguished by their slightly monodelphous stamens, and anthers turned inwards. Their two-celled anthers and not columnar stamens distinguish them from Mallow-worts. The order has been divided into six suborders founded on the following genera: *Examples*—*Lasiopetalum*, *Byttneria*, *Hermannia*, *Dombeya*, *Eriolæna*, and *Philippodendron*.

786. The plants abound in mucilage, and many yield cordage. The seeds of *Theobroma Cacao* are called Cacao-beans, and are the chief ingredient in chocolate, which contains also sugar, arnotto, vanilla, and cinnamon. The seeds by pressure yield a fatty oil, called Butter of Cacao. They contain a crystalline principle analogous to caffeine, called Theobromine. The *Cocoa* of the shops consists generally of the roasted beans, and sometimes of the roasted integuments of the beans, ground to powder.

787. Order 30.—**Tiliaceæ**, the Lime-tree Family. (*Polypet. Hypog.*) Sepals 4-5, with a valvate æstivation. Petals 4-5, entire, rarely wanting. Stamens hypogynous, free, or united by the enlarged border of the stalk of the pistil (fig. 316, 1, 2), usually ∞ ; anthers 2-celled, dehiscing longitudinally or by pores, occasionally some abortive (fig. 316, 2). Disk often large and glandular. Ovary solitary, formed by the union of 2-10 carpels; style 1; stigmas as many as the carpels. Fruit dry or pulpy, either multilocular with numerous seeds, or by abortion unilocular and 1-seeded. Seeds anatropal; embryo erect in the axis of fleshy albumen, with flat, leafy cotyledons (fig. 510).—Trees or shrubs, rarely herbaceous plants, with alternate stipulate leaves (fig. 195).

The principal part of the order is found within the tropics forming weed-like plants, or shrubs, or trees, with handsome, usually white or pink flowers. A small number are peculiar to the northern parts of either hemisphere, where they form timber trees. The order has been divided into two sections:—1. *Tiliæ*, with entire petals or 0, and anthers dehiscing longitudinally. 2. *Elæocarpeæ*, with lacerated petals, and anthers opening at the apex. Lindley enumerates 35 genera, including 350 species. *Examples*—*Tilia*, *Corchorus*, *Grewia*, *Aristolelia*, *Elæocarpus*.

788. The plants possess mucilaginous properties, and many of them furnish excellent materials for cordage. The fruit of some is edible. From the gummy matter they contain, some have been employed as demulcents. The inner bark, the bast or bass, of the Linden or Lime-tree (*Tilia Europæa*), is tough and fibrous, and from it are manufactured Russian mats. The bark of *Luhæa grandiflora* is used in Brazil for tanning leather. An infusion of the flowers is used on the continent as an antispasmodic and expectorant. A species of *Corchorus* is used in India for the manufacture of bags and a coarse kind of linen. The leaves of *Corchorus olitorius*, Jew's-mallow, are used in many countries as a culinary vegetable.

789. Order 31.—**Dipterocarpaceæ**, the Sumatra-Camphor Family. (*Polpet. Hypog.*) Calyx tubular, 5-lobed, unequal, naked, persistent, and afterwards enlarged, with an imbricated æstivation. Petals hypogynous, sessile, often combined at the base, with a twisted æstivation. Stamens indefinite, hypogynous; filaments dilated at the base, either distinct or irregularly cohering; anthers innate, bilocular, subulate, opening by terminal fissures. Torus not enlarged in a disk-like manner. Ovary superior, 3-celled; ovules in pairs, pendulous; style and stigma simple. Fruit coriaceous, unilocular by abortion, 3-valved or indehiscent, surrounded by the calyx, which is prolonged in the form of long wing-like lobes. Seed solitary, exalbuminous; cotyledons often twisted and crumpled; radicle superior.—Trees with alternate leaves, having an involute vernation, and deciduous convolute stipules. They are found in India and especially in the eastern islands of the Indian Archipelago, where, according to Blume, they form the largest trees of the forest. There are about 8 known genera, including 48 species. *Examples*—*Dipterocarpus*, *Vateria*, *Dryobalanops*.

790. The trees belonging to this order are handsome and ornamental, and they abound in resinous juice. A kind of Camphor is procured in Sumatra from *Dryobalanops Camphora* or *aromatica*. It is secreted in crystalline masses naturally into cavities in the wood. It is less volatile than the common camphor of commerce. It supplies this camphor only after attaining a considerable age. In its young state it yields on incision a pale yellow liquid, called the liquid camphor of Borneo and Sumatra, which consists of resin and a volatile oil

having a camphoraceous odour. Indian copal, or Gum animi of commerce, is the inspissated varnish got from *Vateria Indica*. The fruit of this tree yields to boiling water the celebrated butter of Canara, or Pinei tallow. Various species of *Dipterocarpus* yield a substance like Balsam of Copaiva.

791. Order 32.—**Chlænaceæ**, the Chlænad Family. (*Polypet. Hygog.*) Involucre 1-2-flowered, persistent. Sepals 3, small. Petals 5-6, hypogynous, sometimes combined at the base where they are broader. Stamens 10, or indefinite; filaments cohering at the base, and united to the base of the petals; anthers roundish, free or united, bilocular. Ovary single, trilocular; style 1, filiform; stigma trifid. Capsule 3-celled, or by abortion 1-celled. Seeds solitary or numerous, suspended, attached to a central placenta; embryo in the axis of fleshy or horny albumen; cotyledons leafy, undulated.—Trees or shrubs, with alternate stipulate leaves, found in Madagascar. Their properties are unknown. There are four genera enumerated, including probably about 8 or 10 species. *Examples*—*Sarcolæna*, *Leptolæna*.

792. Order 33.—**Ternstræmiaceæ**, the Tea Family. (*Polypet. Hygog.*) Sepals 5 or 7, concave, coriaceous, deciduous, the innermost often the largest; æstivation imbricated (fig. 265 c). Petals 5, 6, or 9, often combined at the base (fig. 265 p). Stamens indefinite, hypogynous; filaments free, or united at the base into one or more parcels; anthers versatile, or adnate, dehiscing longitudinally. Ovary multilocular; styles 2-7. Fruit either a capsule, 2-7 celled, opening by valves, or coriaceous and indehiscent. Seeds attached to the axis, few and large; albumen 0, or in very small quantity; embryo straight or bent, or folded back; radicle next the hilum; cotyledons very large, often containing oil.—Trees or shrubs, with alternate coriaceous, exstipulate leaves, which are sometimes dotted. They abound in South America, and many occur in India, while others inhabit China and North America. There are 33 genera, and 130 species enumerated. *Examples*—*Ternstræmia*, *Gordonia*, *Camellia*, *Thea*.

793. The most important plants of this order are those which yield Tea. Considerable discussion has taken place regarding the Tea plants; some say that there is only one species; others, two; others, three. Fortune visited the black and green tea districts of Canton, Fokien, and Chekiang, and he says that the black and green teas of the northern districts of China are obtained from the same species or variety, viz. that cultivated in Britain under the name of *Thea viridis*; while the black and green teas from the neighbourhood of Canton are made from the species or variety cultivated in this country under the name of *Thea Bohea*. Some make the Assam plant a different species, and thus recognize three, *Thea Cantonensis* or *Bohea*, *Thea viridis*, and *Thea Assamica*. The quality of the tea depends much on the season when the leaves are picked, the mode in which it is prepared, as well as the district

in which it grows. Green Tea contains more essential oil and tannin than Black Tea. The Green Teas include Twankay, Young Hyson, Hyson, Gunpowder, and Imperial; while the black include Bohea, Congou, Souchong, Oolong, and Pekoe. The teas of certain districts, such as Ankoï, have peculiar characters. In some instances teas are dyed by means of *Isatis indigotica*; in other cases, by means of Prussian blue and gypsum. Perfume is communicated to teas by means of *Olea fragrans*, *Chloranthus inconspicuus*, and *Aglaia odorata*. There is a bitter principle in tea called *theine*, which may be procured by adding a slight excess of acetate of lead to a decoction of tea, filtering hot, evaporating, and subliming. According to Dr. Stenhouse,

- 1 lb. of Green Hyson Tea gave 72 grains pure white Theine, and 2 coloured
 = 74 grains or 1·05 p. c.
 8 oz. Black Congou gave 34·5 gr. pure, and 1·5 impure = 36 gr. or 1·02 p. c.
 6 oz. of Black Assam Tea yielded 36 gr. or 1·37 p. c.
 1 lb. of a cheap Green Tea, called Twankay, gave 69 gr. or 0·98 p. c.

In 1846, the Imports of Tea into the United Kingdom were—

Black, about.....	43,000,000 lbs.
Green, about.....	13,000,000 —
Total,.....	56,000,000 —

An interesting account of the Tea plants, and the manufacture of Tea, will be found in Fortune's "Travels in China," in Ball's "Account of the Cultivation and Manufacture of Tea," Royle's "Illustrations of Himalayan Botany," and his "Productive Resources of India." The genus *Camellia* is prized on account of its showy flowers. There are numerous cultivated varieties of *Camellia japonica*, many of which can endure the climate of Britain when trained on a wall with a southern exposure, or slightly protected. In China, *Camellia Sasanqua*, or Sasanqua tea, is cultivated on account of its flowers, which are said to impart fragrance and flavour to other teas. *Camellia oleifera* yields a valuable oil.

794. Order 34.—**Olacaceæ**, the Olax Family. (*Polypet. Hypog.*) Calyx small, gamosepalous, entire or toothed, often becoming finally large and fleshy; æstivation imbricated. Petals 3-6, hypogynous, free, or adhering in pairs by means of the stamens; æstivation valvate. Stamens hypogynous, some fertile, others sterile; the former 3-10, alternate with the petals, the latter opposite to the petals; filaments compressed; anthers innate, bilocular, with longitudinal dehiscence. Ovary 1-3-4-celled; ovules 1-3, pendulous from a central placenta; style filiform; stigma simple. Fruit fleshy, indehiscent, often surrounded by the enlarged calyx, unilocular, monospermal. Seed anatropal, pendulous; albumen copious, fleshy; embryo small, at the base of the albumen.—Trees or shrubs, with simple, alternate, exsti-

pulate leaves, which are, however, sometimes abortive. They are chiefly tropical or subtropical, chiefly found in the East Indies, New Holland, and Africa. One is only known in the West Indies. A few are from the Cape of Good Hope. Little is known in regard to their properties. *Olex zeylanica* has a fœtid wood with a saline taste and is employed in putrid fevers; its leaves are used in a salad. There are 24 genera, and 53 species enumerated. *Examples*—*Olex*, *Opilia*, *Icacina*.

795. Order 35.—**Aurantiaceæ**, the Orange Family. (*Polypet. Hypog.*) Calyx urceolate or campanulate, short, 3-5-toothed, withering. Petals 3-5, broad at the base, sometimes slightly coherent; aestivation imbricated. Stamens equal in number to, or a multiple of, the petals; filaments flattened at the base, distinct or combined into one or more parcels; anthers erect. Thalamus enlarged in the form of a hypogynous disk, to which the petals and stamens are attached. Ovary free, multilocular; style 1; stigma thickish, somewhat divided. Fruit a hesperidium, having a spongy separable rind, and pulpy separable cells (§ 552). Seeds anatropal, attached to the axis, solitary or several, usually pendulous, having the chalaza and raphe usually well marked; perisperm 0; embryo straight; cotyledons thick and fleshy.—Trees or shrubs, usually conspicuous for their beauty, with alternate, often compound leaves, which are articulated with a usually winged petiole (fig. 185). They abound in the East Indies. *Limonia laureola* is remarkable as the only plant of this family found on the tops of cold and lofty mountains, where it is for some months of the year covered with snow. There are 20 genera, and nearly 100 species enumerated. *Examples*—*Citrus*, *Limonia*, *Triphasia*.

796. The plants exhibit in every part receptacles of volatile oil. The oil abounds in the leaves and in the rind of the fruit. It is fragrant and bitter. The fruit has a more or less acid pulp, and the wood is generally compact. The Orange, Lemon, Lime, Citron, Shaddock, and Forbidden Fruit, belong to this order. *Citrus vulgaris*, yields the Bitter or Seville Orange, from the flowers of which an essential oil, called Neroli-oil, is procured, in the proportion of an ounce from 550 pounds of flowers. A similar oil is got from the flower of the Sweet Orange, *Citrus Aurantium*. The rind of the Bitter Orange is used in conserves. In the young state, the fruit is sold under the name of Orangettes or Curaçoa oranges. Orange-flower-water, as obtained from the flowers of the Bitter Orange, is employed as an anodyne. The chief kinds of Sweet Orange are the Common Orange, the Chinese or Mandarin Orange, the Maltese, and St. Michael's. The last are the finest imported into Britain, and are distinguished by their smooth, thin rind. A single tree, it is said, will produce 20,000 good oranges. Their fruit is used medicinally,

on account of the pulp, which contains sugar, mucilage, and citric acid. From the rind of the Sweet Orange, an oil, called Oil of Orange, is procured, which differs from Neroli-oil. A similar oil, but of inferior quality, is procured from the rind of the Seville Orange. Many look on the Bitter and Sweet Oranges as produced by varieties of one species. The Bitter Orange-tree is less than that yielding the Sweet Orange; the petioles are more distinctly foliaceous; the flowers have a sweeter fragrance; the rind of the fruit is darker and more bitter; and its pulp more bitter and less saccharine. The Lemon, Lime, and Citron, are distinguished from oranges by their oblong form, their adherent rind, and a protuberance at the apex. *Citrus Limonum* yields the Lemon, the juice of which is antiscorbutic, and is used for cooling drinks and effervescing draughts, while the peel or rind, on account of the oil it contains, is employed as an aromatic and anthelmintic. A single tree will produce 8,000 lemons. *Citrus medica* furnishes the Citron, which is larger than the Lemon, has a thicker and tuberculated rind, and a less acid pulp. The rind and juice may be applied to the same purposes as those of the Lemon. *Citrus Limetta* is the source of the Bergamot and Lime, which are probably merely varieties. The Bergamot is less than the Lemon in size, and is more pyriform, while its colour is golden. The Lime is about half the size of the Lemon; its rind is thin, dense, and of a greenish-yellow colour, and its taste is more bitter. Oil of Bergamot is the volatile oil of the rind, and 100 fruits are said to yield $2\frac{1}{2}$ ounces. The fruit of *Citrus decumana* is known by the name of the Shaddock, while that of *C. Pompelmos* is the Pomпельmoose fruit. What are called horned oranges and fingered citrons, are produced by a separation or multiplication of the carpels, so that small fruits appear to be enclosed within the large one. *Ægle Marmelos* yields an excellent fruit. From *Feronia elephantum*, a gum, like gum-arabic, is procured.

797. Order 36.—**Hypericaceæ**, the Tutsan or St. John's-wort Family. (*Polypet. Hypog.*) Sepals 4-5, separate or united, persistent, usually with glandular dots, unequal; æstivation imbricated. Petals 4-5, oblique, often with black dots; æstivation contorted. Stamens hypogynous, ∞ ; generally polyadelphous (fig. 315), very rarely 10, and monadelphous or distinct; filaments filiform; anthers bilocular, with longitudinal dehiscence; carpels 2-5, united round a central or basal placenta; styles the same number as the carpels, usually separate; stigmas capitate or simple. Fruit either fleshy or capsular, multilocular, and multivalvular, rarely unilocular. Seeds usually 00, minute, anatropal, usually exalbuminous; embryo usually straight.—Herbaceous plants, shrubs or trees, with exstipulate entire leaves, which are usually opposite and dotted. Flowers often yellow. They are distributed very generally over all parts of the globe, are found in elevated and low, dry and damp situations. They yield a resinous coloured

Hypericaceæ - Opposit. dotted leaves - Andrie. Calyx - twisted Petals - Only ad. 1 flower - 1 stamen -

juice which has purgative properties, and resembles gamboge. In the European species this yellow juice is in small proportion to the essential oil, and the rest of the vegetable matter. They have been used as tonics and astringents. *Hypericum hircinum* is fœtid. A gargle for sore throats is prepared in Brazil from the *Hypericum Connatum*. A decoction of the leaves of *Hypericum laxius acum*, or *Allecim Crabs*, is reputed in the same country to be a specific against the bite of serpents. Lindley places *Parnassia* in this order. There are 15 known genera, and about 270 species. *Examples*—*Hypericum*, *Elodea*, *Vismia*, *Reaumuria*.

798. Order 37.—**Guttiferae**, or **Clusiaceæ**, the Gamboge Family. (*Polypet. Hypog.*) Sepals 2-6 or 8, usually persistent, round, frequently unequal and coloured; æstivation imbricated. Petals hypogynous, equal to, or a multiple of, the sepals. Stamens hypogynous, usually 00, rarely definite, free or variously united at the base; filaments unequal in length; anthers adnate, introrse or extrorse, sometimes very small, occasionally unilocular, and sometimes with porous or circumscissile dehiscence. Thalamus forming a fleshy, sometimes 5-lobed disk. Ovary solitary, 1 or many-celled; ovules either solitary and erect or ascending, or numerous and attached to central placentas; style 0 or very short: stigmas peltate or radiate. Fruit dry or fleshy, 1 or many-celled, 1 or many-seeded, either with septicidal dehiscence or indehiscent. Seeds definite, anatropal, or orthotropal, in a pulp, apterous and often arillate, with a thin and membranous spermoderm; albumen 0; embryo straight; cotyledons usually cohering.—Trees or shrubs, sometimes parasitical, with exstipulate, opposite, coriaceous, entire leaves, having a strong midrib, and lateral veins running directly to the margin. Flowers articulated with the peduncle, often unisexual by abortion. They are natives of tropical regions, more especially of South America; a few are from Madagascar and the continent of Africa. They generally require situations combining excessive heat and humidity. Lindley enumerates 30 genera, including 150 species. *Examples*—*Clusia*, *Garcinia*, *Cambogia*, and *Calophyllum*.

799. The plants of this order yield a resinous juice, which is acrid, purgative, and has a yellow colour. Gamboge is one of the most important products. There are two kinds of gamboge, one called Siam, and the other Ceylon Gamboge. The latter is the product of *Cambogia gutta* or *Hebradendron gambogioides*, and it is probable that the other, which is the only sort in commerce, and which is known by the names of Pipe, Lump, and Coarse Gamboge, is derived from the same source. Another kind of gamboge, called Coorg or Wynaad Gamboge, has been recently described by Christison. Gamboge is a powerful irritant, and in large doses acts as a poison, causing inflammation of the mucous membrane. It is employed medicinally, as a drastic and hydragogue cathartic. It is an excellent pigment. The

resin of Tacamahaca is yielded by *Calophyllum Calaba*. An oil is obtained from the seeds of *Calophyllum inophyllum*. *Pentadesma butyracea* is the Butter and Tallow-tree of Sierra Leone, so called on account of the solid oil which is furnished by the fruit. While an acrid resin is the product of most of the plants of the order, there are some parts in which the resin is either absent or elaborated in small quantity. Thus some of them produce fruits which are used as articles of diet. *Garcinia Mangostana* supplies the East Indian Mangosteen, which is said to be one of the finest known fruits; it resembles a middle-sized Orange, and is filled with a sweet and highly-flavoured pulp. *Mammea americana* gives a drupaceous fruit, called Mammee Apple, or Wild Apricot of South America. Its seeds are anthelmintic; its flowers yield by distillation a stomachic spirit called Eau de Créole; and a wine is obtained by fermenting its sap. *Mesua ferea* yields a hard and durable timber. The *Clusias* are handsome trees, remarkable for the mode in which they send out adventitious roots. The fruit of *Clusia flava*, sometimes called Wild Mango, or Balsam-tree, yields a yellow juice like gamboge.

800. Order 38.—**Margraviaceæ**, the Marcgravia Family. (*Polypet. Hypog.*) Sepals 2-7, usually coriaceous and persistent; æstivation imbricated. Corolla hypogynous, of 5 petals, or gamopetalous, calyptriform, entire or torn at the point. Stamens usually ∞, very rarely 5, hypogynous; filaments dilated at the base; anthers long, erect, introrse. Ovary single, unilocular; style 1; stigma often capitate. Fruit coriaceous, indehiscent, or dehiscing by valves in a loculicidal manner, the placentas being parietal and forming spurious dissepiments. Seeds indefinite, minute, in a pulp, anatropal, exalbuminous; embryo straight.—Trees and shrubs, with alternate, simple, entire, coriaceous, and exstipulate leaves. Flowers furnished occasionally with bracts, which are folded and united so as to form ascidia. They occur chiefly in the warmer parts of America. Their properties are scarcely known. The stem, root, and leaves of Marcgravia Umbellata, are regarded in the West Indies as diuretic and antisiphylitic. There are 4 genera mentioned, and 26 species. *Examples*—Marcgravia, Norantea.

801. Order 39.—**Hippocrateæ**, the Hippocratea Family. (*Polypet. Hypog.*) Sepals 5, very small, united up to the middle, persistent, with an imbricated æstivation. Petals 5, with an imbricated æstivation. Stamens 3, monadelphous; the united filaments forming a tube or a disk-like cup round the ovary; anthers with transverse dehiscence. Ovary free, trilocular; style 1; stigmas 1-3. Fruit consisting either of 3 samaroid carpels, or fleshy and 1-3 celled. Seeds definite, about 4 in each cell, attached to a central placenta, exalbuminous, anatropal, with a straight embryo, and flat somewhat fleshy cotyledons.—Arborescent or climbing shrubs, with opposite, simple, somewhat coriaceous leaves, having small deciduous stipules. They are found principally

in South America; a few are natives of Africa and the East Indies. The fruit of some is eatable. The fruit of *Touteleia* (*Salacia*) *pyriformis*, a native of Sierra Leone, is about the size of a Bergamot Pear; its flavour is rich and sweet. The nuts of *Hippocratea Comosa* are oily and sweet; it is called in the French West Indian Islands, *Amandier du Bois*. Lindley mentions 6 genera, comprehending 86 species.

Examples—*Hippocratea*, *Salacia*.

802. Order 40. **Erythroxylaceæ**, the Erythroxylon Family. (*Polypet. Hypog.*) Sepals 5, united at the base, persistent; æstivation imbricated. Petals 5, hypogynous, broad and with a small scale at the base, slightly contorted in æstivation. Stamens 10, monadelphous; anthers erect, bilocular, with longitudinal dehiscence. Ovary 3-celled, two of which are sometimes abortive; styles 3, distinct or united; stigmas 3; ovule single, pendulous. Fruit a 1-seeded drupe. Seed angular, anatropal; embryo in the axis of firm albumen, rarely exalbuminous; cotyledons linear, flat, and leafy.—Shrubs or trees with alternate stipulate leaves. Flowers arising from numerous, imbricated, scale-like bracts. Found chiefly in the West Indies and South America. The plants of the order have tonic, purgative, and narcotic qualities. The leaves of *Erythroxylon Coca* are used by the miners of Peru as a stimulant like opium. Its leaves are chewed with a small mixture of finely powdered chalk. The wood of some is of a bright red colour, and yields a dye. There are 2 or 3 known genera, and about 80 species. *Examples*—*Erythroxylon*, *Sethia*.

803. Order 41.—**Malpighiaceæ**, the Malpighia Family. (*Polypet. Hypog.*) Sepals 5, slightly united, persistent, often glandular at the base; æstivation imbricated. Petals 5, unguiculate, with convolute æstivation. Stamens usually 10, often monadelphous; anthers roundish, with a projecting process from the connective (fig. 342). Ovary formed by 3 (rarely 2 or 4) carpels, more or less combined; ovules solitary, with a long pendulous cord; styles 3, distinct or united. Fruit dry or fleshy, sometimes winged (fig. 466). Seeds solitary, orthotropal, suspended, exalbuminous; embryo straight or curved in various ways; cotyledons foliaceous or thickish (fig. 506).—Trees or shrubs, sometimes climbing, with simple, opposite, or very rarely alternate, stipulate leaves, without dots. Hairs, when present, peltate. Flowers either perfect or unisexual. They are inhabitants of tropical countries chiefly, and a great number of them are found in South America. Lindley notices 42 genera, including 555 species. *Examples*—*Malpighia*, *Banisteria*, *Hiptage*, *Hiræa*, *Gaudichaudia*.

804. Some of the woody plants of this order exhibit an anomalous formation of the stem, from the absence of annular rings and medullary rays, and the peculiar mode in which the bark is produced. This is shown in figs. 105, 108, and 109. Many of the plants are astringent. Some have stinging hairs (fig. 84). The fruit of *Mal-*

pighia glabra is called Barbadoes Cherry, and is used as an article of dessert. *Nitraria* is a genus doubtfully referred to this order. *N. tridentata*, found in the desert of Soussa, near Tunis, is said by some to be the true Lotus-tree of the ancient Lotophagi.

805. Order 42.—**Aceraceæ**, the Maple Family. (*Polypet. Hypog.*) Calyx divided into 5, rarely into 4 or 9 parts, with an imbricated æstivation. Petals equal in number to the lobes of the calyx, with which they alternate, rarely wanting. Stamens generally 8, inserted on a hypogynous disk. Ovary free, 2-lobed, 2-celled: ovules in pairs; amphitropal, pendulous; style 1; stigmas 2. Fruit, a samara (fig. 465), composed of two winged carpels, each 1-celled with 1-2 seeds. Seeds erect, exalbuminous; embryo curved, with foliaceous cotyledons, and the radicle next the hilum.—Trees with opposite, simple, lobed or palmate, exstipulate leaves. Flowers often polygamous. They are confined chiefly to the temperate parts of Europe, Asia, and North America. They yield a saccharine sap, from which sugar is sometimes manufactured. It is said that their juices become acrid as the season advances. The bark is astringent, and yields the dyer reddish-brown and yellow colours. *Acer saccharinum* is the Sugar Maple of America. *Acer Pseudo-platanus*, the Sycamore or Great Maple (the Plane-tree of Scotland), acts well as a shelter in exposed places, as near the sea. Its sap is slightly saccharine. Its wood is used in machinery and for charcoal. The leaves are often covered with black spots, caused by the attack of a fungus, *Xyloma* or *Rytisma acerinum*. There are 3 known genera, and 60 species. *Examples*—Acer, Negundo, Dobinea.

806. Order 43.—**Sapindaceæ**, the Soapwort Family. (*Polypet. Hypog.*) Sepals 4-5, distinct or cohering at the base; æstivation imbricated. Petals 4-5, occasionally absent, hypogynous, sometimes naked, sometimes with a glandular or scaly appendage inside; æstivation imbricated. Stamens usually 8-10, sometimes 5-6-7, very rarely 20; filaments free, or combined just at the base; anthers introrse. Thalamus forming a fleshy or glandular disc, into which the stamens are often inserted. Ovary trilocular, rarely bi- or quadri-locular; ovules anatropal, definite; style either undivided or 2-3 cleft. Fruit either fleshy and indehiscent, or samaroid, or capsular, and 2-3 valved. Seeds solitary, often arillate, exalbuminous; embryo straight, curved, or spiral; cotyledons incumbent; radicle next the hilum.—Trees or shrubs, sometimes climbing herbaceous plants, with alternate, sometimes opposite, compound (fig. 176), rarely simple leaves, often marked with lines or pellucid dots. They are natives principally of South America and India. Africa contains many of them; they are wanting in the cold regions of the north. None are found wild in Europe. In this order are included the Hippocastaneæ or Horse-chestnuts, which are distinguished by their opposite leaves, and their two ovules in each cell, one ascending, the other suspended (fig. 428). Lindley gives 50

genera, including 380 species. *Examples*—*Sapindus*, *Paullinia*, *Nephelium*, *Æsculus*, *Pavia*, *Dodonæa*, *Meliosma*.

807. In this order are included many plants which yield edible fruits, and others which are poisonous. A saponaceous principle exists in certain species. The fruit of *Sapindus Saponaria*, under the name of Soap-berries, is used as a substitute for soap in the West Indies. The Longan and Litchi are excellent Chinese fruits, the produce of *Nephelium Longan* and *N. Litchi*. The kernel of the Longan powdered, is sometimes made into paper. *Blighia* or *Cupania sapida* yields the Akee fruit, the succulent arillus of which is used as food. Many of the *Paullinias* are poisonous. Martius suggests that the nature of this poison should be inquired into, and experiments made as to whether it may not be advantageously administered in hydrophobia and insanity. From the seeds of *Paullinia sorbilis*, however, the Guarana bread is prepared in Brazil. This Guarana contains a bitter principle, identical with Caffeine. The bark of *Æsculus Hippocastanum*, Horse-chestnut, has been recommended as a febrifuge, and its seeds have been used as a substitute for coffee. The fruit and leaves of *Æsculus ohiotensis*, the Buck-eye or American Horse-chestnut, are said to be poisonous. *Paullinia pinnata*, and some other Sapindaceæ of Brazil, exhibit anomalous exogenous stems (fig. 106). *Ophiocaryon paradoxum*, is the Snake-nut-tree of Demerara, and is so called on account of the embryo, resembling a coiled-up snake.

808. Order 44.—**Rhizobolaceæ**, the Souari-nut Family. (*Polypet. Hypog.*) Sepals 5, more or less combined; æstivation imbricated. Petals usually 5, unequal, thickish. Stamens indefinite, slightly monadelphous, arising from a hypogynous disk, in a double row, of which the inner is often abortive; anthers roundish, with longitudinal dehiscence. Ovary 4-5 celled; ovules solitary, semi-anatropal; styles as many as the cells of the ovary; stigmas simple. Fruit formed of several indehiscent, 1-celled, 1-seeded nuts, with a thick double endocarp. Seed reniform, exalbuminous, with the funiculus dilated into a spongy excrescence; embryo with a very large radicle, which constitutes nearly the whole of the kernel; cotyledons small, lying in a furrow of the radicle (fig. 503).—Trees with opposite, palmately compound, coriaceous, exstipulate leaves. They grow in the warm forests of South America. Some of them furnish oil, others yield edible nuts. Souari nuts are the produce of *Caryocar butyrosum* (*Pekea butyrosa*). Lindley notices 2 genera, and 8 species. *Examples*—*Caryocar*, *Anthodiscus*.

809. Order 45.—**Meliaceæ**, the Melia Family. (*Polypet. Hypog.*) Sepals 4-5, more or less united, with an imbricated æstivation. Petals 4-5, hypogynous, sometimes cohering at the base, with a valvate or imbricated æstivation. Stamens equal in number to the petals, or 2, 3, or 4 times as many; filaments combined in a long tube; anthers


sessile within the orifice of the tube. Disk often large and cup-shaped. Ovary single, multilocular, the cells often equal in number to the petals; ovules usually anatropal, 1-2 in each cell; style 1; stigmas distinct or united. Fruit baccate, drupaceous or capsular, multilocular, or by abortion unilocular, when valves are present opening by loculicidal dehiscence. Seeds not winged; albumen usually absent; embryo straight, with leafy cotyledons.—Trees or shrubs with alternate (occasionally opposite), exstipulate, simple, or pinnate leaves. They are chiefly found in the tropical parts of America and Asia. Under this order, some include Humiriaceæ, which are distinguished by a prolonged fleshy connective (fig. 341), albuminous seeds, and a slender embryo. Arnott includes Cedrelaceæ also under this order. There are about 40 known genera, and upwards of 160 species. *Examples*—*Melia*, *Trichilia*, *Humirium*, *Canella*.

810. The plants of this order possess bitter, tonic, and astringent qualities. *Melia Azedarachta* is used in India as a febrifuge, and its fruit yields an oil which is employed for domestic purposes, and as an antispasmodic. The root of *Melia Azedarach* is bitter and nauseous, and is used in North America as an anthelmintic. Oils are procured also from species of *Trichilia* and *Carapa* (fig. 507). A warm pleasant smelling oil is prepared from the fruit of *Trichilia Speciosa*, which in India is considered as a valuable external remedy in chronic rheumatism and paralytic affections. The bark of *Carapa guianensis* has reputation as an anthelmintic. The fruit called in the Indian Archipelago, *Langsat*, is the produce of a species of *Lansium*. A fragrant balsam, called *Balsam of Umiri*, is got from the trunk of *Humirium floribundum*. *Canella alba*, the plant which furnishes *Canella bark*, has been referred to this order, but great doubts exist as to the propriety of doing so. Lindley makes it provisionally a separate order, *Canellaceæ*, which he places near *Pittosporaceæ*. *Canella bark* is hot and aromatic, and is used as a spice, and medicinally as a stimulant tonic. It was formerly confounded with *Winter's bark*, and the plant was called by Linnæus, *Winterania Canella*, or spurious *Winter's bark*.

811. Order 46.—*Cedrelaceæ*, the Mahogany Family. (*Polypet. Hypog.*) Calyx 4-5-cleft, with imbricated æstivation. Petals 4-5, with imbricated æstivation. Stamens 8-10, united below into a tube, sometimes distinct, inserted into a hypogynous annular disk; anthers bilocular, acuminate, with longitudinal dehiscence. Ovary usually 4 or 5-celled; ovules anatropal, pendulous; style simple; stigma peltate. Fruit a capsule opening septifragally (figs. 450, 451). Seeds winged; albumen thin or 0; embryo straight, erect; cotyledons fleshy.—Trees with alternate, pinnate, exstipulate leaves. They are found in the tropical parts of America and Asia. Lindley enumerates 9 genera, including 25 species. *Examples*—*Cedrela*, *Swietenia*.

812. The plants of this order are bitter, and have an aromatic fragrance. *Swietenia Mahagoni* supplies the well-known mahogany wood. Its bark, as well as that of *Soymida febrifuga*, and of *Cedrela febrifuga*, are used for the cure of intermittents. *Chloroxylon Swietenia* produces satin wood, and also yields a kind of wood-oil.

813. Order 47.—**Ampelideæ** or **Vitaceæ**, the Vine Family. (*Polypet. Hypog.*) Calyx small, nearly entire (fig. 586 c). Petals 4-5, sometimes cohering above (fig. 586 p), inserted outside an annular hypogynous disk (figs. 586, 587 g); æstivation valvate. Stamens 4-5, opposite to the petals (figs. 586, 587 e), inserted on the disk; filaments free, or united at the base; anthers ovate, versatile (fig. 587). Ovary 2-6-celled; ovules erect, anatropal (fig. 588 o); style 1, very short; stigma simple (588 s). Fruit pulpy and globular, not united to the calyx (fig. 589), sometimes 1-celled by abortion. Seeds 1 to



4 or 5, erect (590), with an osseous spermoderm, horny albumen (figs. 591, 592 p), and an erect embryo (fig. 591 e).—Climbing shrubs, having the lower leaves opposite, the upper ones alternate (fig. 221). Flowers in racemes, which are often opposite the leaves; floral peduncles sometimes becoming cirrhose. They inhabit the

Figs. 585-592.—Organs of fructification of *Vitis vinifera*, to illustrate the natural order Vitaceæ or Ampelideæ.

Fig. 585.—Diagram of the flower, showing 5 sepals, 5 petals, 5 stamens opposite the petals on account of the non-development of one staminal row, a disk, and the ovary.

Fig. 586.—Flower showing the petals, p, detached at the base, and remaining united above in a calyptra-like manner. c, Calyx. g, Glands, forming a disk. e, Stamens, the filaments of which only are seen.

Fig. 587.—Flower after the petals have fallen. g, Glands of the disk. e, Stamens, with versatile anthers. p, Pistil.

Fig. 588.—Vertical section of the flower. c, Calyx. p, Petals. e, Filaments. o, Ovary, with 2 cells and their erect anatropal ovules. s, Stigma.

Fig. 589.—Globular pulpy fruit, uva, or grape, differing from a berry, in the calyx not forming part of the pericarp. It is by some called nuculanum.

Fig. 590.—The seed of the grape, with its osseous spermoderm enclosing a hard perisperm.

Fig. 591.—The seed cut vertically. t, The integument or spermoderm. p, Perisperm, or albumen, which is horny. e, Erect embryo, with lanceolate cotyledons.

Fig. 592.—Horizontal section of the seed of the grape, about the middle. t, Integument or spermoderm. p, Perisperm or albumen.

milder as well as the hotter parts of both hemispheres, and abound in the West Indies. There are 7 genera, and 260 species. *Examples*—*Vitis*, *Cissus*, *Leëa*.

814. The plants of this order have generally acid leaves, and their fruit, when ripe, is saccharine. The leaves, on account of their astringency, have been used in diarrhœa. In France the sap or tears of the vine are regarded as a popular remedy for chronic ophthalmia; they are, however, of little value. *Vitis vinifera*, the Grape Vine, belongs to this order. It is said to be a native of the shores of the Caspian, whence it was imported into Europe. The unripe fruit contains a harsh acid juice, called verjuice. It contains free citric, malic, and tartaric acids, along with bitartrate of potass. As grapes ripen, sugar, called Grape-sugar, is formed at the expense of the acids (§ 304). The vessels of the Vine are large (fig. 61), and the sap passes through them with great force and rapidity. When cut in spring the plant bleeds freely. The leaves of *Cissus cordata* and *C. setosa* are said to possess acrid properties, and to prove useful in bringing indolent tumours to suppuration. The berries of the latter are acrid. Both leaves and fruit of *Cistus tinctoria* abound in a green colouring matter, which on exposure to air and light becomes blue, and is highly esteemed as a dye for cotton fabrics.

815. Order 48.—**Geraniaceæ**, the Cranesbill Family. (*Polypet. Hypog.*) Sepals 5, persistent, more or less unequal (figs. 307, 347 cc), one sometimes spurred at the base; æstivation imbricated. Petals 5 (or by abortion 4), unguiculate, with contorted æstivation (figs. 307, 347, pp). Stamens monadelphous, hypogynous (figs. 307, 347 e), twice or thrice as many as the petals, some occasionally abortive. Ovary of 5 carpels, placed round an elongated axis (fig. 307 t); ovules pendulous, solitary; styles 5, cohering round the axis (fig. 307 o). Fruit formed of 5 one-seeded cocoons, terminated each by an indurated style, which curls upwards, carrying the coccus or pericarp with it (fig. 455). Seeds exalbuminous, solitary, with a curved folded embryo, and leafy, convolute, and plaited cotyledons (fig. 511.)—Herbs or shrubs with simple, stipulate leaves, which are either opposite, or alternate with peduncles opposite to them. They are distributed over various parts of the world. The species of *Pelargonium* abound at the Cape of Good Hope. Lindley mentions 4 genera, including, after separating hybrids, about 500 species. *Examples*—*Geranium*, *Pelargonium*.

816. The name Cranesbill is derived from the long beak-like prolongation of the axis, or what is called the carpophore (§ 437). The plants of this order are astringent and aromatic. The root of *Geranium maculatum*, or Alum root, is a most powerful astringent, and is said to contain more tannin than Kino. The tuberous or moniliform roots of some, such as *Pelargonium triste* (fig. 123), are eatable.

Geraniaceæ *Spurred Stamens* - *Opposite Leaves* *Monadelphous*
Stamens - *Roots* *Plataneous* *Fruit*

The species of *Pelargonium* are remarkable for the beauty of their flowers. By the art of the gardener, and by hybridization, many fine varieties of *Pelargonium* have been produced.

817. Order 49.—**Linaceæ**, the Flax Family. (*Polypet. Hypog.*) Sepals 3, 4, or 5, persistent, with an imbricated æstivation. Petals 3, 4, or 5, fugitive, unguiculate, hypogynous, with a twisted æstivation. Stamens equal to the petals and alternate with them (with intermediate teeth or abortive stamens), arising from a hypogynous annular disk; anthers ovate, erect. Ovary with as many cells and styles as sepals, seldom fewer; stigmas capitate; ovules anatropal, pendulous. Fruit a multilocular capsule, pointed generally with the indurated base of the styles; each loculament or cell more or less completely divided by a spurious dissepiment, arising from the dorsal suture, and opening by two valves at the apex. Seeds solitary in each spurious cell, compressed, pendulous; albumen usually in small quantity, sometimes 0; embryo straight; cotyledons flat; radicle next the hilum.—Annual and perennial plants, with exstipulate, simple, entire leaves, which are usually alternate. They are scattered over the globe, but are said to be most abundant in Europe, and in the north of Africa. By some authors the order is associated with Geraniaceæ, from which it differs in its unbeaked fruit and exstipulate leaves, as well as the absence of joints in the stem. There are 3 genera mentioned by Lindley, comprising 90 species. *Examples*—*Linum*, *Radiola*.

818. The plants yield mucilage and fibre. Flax, which consists of woody fibre (fig. 44), is procured from the inner bark of the stalk of *Linum usitatissimum*, by the process of steeping and stripping off the bark. Linen and cambric are prepared from it. The flax plant is supposed to have been originally a native of Egypt, and mummy cloth has been shown to be formed of linen. The integument of the seeds is mucilaginous, and an infusion of them in boiling water is used as a demulcent and diuretic. The cotyledons of the seeds are oleaginous, and by expression yield Linseed oil, which has the property of drying and hardening into an elastic varnish, on exposure to the air. It is used medicinally for burns mixed with lime water. After expressing the oil a cake remains, called oil-cake, which is used for fattening cattle. The powdered cake receives the name of Linseed meal, and is commonly used for poultices. Another species of *Linum*, called *L. catharticum*, has purgative properties, which seem to depend on the presence of an acrid bitter matter, called Linin. *Linum selaginoides* is considered in Peru bitter and aperient.

819. Order 50.—**Balsaminaceæ**, the Balsam Family. (*Polypet. Hypog.*) Sepals 5, irregular, deciduous, the two inner and upper connate, coloured, the lower (odd) sepal spurred (fig. 541); æstivation imbricated. Petals alternate with the sepals, usually 4, in conse-

Linaceæ. *Imbricated Calyx* - *Twisted Petals* - *Stems* *Cells* - *Seeds* - *the number of the Sepals*

quence of 1 being abortive, often more or less irregularly united; æstivation convolute. Stamens 5. Ovary 5-celled; ovules usually numerous; stigma sessile, more or less 5-lobed. Fruit a 5-celled capsule, opening septifragally, by 5 elastic valves. Seeds usually numerous; suspended, exalbuminous, with a straight embryo, and radicle next the hilum. Succulent herbaceous plants with watery juice, having simple, opposite, or alternate, exstipulate leaves, and axillary irregular flowers. They inhabit chiefly the East Indies, and are remarkable for the force with which the seed-vessels open when ripe. The valves give way on account of the exosmose which goes on in the cells, and they then curl up in a peculiar manner (§ 27, 606). They have usually showy flowers, but their properties are unimportant. Lindley mentions 3 genera, including 110 species. *Examples*—*Impatiens*, *Hydrocera*.

820. Order 51.—*Oxalidaceæ*, the Wood-sorrel Family. (*Polypet. Hypog.*) Sepals 5, equal, sometimes cohering slightly at the base, persistent, imbricate in æstivation. Petals 5, equal, unguiculate, hypogynous, with a twisted æstivation. Stamens 10, more or less monadelphous, in 2 rows; those opposite the petals being longer than those in the outer row; anthers erect, bilocular. Ovary usually quinquelocular; styles filiform, distinct; stigmas capitate or slightly bifid. Fruit capsular, membranous or fleshy, usually 5-celled, and when dehiscent 5-10 valved. Seeds few, anatropal, albuminous, attached to a central placenta, sometimes with a peculiar elastic integument; embryo straight, as long as the fleshy albumen, with a long radicle and leafy cotyledons.—Herbs, undershrubs, or trees, with alternate, rarely opposite compound (occasionally simple) leaves, which are generally without stipules. They are found in the hot as well as the temperate parts of the world, and are abundant in North America and at the Cape of Good Hope. The shrubby species are confined to the hotter parts of the world. In some cases phyllodia, or winged petioles, occupy the place of leaves. There are about 6 known genera, and upwards of 320 species. *Examples*—*Oxalis*, *Averrhoa*, *Hugonia*.

821. They are often acid in their properties. Some of them yield esculent roots. *Oxalis Acetosella*, common Wood-sorrel, receives its name from its acid taste. It contains binoxalate of potash, which is sometimes called the salt of sorrel, and at other times the essential salt of lemons. The plant has been used as a refrigerant and antiscorbutic. Its leaves are trifoliate, and some have considered it to be the true Shamrock, in consequence of being in flower about the period of the year when St. Patrick's day occurs. Some of the oxalises, as *O. sensitiva*, have sensitive leaves, and experiments have been made in regard to their closing and opening by Morren (§ 660). *Oxalis crenata*, *esculenta*, and *Deppei*, yield tubers, which have been used as a substi-

tute for potatoes. The acid fruit of *Averrhoa Bilimbi* and *Carambola* is used in the East Indies as food.

822. Order 52.—**Tropæolaceæ**, the Indian Cress Family. (*Polypet. Hypog.*) Sepals usually 5, the upper spurred (fig. 275); æstivation slightly imbricate. Petals often 5, hypogynous, more or less unequal, sometimes abortive (fig. 542); æstivation convolute. Stamens 8 or 10, seldom fewer, free, almost perigynous; anthers bilocular, innate. Ovary triquetrous, composed of 3-5 carpels, with a single style, and 3-5 acute stigmas; ovules solitary, often pendulous. Fruit indehiscent, usually composed of 3 pieces. Seeds exalbuminous, with a large embryo, which has thick, often united, cotyledons, and a radicle next the hilum.—Herbaceous trailing or twining plants, having a delicate texture, with alternate exstipulate leaves, and axillary, often gay, flowers. They are natives of the temperate parts of America, and are extensively cultivated on account of their showy yellow, orange, scarlet, and occasionally blue flowers. They have more or less pungency in their fruit, which is used as a cress. The unripe fruit of *Tropæolum majus*, common Indian Cress, has been pickled and used as capers. Their roots are sometimes eaten. Lindley includes *Limnanthæ* in this order, and enumerates 6 genera, including 44 species. *Example*—*Tropæolum*.

823. Order 53.—**Pittosporaceæ**, the Pittosporum Family. (*Polypet. Hypog.*) Sepals 4 or 5, deciduous, distinct or partially united; æstivation imbricated. Petals 4 or 5, sometimes slightly cohering, with imbricated æstivation. Stamens 5, distinct, alternate with the petals. Ovary single, 2-5 celled; style 1; stigmas 2-5, equal in number to the placentas. Fruit capsular or berried, with many-seeded cells, which are sometimes incomplete; dehiscence loculicidal. Seeds often enveloped in a glutinous or resinous pulp, anatropal, with a minute embryo lying in fleshy albumen; radicle long; cotyledons very short.—Trees or shrubs, with simple, alternate, exstipulate leaves and flowers occasionally polygamous. They are found chiefly in Australasia. Many of them are resinous, and, in some instances, the berries are eaten. Lindley mentions 12 genera, including 78 species. *Examples*—*Pittosporum*, *Billardiera*, *Sollya*.

824. Order 54.—**Brexia**, the Brexia Family. (*Polypet. Hypog.*) Calyx small, persistent, of 5 coherent sepals, with an imbricated æstivation. Petals 5, with twisted æstivation. Stamens 5, alternate with the petals, arising from a narrow cup or disk, which is to the style

principally in Madagascar. Lindley associates some perigynous genera with *Brexia*, and places the order near *Saxafragaceæ*. He enumerates 4 genera, including 6 species. *Example*—*Brexia*.

825. Order 55.—**Zygophyllaceæ**, the Guaiacum Family. (*Polypet. Hypog.*) Calyx 4-5 parted, with convolute æstivation. Petals alternate with the calycine segments, with imbricated æstivation. Stamens twice as many as the petals; filaments dilated at the base, usually arising from scales (fig. 313). Ovary simple, 4-5 celled; divisions occasionally formed by spurious dissepiments (figs. 438, 439). Ovules 2 or more in each cell, usually pendulous; style simple, 4-5 furrowed; stigma simple, or 4-5-lobed. Fruit capsular or rarely fleshy, with 4-5 angles or wings, 4-5-valved, either opening by loculicidal dehiscence, or indehiscent. Seeds few, usually with whitish albumen, sometimes exalbuminous; embryo green, with foliaceous cotyledons, and a superior radicle.—Herbs, shrubs, or trees, with opposite, stipulate, usually compound leaves, which are not dotted, and hermaphrodite flowers. They occur in various parts of the world, chiefly in warm extra-tropical regions, as in the south of Europe, America, Africa, and India. The order has been divided into two sections:—1. *Zygophyllæ*, having albuminous seeds. 2. *Tribulleæ*, having exalbuminous seeds. Lindley mentions 7 genera, comprising 100 species. *Examples*—*Zygophyllum*, *Guaiacum*, *Tribulus*.

826. Some of the plants abound in a stimulant resin, which pervades the wood and bark; others are bitter and acrid. The medicinal species are used as sudorifics. *Zygophyllum Fabago* is called the Bean-caper, on account of its flowers being used as a substitute for capers. The plant is said to act as a vermifuge. *Guaiacum officinale* is a beautiful West Indian tree, the wood of which, commonly called *lignum-vitæ*, is prized for its hardness. The alburnum is of a grayish-yellow colour, while the duramen is greenish-black. The fibres of the wood are remarkable for their direction, being cross-grained, in consequence of one layer crossing another diagonally. It yields a resinous matter known as the resin of Guaiac, or Gum-guaiac. This resin exudes spontaneously, or it may be procured by incisions, or by the application of heat. A solution of the resin in alcohol, when applied to the fresh cut surface of a potato, gives rise to a blue colour. Both the wood and the resin are used medicinally on account of their stimulant diaphoretic properties. In decoction and tincture they are administered in cutaneous and syphilitic diseases. *Guaiacum sanctum* from Mexico has similar properties, and is sometimes used medicinally on the continent.

827. Order 56.—**Rutaceæ**, the Rue Family. (*Polypet. Hypog.*) See figs. 532, 533. Calyx having 4-5 segments, with an imbricated æstivation. Petals alternate with the divisions of the calyx, distinct, or cohering below into a spurious gamopetalous corolla, rarely want-

ing; æstivation either contorted or valvate. Stamens equal in number to the petals, or twice or thrice as many (rarely fewer by abortion or non-development) (fig. 534), usually hypogynous, but in some instances perigynous. Between the stamens and ovary there is a more or less complete cup-shaped disk, which is either free or united to the calyx. Ovary sessile or supported on a gynophore (fig. 382), its carpels equal to the petals in number or fewer; ovules 2, rarely 4 or fewer in each carpel; styles adherent above (fig. 382); stigma simple or dilated. Fruit capsular, its parts either combined completely or partially; seeds solitary or in pairs, albuminous or exalbuminous; embryo with a superior radicle.—Trees or shrubs, with exstipulate, opposite, or alternate leaves, usually covered with pellucid resinous dots (figs. 87, 89), and hermaphrodite flowers. The order has been subdivided into two suborders:—1. Ruteæ, with albuminous seeds, and the fruit with sarcocarp and endocarp combined. 2. Diosmeæ, with exalbuminous seeds, and a 2-valved endocarp, which dehisces at the base, and when the fruit is ripe separates from a 2-valved sarcocarp. Ruteæ are found chiefly in the southern part of the temperate zone, as in the south of Europe, while Diosmeæ abound at the Cape of Good Hope, and in New Holland. Lindley mentions 48 genera, and 400 species. *Examples*—*Ruta*, *Dictamnus*, *Diosma*, *Barosma*, *Correa*, *Boronia*.

828. The plants are remarkable for their peculiar odour, which is very powerful and penetrating. Many have antispasmodic properties, while others are bitter, and act as febrifuges and tonics. The leaves and unripe fruit of *Ruta graveolens*, common or garden Rue, are used in medicine as stimulants, antispasmodics, anthelmintics, and emmenagogues. They emit when bruised a strong and peculiar oppressive odour, and have a bitter and acrid taste. By distillation with water, they yield a yellow acrid volatile oil, which is their active constituent. The leaves of various species of *Barosma*, especially *B. crenata* or *crenulata*, and *serratifolia*, are used in medicine under the name of Bucku or Buchu. They contain a yellowish volatile oil, having a powerful odour, and they have been used as stimulants and antispasmodics. They are prescribed in cases of irritation and catarrh of the bladder in the form of infusion and tincture. *Galipea officinalis*, a plant found in Columbian Guiana, supplies the Angustura bark, which is used as a tonic and febrifuge. It is probable that *Galipea Cusparia*, sometimes called *Bonplandia trifoliata*, also furnishes a variety of Angustura bark. On the continent, Angustura bark is sometimes adulterated with the poisonous bark of *Strychnos Nux vomica*. Some of the species of *Dictamnus*, such as *D. Fraxinella*, False Dittany, abound in volatile oil to such a degree that the atmosphere around them becomes inflammable in hot, dry, and calm weather. The *Correas* are remarkable for their gamopetalous corolla. The leaves of some of the species have been used for tea in New Holland.

829. Order 57.—**Xanthoxylaceæ** or **Zanthoxylaceæ**, the Xanthoxylon Family. (*Polypet Hypog.*) Flowers unisexual. Calyx in 3, 4, or 5 segments, with imbricated æstivation. Petals the same number, rarely 0, usually larger than the calyx; æstivation imbricated or convolute. Stamens as many or twice as many as the petals, not developed in the female flowers. Ovary consisting of as many carpels as there are petals (sometimes fewer), the carpels being either completely or partially united (fig. 380); ovules 2, rarely 4, in each carpel; styles more or less combined (fig. 380 s). Fruit baccate or membranous, sometimes of 2-5 cells, sometimes of several drupes, or 2-valved capsules, of which the fleshy sarcocarp is partly separable from the endocarp. Seeds solitary or in pairs, pendulous; embryo lying within fleshy albumen; radicle superior; cotyledons ovate, flat.—Trees or shrubs, with exstipulate, alternate, or opposite leaves, having pellucid dots. They exist chiefly in the tropical parts of America. Lindley enumerates 20 genera, including 110 species.

830. The plants yield a volatile oil, which is aromatic and pungent. Some are diaphoretic in their properties, others are febrifugal and tonic. The pungency of species of *Xanthoxylon* has caused them sometimes to be denominated peppers. *Xanthoxylon fraxineum*, or prickly ash, acts as a sialagogue. *X. caribæum* has a bitter and febrifugal bark. The bitter principle secreted by many of the plants of this order is called Xanthopirine, and is said to have been used with some success as a substitute for hops.

831. Order 58.—**Simarubaceæ**, the Quassia and Simaruba Family. (*Polypet. Hypog.*) Flowers usually hermaphrodite. Calyx in 4 or 5 divisions; æstivation imbricated. Petals 4 or 5, spreading or connivent into a kind of tube; æstivation twisted. Stamens twice as many as the petals; filaments arising from scales. Ovary 4-5-lobed, 4-5-celled, supported on a gynophore; ovules solitary; style simple; stigma 4-5-lobed. Fruit indehiscent, consisting of 4 or 5 drupes arranged round a common receptacle. Seeds anatropal, pendulous; embryo exalbuminous.—Trees or shrubs, with exstipulate, alternate, usually compound leaves without dots. They are found in the tropical parts of America, Asia, and Africa. Lindley gives 10 genera, and 35 species. *Examples*—Simaruba, Quassia, Picræna.

832. All the plants of the order are intensely bitter. Quassia wood was originally the product of *Quassia amara*, a tall shrub, never above 15 feet in height, inhabiting Surinam, Guiana, and Colombia. It is a very ornamental plant, and has remarkable pinnate leaves, with winged petioles. In their early state, the leaves seem to be simple, but in the progress of growth two or more contractions take place, at each of which two leaflets appear, the pairs being separated by a winged midrib,—a continuation of the petiole. This Surinam Quassia does not appear to be exported now, and it is not met with in English

trade. A thriving specimen was recently sent to the Edinburgh Botanic Garden by Professor Syme. The Quassia of the shops is the wood of *Simaruba* or *Picræna excelsa*, a very large forest tree, attaining a height of nearly 100 feet, growing in Jamaica and other West Indian islands, where it is called Bitter Ash, and Bitter Wood. Quassia is used medicinally, in the form of infusion and tincture as a tonic and anthelmintic. It acts as a narcotic poison on flies and other insects. Although prohibited by law, it is frequently employed by brewers as a substitute for hops. The bitterness of Quassia is said to be owing to a crystalline principle called Quassin. The bark of the root of *Simaruba amara* or *officinalis*, a tree found in Cayenne and in the West Indies, is used as a bitter tonic and astringent, more especially in the advanced stages of diarrhœa and dysentery. *Brucea antidysenterica* was at one time erroneously supposed to furnish false Angustura bark. It has properties similar to those of Quassia. Malombo bark is probably yielded by a species of Quassia.

833. Order 59.—**Ochnaceæ**, the Ochna Family. (*Polypet. Hypog.*) Sepals 5, persistent, imbricated in æstivation. Petals equal to, or twice as many as the sepals, deciduous, spreading, imbricated in æstivation. Stamens 5, opposite the sepals, or 10, or indefinite; filaments persistent, attached to a hypogynous disk; anthers bilocular, innate, opening by pores, or longitudinally. Carpels as many as the petals, seated on an enlarged gynobase (thecaphore); ovule erect or pendulous, styles united into one. Fruit gynobasic, consisting of several succulent, indehiscent, monospermous carpels. Seeds anatropal, usually exalbuminous; embryo straight; radicle short; cotyledons thick.—Undershrubs or trees, with alternate, simple, stipulate leaves, and pedicels articulated in the middle. They grow in tropical countries, and are remarkable for the large succulent prolongation of the receptacle to which the carpels are attached. They are generally bitter, and some of them are used as tonics. Lindley enumerates 6 genera, comprehending 82 species. *Examples*—Ochna, Gomphia, Castela.

834. Order 60.—**Coriariaceæ**, the Coriaria Family. (*Polypet. Hypog.*) Flowers unisexual. Calyx campanulate, 5-parted; æstivation imbricate. Petals alternate with the calycine segments, very small, fleshy, with a keel on the internal surface. Stamens 10 (fig. 537); filaments filiform, distinct; anthers dithecal, oblong. Ovary composed usually of 5 carpels, attached to a thickened receptacle or gynobase, 5-celled; ovules solitary, pendulous; style 0; stigmas 5, long and glandular. Fruit, consisting of 5, monospermous, indehiscent crustaceous carpels, enclosed by the enlarged petals. Seeds pendulous, anatropal, exalbuminous; embryo nearly straight; cotyledons fleshy; radicle short and blunt.—Shrubs with opposite square branches, opposite, simple, ribbed leaves, and scaly buds. They are found in small numbers in the south of Europe, South America, India, and New

Zealand. Some of them are poisonous. The leaves of *Coriaria myrtifolia* have been employed to adulterate Alexandrian Senna on the continent. The leaves are known from those of true Senna by being 3-ribbed. The leaves are used for dyeing black, and an infusion of them gives a dark-blue with sulphate of iron. Lindley mentions the genus *Coriaria* only as belonging to this family, the position of which he considers as doubtful. There are 8 species of the genus. *Example*—*Coriaria*.

Subclass II.—CALYCIFLORÆ.

835. In this division are included the Polypetalous orders of Jussieu, in which the stamens are not hypogynous, as well as some declinuous orders. A calyx and corolla are present, in other words, the plants are Dichlamydeous; the petals are distinct and the stamens are attached to the calyx,—being thus more or less Perigynous. This subclass, along with Thalamifloræ, comprises the Dialypetalæ of Endlicher. It has been already stated that De Candolle included in this division gamopetalous plants, in which the ovary is inferior.

836. Order 61.—**Stackhousiaceæ**, the Stackhousia Family. (*Polypetal. Perigyn.*) Calyx 5-cleft, equal, with an inflated tube. Petals 5, equal, inserted at the top of the tube of the calyx, claws of the petals united, limb narrow and stellate. Stamens 5, unequal, attached to the tube of the calyx. Ovary superior, 3-5-celled, cells partially distinct; ovules solitary, erect; styles 3-5, sometimes united at the base; stigmas simple. Fruit consisting of 3-5 indehiscent pieces, which are sometimes winged, and are attached to a central persistent column. Seeds anatropal; embryo long, erect, in the axis of fleshy albumen.—Shrubs with simple, entire, alternate, stipulate, leaves, found in New Holland, and not possessing any marked properties. Lindley notices 2 genera, and 10 species. *Example*—*Stackhousia*.

837. Order 62.—**Celastraceæ**, the Spindle-tree Family. (*Polypetal. Perigyn.*) Sepals 4-5, imbricated in æstivation. Petals 4-5, with a broad base, and an imbricated æstivation, rarely wanting. Stamens alternate with the petals; anthers erect. Disk large, flat, and expanded, surrounding the ovary to which it adheres. Ovary superior, 2-5-celled; ovules ascending, one or numerous, attached to the axis by a short funiculus. Fruit either a 2-5-celled capsule, with loculicidal dehiscence, or drupaceous. Seeds one or many in each cell, anatropal, usually ascending, and sometimes arillate (figs. 481, 482); albumen fleshy; embryo straight, with flat cotyledons and a short radicle.—Small trees or shrubs, with simple, alternate, rarely opposite leaves, and small deciduous stipules. They inhabit the warm parts of Europe,

North America, and Asia, and many are found at the Cape of Good Hope. The order contains 24 known genera, and 260 species. It has been divided into two tribes:—1. *Euonymææ*, with capsular fruit, 2. *Elæodendreæ*, with drupaceous fruit. *Examples*—*Celastrus*, *Euonymus*, *Elæodendron*.

838. The plants of the order have subacid properties, and the seeds of some yield a useful oil. That of *Celastrus nutans*, or *paniculatus*, is said in India to be of a stimulant nature, and to be used in medicine in the disease called *Berriberri*. Some of the species of *Celastrus*, as *C. venenatus*, are reckoned poisonous. The seeds of *Euonymus*, Spindle-tree, are surrounded by an aril, or rather arillode, which is considered as a prolongation from the exostome (figs. 481, 482). In some of the species the capsules are crimson, and with the bright scarlet arillodes, they present a very showy appearance when the fruit is ripe. The bark of *Euonymus tingens* furnishes a yellow dye. It is used to mark the tika on the forehead of the Hindoos. It is also considered useful in diseases of the eye. The young shoots of *Euonymus europæus*, when charred, are used to form a particular kind of drawing pencil; its fruit and inner bark are said to be purgative and emetic.

839. Order 63.—*Staphyleaceæ*, the Bladder-nut Family. (*Polypet. Perigyn.*) (Fig. 539.) Sepals 5, united at the base, coloured, imbricated in æstivation. Petals 5, alternate, with an imbricated æstivation. Stamens 5, alternate with the petals. Disk large and urceolate. Ovary 2-3-celled, superior; ovules usually ascending; styles 2-3, cohering at the base. Fruit membranous or fleshy, indehiscent or opening internally, often partly abortive. Seeds anatropal, roundish, truncate at the hilum, with a bony testa; albumen generally 0; embryo straight, with thick cotyledons and a small inferior radicle.—Shrubs, with opposite, pinnate leaves, having stipules and stipels (§ 161). By many authors they are included under the last order. The plants are irregularly scattered over the globe, and are found in Europe, America, and Asia. Some of them appear to be subacid, while others are bitter and astringent. The species of *Staphyllea* receive the name of bladder-nut, on account of their inflated bladder-like pericarp. They are cultivated as handsome shrubs. Three known genera are enumerated, and 14 species. *Example*—*Staphylea*.

840. Order 64.—*Rhamnaceæ*, the Buckthorn Family. (*Polypet. Perigyn.*) Calyx 4-5-cleft, valvate in æstivation. Petals distinct, hooded or convolute, inserted into the throat of the calyx, sometimes 0. Stamens definite, opposite the petals. Disk large, fleshy, flat or urceolate. Ovary superior or half superior, 2-3 or 4-celled; ovules solitary, erect, anatropal. Fruit fleshy and indehiscent, or dry and separating into three parts. Seeds erect; albumen fleshy, rarely 0; embryo about as long as the seed, with a short inferior radicle, and large flat cotyledons.—Trees or shrubs, often spiny, with simple, alter-

nate, rarely opposite leaves, and minute stipules. They are generally distributed over the globe, and are found both in temperate and tropical regions. There are 42 genera, and 250 species enumerated. *Examples*—*Rhamnus*, *Ceanothus*, *Phytica*, *Pomaderris*.

841. Many of the plants of the order have active cathartic properties. Some, however, yield edible fruit, and others are tonic and febrifugal. *Rhamnus catharticus*, common or purging Buckthorn, is a European shrub, the black succulent fruits or berries of which are used as a hydragogue cathartic in cases of dropsy. The greenish juice becomes gradually red by the formation of acetic acid in it. It may be preserved unchanged in the form of syrup. When mixed with lime and evaporated to dryness, it forms the colour called sap-green. The fruit of *Rhamnus Frangula*, Black Alder, is emetic and purgative. The berries of *Rhamnus infectorius*, as well as those of other species, are known by the name of French berries. They have been used for dyeing yellow. The fruit of many species of *Zizyphus* is used for food; *Zizyphus Jujuba* supplies the fruit called Jujube; and the Lotus, or Lote-bush of the classics, whence the Lotophagi were named, is *Zizyphus Lotus*. A kind of Scinde lac is found on *Zizyphus Jujuba*. *Paliurus aculeatus*, Christ's-thorn, is common in the hedges of Judæa. *Ceanothus Americanus* is used in America as an astringent, and its leaves, under the name of New Jersey Tea, have been used as a substitute for tea. The leaves of *Seegeretia theezans* are used for the same purpose by the poorer classes in China.

842. Order 65.—**Anacardiaceæ**, the Cashew-nut Family. (*Polypet. Perigyn.*) Flowers usually unisexual. Calyx usually small and persistent, with 5, or sometimes 3-4-7 divisions. Petals equal in number to the calycine divisions, perigynous, sometimes 0; imbricated in æstivation. Stamens either equal to the petals in number and alternate with them, or twice as many or more; filaments distinct or cohering at the base, usually perigynous. Disk fleshy, annular or cup-shaped, sometimes inconspicuous. Ovary single, rarely 5 or 6, free or adhering to the calyx, 1-celled; ovule solitary, attached by a funiculus to the bottom or along the side of the cell; styles 1-3, occasionally 4; stigmas 1-3 or 4. Fruit usually drupaceous and indehiscent. Seed ascending or frequently pendulous, from the adherence of the funiculus to the angle of the cell, exalbuminous; radicle inferior or superior, sometimes curved suddenly back; cotyledons thick, fleshy, or leafy.—Trees or shrubs, with a resinous, often caustic juice, and alternate leaves without dots. The order is a subdivision of the Terebinthaceæ of Jussieu. The plants inhabit chiefly the tropical parts of America, Africa, and India; some, however, are found in Europe. The order is unknown in New Holland. There are 41 known genera, and 95 species. *Examples*—*Anacardium*, *Rhus*, *Mangifera*, *Spondias*.

"Found about the size of a Plum" Also the text

843. The order is characterized by the presence of an acrid resinous juice. In some cases, however, the fruit of the plant is edible. Many of them supply varnishes. *Anacardium occidentale* furnishes the Cashew-nut, which is remarkable for its large succulent peduncle supporting the fruit or nut (fig. 227). The pericarp has the acrid properties which pervade the order, while the seed is eatable. The fleshy peduncle is acid and edible, and a bland gum exudes from the bark. *Pistacia vera* is the Pistacia or Pistachio nut-tree, which extends from Syria to Bokhara and Caubul, and is cultivated in the south of Europe. It has green-coloured oily kernels, which are used as articles of diet. *P. Terebinthus* is a native of the southern part of Europe, and the northern part of Africa, and yields a liquid resinous exudation, known as Chian turpentine. The turpentine receives its name on account of being collected in the island of Chio or Scio, where the plant thrives. The tree attains a height of 30 or 35 feet, and one tree will yield ten ounces of the liquid resinous matter, which thickens on exposure to air, by the loss of volatile oil. Like other turpentine, it has diuretic and excitant properties. *Pistacia Lentiscus*, a native of the coasts and islands of the Mediterranean, furnishes the concrete resinous exudation called Mastiche or Mastic. It is a bush of about 10 or 12 feet in height, which is cultivated abundantly in the island of Chios. Mastic is used as a masticatory for consolidating the gums and cleansing the teeth. It has also been employed as an antispasmodic, and it enters into the composition of varnishes. *Rhus Toxicodendron*, Poison-oak, is a shrub found in Canada and the United States, the leaves of which have been used as stimulants in cases of palsy. Like the other species of this genus, it yields an acrid milky juice, which becomes black on exposure to the air. *Rhus radicans*, Poison-ivy, or Poison-vine, is probably another name of the same species. *Rhus venenata*, Poison-sumach, or Poison-elder, has acrid poisonous properties, and contact with it, in some instances, gives rise to inflammation of the skin. Cases are related of persons who are peculiarly liable to be thus affected, and in whom the irritation caused by the juice of the poisonous species of *Rhus* is very great, and even alarming. *Rhus coriaria*, *R. typhina*, and *R. glabra* are used for tanning, and their fruit is acid. *Rhus Cotinus* is called *Arbre à perruque* in France, on account of the hairy appearance presented by its abortive pedicels. Many of the plants in this order furnish varnishes and marking ink. *Semecarpus Anacardium*, commonly called the Marking-nut tree, supplies the Sylhet varnish, while *Melanorrhæa usitatissima* furnishes that of Martaban. *Stagmaria verniciiflua* is the source of the hard black varnish called Japan Lacquer. The leaves of many of the species of *Schinus*, as *S. Molle*, when broken and thrown on the surface of water, send out a resinous matter with great force, so as to cause a sort of spontaneous motion by the recoil.

Although a resinous principle pervades the plants of this order, yet in some cases it is not developed in the fruit, which becomes catable. Of this an illustration is furnished by the Mango, the produce of *Mangifera indica*. The Hog-plums of the West Indies are furnished by various species of *Spondias*, as *S. purpurea* and *Mombin*.

844. Order 66.—**Amyridaceæ**, the Amyris Family. (*Polypet. Perigyn.*) Flowers usually bisexual, sometimes unisexual by abortion. Calyx persistent, regular or nearly so, with 2 to 5 divisions. Petals 3-5, inserted at the base of the calyx; æstivation valvate or imbricated. Stamens twice or four times as many as the petals, perigynous. Disk covering the base of the calyx often in a ring-like manner. Ovary superior, sessile, 1-5 celled; ovules in pairs, anatropal, pendulous or suspended; style 1 or none; stigma simple or lobed, sometimes capitate. Fruit dry, 1-5-celled, indehiscent, or its epicarp splitting into valves. Seeds solitary, exalbuminous, with a superior radicle next the hilum, and cotyledons, which are fleshy or wrinkled.—Trees or shrubs, abounding in resin, with opposite or alternate compound leaves, which are frequently stipulate and dotted. They are natives of tropical regions. There are two suborders:—1. Amyrideæ, with an unilocular ovary. 2. Bursereæ, with a 2-5-celled ovary. Some look upon the stamens of Amyrideæ as truly hypogynous, and consider the order as allied to Aurantiaceæ. Lindley gives 22 genera, and 45 species. *Examples*—Amyris, Boswellia, Bursera, Balsamodendron.

845. The plants yield a fragrant balsamic and resinous juice, which, in a dry state, is often used as frankincense, and is employed medicinally as a stimulant or expectorant. The resinous substance called Elemi seems to be obtained from one of the plants of this order. Linnæus referred it to *Amyris elemifera*, under which name several species seem to be included. Some authors think that *Icica Icicariba* furnishes elemi in Brazil, while Royle refers a Mexican kind to *Elaphrium elemiferum*; and other varieties, according to Christison, are probably the produce of *Canarium commune* and *balsamiferum*. The resin contains a stimulant volatile oil. *Boswellia serrata*, a large Indian tree, supplies the gum-resin called olibanum, or the true frankincense of the ancients, the לִבְנָן of the Scriptures. It contains a volatile oil, and has been used as a stimulant, and as a material for fumigation. *Balsamodendron* (*Protium*?) *Myrrha*, a shrub growing in Abyssinia, appears to be the source of the officinal myrrh, the מוֹר of the Bible (מֹר). It is a bitter aromatic gum-resin, containing volatile oil, and was used in ancient times as frankincense. It is a heating stimulant, and is employed medicinally as an emmenagogue and diaphoretic, as well as for arresting various mucous discharges. The resin called Bdellium is procured from various species of *Balsamodendron*, as *B. africanum* and *Roxburghii*. The celebrated balsam called Balm of Gilead, is an exudation from *Balsamodendron gileadense*. Tecamahac

is procured from *Elaphrium tomentosum*. Various other balsams and resins are yielded by plants of this order. *Amyris toxifera* is said to be poisonous.

846. Order 67.—**Connarace**, the Connarus Family. (*Polypet. Perigyn.*) Flowers bisexual, rarely unisexual. Calyx 5-partite, regular, persistent; æstivation imbricate or valvate. Petals 5, inserted at the base of the calyx. Stamens twice as many as the petals, inserted with them, and doubtfully hypogynous; filaments united at the base. Ovary consisting of one or more separate carpels, each having a terminal style, and a dilated stigma; ovules in pairs, collateral, ascending, orthotropical. Fruit follicular, dehiscing along the ventral suture. Seeds solitary or in pairs, erect, with or without albumen, sometimes arillate; embryo with a superior radicle, remote from the hilum, and cotyledons, which are either fleshy or leafy.—Trees or shrubs, with compound, alternate, exstipulate leaves, which are not dotted. They are tropical plants. According to Endlicher they are common in America; some of them have febrifuge properties. *Omphalobium Lamberti* is said to furnish Zebra-wood. This order, as well as the orders Anacardiaceæ and Amyridaceæ, are by many considered truly hypogynous, and as belonging to Thalamifloræ. Lindley includes them in his Rutal alliance. He notices 5 genera, and 41 species. *Examples*—Connarus, Omphalobium, Cnestis.

847. Order 68.—**Leguminosæ** (Fabaceæ of Lindley), the Pea and Bean Tribe. (*Polypet Perigyn.*) Calyx 5-partite, toothed, or cleft (figs. 593, 594 c c), with the odd segment anterior (¶ 357); segments often unequal and variously combined. Petals 5 (figs. 593, 594), or by abortion, 4, 3, 2, 1, or 0, inserted into the base of the calyx, sometimes equal, but usually unequal, often papilionaceous, with the odd petal superior (fig. 594 e). Stamens definite or indefinite, usually perigynous, distinct, or monadelphous or diadelphous (fig. 594 t), or rarely triadelphous; anthers bilocular, versatile. Ovary superior, 1-celled, consisting usually of a solitary carpel (fig. 594 o), sometimes of 2-5; ovules 1 or many; style simple, proceeding from the upper or ventral suture; stigma simple (fig. 594 s). Fruit a legume (figs. 440, 469, 595), or a drupe. Seeds solitary or several (fig. 595), sometimes arillate, often curved (fig. 596); embryo usually exalbuminous, straight, or with the radicle bent upon the edges of the cotyledons (figs. 429, 516), which are either epigeal or hypogeal (¶ 629) in germination (fig. 597), and leafy (Phyllolobæ) or fleshy (Sarclobæ).—Herbaceous plants, shrubs, or trees, with alternate, usually compound leaves, having two stipules at the base of the petiole (fig. 193), and two at the base of each leaflet in the pinnate leaves. Pedicels usually articulated. The flowers are frequently papilionaceous (fig. 292), and the fruit is commonly leguminous (figs. 460, 469, 470, 471), and by the presence of one or other of these characters the order may be recognized. It is remarkable that one or other of these dis-

Fruit. Legum. Seed. Sepal. Petal. Add. Sepal. Petal. Stamens. Stip. Pale. Leaves. Leguminosæ.

tinctions disappears in a great number of cases. *Cæsalpinieæ* have irregular flowers with spreading petals and stamens adhering to the



calyx; others have no petals at all, or some number less than five; while *Mimoseæ* have perfectly regular flowers and indefinite hypogynous stamens. *Detarium* and other plants of this family bear fruits not to be distinguished from a drupe. Leguminous plants and *Rose-worts* have in some instances so many features in common that it may be affirmed no positive character has been discovered to distinguish the one order from the other.

848. The plants of the order are very generally distributed over the globe, but many genera are very limited in their range. De Candolle gives the following geographical distribution of the 3600 species known in his day:—

Figs. 593-597.—Organs of fructification of *Lathyrus odoratus*, Sweet-pea, a papilionaceous flower, showing the structure of the natural order Leguminosæ.

Fig. 593.—Diagram of the flower, showing five divisions of the calyx, 5 petals, consisting of 2 parts forming the carina, 2 alæ, and the vexillum, which is superior, 10 stamens in 2 rows, diadelphous; ovary 1-celled, formed by a single carpel; one of the ovules shown with its funiculus attached to the ventral suture.

Fig. 594.—Longitudinal section of the flower of *Lathyrus odoratus*. c c, Calyx, with five segments. e, Vexillum or standard, being the superior or posterior odd petal. a, One of the alæ, or wings. c, a, One-half of the carina, or keel. f, Tube of the stamens, the filaments being united in two bundles, or diadelphous. o, Ovary laid open, showing the ovules attached to the placenta, on the ventral or upper suture. s, Stigma, at the apex of the style, which is continuous with the ventral suture.

Fig. 595.—Fruit, a Legume or Pod, opening by two valves, and dehiscing by the ventral and dorsal suture. Seeds attached on each side of the ventral suture, curved upon themselves, having a marked hilum and funiculus (podosperm or umbilical cord).

Fig. 596.—A seed separated. f, Funiculus. c, Chalazal unit to the funiculus by the raphe. m, Micropyle or foramen.

Fig. 597.—Embryo, which occupies the entire seed after the spermoderm is removed. c, c, Two cotyledons separated; they are fleshy and hypogæal, i.e. remain under ground during germination. g, Gemmule or plumule. r, Radicle.

Equinoctial America.....	605
Basin of the Mediterranean.....	468
East Indies.....	452
Cape of Good Hope.....	353
Levant.....	250
New Holland.....	229
West Indies.....	221
Europe, excepting the Mediterranean.....	184
United States.....	183
Mexico.....	152
Equinoctial Africa.....	130
Siberia.....	129
Arabia and Egypt.....	87
China, Japan, Cochin-China.....	77
Isles of Southern Africa.....	42
South America, beyond the tropics.....	29
Canaries.....	21
South Sea Islands.....	13

It is said that no native species occur in the islands of Tristan d'Acugna and St. Helena.

The order has been divided into three suborders:—1. Papilionaceæ; papilionaceous flowers, petals imbricated in æstivation, and upper one exterior. This suborder is subdivided into the tribes Podalyrieæ, Lotææ, Viciææ, Hedysareæ, Phaseoleæ, Dalbergiææ, Sophoreæ; according to the nature of the filaments, whether free or variously united, the form and dehiscence of the legume, the cotyledons whether fleshy or leafy, and the simple or compound nature of the leaves. *Examples*—Podalyria, Lotus, Cytisus, Pisum, Hedysarum, Phaseolus, Dalbergia. 2. Cæsalpinieæ; flowers irregular but not papilionaceous, petals spreading, imbricated in æstivation, upper one interior. *Examples*—Hæmatoxylon, Cæsalpinia, Cassia, Swartzia, Amherstia, Bauhinia, Copaifera, and Ceratonia. 3. Mimoseæ; flowers regular, petals valvate in æstivation. *Examples*—Parkia, Mimosa, Acacia.

The number of known genera at the present day, according to Lindley, is 467, comprehending 6500 species. The following is the estimate of species in the different suborders and tribes, considered in reference to the flora of the globe and the flora of Britain (Bentham and Henslow):—

Suborders.	Tribes.	Species.	British species.
1. Papilionaceæ.....	1. Podalyrieæ.....	350	0
	2. Lotææ with } Viciææ.....	3000	48 }
	3. Hedysareæ.....	500	4
	4. Phaseoleæ.....	650	0
	5. Dalbergiææ.....	250	0
	6. Sophoreæ.....	50	0
2. Cæsalpinieæ.....		700	0
3. Mimoseæ.....		1000	0
		6500	75

849. This is a very extensive and a very important natural order. It embraces many valuable medicinal plants, such as those yielding Senna, Gum-arabic, Tragacanth, Catechu, and Kino; important dyes, as Indigo and Logwood; many valuable timber trees, as Locust-tree and Rosewood; plants furnishing nutritious food, such as the Bean and Pea. The properties of the order may be considered in general as wholesome, although it contains some poisonous plants. Lindley, however, says that the order must be considered upon the whole as poisonous, and that the plants used for food are exceptions to the general rule; the injurious juices of the order not being in such instances sufficiently concentrated to prove injurious, being replaced to a considerable extent by either sugar or starch.

850. Suborder *Papilionaceæ*. The plants in this section have frequently beautiful showy flowers; for example, *Robinia*, *Laburnum*, *Wisteria*, *Lupinus*, *Clianthus*, *Erythrina*, or Coral-flower, *Hovea*. They are often nutritious. The various kinds of Clover, Beans, Peas, and Pulse belong to it. The common red Clover is *Trifolium pratense*. White or Dutch Clover (*T. repens*) springs up frequently on ground recently cleared. The Shamrock is generally considered as a species of Trefoil. Various species of Medick and Lucerne (*Medicago*, fig. 471), of Saintfoin (*Onobrychis*), and Melilot (*Melilotus*), are cultivated as food for cattle. Many are used for their medicinal qualities. *Glycyrrhiza glabra*, or *Liquiritia officinalis*, is the plant which yields liquorice-root. This plant is a native of the southern part of Europe, and it has been occasionally cultivated with success in Britain, especially at Pontefract in Yorkshire, and at Mitcham in Surrey. An extract is prepared from the root or underground stem by decoction in water, and subsequent inspissation. It owes its sweetness to a peculiar principle called Glycion, or Glycyrrhizin, which appears also to be present in the root and leaves of other papilionaceous plants, as *Glycyrrhiza echinata* and *glandulifera*, *Trifolium alpinum*, and *Abrus precatorius*. Liquorice is used medicinally as a demulcent. A sweet secretion (a kind of Manna) is produced by *Alhagi Maurorum*. *Astragalus verus*, *creticus*, *aristatus*, *gummifer*, and other species, yield an exudation known by the name of Gum Tragacanth. *A. verus* seems to be the chief source of the European tragacanth. It is a shrub found in Asia Minor and Persia, and the gum is procured by exudation or incision. Tragacanth forms with cold water a bulky jelly, while it is soluble in boiling water. It contains both Arabin and Bassorin in its composition, and is used as a demulcent. *Myrospermum*, or *Myroxylon peruiferum*, yields the Balsam of Peru, while *Myrospermum*, or *Myroxylon toluiferum*, is the source of the Balsam of Tolu. These balsams are procured chiefly by making incisions in the trees. They consist of resinous and oily matter, with cinnamic acid, and they are used as stimulant expectorants. *Pterocarpus Marsupium*, a tree of the Indian forests, furnishes the concrete exuda-

tion called Kino. *Butea frondosa*, or the Dhak tree of the East Indies, yields a similar product, its flowers discharging a beautiful yellow or orange dye. African Kino is procured from *Pterocarpus erinaceus*. Kino is used as a powerful astringent, and is administered in the form of powder and tincture. Broom-tops, procured from *Cytisus* (*Sarothamnus*) *Scoparius*, are used as a diuretic. The hairs from the legumes of *Mucuna pruriens* in the West Indies, and of *M. Prurita* in the East, under the name of Cowhage, or Cowitch, have irritating properties, and, mixed with syrup, they are used in the treatment of intestinal worms. The leaves of *Colutea arborescens*, Bladder-Senna (fig. 470), are purgative, and are used abroad to adulterate the obovate or blunt-pointed Senna. The leaves of *Tephrosia apollinea* are also purgative, and are occasionally mixed with Alexandrian Senna. The bark of *Andira inermis*, the Cabbage-tree of the West Indies, acts as a purgative and anthelmintic. The fruit of *Geoffroya superba*, Umari, is much used by the inhabitants of Brazil on the banks of the Rio San Francisco; the fruit is a drupe.

Besides the plants which have active medicinal qualities, there are others which are valuable in commerce and the arts, as furnishing food, dyes, fibres, timber. Various species of *Indigofera*, as *I. tinctoria* and *cærulea*, furnish the Indigo of commerce. *Pterocarpus santalinus* yields red Sandal-wood, which is used as a dye. *P. Draco* yields Gum-Dragon, and *P. Dalbergioides* is said to yield Andaman redwood, and to be valuable both as a dye and as timber. *Baptisia tinctoria* gives a blue dye, and is the wild Indigo of the United States. *Dalbergia Sissoo* is an Indian forest tree, which is valued on account of its wood. *Crotalaria juncea* supplies fibres, which are known as Sun or Bengal Hemp. The fragrant seeds of *Dipterix odorata* are known as Tonka-beans. A similar fragrance is given out by some species of Melilot, the flowers and seeds of which are employed to give the peculiar odour to Gruyere cheese. *Arachis hypogæa* produces its legumes under ground, and receives the name of under-ground Kidney-bean, or Ground-nut. *Erythrina monosperma* yields Gum lac. The roots of *Glycine Apios*, or *Apios tuberosa*, are used as an article of food in America. *Robinia pseudo-acacia* is often cultivated in Britain as the Locust-tree. The tree attains in England a height varying from forty-five to eighty feet, and sometimes has a diameter of three feet. Its wood is durable. According to Bertoloni, a kind of Ebony is the produce of *Fornasinia ebenifera*, a papilionaceous plant, found in Caffraria, near Mozambique. Rosewood is said to be the timber of two or three species of *Trioptolomea*. It is rare to find papilionaceous plants produce double flowers. The Whin is one of the plants which exhibits this monstrosity. *Desmodium* or *Hedysarum gyrans* exhibits a remarkable irritability in its leaves (¶ 660). There are certain poisonous plants in this suborder. The seeds and bark of *Cytisus*

Laburnum are narcotic; such is said also to be the case with those of *Lathyrus Cicera*, and *L. Aphaca*. The roots of many species of *Phaseolus*, as *P. multiflorus*, the Scarlet-runner, and *P. radiatus*, are poisonous. The branches and leaves of *Tephrosia toxicaria*, and the bark of the root of *Piscidia erythrina*, Jamaica Dogwood, are employed as fish poisons. They yield a narcotic and diaphoretic tincture. A species of *Gompholobium* has poisoned sheep in the Swan River colony. *Coronilla varia* acts as a narcotic poison. The leaves of it and of *Coronilla Emerus* are sometimes used to adulterate Senna.

851. Suborder *Cæsalpinieæ*. In this section there are many plants which furnish purgative remedies. Among these may be noticed various species of *Cassia*. *C. lanceolata*, *acutifolia*, *elongata*, *obtusata*, and *obovata*, supply the various kinds of Senna known as Alexandrian or Egyptian, Tripoli, and East Indian Senna. Other species also, as *Cassia marilandica*, *Absus*, *corymbosa*, *biflora*, *tomentosa*, *alata*, and *Porturegalis*, have purgative leaves. *Cassia Fistula*, called also *Cathartocarpus Fistula*, has an indehiscent pod, divided by numerous transverse phragmata (fig. 395), and containing a laxative pulp, which is a secretion from the endocarp. A pulp having similar properties is procured from the pericarp of *Tamardinus indica*, the Tamarind-tree. The pod of *Ceratonia Siliqua* is known as the Algaroba-bean, and is used occasionally for feeding horses. The tree is denominated Carob-tree, and sometimes Locust-tree, or St. John's Bread, from a tradition that the pulpy matter surrounding the seeds supplied food to St. John in the wilderness. The pods of *Hymenæa Courbaril*, the West Indian Locust-tree, supply a nutritious matter; its inner bark is anthelmintic, and the plant yields a kind of resin called Anime. The bark of *Guilandina Bonduc*, the Nicker-tree, is bitter, tonic, and its seeds are said to be emetic. The curved pods of *Cæsalpinia coriaria*, under the name of Dividivi, are used for tanning. *Cæsalpinia brasiliensis* yields the Brazil-wood of commerce, and the Mora-wood of Guiana is yielded by a large tree called *Mora excelsa*. Many dyes are furnished by the plants of this suborder. *Hæmatoxylon campeachianum* gives Logwood or Campeachy-wood, which is employed both as a dye and as an astringent. The inner wood is the part employed both in the arts and officinally. The alburnum is of a yellowish colour, and is not imported. The red colouring principle is Hæmatoxylin. *Cæsalpinia echinata* furnishes Pernambuco-wood; *C. sappan*, Sappan-wood, the Wukkum or Bukkum-wood of Scinde; *Baphia nitida*, Camwood. Various species of *Copaifera*, as *C. Jacquinii*, *Langsdorffii*, *biguga*, *multiguga*, *Martii*, *Guianensis*, *coriacea*, &c., furnish the Baslam of Copaiva, of which two kinds are distinguished—the West Indian and Brazilian. The balsam contains a resin and volatile oil. It is used in medicine as a stimulant, cathartic, and diuretic, and is especially employed in the treatment of mucous inflammations. *Cassia Chamæ-*

crista, *marilandica*, and *nititans*, all exhibit, according to Bromfield, a high degree of irritability; the leaflets close together almost as soon as gathered, and even when rudely handled or brushed by the feet in walking through the herbage.


852. Suborder *Mimoseæ*. The plants of this section yield Gum in large quantity, and their bark is frequently astringent. *Acacia Ehrenbergii*, *tortilis*, *Seyal*, *arabica*, *vera*, *gummifera*, *Adansonii*, *Verek*, *albida*, and various other species, yield the gummy substances known as Gum Arabic, Gum Senegal, Barbary Gum, and East Indian Gum. A kind of gum is procured at the Cape of Good Hope from *Acacia Karroo*, and in New Holland, *A. decurrens* yields another variety. A variety of Indian gum procured from *A. arabica*, is called Babul, or Babool-Gum; Babul-wood is used for tanning in Scinde. These gums are all more or less nutritive and demulcent, and are administered in the form of mucilage, emulsion, or lozenges. The Wattles of Australia are species of *Acacia*, which furnish astringent barks. An extract made from them has been imported for the purpose of tanning. The inner wood of *Acacia Catechu*, an Indian shrub, furnishes a kind of Catechu, or Cutch, which contains much tannin, and is used for tanning, and as a powerful astringent. Some of the New Holland Acacias are remarkable for the peculiar development of the petiole, which assumes the form of a phyllodium (fig. 188 p). The large seeds of *Entada scandens* are sometimes carried by the winds and tides from the West Indies to the shores of the outer Hebrides. Some say that a species of *Desmanthus* is the source of the cellular substance called Rice-paper; others refer it to the genus *Æschynomene*. Some of the plants in this suborder display peculiar irritability in their pinnate leaves. This is particularly the case with *Mimosa sensitiva* and *pudica*, which are commonly called sensitive plants (¶ 659). Almost all of the pinnate-leaved Leguminous plants close their leaves in a marked way during darkness.

853. Order 69.—**Moringaceæ**, the Moringa Family. (*Polypet. Perigyn.*) Calyx 5-partite; æstivation slightly imbricated. Petals 5, rather unequal, upper one ascending. Stamens 8 or 10, perigynous; filaments slightly petaloid, callous, and hairy at the base; anthers simple, 1-celled, with a thick convex connective. Disk lining the tube of the calyx. Ovary superior, stipitate, 1-celled; ovules anatropal, attached to parietal placentas; style filiform; stigma simple. Fruit a pod-like capsule, 1-celled, 3-valved, opening by loculicidal dehiscence. Seeds numerous, half buried in the spongy substance of the valves, sometimes winged, exalbuminous; embryo with a superior, straight, small radicle, and fleshy cotyledons.—Trees, with bi- or tri-pinnate, stipulate leaves, natives of the East Indies and Arabia. Some of them are pungent and aromatic. The seeds of *Moringa pterygosperma*, Horse-radish tree, are winged, and are called Ben-nuts. From

Fruit, Mithradatum, Follicle, Pome or Pomme—

them is procured a fluid oil, used by watchmakers, and called Oil of Ben. The root is pungent and stimulant, and resembles Horse-radish in its taste. It is used as a stimulant in paralytic affections and in intermittent fever. It is also a rubefacient. Lindley places this order in his Violal alliance. He mentions 1 genus, and 4 species. *Example*—Moringa.

854. Order 70.—**Rosaceæ**, the Rose Family. (*Polypet. Perigyn.*) (Figs. 226, 235, 236, 276, 289, 385, 598). Calyx 4-5-lobed (fig. 599 *c c*), the fifth lobe superior. Petals as many as the divisions of



the calyx, often 5 (fig. 599 *p e*), sometimes wanting, perigynous, generally regular; æstivation quincuncial (fig. 598). Stamens inserted with the petals (fig. 599 *e*), definite or indefinite; filaments incurved in æstivation; anthers bilocular (fig. 600) dehiscing longitudinally

Figs. 598-605.—Organs of fructification of *Rubus strigosus*, illustrating the natural order Rosaceæ.

Fig. 598.—Diagram of the flower, with five divisions of the calyx, 5 quincuncial petals, indefinite perigynous stamens, and numerous succulent carpels.

Fig. 599.—The flower cut vertically. *c c*, Calyx. *pe*, Petals. *e*, Stamens. *d*, Disk, lining the base of the calyx, upon which the stamens are inserted. *pi*, Plstil composed of several carpels.

Fig. 600.—Bilocular anther separated with the upper part of the filament, seen on the outside.

Fig. 601.—Ovary, *o*, cut vertically. *g*, Exalbuminous, suspended seed. *s*, Lateral style.

Fig. 602.—Fruit *f*, Fleshy carpels accompanied with the persistent calyx, *c*, connected with the base of the calyx.

Fig. 603.—Vertical section of a carpel. *s*, Lateral Style. *m*, Fleshy mesocarp or sarcocarp. *e*, Endocarp. *g*, Seed.

Fig. 604.—Horizontal section of the exalbuminous seed. *t*, Integument (spermoderm). *c*, Cotyledons of the embryo.

Fig. 605.—Embryo isolated. It fills the entire seed.

Rosaceæ - Stipitate Leaves - Old Sepal next Axis - Numerous Stamens -

(fig. 322). Ovaries superior, either solitary or several, unilocular (fig. 601), sometimes uniting so as to form a many-celled pistil; ovules 1, 2, or more, anatropal, suspended (figs. 373 *g*, 601 *g*), rarely erect; styles lateral (figs. 400, 601, 603); stigmas usually simple. Fruit either achænia (fig. 270) or drupes (figs. 373, 602), or follicles or pomes (fig. 472). Seeds erect or inverted, usually exalbuminous; embryo straight, with the radicle next the hilum (figs. 603, 605,) and leafy or fleshy cotyledons (figs. 501, 604).—Herbaceous plants, or shrubs, or trees, with simple or compound, alternate, stipulate leaves (fig. 191), and the flowers sometimes unisexual. They are found chiefly in the cold and temperate climates of the northern hemisphere. Some are found on high mountains within the tropics, and a few occur in warm regions. One species only occurs in the West Indies, viz., *Rubus jamaicensis*.

The order has been divided into the following suborders:—1. *Chrysobalanææ*, petals and stamens more or less irregular; ovary stipitate, its stalk adhering on one side to the calyx, style basilar (fig. 401), fruit a 1-2-celled drupe. 2. *Amygdalææ* (*Drupacææ* of Lindley), tube of calyx lined with a disk, styles terminal, fruit a drupe (figs. 308, 371-373). 3. *Spiræææ*, calyx-tube herbaceous, lined with a disk, fruit consisting of numerous follicles, seeds apterous. 4. *Quillaieææ*, flowers unisexual, calyx-tube herbaceous, fruit capsular, seeds winged at the apex. 5. *Sanguisorbææ*, petals 0, tube of calyx thickened and indurated, lined with a disk, stamens definite, not solitary, enclosed in the calycine tube. 6. *Potentilleææ*, calyx-tube herbaceous, lined with a disk which sometimes becomes fleshy, fruit consisting of numerous achænia. 7. *Rosææ*, calyx-tube contracted at the mouth, becoming fleshy, lined with a disk, and covering numerous hairy achænia (figs. 270, 289). 8. *Pomeææ* (*Pomacææ* of Lindley), tube of calyx more or less globose, ovary fleshy and juicy, lined with a thin disk, fruit a 1-5-celled (fig. 472) or spuriously 10-celled pomum. There are 82 known genera, and about 1000 species. *Examples*—*Chrysobalanus*, *Amygdalus*, *Prunus*, *Spiræa*, *Quillaia*, *Sanguisorba*, *Potentilla*, *Rubus*, *Fragaria*, *Rosa*, *Pyrus*.

855. Many of the plants of the order yield edible fruits, such as Raspberries, Strawberries, Brambles, Plums, Apples, Pears, Quinces, Cherries, Almonds, Peaches, Nectarines, and Apricots. Some are astringent, others yield hydrocyanic acid. Those belonging to the suborder *Chrysobalanææ*, are principally natives of the tropical parts of Africa and America. Many of them furnish edible fruits. The drupes of *Chrysobalanus Icaco*, are eaten in the West Indies under the name of Cocoa-plums. The root and bark are used as astringents.

The plants in the suborder *Amygdalææ*, are chiefly remarkable on account of the presence of hydrocyanic acid in their kernels, leaves,

or flowers. *Amygdalus communis*, the Almond-tree, grows naturally in Barbary and in Asia, from Syria to Affghanistan. It is extensively cultivated in the south of Europe. There are two varieties of the tree,—*α. dulcis*, yielding the sweet Almond, and *β. amara*, yielding the bitter Almond. In the former, the style is much longer than the stamens, and there are glands on the base of the leaf; while in the latter, the style is equal in length to the stamens, and the glands are situated on the petioles. The chief kinds of sweet Almonds are the Valentia, the Italian, and the Jordan Almonds; the latter come from Malaga. Under the name of shell Almonds, they are often sold with the brittle endocarps on them. They consist chemically of a bland fixed oil, and a kind of vegetable albumen called Emulsin or Synaptase. Bitter Almonds are imported from Mogadore. Besides a fixed oil and synaptase, they contain a bitter azotised principle called Amygdalin, which, when brought into contact with a solution of Emulsin, produces a volatile oil, containing hydrocyanic acid. This gives rise to the peculiar aroma of bitter Almonds, when mixed with water. Sweet Almonds are used medicinally, in the form of Emulsion, as demulcents. The hydrocyanated essential oil of bitter Almonds is sedative, and has been used as a substitute for Prussic acid. They sometimes produce derangement of the digestive functions, and give rise to nettle-rash. The leaves of *Amygdalus persica* (*Persica vulgaris* of some), the Peach, contain a similar oil, and have been employed as sedative and vermifuge. The flowers of the Peach exhale the odour of bitter Almonds. Peaches are divided into Freestone and Clingsone, according as the pulp (sarcocarp) separates easily from the endocarp or adheres to it. The fruit of *Prunus domestica*, the Plum-tree and its varieties, when dried, constitute Prunes, which are used medicinally, on account of their nutritive and laxative qualities. The leaves of *Prunus* or *Cerasus Laurocerasus*, Cherry Laurel, or common Bay Laurel, have been used medicinally, as anodyne and hypnotic remedies. The water distilled from them has poisonous properties, owing to the presence of a hydrocyanated oil, which seems to be developed in a similar manner as in the case of bitter Almonds. The oil exists in largest quantity in the young leaves. *Prunus Lusitanica* is the Portugal Laurel, which is extensively cultivated in Britain as an evergreen. The leaves of *Prunus spinosa*, the Sloe, have been used as a substitute for, as well as an adulteration of Tea. The fruit of a variety of *Cerasus avium*, the Cherry, is used in the manufacture of Kirschenwasser. The kernel of *Cerasus occidentalis* is used for flavouring Noyau. The flavour of Ratafia, Cherry-brandy, and Maraschino, are due to the kernels of *Cerasus*.

The suborder *Pomeæ* supplies many edible fruits, as Apples, Pears, Medlars (fig. 472), and Quinces. The seeds and occasionally the flowers and bark of some, yield hydrocyanic acid. All the cultivated

varieties of Apple are derived by grafting from the native species, *Pyrus Malus*; while Pears have their origin in *Pyrus communis*. The seeds of *Cydonias vulgaris* (*Pyrus Cydonia*), the Quince, when boiled in water, yield a mucilaginous decoction, which has been used as a demulcent. Malic acid is found in some of the fruits of this suborder. *Eriobotrya japonica* yields the Loquat, a Japan fruit.

The other suborders contain plants which are distinguished by astringent and tonic qualities. *Geum urbanum* and *rivale* (Avens) have been employed as tonics and astringents, as also the root of *Potentilla Tormentilla* (Tormentil). Cusso or Cabotz, known as Brayera anthelmintica, one of the most powerful remedies for the cure of tænia, belongs to this family. The properties of this plant being first described by Brayer, it has been named after him. The varieties of Scotch Roses are derived from *Rosa spinosissima*. The fruit of *Rosa canina*, the Dog-rose, which consists of the enlarged fleshy calyx enclosing numerous achænia (fig. 270), is beat into a pulp with sugar, and used as an acidulous refrigerant and astringent. The petals of *Rosa gallica*, Red, French, and Provins Rose, are employed in the form of infusion, as a tonic and slightly astringent remedy. The petals of *Rosa centifolia*, the Hundred-leaved or Cabbage-rose (fig. 88), and its varieties, *R. damascena*, Damask-rose, *R. moschata*, Musk-rose, &c., are employed in the preparation of Rose-water, and of the oil or attar of Roses. It is stated by Dr. Christison, that 100,000 roses, the produce of 10,000 bushes of *Rosa damascena*, yield at Ghazeepore only 180 grains of attar. The bark of many species of *Quillaia*, as *Q. saponaria*, are used as a substitute for soap.

856. Order 71.—**Calycanthaceæ**, the Calycanthus Family. (*Polypet. Perigyn.*) Sepals and petals confounded, indefinite, combined in a fleshy tube; æstivation imbricated. Stamens ∞ , perigynous; anthers adnate, extrorse, with longitudinal dehiscence. Ovaries several, 1-celled, adhering to the tube of the calyx; ovules solitary or two, one above the other, anatropal; style terminal. Fruit consisting of achænia enclosed in the fleshy tube of the Calyx. Seed exalbuminous; embryo straight; cotyledons convolute; radicle inferior.—Shrubs, with square stems, consisting of a central woody mass, with four smaller ones around (¶ 90); leaves opposite, simple, scabrous, exstipulate. They are natives of North America and Japan. Their flowers are aromatic; the bark of some is used as a carminative. *Calycanthus floridus* is called Carolina or common American Allspice. The order includes 2 genera, and 6 species. *Examples*—Calycanthus, Chimonanthus.

857. Order 72.—**Lythraceæ**, the Loosetrife Family. (*Polypet. Perigyn.*) Calyx tubular, lobed, the lobes sometimes with intermediate lobes or teeth; æstivation valvate. Petals alternate with the primary lobes of the calyx, very deciduous, sometimes 0. Stamens inserted

into the tube of the calyx a little below the petals, equal in number to them, or two, three, or four times as many; anthers adnate, dithecal, introrse, with longitudinal dehiscence. Ovary superior, 2-6-celled; ovules numerous, anatropal; style filiform; stigma usually capitate. Fruit a dehiscent membranous capsule, surrounded by the calyx, but not adherent to it, sometimes 1-celled by the obliteration of the dissepiments. Seeds numerous, small, apterous, or winged, exalbuminous, attached to a central placenta; embryo straight; cotyledons flat and foliaceous; radicle next the hilum.—Herbs and shrubs, with branches which are usually tetragonal, and with opposite, rarely alternate, entire, exstipulate leaves without glands. They are natives of Europe, North and South America, and India. The order is divided into two suborders:—1. *Lythreæ*, with apterous (wingless) seeds. 2. *Lagerströmiæ*, with winged seeds. Lindley gives 35 genera, including 300 species. *Examples*—*Lythrum*, *Cuphea*, *Lagerströmia*.

858. Many of the plants of the order are distinguished by astringent properties, and some are used for dyeing. *Lythrum Salicaria*, Purple Loosestrife, or Willowstrife, a European plant, found also in Australia, has been used in cases of diarrhœa, on account of the tannin in its composition. The flowers of *Grislea tomentosa* are employed in India, mixed with morinda, for dyeing, under the name of *Dhaec*. *Heimia Salicifolia* is said to possess diaphoretic properties, and is considered by the Mexicans as a potent remedy for venereal diseases. The Henna, or Alhenna of the Arabs, which is used in Egypt for dyeing orange, is the product of *Lawsonia inermis*. The *Cupheas* are remarkable for the mode in which the placenta bursts through the ovary and floral envelopes, so as to expose the seeds.

859. Order 73.—**Rhizophoraceæ**, the Mangrove Family. (*Polypet. Epigyn*). Calyx adherent, 4-12-lobed; æstivation valvate, or sometimes calyptriform. Petals arising from the calyx, alternate with the lobes, and equal to them in number. Stamens inserted with the petals, twice or thrice their number; filaments distinct, subulate; anthers erect. Ovary 2-3-4-celled; ovules 2 or more in each cell, anatropal. Fruit indehiscent, adherent to the calyx, and crowned by it, unilocular, monospermous. Seed solitary, pendulous, exalbuminous; cotyledons flat; radicle long, piercing the fruit.—Trees or shrubs, with simple opposite leaves, and deciduous interpetiolar stipules. They are found on the muddy shores of the tropics. There are 5 genera, and 20 species known. *Examples*—*Rhizophora*, *Kandelia*.

860. The plants of the order have frequently an astringent bark, which is in some cases used for dyeing black. *Rhizophora Mangle*, the Mangrove, forms thickets at the muddy mouths of rivers in tropical countries, and sends out adventitious roots which often raise the main trunk much above its original level, and give the tree the appearance

of being supported upon stalks. The fruit is sweet and eatable. The embryo germinates before the fruit falls, and the radicle is much elongated before the seed drops into the mud. The anther consists of numerous cells containing pollen.

861. Order 74.—**Vochysiaceæ**, the Vochysia Family. (*Polypet. Perigyn.*) Sepals 4-5, united at the base, unequal, the upper one largest and spurred; æstivation imbricated. Petals 1, 2, 3, or 5, alternate with the divisions of the calyx, and inserted into its base, unequal. Stamens 1-5, opposite to, or alternate with the petals, perigynous, one having an ovate, fertile, 4-celled anther, the rest being sterile. Ovary free, or partially adherent to the calyx, 3-celled; ovules solitary or in pairs, rarely numerous, amphitropal or anatropal; style and stigma one. Fruit a triquetrous, 3-celled and 3-valved capsule, usually with loculicidal dehiscence. Seeds usually 1-2 in each cell, erect, exalbuminous, attached to a central placenta; embryo straight; cotyledons large and leafy; radicle short and superior.—Trees or shrubs, with opposite, entire, stipulate leaves. They inhabit the warmer parts of America. Their properties are little known. Their flowers are reputed to be very sweet, and some are said to have a resinous juice. There are 8 genera enumerated, including 51 species. *Examples*—Vochysia, Qualea.

862. Order 75.—**Combretaceæ**, the Myrobalan Family. (*Polypet. Epigyn.*) Calyx 4-5-lobed, lobes deciduous. Petals arising from the orifice of the calyx, alternate with the lobes, or wanting. Stamens epigynous, twice as many as the lobes of the calyx, rarely equal in number, or thrice as many; filaments distinct, subulate; anthers dithecal, dehiscing longitudinally or by recurved valves. Ovary adherent to the tube of the calyx, unilocular; ovules 2-4, pendulous; style 1; stigma simple. Fruit succulent or nut-like, inferior, unilocular, indehiscent, often winged. Seed solitary, pendulous, exalbuminous; cotyledons leafy, usually convolute, sometimes plicate; radicle turned towards the hilum.—Trees or shrubs, with alternate or opposite exstipulate, entire leaves. They are natives of the tropical regions of Asia, Africa, and America. The general property of the order is astringency. Many are used for tanning, and some for dyeing. The fruit of *Terminalia Belerica*, and of *T. Chebula*, under the name of Myrobalans, is used as an astringent. The seeds of *Terminalia Catappa* are eaten like almonds. The order has been divided into three suborders:—1. *Terminaliæ*, petals 0, cotyledons convolute. 2. *Combretæ*, petals present, cotyledons plicate. 3. *Gyrocarpæ*, petals 0, cotyledons convolute, anthers dehiscing by recurved valves. There are 22 genera enumerated by Lindley, including 200 species. *Examples*—*Terminalia*, *Combretum*, *Gyrocarpus*.

863. Order 76.—**Melastomaceæ**, the Melastoma Family. (*Polypet. Perig. or Epigyn.*) Calyx with 4, 5, or 6 divisions, which are more

or less deep, or are sometimes united and separate from the tube like a lid. Petals equal to the segments of the calyx, perigynous, æstivation twisted. Stamens equal in number to the petals and alternate with them, usually with intermediate sterile ones; filaments curved downwards in the young state; anthers long, often beaked, bilocular, dehiscing by two terminal pores or longitudinally. Ovary more or less adherent to the calyx, multilocular; ovules usually 00; style 1; stigma simple, either capitate or minute. Fruit multilocular, either capsular, with loculicidal dehiscence, or succulent, combined with the calyx and indehiscent. Seeds ∞ , minute, attached to central placentas, exalbuminous; embryo straight or curved; cotyledons sometimes unequal, flat, or convolute.—Trees, shrubs, or herbs, with opposite, undivided, usually entire, often 3-9-ribbed leaves, not dotted. They are found chiefly in warm climates. Many are natives of America and India. There are no unwholesome plants in the order, and the succulent fruit of several is edible. A slight degree of astringency pervades all the plants of the order, and hence some are used medicinally in cases of diarrhœa. The name *Melastoma* (μέλας, black, and στόμα, mouth), is derived from the circumstance that the fruit of some dyes the lips black. There are two suborders:—1. *Melastomææ*, with ribbed leaves and flat cotyledons. 2. *Memecyleæ*, with ribless leaves and convolute cotyledons. Lindley notices 118 genera, comprising 1200 species. *Examples*—*Melastoma*, *Osbeckia*, *Lasiandra*, *Rhexia*, *Lavoisiera*, *Miconia*, *Charianthus*, *Memecylon*, *Mouriria*.

864. Order 77.—*Alangiaceæ*, the *Alangium* Family. (*Polypet. Epigyn.*) Calyx campanulate, adherent, 5-10-toothed. Petals 5-10, linear, reflexed, inserted into a fleshy disk, which is adherent to the calyx, and ovary; æstivation twisted. Stamens long, exserted, 2 or 4 times as many as the petals; filaments distinct, villous at the base; anthers bilocular, adnate, introrse, often sterile. Ovary globose, unilobular; ovules solitary, anatropal; style filiform; stigma capitate or conical. Fruit oval, fleshy, coherent with the tube of the calyx, and somewhat crowned by its limb, slightly ribbed; endocarp sometimes osseous, with a foramen at the apex. Seed solitary, anatropal; albumen fleshy, brittle; embryo straight; cotyledons flat, foliaceous; radicle long, superior.—Trees or shrubs, with alternate, exstipulate leaves, which are not dotted. They are found chiefly in India, some are natives of America; some of the plants yield edible fruits, others are purgative. Lindley enumerates 3 genera, comprehending 8 species. *Examples*—*Alangium*, *Nyssa*.

865. Order 78.—*Philadelphaceæ*, the *Syringa* Family. (*Polypet. Epigyn.*) Calyx with a 4-10-divided, persistent limb. Petals alternate with the divisions of the calyx, and equal to them in number; æstivation convolute, imbricate. Stamens ∞ (rarely 10), in one or two rows, arising from the orifice of the calyx. Ovary adherent to

the tube of the calyx; styles distinct, or united into one; stigmas 4-10; ovules ∞ , attached to a central placenta. Fruit a 4-10-celled capsule, free above. Seeds ∞ , scrobiform, subulate, smooth, pendulous, with a loose membranous arillus; albumen fleshy; embryo straight, about as long as the albumen; cotyledons flat; radicle next the hilum, obtuse.—Shrubs with deciduous, opposite, exstipulate leaves without dots; flowers usually in trichotomous cymes. They are natives of the South of Europe, of North America, Japan, and India. They have no marked properties. The flowers of *Philadelphus coronarius*, *Syringa*, have a peculiar sweetish odour, which to some persons is overpowering and disagreeable. The smell is due to the presence of an oil. *Deutzia scabra* has a scurfy matter on its leaves, which, under the microscope, is seen to consist of beautiful stellate hairs. The leaves are in consequence used in Japan by polishers. Its inner bark is used for poultices. There are 3 genera enumerated, including 25 species. *Examples*—*Philadelphus*, *Deutzia*, *Decumaria*.

866. Order 79.—**Myrtaceæ**, the Myrtle Family. (*Polypet. Epigyn.*) Calyx 4-5-6-8-cleft, the limb sometimes cohering at the apex, and falling off like a lid; æstivation valvate. Petals attached to the calyx, alternating with its segments, and equal to them in number, with a quincuncial æstivation, rarely 0. Stamens inserted with the petals, twice as many as the petals, or ∞ ; filaments distinct, or united in one or more parcels, curved inwards in the bud; anthers ovate, dithecal, with longitudinal dehiscence. Ovary adherent to the tube of the calyx, 1-6-celled; style and stigma simple; ovules anatropal, pendulous or erect. Fruit dry or fleshy, dehiscent or indehiscent. Seeds usually ∞ , attached to a central placenta; mostly exalbuminous; embryo straight or curved; cotyledons distinct (fig. 514), or consolidated with the radicle, which is next the hilum.—Trees or shrubs, with opposite, rarely alternate leaves, which are usually entire and dotted, and frequently have an intramarginal vein. They are natives chiefly of warm countries, as South America and the East Indies. Many, however, are found in more temperate regions. Some of the genera are peculiar to Australia. The order has been divided into the following suborders:—1. *Chamælaucieæ*, heath-like plants with a 1-celled ovary and capsule, and opposite dotted leaves. 2. *Leptospermeæ*, having a multilocular capsule, and opposite or alternate, usually dotted leaves. 3. *Myrteæ*, having a baccate fruit, distinct stamens, opposite dotted leaves. 4. *Barringtonieæ*, having a fleshy 1-celled fruit, monadelphous stamens, albuminous seeds, opposite or verticillate leaves, not dotted. 5. *Lecythideæ*, having a multilocular woody capsule, which either remains closed or opens by a lid, monadelphous stamens, alternate, not dotted leaves. Several of these suborders are made separate orders by Lindley and others. There are 77 known genera, and upwards of 1400 species. *Examples*—*Chamælaucium*,

Calytrix, Leptospermum, Melaleuca, Metrosideros, Eucalyptus, Myrtus, Psidium, Eugenia, Caryophyllus, Barringtonia, Gustavia, Lecythis, Bertholletia.

867. Many of the plants of the order yield an aromatic volatile oil. This is particularly the case with those having pellucid dots in their leaves. Many yield edible fruits, others furnish astringent and saccharine substances. The leaves of some species of *Leptospermum* and *Melaleuca* are used as tea in Australia. The leaves of *Melaleuca minor* (*Cajuputi* of some), a native of the Moluccas, yield the volatile oil of Cajeput. It is a very liquid oil, of a grass-green colour, having a pungent camphoraceous odour, and capable of dissolving caoutchouc. It is used medicinally as a stimulant and antispasmodic. Species of *Eucalyptus* constitute the gigantic gum-trees of Australia, some of which attain a height of two hundred feet. They are remarkable for their operculate calyx, which may be considered as formed by several jointed leaves (like those of the orange), united throughout, and separating at the articulation in the form of a lid (§ 366). Their bark also separates remarkably in layers. They yield an astringent matter, which has been used for tanning. *Eucalyptus resinifera*, Brown Gum-tree of New Holland, furnishes Botany-Bay Kino, an astringent resinous-like substance, which exudes in the form of red juice from incisions in the bark. A single tree will yield sixty gallons. *E. mannifera* gives a saccharine exudation resembling manna. A saccharine substance, mixed with cellular hairs, which arise from a cup-like body, has been sent to this country by Mr. Cay, found upon the leaves of *E. dumosa*. It is called Layurp by the natives, and is thought by Mr. Newport to be the produce of an insect of the tribe Coccidæ. The wood of many species of *Metrosideros* is hard and dark-coloured. The flower-buds of *Caryophyllus aromaticus* (*Eugenia caryophyllata*), a tree which was originally a native of the Moluccas, but is now cultivated in the East and West Indies, constitute the Cloves of commerce. They have the form of a nail, and, when examined, are seen to consist of the tubular calyx with a roundish projection formed by the unopened petals. They contain a volatile oil, associated with resinous, gummy, and astringent matter. The oil is aromatic and acrid, and has been used as a condiment and a stimulant carminative. Pimento, Allspice, or Jamaica Pepper, is the berried fruit of *Eugenia Pimenta* (*Myrtus Pimenta*), a tree which is a native of the West Indies and Mexico. The fruit has an aromatic odour, and its taste combines that of cinnamon, nutmeg, and cloves; hence the name Allspice. It contains an acrid volatile oil, to which its properties are due. Medicinally Pimento is sometimes employed as a stimulant and carminative. The fruit of *Eugenia acris* is used for Pimento. Among the pulpy edible fruits of the order may be noticed Guavas and Rose-apples. The former are the produce of various species of *Psidium*, such as *P.*

pyriferum, *pomiferum*, and *Catleyanum*; the latter are procured from species of *Eugenia*, as *E. Jambos*, and *Malaccensis*. The berries of the common Myrtle (*Myrtus communis*) are also used as food. *Punica Granatum*, the Pomegranate-tree, is a native of the warmer parts of Asia and Northern Africa, whence it was introduced into Europe. It is the רמון of Scripture. It produces dark scarlet flowers, formerly called Balaustia, which have been used as an astringent. The fruit of the Pomegranate has given rise to much difference of opinion among botanists. It is composed, in the young state, of two rows of carpels, some of which are placed round the axis, and adhering to the bottom of the calycine tube, while others are placed outside, and adhere to the upper part of the tube. The subsequent contraction of the tube of the calyx, and the peculiar adhesion of the placentas, according to Lindley, account for the anomaly in the fruit (Balausta, ¶ 550). The rind of the fruit (malicorium), and the bark of the root, are used as anthelmintics, especially in cases of tape-worm. *Lecythis ollaria*, a large Brazilian tree, yields the woody capsules called Monkey-pots, which open by circumscissile dehiscence. These seed-vessels seem to be formed in the same way as the calyx of Eucalyptus, the part where the lid separates indicating the articulations of the carpellary leaves. The seeds are eatable, and are relished by monkeys. The bark of the tree may be separated into numerous thin layers. *Bertholletia excelsa* is the source of the Brazil nuts.

868. Order 80.—**Onagraceæ**, the Evening-Primrose Family. (*Polypet. Epigyn.*) Calyx tubular, the limb having usually 4 (fig. 399 *l*), sometimes 2, 3, or 6 divisions (fig. 531), which cohere in various ways; æstivation valvate. Petals usually equal in number to the calycine segments, regular (rarely irregular), inserted into the tube of the calyx (fig. 399 *p*); æstivation twisted. Stamens usually 4 or 8 (rarely 1 or 2, fig. 531), epigynous (fig. 399 *e*); filaments distinct; pollen triangular, usually cohering by threads. Ovary 2-4-celled (figs. 384, 531), adherent (fig. 399 *o*), usually with an epigynous disk; style filiform; stigma capitate (fig. 399 *s*) or 4-lobed; ovules (figs. 384 *o*, 399 *g*) indefinite, rarely definite, anatropal. Fruit succulent or capsular, dehiscent or indehiscent, 1-2-4-celled. Seeds usually ∞ , exalbuminous; embryo straight, with a long slender radicle pointing to the hilum, and short cotyledons.—Herbs or shrubs, with alternate or opposite, simple, not dotted leaves, and with the parts of the flower usually tetramerous. They inhabit chiefly temperate regions, and are found abundantly in Europe, Asia, and America, and sparingly in Africa. Some yield edible fruits, as *Fuchsia*, others furnish edible roots, as *Enothera biennis*. Many of them have mucilaginous properties, while a few are astringent. *Trapa* has unequal cotyledons. *T. natans*, Water Chestnut, and *T. bicornis*, remarkable for its horned fruit, both supply edible seeds. There are about 30 known genera,


and upwards of 450 species. *Examples*—*Oenothera*, *Epilobium*, *Jussiaea*, *Montinia*, *Fuchsia*, *Circæa*, *Gaura*, *Trapa*.

869. Order 81.—**Halorageæ**, the Mare's-Tail Family. (*Polypet. Epigyn.*) Calyx with a minute limb, which is either 3-4-divided or entire; it is sometimes reduced to a mere rim. Petals epigynous or 0. Stamens epigynous, equal in number to the petals, or twice as many, rarely fewer; when the petals are wanting, stamens often 1 or 2. Ovary cohering with the tube of the calyx, with 1 or more cells, sometimes tetragonal or compressed. Style 0, what is frequently called the styles being the papulose stigmas, which are equal in number to the cells; ovules pendulous, anatropal. Fruit dry, indehiscent, membranous or bony, with 1 or more cells. Seed solitary or in pairs, pendulous; albumen fleshy or thin; embryo straight, or slightly curved, in the axis of the albumen; cotyledons minute; radicle superior, long.—Herbs or undershrubs, often aquatic, with large air cavities, having alternate, opposite, or whorled leaves, and axillary, sessile flowers, which are occasionally unisexual. They are found in ditches and lakes in various parts of the world. They have no properties of importance. There are 8 known genera, and about 70 species. *Examples*—*Hippuris*, *Myriophyllum*, *Haloragis*, *Callitriche*.

870. Order 82.—**Loasacæ**, the Chili-Nettle Family. (*Polypet. Epigyn.*) Calyx 4-5-parted, persistent, spreading in æstivation. Petals 5, cucullate, epigynous, alternate with the segments of the calyx, sometimes with an inner row of 5, which are either similar to the outer or dissimilar; æstivation inflexed, valvate, or twisted. Stamens ∞ , in several rows, distinct, or polyadelphous, each parcel being opposite the outer petals; filaments subulate, unequal, the outer ones often sterile. Ovary inferior, 1-celled, with parietal placentas; ovules anatropal; styles combined into 1; stigma 1 or several. Fruit capsular or succulent, 1-celled. Seeds without an arillus; embryo straight, in the axis of fleshy albumen; cotyledons small, flat; embryo pointing to the hilum.—Herbaceous plants, hispid with stinging hairs, having opposite or alternate exstipulate leaves, and axillary 1-flowered peduncles. They are American plants, chiefly distinguished for their stinging qualities, and hence the name of Chili-Nettle. The roots of *Mentzelia hispida*, a Mexican herb, are said to possess purgative qualities. There are 15 genera enumerated by Lindley, including 70 species. *Examples*—*Loasa*, *Mentzelia*, *Gronovia*.

871. Order 83.—**Cucurbitacæ**, the Cucumber Family. (*Polypet. or Monopet. Epigyn. and Diclinal.*) Calyx 5-toothed (figs. 396 l, 606 c), sometimes obsolete. Petals 5, distinct, or more or less united, sometimes scarcely distinguishable from the calyx, strongly marked with reticulated veins (figs. 396 p, 606 p), sometimes fringed. Stamens 5, distinct or united in one or three parcels, attached to the petals (fig. 606 e); anthers bilocular, sinuous (figs. 332, 607, a); ovary (figs.

396 o, 608 c o) adhering to the tube of the calyx, 1-celled, formed by 3 carpels, and having 3 parietal placentas (fig. 609), which sometimes project so as to join in the centre, the ovules remaining attached to the curved free edges; ovules solitary or indefinite (fig. 609), anatropal; styles short; stigmas very thick, velvety or fringed (fig. 608 s).



Fruit a pepo (§ 551). Seeds flat and ovate (fig 610), enveloped in a juicy or dry and membranous covering; testa coriaceous; albumen 0; embryo straight (figs. 610 e, 611); cotyledons leafy and veined; radicle next the hilum.—Herbaceous plants, with succulent stems, climbing by means of lateral tendrils, which are transformed stipules; leaves alternate and palmate, covered with asperities; flowers generally unisexual. They are natives of warm climates chiefly, and abound in India. A few are found to the north in Europe and North America, and several are natives of the Cape of Good Hope. Those which are annuals readily submit to the climate of northern latitudes during the

Figs. 606-611.—Organs of fructification of Cucurbitaceæ.

Fig. 606.—Male flower of *Cucumis sativus*, Common Cucumber, laid open to show the interior of it. c, 5-divided calyx. p, United petals, by some considered as being an internal coloured calyx. e, Epigynous stamens.

Fig. 607.—Stamens separated. f, Filament. a, Long sinuous anther.

Fig. 608.—Female flower. c o, Calyx adherent to the ovary. p, United petals. s, Thick velvety stigmas.

Fig. 609.—Horizontal section of the ovary, showing its division into three, by projections from the parietal placentas, to which the indefinite ovules are attached.

Fig. 610.—Anatropal seed cut vertically. t, Spermoderm swollen at the chalaza. e, Embryo.

Fig. 611.—Embryo separated. r, Radicle. c, Cotyledons.

summer, and thus though of tropical origin, they grow well in European gardens. There are nearly 60 known genera, and about 300 species. *Examples*—*Cucurbita*, *Cucumis*, *Momordica*, *Bryonia*, *Telfairia*.

872. A certain degree of acridity pervades the order, and many of the plants are drastic purgatives. In some cases, however, more especially under cultivation, the fruits are eatable. Instances of edible fruits are seen in the case of the Melon, Cucumber, Gourd, Pumpkin, Squash, and Vegetable Marrow. The genus *Cucumis* contains the Melon and Cucumber, with edible fruits, and the *Colocynthis* with purgative fruit. Much discussion has taken place in regard to the structure of the fruit in this genus, and in *Cucurbitaceæ* in general. Some have considered it an anomaly in vegetable structure, from the apparent formation of the placenta and ventral suture, externally, as if the usual position of the carpels were reversed. It would appear, however, as shown by Lindley, that the placentas follow the ordinary law. They are parietal, and curve in a peculiar way, bearing the seeds on their curvature; at the same time prolongations are sent inwards, which often meet in the centre. Stocks and others consider the carpels as being involute, and they trace this involution particularly in *Luffa pentandra*. *Cucumis Colocynthis*, or *Citrullus Colocynthis*, yields a globular fruit called Coloquintida, or Bitter Apple, the pulp of which constitutes the medicinal *Colocynthis*. It is imported from the Levant and the coasts of the Mediterranean. It is used in the form of powder and extract as an irritant cathartic. The plant is supposed to be the *rypa*, or Wild Gourd of Scripture. *Momordica Elaterium* or *Ecbalium agreste*, the Wild or Squirting Cucumber, is so called on account of the force with which its seeds are expelled when ripe. The fruit, by a process of endosmose going on in the cells, becomes distended, and ultimately gives way at the weakest part, where the peduncle is united to it. In separating from the stalk, the elasticity of the parietes comes into play, so as to discharge the brown seeds and slimy juice through the aperture at the base of the fruit. The feculence which subsides from the juice constitutes the medicinal *Elaterium*, which is used in small doses of $\frac{1}{8}$ — $\frac{1}{2}$ a grain, as a violent cathartic, especially in dropsical cases. The active principle is *Elaterin*. The roots of *Bryonia alba* and *dioica* are also powerful purgatives. The fruit of various species of Gourd, as *Cucurbita Pepo*, the White Gourd, and *C. maxima* the Red Gourd, *C. ovifera*, the Vegetable Marrow, are used as potherbs; while *C. Citrullus*, the Water Melon, is prized for its cool refreshing juice. The fruit of *Lagenaria vulgaris*, in consequence of having a hard outer covering, is used as a vessel for containing fluid, after the pulp and seeds are removed. It is hence called Bottle Gourd. It is stated that poisoning has followed on the drinking of beer that had been standing in a flask made of one

of those Gourds. Dr. Royle mentions that symptoms of cholera have been induced by eating the bitter pulp. The seeds of the plants in this order frequently supply a bland oil. The seeds of *Telfairia pedata* are as large as Chestnuts, and are used as food.

873. Order 84.—**Papayaceæ**, the Papaw Family. (*Monopet. Polypet. Epigyn. and Declines.*) Calyx minute, 5-toothed. Corolla monopetalous, inserted into the base of the calyx; in the male, tubular and 5-lobed; in the female, divided nearly to the base into 5 segments. In the section Pangieæ the sepals and petals are distinct. Stamens 10, inserted into the throat of the corolla; anthers bilocular, introrse, innate, dehiscing longitudinally. Ovary free, 1-celled; ovules indefinite, attached to 5 parietal placentas; stigma 5-lobed, lacerated. Fruit usually succulent and indehiscent, sometimes capsular and dehiscent, 1-celled. Seeds ∞ , enveloped in a loose mucous coat, parietal; spermoderm brittle, pitted; embryo in the axis of fleshy albumen; cotyledons flat; radicle slender, turned towards the hilum.—Trees or shrubs, not branching, with alternate lobed leaves, supported on long slender petioles, and with unisexual flowers. They are found in South America, and in other warm countries. One of the most important plants of the order is *Carica Papaya*, the Papaw-tree, which yields an acrid milky juice, and an edible fruit. Its chief peculiarities are, that the juice of the unripe fruit is a powerful anthelmintic, (the powder of the seeds possessing the same properties). A constituent of this juice is fibrine, a principle formerly supposed peculiar to the animal kingdom and to Fungals. This tree is said to have the property of rendering meat tender, by causing a separation of the muscular fibre. Newly-killed meat suspended among the leaves, and even old animals when fed on the leaves and fruit, become tender in a few hours. The fruits of this plant, though handsome, scentless, and insipid, are untouched by birds or other animals except an ant belonging to the genus *Atta*. The order has been divided into three sections:—1. Cariceæ, corolla monopetalous, fruit succulent and indehiscent. 2. Modecceæ, corolla monopetalous, fruit capsular and dehiscent. 3. Pangieæ, corolla polypetalous. There are 11 known genera, including 29 species. *Examples*—*Carica*, *Modecca*, *Pangium*.

874. Order 85.—**Belvisiaceæ**, the Belvisia Family. (*Monopet. Epigyn.*) Calyx gamosepalous, persistent, limb divided into 5 thick ovate segments; æstivation valvate. Petals inserted in the tube of the calyx, united more or less, and forming 3 verticils, the innermost of which may be considered as an altered staminal row; the outer petaline verticil consists of 5 plaited lobes, each of which are 7-toothed, and have 7 feathered ribs; the second petaline verticil is cut into a number of narrow segments; while the third is an inconspicuous cup-like ring, with its edge minutely divided. Stamens ∞ , united at

their base so as to be monadelphous, or unequally polyadelphous; filaments curved inwards; anthers dithecal, oblong. Ovary surrounded by a fleshy disk, and adherent to the tube of the calyx, 5-celled; ovules 2 in each cell, attached to a central placenta, nucleus curved; style 5-angled; stigma broad, flat, pentagonal. Fruit a large fleshy rounded berry, crowned by the lobes of the calyx. Seeds large, kidney-shaped; cotyledons plano-convex; radicle and plumule immersed in their substance.—Shrubs, with alternate, simple, coriaceous, exstipulate leaves; and axillary flowers often in sets of three. They are tropical, chiefly African. The pulp of the fruit is eatable. The rind contains much tannin, and has been used for making ink. The wood is soft, and presents numerous dotted vessels. Their place in the natural system is not well determined; some placing the order next Passifloraceæ, others near Symplocaceæ, and Lindley recognizing its affinity to Rhizophoraceæ. There are 2 genera and 4 species. *Examples*—*Bervisia* (Napoleona), *Asteranthos*.

875. Order 86.—**Passifloraceæ**, the Passion-flower Family. (*Polypet. Perigyn.*) Sepals 5, combined below into a more or less elongated tube. Petals 5, perigynous, often with filamentous or annular processes on their inside, which appear to be an altered whorl or whorls of petals, occasionally wanting, imbricated in æstivation. Stamens 5, monadelphous, surrounding the gynophore when present, rarely ∞ , usually with processes from the thalamus, interposed between them and the petals; anthers dithecal, extrorse, versatile, dehiscing longitudinally; pollen-grains sometimes bursting by opercula (fig. 354). Ovary 1-celled, often with a gynophore (¶ 437); ovules anatropal, ∞ ; styles 3; stigmas dilated. Fruit often stipitate, 1-celled, sometimes 3-valved, opening by loculicidal dehiscence, or succulent and indehiscent. Seeds ∞ , attached to parietal placentas, arillate, or strophiolate; spermoderm brittle and sculptured; embryo straight in the midst of thin fleshy albumen; radicle pointing to the hilum.—Herbs or shrubs, often climbing, with alternate, stipulate, or exstipulate leaves. The order has been divided into three suborders:—1. *Paropsiæ*, plants not climbing, with a sessile ovary, arillate seeds, and exstipulate leaves. 2. *Passifloreæ*, climbing plants with a stalked ovary, arillate seeds, stipulate leaves, and glandular petioles. 3. *Malesherbieæ*, plants not climbing, with a stalked ovary, style below the apex of the ovary, strophiolate seeds, and exstipulate leaves. They are natives chiefly of warm climates, and are found in America, the East and West Indies. There are 14 known genera, and 215 species. *Examples*—*Paropsia*, *Smeathmannia*, *Passiflora*, *Tacsonia*, *Malesherbia*.

876. Considerable discussion has taken place as to the true nature of the calyx and corolla in Passifloraceæ. Lindley supports the view here given. Others consider the calyx as consisting of ten sepals in two rows, the inner more or less petaloid, and they look on the petals

as either wanting, or existing in the form of filamentous or annular processes. The name Passion-flower was given on account of a fancied resemblance in the flowers to the appearances presented at Calvary. In the five anthers the superstitious monks saw a resemblance to the wounds of Christ; in the triple style, the three nails on the cross; in the central gynophore, the pillar of the cross; and in the filamentous processes, the rays of light round the Saviour, or the crown of thorns. Many of the plants, such as *Passiflora quadrangularis* and *edulis*, Grenadillas, yield edible fruits, the pulp or succulent arillus being fragrant and cooling. The root of *Passiflora quadrangularis* is said to be emetic and powerfully narcotic, on which account Mr. Burnett declares it is cultivated in several French settlements. It is said to owe its activity to a peculiar principle called Passiflorine. Others are bitter and astringent, and some are said to be narcotic.

877. Order 87.—**Turneraceæ**, the Turnera Family. (*Polypet. Perigyn.*) Calyx with 5 equal lobes; æstivation imbricated. Petals 5, perigynous, equal; æstivation twisted. Stamens 5, perigynous, alternating with the petals; filaments distinct; anthers dithecal, innate, oblong. Ovary free, 1-celled, with 3 parietal placentas; ovules ∞ , anatropal; styles more or less cohering, or forked; stigmas multifid. Fruit a 1-celled, 3-valved capsule, dehiscent only half-way down, in a loculicidal manner. Seeds crustaceous, reticulated, arillate on one side; embryo slightly curved, in the midst of fleshy albumen; cotyledons plano-convex; radicle pointing to the hilum.—Herbaceous or somewhat shrubby plants, occasionally with stellate pubescence, having alternate, exstipulate leaves, and frequently two glands at the apex of the petiole. They are natives of the West Indies and South America. They are not put to any important use. *Turnera opifera* is astringent, and is employed in Brazil against dyspepsia. *Turnera ulmifolia* is considered a tonic and expectorant. Lindley gives 2 genera, including 60 species. *Examples*—*Turnera*, *Piriqueta*.

878. Order 88.—**Portulacaceæ**, the Purslane Family. (*Polypet. Perigyn.*) Sepals 2, cohering at the base. Petals usually 5, rarely wanting, distinct or cohering at the base, sometimes hypogynous. Stamens usually perigynous, variable in number, all fertile, sometimes opposite the petals; filaments distinct; anthers versatile, bilocular, with longitudinal dehiscence. Ovary free or partially adherent, 1-celled, formed by 3 united carpels; style single or 0; stigmas several. Fruit capsular, 1-celled, opening by circumscissile dehiscence, or by 3 valves, occasionally monospermous and indehiscent. Seeds numerous or definite, or solitary, attached to a central placenta; albumen farinaceous; embryo peripheral; radicle long.—Succulent shrubs or herbs, with alternate, seldom opposite, entire, exstipulate leaves, often having hairs in their axils. They are found in various parts of the world, chiefly, however, in South America and at the Cape of Good

Hope. They always inhabit dry parched places. They have a great affinity to Caryophyllaceæ, from which they are chiefly distinguished by their bisepalous calyx, perigynous stamens, and transversely dehiscent capsule. Lindley places this and the succeeding order among hypogynous Exogens, next to Caryophyllaceæ. The plants belonging to the order have few properties of importance. They are insipid and destitute of odour. *Portulaca oleracea*, common Purslane, is used as a potherb on account of its cooling and antiscorbutic qualities; the ancients thought the seeds, steeped in wine, to be an emmenagogue. The tuberous roots of *Claytonia tuberosa*, a Siberian plant, are eaten; and those of *Mellocca (Ullucus) tuberosa*, a native of Peru, have been recommended as a substitute for the potato. There are 12 known genera, and 184 species. *Examples*—*Portulaca*, *Talinum*, *Calandrinia*, *Claytonia*, *Montia*.

879. Order 89.—**Paronychiaceæ**, the Knotwort Family. (*Polypet. Perigyn.*) Sepals 4-5, distinct or cohering. Petals perigynous, between the divisions of the calyx, usually inconspicuous, sometimes 0. Stamens usually perigynous, sometimes hypogynous, opposite to the sepals when equal to them in number, some of them occasionally wanting; filaments distinct, rarely united; anthers bilocular. Ovary superior, with one or more ovules; styles 2-3, distinct or combined. Fruit unilocular, either a utricle covered by the calyx, or a 3-valved capsule. Seeds either numerous, attached to a free central placenta, or solitary and pendulous from a long funiculus arising from the base of the fruit. Embryo more or less curved, lying on one side of the farinaceous albumen, or surrounding it.—Herbaceous or somewhat shrubby plants, with opposite or alternate, sometimes setaceous and clustered leaves, which are either exstipulate or have scarious stipules. Found in barren places in various parts of Europe, Asia, and North America. A slight degree of astringency pervades this order, and is the only sensible property that it is known to possess. The order has been divided into two sections:—1. Illecebreæ, with the embryo lying on one side of the albumen, and stipulate leaves. 2. Scleranthæ, with a peripheral embryo, and exstipulate leaves. There are 28 known genera, and nearly 120 species. *Examples*—*Paronychia*, *Illecebrum*, *Polycarpon*, *Corrigiola*, *Scleranthus*.

880. Order 90.—**Crassulaceæ**, the Houseleek Family (figs. 535, 536). (*Polypet. Perigyn.*) Sepals 3-20, more or less united at the base (fig. 258 c c). Petals equal to the sepals in number, inserted in the bottom of the calyx (fig. 258 p p), either distinct or cohering in a gamopetalous corolla. Stamens inserted with the petals, either equal to them in number, and alternate with them (fig. 258 e e), or twice as many, those opposite the petals being shortest; sometimes one or two rows of abortive stamens; filaments distinct, or united, subulate, anthers bilocular, dehiscing longitudinally or transversely. Abortive

Crassulaceæ - Succulent leaves - Floral Organs numerous - lower throughout - Ovaries equal to Petals - Gillian's Paronychia.

stamens or scales (sometimes obsolete), at the base of each carpel (fig. 258 *a a*). Carpels equal in number to the petals and opposite to them, 1-celled (fig. 258 *o o*), sometimes consolidated; styles several or combined; stigmas pointed or 4-cornered; ovules 00, or definite, anatropal. Fruit consisting of several follicles, dehiscing by the ventral suture, sometimes by the dorsal suture. Seeds variable in number; embryo straight, in the midst of fleshy albumen; radicle pointing to the hilum.—Herbaceous plants or shrubs, often succulent, with simple, entire, or pinnatifid, exstipulate leaves. They are found in the driest situations, as on rocks, walls, and sandy plains, in various parts of the world. Some of them are acrid, as *Sedum acre*, Biting Stonecrop; others are refrigerant, from the presence of an acid, such as malic acid. *Sempervivum tectorum* is commonly known as the Houseleek. The fishermen of Madeira rub their nets with the fresh leaves of the *Sempervivum glutinosum*, by which the nets are rendered as durable as if tanned, provided they are steeped in some alkaline liquor. *Bryophyllum calycinum* is remarkable for the property of producing germinating buds at the edges of its leaves (§ 199). There are two suborders:—

1. Semperviveæ, with carpellary scales, numerous separate carpels in the pistil.
2. Galacieæ or Francoeæ, without true scales, pistil consolidated.

There are 25 genera, and about 460 species. *Examples*—*Crassula*, *Sempervivum*, *Cotyledon*, *Sedum*, *Penthorum*, *Galax*, *Francoa*.

881. Order 91.—**Ficoideæ**, or **Mesembryaceæ**, the Ficoid or Mesembryanthemum Family. (*Polypet. Perigyn.*) Sepals definite, usually 5, but varying from 4-8, more or less combined at the base, adherent to the ovary or distinct from it, equal or unequal; æstivation valvate or imbricate. Petals indefinite, coloured, sometimes 0. Stamens perigynous, distinct, definite or indefinite; anthers oblong, incumbent. Ovary usually many-celled; stigmas several, distinct; ovules 00, anatropal or amphitropal, attached by cords to the placenta, which is either central or parietal. Fruit a many-celled capsule, opening in a stellate or circumscissile manner at the apex, or an indehiscent nut. Seeds 00, rarely definite or even solitary; embryo curved or spiral on the outside of mealy albumen; radicle next the hilum.—Herbaceous or shrubby succulent plants, with opposite or alternate simple leaves. They are found in warm regions chiefly. The greater part of them grow at the Cape of Good Hope. The order has been divided into three sections:—

1. Mesembryeæ, numerous conspicuous petals, many-celled capsule, with stellate dehiscence.
2. Tetragonieæ, petals 0, fruit woody and indehiscent.
3. Sesuveæ, petals 0, capsule with circumscissile dehiscence.

There are 16 known genera, and 440 species. *Examples*—*Mesembryanthemum*, *Tetragonia*, *Aizoon*, *Sesuvium*.

Some of them are used as articles of diet, as the leaves of *Mesembryanthemum edule*, Hottentot's Fig, and *Tetragonia expansa*, New

Zealand Spinach. Others yield soda, and have been employed in the manufacture of glass. *Mesembryanthemum crystallinum*, the Ice-plant, is remarkable for the watery vesicles which cover its surface, and which have the appearance of pieces of ice. Its juice is said to be diuretic, and has been prescribed in dropsy and liver complaints. The seed-vessels of some species of *Mesembryanthemum*, as *M. Tripolium*, have the property of expanding in a star-like manner when put into water, and closing when dry. The flowers of many of the plants of the order exhibit the phenomenon of opening only under the influence of sunshine and closing in dull weather (§ 483).

882. Order 92. — **Cactaceæ**, the Cactus or Indian Fig Family. (*Polypet. Epigyn.*) Sepals numerous, usually ∞ , and confounded with the petals, adherent to the ovary. Petals numerous, usually indefinite, sometimes irregular, inserted at the orifice of the calyx. Stamens indefinite, cohering more or less with the petals and sepals; filaments long, filiform; anthers ovate, versatile. Ovary fleshy, inferior, unilocular; style filiform; stigmas numerous; ovules ∞ , attached to parietal placentas equal in number to the stigmas. Fruit succulent, 1-celled. Seeds ∞ , parietal, or, after losing their adhesion to the placenta, nestling in pulp, ovate or obovate; albumen 0; embryo straight, curved or spiral; cotyledons thick, leafy, sometimes nearly obsolete; radicle thick, obtuse, next the hilum.—Succulent shrubs, with peculiar angular or flattened stems, having the woody matter often arranged in wedges. Leaves usually absent; when present, fleshy, smooth, entire, or spinous. Flower sessile, sometimes showy. They grow in hot, dry, and exposed places, and are natives chiefly of the tropical parts of America. Some grow rapidly on the lava in volcanic countries. There are 16 known genera, and about 800 species. *Examples*—*Opuntia*, *Melocactus*, *Mammillaria*, *Echinocactus*, *Cereus*, *Epiphyllum*, *Pereskia*, *Rhipsalis*.

883. The plants of this order are remarkable for their succulence, for the great development of their cellular tissue, and the anomalous forms of their stems, some of which attain a great size. In their structure numerous spiral cells are met with, and in many cases the fibre in these cells is interrupted so as to present thickened rings united by membrance. These rings, when the cells are macerated, can be obtained in a free state. Many of the plants in this order show a remarkable tendency to spiral development. The setæ, spines, and hairs, are sometimes arranged spirally, and in *Cereus flagelliformis* the cells of the setæ have this tendency. Many of them yield an edible fruit, which is sometimes refreshing and agreeable, at other times insipid. The fruit of *Pereskia aculeata*, under the name of Barbadoes Gooseberry, is used in the West Indies as an article of diet. That of *Opuntia vulgaris* is known under the name of Prickly Pear. The juice of the fruit of some species is subacid, and has sometimes been used as a refrigerant.

Cattle sometimes feed on the succulent stems in dry seasons. Some of the plants are noted as night-flowering (§ 484). *Cereus grandiflorus* expands its large white blossoms about 10 P.M. in our hothouses, and their beauty lasts only for the night. Such is also the case with *Cereus nycticalus*. A plant of the latter species, in the Glasgow Botanic Garden, began to open its flowers between 7 and 8 P.M., and they were fully opened at 10. The following were the numbers and sizes of the various parts:—

Length of the tube of the calyx.....	7 inches.
Length of the petals.....	4½ —
Length of the style.....	10 —
Breadth of flower when fully expanded.....	11 —
Number of long sepals.....	75
Number of short sepals.....	20
Number of petals.....	25
Number of stamens.....	400
Number of stigmas.....	15

The size to which some of the Cactus family grow may be illustrated by a specimen of *Echinocactus Viznaga*, imported into Kew gardens from the mountains of San Luis, Potosi:—

Weight of the plant.....	713 lbs.
Height from surface of the earth.....	4½ feet.
Measured over the top from the ground on each side.....	10 feet 9 inches.
Circumference at 1 foot from the ground.....	8 feet 7 inches.
Number of deep angles or costæ.....	44
Number of spines.....	8800


In Brazil, some epiphytic Cactuses are met with; and there are some species described by Gardner as attaining a height of thirty feet, with a circumference of three feet. *Opuntia cochinellifera*, and other species, are infested by the Coccus Cacti, or the cochineal insect, which feeds upon them. The plants are cultivated in what are called nopaleries, for the sake of the insect, the females of which, when dried, constitute the cochineal of commerce.

884. Order 93.—**Grossulariaceæ**, the Gooseberry and Currant Family. (*Polypet. Epigyn.*) Calyx 4-5-cleft, regular, coloured. Petals minute, perigynous, equal in number to the segments of the calyx, and alternate with them. Stamens 4-5, alternate with the petals, and inserted into the throat of the calyx; filaments short; anthers dithecal. Ovary unilocular, adherent to the tube of the calyx; ovules ∞, anatropal, attached to two opposite parietal placentas; style single, 2-4-cleft. Fruit a 1-celled berry, crowned with the remains of the flower. Seeds ∞, immersed in pulp, and attached to the placentas by long thread-like funiculi; spermoderm gelatinous externally; albumen horny; embryo straight, minute; radicle pointing to the hilum.—Shrubs, with alternate lobed leaves, having a

plicate vernation. They are natives of temperate regions, and are found in Europe, Asia, and America. Many yield edible fruits, which sometimes contain malic acid. The various kinds of Gooseberry (*Ribes Grossularia*), and Currant (*Ribes rubrum* and *nigrum*) belong to this order. The black currant possesses tonic and stimulant properties. On the under surface of its leaves and flowers fragrant glands may be perceived. It contains 2 or 3 genera, and nearly 100 species.
Example—*Ribes*.

885. Order 94.—**Saxifragaceæ**, the Saxifrage Family. (*Polypet. Perigyn.*) Calyx superior, or more or less inferior (fig. 397 *cc*); sepals usually 5, more or less cohering at the base. Petals usually 5, perigynous, alternate with the lobes of the calyx (fig. 397 *pp*), rarely 0. Stamens perigynous (fig. 397 *e*), 5-10 or ∞ , in 1 or more rows; anthers bilocular, with longitudinal or porous dehiscence. Disk often present, either annular or scaly. Ovary more or less completely united to the tube of the calyx, consisting usually of two carpels, cohering by their face (figs. 397, 398 *o*), but distinct and diverging at the apex; styles as many as the carpels, distinct (fig. 398 *t*) or combined; stigmas capitate (fig. 398 *s*) or clavate. Placentas (fig. 398 *p*) marginal (basal or apicular), rarely central. Fruit generally a 1-2-celled capsule, the cells dehiscing at the ventral suture, and often divaricating when ripe. Seeds usually ∞ , rarely definite; spermoderm often reticulated; embryo small, in the axis of fleshy albumen; radicle pointing to the hilum.—Shrubs or trees, or herbs, with alternate or opposite, usually exstipulate leaves. They are generally natives of temperate climates, and some of them characterize alpine districts. The order has been divided into the following suborders:—1. Escalloniæ, petals and stamens 5; ovary inferior; style simple; albumen oily; evergreen shrubs, with alternate, simple, exstipulate leaves, found in the temperate regions of South America, often at a great elevation. 2. Cunoniæ, petals 4-5 or 0; stamens 8-10 or ∞ ; ovary half inferior; styles 2, distinct or combined; trees or shrubs, with opposite leaves, having interpetiolar stipules; found in South America, the East Indies, south of Africa, and Australasia. 3. Hydrangeæ, petals 4-6; stamens 8-12 or ∞ ; anthers sometimes biporose; ovary more or less inferior; styles 2-5, usually distinct; shrubs with opposite, sometimes whorled, exstipulate leaves, and flowers, frequently cymose, with the exterior flowers sterile and dilated; found chiefly in the temperate parts of Asia and America. 4. Saxifrageæ, petals 5 or 0; stamens 5-10; ovary more or less adherent; styles usually 2, and distinct; herbs, with alternate, usually exstipulate leaves, found in the mountainous regions of Europe, &c. Few of the plants are put to any use. Some of them are astringent, and used for tanning; others have bitter tonic properties. The glutinous exudation of a few of them is acrid. In the entire order there are 57 known genera, and upwards of 900

traversed by five primary longitudinal ridges (jugæ), and often by four alternating secondary ones, the ridges being separated by channels



(valleculæ). In the substance of the pericarp there are frequently vittæ containing oil, which are usually opposite the channels. Seeds pendulous (fig. 616 *g*), usually adherent to the pericarp, rarely loose; embryo minute, at the base of abundant horny albumen (fig. 616 *e*); radicle pointing to the hilum.—Herbaceous plants, often with hollow and furrowed stems, with alternate, rarely opposite, variously divided, sheathing leaves, (which sometimes assume the appearance of phyllodia), and with umbellate, involucrate flowers (fig. 241). They are found chiefly in the northern parts of the northern hemisphere. In warm countries they occur at high elevations, The order has been divided according to the number and size of the pericarpial ridges, the presence or absence of vittæ, and the form of the albumen. The following are the several sections, which, however, are not considered altogether satisfactory:—1. Orthospermæ (ὀρθός, straight, and σπέρμα, seed), albumen flat on the inner face, neither involute nor convolute.

Figs. 612-616.—Organs of fructification of *Daucus Carota*, common Carrot, to illustrate the natural order Umbellifereæ.

Fig. 612.—Diagram of the flower, with a 5-toothed calyx, 5 inflexed petals, 5 stamens, and fruit formed by 2 carpels, with primary and secondary ridges, valleculæ, commissure, and flat albumen.

Fig. 613.—The flower viewed from above, showing the petals with inflexed points and 5 stamens. *g e*, Epigynous disk or stylopod.

Fig. 614.—Vertical section of the flower. *p*, Petals with inflexed points. *e*, Stamens, one incurved at the apex. *o*, Ovary formed by two carpels, adherent to the calyx throughout. *s*, Styles and stigmas. *g e*, Epigynous disk or stylopod.

Fig. 615.—Horizontal section of the fruit (cremocarp) with bristly ridges.

Fig. 616.—Vertical section of the cremocarp. *f*, Pericarp. *g*, Seed. *p*, Flat perisperm. *e*, Embryo.

2. *Campylospermæ* (καμπύλος, inflected), albumen curved at the margins, so as to form a longitudinal furrow. 3. *Cœlospermæ* (κοίλος, concave), albumen curved at the ends (from base to apex). Lindley enumerates 267 genera, including 1500 species. *Examples*—*Hydrocotyle*, *Eryngium*, *Apium*, *Bupleurum*, *Æthusa*, *Angelica*, *Heracleum*, *Daucus*, *Myrrhis*, *Conium*, *Coriandrum*.

889. The properties of the plants of this order are various. Some yield articles of diet, others gum-resinous and oily substances, while others are highly poisonous. According to their qualities, the species have been divided into—1. Those which are harmless, and are used as esculent vegetables. 2. Those producing a gum-resin, often having a fetid odour from the presence of a sulphur-oil, and which are used as antispasmodics and stimulants. 3. Those yielding a volatile oil, which renders them carminative and aromatic. 4. Those which are poisonous, in consequence of the presence of an acrid and narcotic juice.

890. Among esculent species may be noticed—*Daucus Carota* (Carrot), *Pastinaca sativa* (Parsnip), *Apium graveolens* (Celery), *Feniculum vulgare* (Fennel), *Petroselinum sativum* (Parsley), *Anthriscus Cerefolium* (Chervil), *Sium Sisarum* (Skirret), and *Archangelica officinalis* (Angelica). *Criihum maritimum* is the Samphire, which grows abundantly on rocks near the sea, and is used as a pickle. The roots of *Arracacha esculenta*, a native of Grenada, have been recommended as a substitute for the potato; they are large and esculent, resembling a Parsnip in quality. The root of *Eryngium campestre* and *maritimum*, or Eryngo, is sweet, aromatic, and tonic. Boerhaave reckons it as the first of aperient diuretic roots. The tubers of *Bunium Bulbocastanum* and *flexuosum* are eaten under the name of Pig-nuts or Earth-nuts. *Prangos pabularia*, a plant of Southern Tartary, is highly recommended as fodder for cattle.

891. Many species yield milky juices, which concrete into a fetid gum-resin. Assafœtida is procured from the *Ferula Assafœtida* of Linnæus, which has recently been described by Dr. Falconer under the name of *Narthex Assafœtida*. The plant is found in Persia and Afghanistan, and seeds of it have been sent to this country by Dr. Falconer, some of which have germinated in the Edinburgh Botanic Garden, and are now vigorous thriving plants of ten years' growth. The fruit of the plant is distinguished by divided and interrupted vittæ, which form a network on the surface, and its leaves have a resemblance to those of a Pæony. It would appear that *Ferula persica*, a plant with very much divided leaves, yields an inferior sort of assafœtida. The assafœtida is procured by taking successive slices off the top of the root, and collecting the milky juice which is allowed to concrete in masses. It consists of resinous and gummy matter, with a sulphur-oil similar to that of Garlic, which is probably its active ingredient. It is employed medicinally in substance or tincture, as a

stimulant, antispasmodic, and anthelmintic. Ammoniac, another fetid gum-resin, is the produce of *Dorema Ammoniacum* (probably *Diseroneston gummiferum* of Jaubert and Spach), a native of Persia. It contains resin, gum, and volatile oil, and is used medicinally as a stimulant, antispasmodic, and expectorant. Galbanum, which seems to be the גלגלית of Scripture, is procured, in all probability, from *Opoidea galbanifera* of Lindley. There are doubts, however, as to its botanical source. Don referred it to *Galbanum officinale*. It consists of resin, gum, and volatile oil, and is used as an antispasmodic and emmenagogue. Opoponax is another gum-resin, procured from *Opoponax Chironum* (*Pastinaca Opoponax*), a native of the southern parts of Europe. Sagapenum seems to be the produce of a species of *Ferula*.

892. There are other species which supply a carminative and aromatic oil. From the fruits of *Carum Carui*, which are commonly called Caraway seeds, a volatile oil of this nature is procured. Oils of a similar kind are obtained from the fruit of *Pimpinella Anisum* (Anise), from that of *Faniculum vulgare*, or *F. dulce* (common Fennel), *Anethum graveolens* (common Dill), *Coriandrum sativum* (Coriander), the כוסברה of the Bible, *Cuminum Cyminum* (Cumin), *Archangelica officinalis* (Garden Angelica), and *Daucus Carota* (Carrot).

893. In regard to the poisonous species of this order, there is still much to be learned. They appear to vary according to the soil and climate in which they grow. Some species, generally reputed poisonous, have been found by Dr. Christison to be quite innocuous when gathered from localities in the neighbourhood of Edinburgh. The most important plant of this section is *Conium maculatum* (Hemlock), the κώνειον of the Greeks. It is a biennial plant, found abundantly in Britain, and distinguished by its undulated ridges, smooth purple-spotted stem, and the peculiar mouse-like odour of its leaves, when being dried. Every part of the plant, especially the fresh leaves and green fruit, contain a volatile oleaginous alkali, called Conia, which acts as an energetic poison. To this substance the effects of hemlock on the animal frame are due, and care is required in the preparation of the leaves and fruit in order to retain this active principle. A few drops of Conia will kill a small animal. It acts on the spinal cord, producing paralysis with slight convulsive twitches, and its fatal effects are attributed to asphyxia, produced by palsy of the muscles of respiration, without convulsions or coma. Hemlock has been employed medicinally to allay pain, more especially in cancerous and neuralgic affections. *Ceanothe crocata* (Hemlock-Dropwort, or Dead-tongue), and a variety called *apiifolia*, have been long looked upon as poisonous. The roots have been mistaken for parsnips, and fatal effects have been thus produced. It would appear, however, that these poisonous qualities are not invariably present, for Dr. Christison found that the

roots of this plant, when growing in a sea-side locality, near Edinburgh, were innocuous. It remains to be determined if the climate and locality have any effect in modifying the properties of the plant. The same remarks may be made in regard to *Enanthe Phellandrium* (Water Dropwort), and *Cicuta virosa* (Water Hemlock or Cowbane), which seem to vary as regards their poisonous properties. *Æthusa Cynapium* (Fool's Parsley), is another plant of the order reputed poisonous. It has been stated that the roots of Parsnip, during the spring of the second year, on the approach of the flowering season, occasionally produce a poisonous matter.

894. Order 98.—**Araliaceæ**, the Ivy Family. (*Polypet. Epigyn.*) Calyx entire or toothed (fig. 309 c). Petals definite (fig. 309 p), 2, 5, 10, deciduous, occasionally 0; æstivation valvate. Stamens as many as the petals, or twice as many, inserted below the margin of an epiginous disk (fig. 309 e e). Ovary adherent to the tube of the calyx, 2 or more celled (fig 309 o); ovules solitary, pendulous (fig. 309), anatropal; styles 2 or more, distinct or connate (fig. 309 s); stigmas simple. Fruit usually succulent, 2-15-celled, covered by the calycine limb. Seeds solitary, pendulous, adhering to the endocarp; albumen fleshy; embryo small; radicle pointing to the hilum.—Trees, shrubs, or herbaceous plants, with alternate exstipulate leaves, and umbellate (fig. 240) or capitate flowers. They are found both in tropical and in cold regions. Lindley enumerates 21 genera, including 160 species. *Examples*—*Aralia*, *Panax*, *Adoxa*, *Hedera*.

895. They have generally aromatic and stimulant properties. They are allied to Umbelliferæ, but do not possess poisonous qualities in a marked degree, nor does their fruit usually yield volatile oil. A species of *Panax* yields the famous Ginseng root of the Chinese, which is used as a stimulant. The celebrated Rice Paper of the Chinese is ascertained to be prepared from the pith of a plant of this order, called by Sir William Hooker, *Aralia papyrifera*. *Panax quinquefolium* possesses qualities resembling those of ginseng. Some species of *Aralia* yield an aromatic gum-resin. *Aralia nudicaulis*, a native of North America, has fragrant and aromatic roots, which are used as a substitute for sarsaparilla. *A. spinosa*, called toothache-tree in North America, is a stimulant diaphoretic. The succulent fruit of *Hedera Helix*, the Ivy, is emetic and purgative.

896. Order 99.—**Cornaceæ**, the Cornel Family. (*Polypet. Epigyn.*) Calyx 4-lobed. Petals 4, oblong, broad at the base, regular, inserted into the upper part of the calycine tube; æstivation valvate. Stamens 4, inserted along with the petals, and alternate with them; anthers dithecal. Ovary adherent to the tube of the calyx, 2-celled, crowned by a disk; ovules solitary, pendulous, anatropal; style filiform; stigma simple. Fruit fleshy, crowned by the limb of the calyx, 2-celled, rarely 1-celled by abortion; endocarp bony (fig. 473). Seeds solitary,

pendulous; embryo straight, long, in the axis of fleshy albumen; radicle superior, shorter than the oblong cotyledons.—Trees, shrubs, or herbs, with opposite, very rarely alternate, exstipulate leaves, and capitate, umbellate, or corymbose flowers. They inhabit the temperate climates of Europe, Asia, and America. The bark of *Cornus florida* and *sericea* is used in America as a tonic and febrifuge. The young branches of *Cornus florida*, stripped of their bark, and rubbed with their ends against the teeth, acts as a dentifrice. The fruit of *Cornus mascula* has been used as food, and the seeds of *Cornus sanguinea* furnish oil. The fruit of *Cornus suecica*, a species found on the Scotch mountains, is reputed tonic. *Aucuba japonica* has leaves which exhibit a variegated aspect. Lindley gives 9 genera, and 40 species. *Examples*—*Cornus*, *Aucuba*.

Subclass III.—COROLLIFLORÆ.

897. Calyx and corolla present; petals united, bearing the stamens. This subclass includes the Monopetalous orders of Jussieu, and the Gamopetalæ of Endlicher. De Candolle restricted this subclass to those orders in which the corolla was hypogynous, placing the other orders with united petals in the subclass Calycifloræ.

898. Order 100.—**Loranthaceæ**, the Mistletoe Family. (*Monopet. Epigyn.*) Calyx arising from a tube, or rim, which some regard as an expansion of the pedicel, often bracteated. Petals (or according to others, sepals) 4-8, distinct, or more or less united; æstivation valvate. Stamens equal in number to the petals, and opposite to them; filaments more or less united to the petals; anthers 1- 2- or many-celled (¶ 405). Ovary unilocular, adherent to the calycine tube or the expanded pedicel; ovules with a naked nucleus, erect or suspended; style filiform or 0; stigma simple. Fruit succulent, crowned by the calyx, 1-celled. Seed solitary, pendulous; embryo straight, in the axis of fleshy albumen; cotyledons either minute or numerous; radicle superior.—Shrubs, usually parasitical, with opposite or alternate, fleshy exstipulate leaves. Many of the plants have showy flowers, which hang from the trunks and branches of trees in the equinoctial parts of Asia and America. Some occur in temperate regions. Lindley gives 23 genera, and 412 species. *Examples*—*Loranthus*, *Viscum*, *Myzodendron*.

899. Disputes have taken place as to the structure of the flowers in this order, some considering the petals as being in reality sepals, and regarding the calycine rim as being an expansion of the pedicel only. The wood of some of the plants is arranged in separate wedges, and their vessels are either annular or scalariform. The fruit contains a viscid matter, like bird-lime, by means of which the seeds adhere to

trees. The seeds in germinating send their radicles into the plant to which they are attached, and grow afterwards as true parasites, selecting certain chemical ingredients in preference to others. The bark is usually astringent. Griffith has carefully described the nature of the parasitism of those plants. He states that in *Loranthus*, the ripe seeds adhere firmly to the substance on which they are applied by means of their viscid coating, which hardens into a transparent glue. In two or three days after application, the radicle curves towards its support, and as soon as it reaches it becomes enlarged and flattened. By degrees a union is established between the woody system of the parasite and stock, after which the former lies exclusively on the latter, the fibres of the sucker-like root of the parasite expanding on the wood of the support. Before this occurs the parasite is nourished by its own albumen, which is gradually absorbed. "As soon as the young parasite has acquired the height of one or two inches, when an additional supply of nourishment is perhaps required, a lateral shoot is sent out, which is, especially towards the point, of a green colour. This at one, or two, and subsequently at various points, adheres to the support by means of sucker-like productions, which are precisely similar in structure and mode of attachment to the original seminal one." The fibres of the parasite never penetrate beyond their original attachment; in the adult the sucker-bearing shoots frequently run to a considerable distance. "I have seen," says Mr. Griffith, "such shoots which had taken their course *along a decayed branch* become replaced, and *return in quest*, as I may express it, *of a part capable of affording some nourishment*." *Viscum album*, Mistletoe, was called by the Druids the Mistletoe of the Oak, on which, however, it is rarely found parasitic. It grows well on the apple-tree. The formation of the ovule in the Mistletoe, according to Schleiden, is described at ¶ 463. *Loranthus tetrandrus* is used in Chili to dye black.


900. Order 101.—**Caprifoliaceæ**, the Honeysuckle Family. (*Monopet. Epigyn.*) Calyx with its limb 4-5 lobed, usually bracteated. Corolla superior, lobed, usually regular and gamopetalous, sometimes irregular. Stamens epicorolline, equal in number to the lobes of the corolla, and alternate with them. Ovary adherent to the tube of the calyx, usually 3-celled, rarely 4-5-celled; ovules few in each cell, pendulous; style one or none; stigmas 3-5. Fruit fleshy or dry, crowned by the limb of the calyx, indehiscent, uni- or multilocular; endocarp sometimes bony. Seeds solitary, or several in each cell, pendulous; spermoderm often bony; embryo small in the centre of fleshy albumen; radicle next the hilum.—Shrubs or herbs, with opposite exstipulate leaves, and corymbose flowers. Chiefly found in the northern parts of Europe, Asia, and America; found very sparingly in northern Africa, and little known in the southern hemisphere. The order has been divided into two suborders:—1. *Lonicereæ*, the

Caprifoliaceæ
Epigyn.
Corolla superior
Stamens equal
Ovary 3-celled
Fruit fleshy or dry
Indehiscent
Uni- or multilocular
Endocarp sometimes bony
Seeds solitary or several
Embryo small
Radicle next hilum
Shrubs or herbs
Opposite exstipulate leaves
Corymbose flowers
Chiefly found in northern parts of Europe, Asia, and America
Found very sparingly in northern Africa
Little known in the southern hemisphere

true Honeysuckles, with a regular rotate or tubular corolla, three sessile stigmas, and a raphe on the inner side of the ovule. 2. *Sambuceæ*, the Elder Tribe, with a corolla more or less tubular, often irregular, a filiform style, and a raphe on the outside of the ovule. Lindley gives 14 genera, and 220 species. *Examples*—*Lonicera*, *Caprifolium*, *Leycesteria*, *Linnaea*, *Sambucus*, *Viburnum*.

901. Many of the plants, such as the Honeysuckle and Elder, have odoriferous flowers. Some possess emetic and purgative properties. The fruit of *Sambucus nigra*, the Common Elder, is used in the manufacture of a kind of wine. The flowers contain a small quantity of concrete volatile oil, and a minute portion of a volatile odoriferous oil. The inspissated juice of the fruit, and the inner bark, possess purgative qualities. *Viburnum Opulus*, the Guelder Rose, is often cultivated in gardens. *Viburnum Lantana* has an acrid bark. *Linnaea borealis* is a delicate northern plant, named after Linnæus. *Symphoricarpos racemosa* yields the Snowberry.

902. Order 102.—**Rubiaceæ**, the Madder and Peruvian Bark Family. (*Monopet. Epigyn.*) Calyx adherent with the ovary, the limb with a definite number (usually 4-5) of divisions, sometimes obsolete (fig. 619 *c*). Corolla gamopetalous, regular, tubular, or rotate (fig. 619 *p*), inserted into the calyx, usually with 4-5 divisions (fig.



618); æstivation valvate or imbricate. Stamens more or less adherent to the corolline tube, as many as the lobes of the corolla, and alternate

Figs. 617-622.—Illustrations of the natural order Rubiaceæ.

Fig. 617.—Diagram of the flower of *Galium Mollugo*, belonging to the section *Stellateæ*. Calyx nearly obsolete, corolla rotate, 4-lobed, 4 stamens, and didymous ovary.

Fig. 618.—Flower entire.

Fig. 619.—Flower cut vertically. *c*, Calyx adherent to the ovary, *o*, which is 2-celled. *p*, Corolla. *e e*, Stamens surrounding the style and stigmas.

Fig. 620.—Fruit of *Rubia tinctoria*, Madder.

Fig. 621.—The same, showing the separation of the two carpels.

Fig. 622.—The seed cut vertically. *p*, Perisperm. *e*, Curved embryo.

with them (fig. 617). Ovary inferior, usually bilocular (fig. 619 o), sometimes multilocular, crowned with a fleshy disk; ovules numerous or solitary, anatropal or amphitropal; style single, sometimes partly divided; stigmas usually 2, more or less distinct (fig. 619). Fruit inferior, 2- or many-celled, dry or succulent, either indehiscent or splitting into two mericarps (figs. 620, 621). Seeds 1 or many in each cell, in the former case erect or ascending (fig. 619), in the latter attached to a central placenta; albumen copious, horny or fleshy (fig. 622 p); embryo small, straight, or slightly curved (fig. 622 e); cotyledons leafy; radicle turned to the hilum.—Trees, shrubs, or herbs, with simple, entire, opposite, or verticillate leaves, which have either interpetiolar stipules (fig. 190), or are exstipulate. The order has been divided into two suborders:—1. Cinchoneæ, with rounded stems, usually opposite leaves, and interpetiolar stipules (fig. 190), natives of the hotter parts of the world. 2. Galieæ, or Stellatæ, with square stems, verticillate leaves, and no stipules; natives of northern and colder regions. Lindley considers these divisions as separate natural orders. Some authors think that the verticillate leaves of Stellatæ consist partly of true leaves, and partly of stipules. The order includes nearly 280 genera, and upwards of 2800 known species. *Examples*—*Cinchona*, *Gardenia*, *Hedyotis*, *Isertia*, *Hamelia*, *Guettarda*, *Pæderia*, *Coffea*, *Cephælis*, *Psychotria*, *Spermacoce*, *Anthospermum*, *Opercularia*, *Galium*, *Rubia*.

903. The properties of the order, in general, are tonic, febrifuge, and astringent. Many important articles of materia medica are furnished by the plants in the suborder Cinchoneæ. Peruvian or Jesuits' Bark, known under the name of Pale, Yellow, and Red Bark, is procured from various species of *Cinchona*, which grow abundantly in the district of Upper Peru. The *Cinchona* trees seem to be confined exclusively to the Andes, within the boundaries of Peru, Columbia, and Bolivia, from 11° north lat. to 20° south lat., chiefly growing at elevations varying from 1,200 to 10,000 feet above the level of the sea, and in a dry rocky soil. There are at least twelve species which are supposed to furnish the barks of commerce. Great obscurity prevails as to the species whence the various kinds of *Cinchona* bark are derived. The names of Yellow, Red, and Pale Bark have been very vaguely applied, and are by no means well defined. The barks are met with either in thick, large, flat pieces, or in thinner pieces, which curl inwards during drying, and are called quilled. Britain is said to import from 225,000 to 556,000 lbs. annually, and to retain 120,000 lbs. for home consumption. The chief officinal kinds of bark are: 1. Crown-bark, China-loxa, a pale bark in quills 6 to 15 inches long, the produce of *Cinchona Condaminea*, which inhabits the mountains in the vicinity of Loxa, at an elevation of from 5,700 to 7,500 feet. 2. Gray bark, Silver bark, or Huanuco bark, China-Huanuco, another

variety of quilled pale bark, obtained from *Cinchona micrantha*, found in the vicinity of Huanuco, in Peru. 3. Yellow bark, China-regia, or Calisaya bark, partly flat, partly quilled, procured from *Cinchona*, Calisaya, which grows around Apolobamba. 4. Red bark, China-rubra, partly flat and partly quilled, procured from an unknown species of *Cinchona*. Besides these, there are various inferior kinds of bark met with in commerce, such as Ash bark, China-Jaen, from *Cinchona ovata*; hard Carthagena bark, China-flava-dura, from *Cinchona cordifolia*; Rusty bark, China-Huamalies, from *Cinchona pubescens*; Orange-bark, from *Cinchona lancifolia*; and Red bark of Santa-Fé, from *Cinchona magnifolia*. The following is the arrangement adopted by Pereira:—

A. True Cinchonas with a Brown Epidermis.

I. Pale Barks.

1. Crown or Loxa Bark.....*Cinchona Condaminea*, *H.* and *B.*
2. Gray, or Silver, or Huanuco Bark.....*Cinchona micrantha*, *R.* and *P.* (*C. scrobiculata*, *H.* and *B.*)
3. Ash or Jaen Bark.....*Cinchona ovata*, *Fl. Peruv.*
4. Rusty or Huamalies Bark.....*C. pubescens*, *Vahl.* (*C. purpurea*, *R.* and *P.*)

II. Yellow Barks.

5. Royal, Yellow, or Calisaya Bark.....*Cinchona* sp.?

III. Red Barks.

6. Red Bark.....*Cinchona* sp.?

B. True Cinchonas with a White Epidermis.

I. Pale Barks.

7. White Loxa Bark.....*Cinchona* sp.?

II. Yellow Barks.

8. Hard Carthagena Bark.....*Cinchona cordifolia*, *Mutis.*
9. Fibrus Carthagena Bark.....Perhaps *C. cordifolia*.
10. Cuzco Bark.....*Cinchona* sp.?
11. Orange Bark of Santa-Fé.....*C. lancifolia*, *Mutis* (*C. angustifolia*, *P.*)

III. Red Barks.

12. Red Bark of Santa-Fé.....*C. oblongifolia*, *Mutis* (*C. magnifolia* *R.* and *P.*)

Cinchona bark contains two alkaloids, Cinchonina and Quina, to which its active properties are due; the former is best obtained from Gray bark, the latter from Yellow bark. In combination with these, there exists an acid called Cinchonic or Kinic acid. *Cinchona* bark is used medicinally as a tonic and antiperiodic, in cases of dyspepsia, neuralgia, and intermittent fever. It has been administered in the form of infusion and tincture; but, at present, the disulphate of Quina is the

preparation used. The genus *Exostemma* yields various kinds of Cinchona bark, which do not contain the Cinchona alkalis. In genus the stamens are exserted, whereas in Cinchona they are ed. The following are some of the false barks noticed by a:—

1. St. Lucie or Piton Bark.....*Exostemma floribundum*.
2. Jamaica Bark.....*E. caribæum*.
3. Pitaya Bark.....*Exostemma* sp. ?
4. False Peruvian Bark.....*E. peruvianum*.
5. Brazilian Bark.....*E. souzianum*.

Pinckneya pubens yields the fever-bark of Carolina.

904. Some of the plants of this order have emetic and purgative qualities. *Cephælis Ipecacuanha* yields the Ipecacuanha of the Pharmacopœia. The plant is found in the woods of several Brazilian provinces, as Pernambuco, Bahia, and Rio Janeiro. The roots which are the officinal part, are contorted, knotty, and annulated, and about the thickness of a goose-quill. They are used as emetic and diaphoretic remedies, in the form of powder or wine. Their active ingredient is an alkaloid called Emetia or Emetine. Besides this brown or gray annulated Ipecacuanha, there are spurious kinds, such as striated or Black Peruvian Ipecacuanha, the produce of *Psychotria emetica*, and white or amylaceous Ipecacuanha, furnished by *Richardsonia scabra* (*brasiliensis*), a native of the provinces of Rio Janeiro and Minas Gerães. Some of the species of *Psychotria*, *Cephælis*, and *Randia*, are said to act so violently as to produce poisonous effects.

905. Among the astringent plants of the order may be noticed *Uncaria Gambir*, which supplies a kind of Catechu, known by the name of Gambeer. Of the plants furnishing articles of diet, the most important is *Coffea arabica*, a native of Arabia and of the borders of Abyssinia, which furnishes the Coffee of commerce. The fruit is succulent, and the horny albumen of the seed is the part used as a beverage. It contains a bitter principle, denominated Caffein, which is identical with that got from Tea. The import of Coffee into the United Kingdom, in 1847, was 19,783 tons, and in 1848, it was 24,553 tons. The seeds of some other plants of the order, as species of *Galium*, have been used as substitutes for Coffee. Among the plants yielding dye, the most interesting is *Rubia tinctoria*, the root of which is the Madder of commerce. It contains three volatile colouring matters—madder purple, orange, and red. The latter is in the form of crystals, having a fine orange-red colour, and called Alizarine. This is the substance which yields the turkey-red dye. *Rubia Munjista*, (*cordifolia*), Munjeet, is also used for a similar purpose. *Oldenlandia umbellata* is employed in the East Indies as a substitute for Madder, and so is the root of *Morinda citrifolia*, under the name of Sooranjee. The latter yields a peculiar colouring matter, called by Dr. Anderson, Morindine. It

is extracted from the bark of the root, and is procured in the form of minute acicular crystals of a fine yellow colour. It is incapable of producing colours with alum and iron mordants, but with turkey-red mordant it produces a dark red. Many of the plants of this order, especially in the section Cinchoneæ, have very showy and fragrant flowers. The species of *Mussaenda* and *Calceophyllum*, are remarkable on account of one of their sepals becoming large and showy. *Asperula odorata*, Wood-ruff, gives out its fragrance when dried.

906. Order 103.—**Valerianaceæ**, the Valerian Family. (*Monopet. Epigyn.*) Calyx superior, its limb being either membranous or pap-pose. Corolla gamopetalous, inserted into the top of the ovary, tubular, 3-4-5-lobed, sometimes gibbous or spurred at the base. Stamens 1-5, adherent to the corolla and alternate with its lobes. Ovary inferior, 1-3-celled; ovule solitary, pendulous, style filiform; stigmas 1-3. Fruit dry, indehiscent, crowned by the limb of the calyx, 1-celled, in consequence of 2 cells being abortive. Seed solitary, pendulous, exalbuminous; embryo straight; radicle superior.—Herbs, with opposite exstipulate leaves, and cymose inflorescence. They are found in temperate climates. Lindley gives 12 genera, and 185 species. *Examples*—*Patrinia*, *Valeriana*, *Centranthus*.

907. The plants belonging to the order are strong scented or aromatic, and some of them have been used as bitter tonics, anthelmintics, and antispasmodics. The root of *Valeriana officinalis* is the common medicinal Valerian. It has a bitter acrid taste, and a peculiar odour, which is fetid and disagreeable in the dry state. In the form of tincture and infusion, it is prescribed in cases of hysteria. Other species of Valerian, as *V. celtica*, *Phu*, *sitchensis*, and *Saliunca*, have similar properties. Valerian is known to have a peculiar effect on cats, causing a species of intoxication. *Nardostachys Jatamansi* is the नर्द, *ναρδος*, or spikenard of the ancients, which was highly prized on account of its perfume. *Valeriana olitoria*, Lamb's lettuce, has been used as a salad. Many of the plants in the order secrete a peculiar volatile oil, to which these properties are due.


908. Order 104.—**Dipsacaceæ**, the Teazel Family. (*Monopet. Epigyn.*) Calyx superior, with an entire, or toothed, or pap-pose limb (fig. 278). Corolla gamopetalous, tubular, inserted on the calycine tube, with an oblique 4-5-lobed limb; æstivation imbricated. Stamens 4, attached to the tube of the corolla, and alternate with its lobes; anthers dithecal, distinct. Ovary cohering with the tube of the calyx, either closely or only at the apex, unilocular; ovule solitary, pendulous, anatropal; style filiform; stigma simple. Fruit dry, indehiscent, crowned by the limb of the calyx, covered by an epicalyx or involucrellum, 1-celled. Seed solitary, pendulous, albuminous; embryo straight; radicle superior.—Herbs or undershrubs, with opposite or verticillate leaves, and capitate or verticillate flowers,

surrounded by a many-leaved involucre (figs. 232, 244). They are found in the south of Europe, the Levant, and at the Cape of Good Hope. The properties of the order are unimportant. The name *Dipsacus* is derived from $\delta\iota\psi\alpha$, thirst, in consequence of the bases of the leaves of some of the species being connate, in such a way as to enclose a cavity which contains water ready to allay thirst. This plant has obtained the name of Venus's Bath; the water contained in which was considered good for bleared eyes. The heads of *Dipsacus fullonum*, Fullers' Teazel, on account of their spiny bracts, are used in dressing cloth. Some of the species are reputed febrifugal. *Scabiosa semisa* is said to yield a green dye, and has from its astringent qualities attracted the attention of tanners. Lindley mentions 6 genera, including 150 species. *Examples*—Morina, Scabiosa, Dipsacus.

909. Order 105.—**Calyceraceæ**, the Calycera Family. (*Monopet. Epigyn.*) Calyx superior, with a limb of 5 unequal segments. Corolla regular, infundibuliform, with a long slender tube, and a 5-lobed limb, the lobes having each three principal veins. Stamens 5, attached to the tube of the corolla, with as many alternating glands below them; filaments monadelphous; anthers partially united. Ovary inferior, 1-celled; ovule solitary, pendulous; style single, smooth; stigma capitate. Fruit an achæmium, crowned by the rigid spiny segments of the calyx, sometimes covered with papillæ, which emit spiral tubes when placed in water. Seeds solitary, pendulous; embryo in the axis of fleshy albumen; radicle superior.—Herbaceous plants, with alternate, exstipulate leaves, and sessile capitate flowers, surrounded by an involucre. They inhabit South America, rarely occurring in the tropical districts, but more plentiful in South Chili. Their properties are unknown. There are 5 known genera according to Lindley, and 10 species. *Examples*—Calycera, Boopis.

910. Order 106.—**Compositæ** (Asteraceæ of Lindley, and Synantheræ of other authors), the Composite Family. (*Monopet. Epigyn.*) (Figs. 623–634). Calyx superior, its limb either wanting or membranous, or divided into bristles, paleæ, or hairs, and called pappus (figs. 277, 279, 626 a). Corolla gamopetalous, ligulate (figs. 301, 624) or tubular (fig. 626 p), in the latter case usually 5-toothed, sometimes bilabiate (fig. 625); two marginal veins, containing spiral cells, run along each of the corolline divisions, and afterwards proceed along the axis of these divisions; æstivation valvate. Stamens usually 5, alternate with the teeth of the corolla (fig. 626 e); filaments distinct; anthers (figs. 301 a, 624, 625, 626 e) cohering into a cylinder (synantherous or syngenesious). Ovary inferior, closely adherent to the tube of the calyx (figs. 624, 625, 626 a, 634), and undistinguishable from it, 1-celled; ovule solitary, erect (figs. 423, 626, 634); style simple, sometimes with collecting hairs (fig. 627); stigmas two, distinct (figs. 404, 627–633) or united. Fruit, an achæmium


Dipsacus Calyx -
 Scabiosa Flowers -
 Morina Stamens -
 Calycera inflorescence -
 Boopis Capitate -
 Composite -



634



624



625



626



633



623



627

629

628

631

630

632

(Cypsela, ¶ 542), crowned with the limb of the calyx (fig. 634). Seed solitary, erect, exalbuminous (fig. 634); radicle inferior.—Herbs or shrubs, with alternate or opposite, exstipulate leaves, and capitula of flowers (called florets), which are either hermaphrodite or unisexual, and are surrounded by bracts in the form of an involucre (figs. 242, 243). Bractlets are sometimes interspersed with the flowers on the receptacle, and are then called paleæ. Some of the flowers belong to the cyanic, others to the xanthic series (¶ 679). In the same head the flowers are sometimes homochromous (ὁμοῖος, similar, and χρώμα, colour), belonging to the same series; at other times they are heterochromous (ἑτεροῖος, diverse), belonging to different series,—the ligulate to the cyanic, and the tubular to the xanthic.

911. This is one of the largest, and, at the same time, one of the most important natural families in the vegetable kingdom. The plants were all included by Linnæus in his class Syngenesia, and were divided into five orders according to the sexes of the florets, and the nature of the involucre. These divisions are given at page 356, under the names Polygamia, Æqualis, Superflua, Frustranea, Necessaria, and Segregata. The following series of terms have also been employed to express the nature of the capitula, as regards stamens and pistils:—

1. Homogamous (ὁμοῖος, alike, and γάμος, marriage), flowers all hermaphrodite (♂).
2. Heterogamous (ἑτεροῖος, diverse), the flowers of the disk (centre) hermaphrodite, those of the ray (circumference) either pistillate (female) only, or neutral, *i. e.* destitute both of stamens and pistils.
3. Monœcious, ♂ — ♀, male and female flowers in the same capitulum.

Figs. 623-634.—Organs of fructification of Compositæ.

Fig. 623.—Diagram of the flower of a Senecio. The outer dotted circle indicates the pappose limb of the calyx, within it is the tubular corolla with five divisions, next five stamens with united anthers, and in the centre the 1-celled, 1-seeded ovary.

Fig. 624.—One of the ligulate flowers or florets of Cichorium Intybus, Succory or Chicory, belonging to the section Cichoraceæ. *o*, Ovary completely adherent to the tube of the calyx, the limb of the calyx forming a crown surrounding the base of the ligulate (strap-shaped) corolla, which has five apical divisions. *e*, Cylinder formed by the anthers (synantherous), traversed by the style with its bifid stigma, *s*.

Fig. 625.—Flower of Chaetanthera linearis, belonging to the section Labiatifloræ. *o*, Ovary with adherent calycine tube. *t*, Tube of the gamopetalous bilabiate corolla. *ls*, Upper lip of the corolla. *li*, Lower lip of the corolla. *e*, Tube of the anthers. *s*, The bifid stigma at the apex of the style.

Fig. 626.—Tubular (flosculus) flower of Aster rubricaulis, belonging to the section Corymbiferae, cut longitudinally, to show the erect ovule, *o*, enclosed in the pericarp, consisting of the walls of the ovary, and the calycine tube incorporated. *p*, United petals. *a*, Pappus, consisting of the altered limb of the calyx. *e*, Stamens with their united anthers, attached to the corolla. *s*, Style traversing the antherine tube.

Figs. 627-633.—Summits of the styles of plants belonging to different tribes of Compositæ. Two stigmatic bands are seen bordering the internal surface of the two branches, which terminate each of these styles. Several have collecting hairs at different parts.

Fig. 627.—Summit of the style of Cichorium Intybus, one of the Cichoraceæ.

Fig. 628.—Chaetanthera linearis, one of the Labiatifloræ.

Fig. 629.—Thevenotia, one of the Cynareæ.

Fig. 630.—Senecio Doria, one of the Senecionideæ.

Fig. 631.—Aster adulterinus, one of the Asteroideæ.

Fig. 632.—Stevia purpurea, one of the Eupatoriaceæ.

Fig. 633.—Vernonia angustifolia, one of the Vernoniaceæ.

Fig. 634.—Ripe fruit (Cypsela) of a Senecio, cut vertically. *e*, Exalbuminous embryo, with inferior radicle. *t*, Spermoderm or covering of the erect seed. *p*, Pericarp consisting of ovarian parietes with the closely-adherent calycine tube. *s*, Style.

4. Heterocephalous (*ἑτεροκεφαλῆς*, diverse, and *κεφαλῆς*, a head), some capitula entirely male, others entirely female, in the same plant.
5. Dioecious, ♂ : ♀, some plants with male capitula only, others with female capitula only.

The following series of terms have been used to express the nature of the capitula, as regards the form and arrangement of the flowers:—

1. Discoid or Flosculous, corollas all tubular.
2. Ligulate or Semiflosculous, corollas all ligulate.
3. Radiate, corollas of the margin or ray ligulate, those of the centre or disk tubular.
4. Falsely-discoid, corollas all bilabiate.
5. Falsely-radiate, or radiatiform, corollas of the margin ligulate, those of the centre bilabiate.

912. Jussieu divided the order into three sections:—1. *Cynarocephalæ* (*cynara*, the artichoke), having the flowers all flosculous (tubular); involucre hard, conical, and often spiny. 2. *Corymbiferæ* (*corymbus*, a corymb, and *fero*, I bear), having flosculous (tubular) florets in the disk (centre), and ligulate (semiflosculous) in the ray (circumference); involucre hemispherical, leafy, or scaly, seldom spiny. 3. *Cichoraceæ* (*cichorium*, succory), having the florets all ligulate. Another section was subsequently added, containing bilabiate flowers.

913. De Candolle made the following divisions, which are now pretty generally adopted:—1. *Tubulifloræ*, hermaphrodite flowers tubular, regularly 5- rarely 4-toothed. Under this section he included several tribes, in which the distinctions are founded on the nature of the style and stigma in the hermaphrodite flowers. These characters are shown in figs. 629-633, which illustrate the tribes *Vernoniaceæ* (fig. 633), *Eupatoriaceæ* (fig. 632), *Asteroidæ* (fig. 631), *Senecionidæ* (fig. 630), and *Cynareæ* (fig. 629). 2. *Labiatifloræ*, hermaphrodite flowers, or at least the unisexual ones, divided into two lips (fig. 625). The subdivisions of this section are also founded on the style and stigma (fig. 628). 3. *Ligulifloræ* (*cichoraceæ*), all the flowers hermaphrodite and ligulate (fig. 624). The form of the style and stigma is seen in fig. 627.

914. Henslow gives the following tabular view of these various divisions—the letter *l* meaning ligulate flowers; *f*, flosculous; *H*, hermaphrodite; *F*, female; *N*, neuter; *M*, male; the relative position of the letters indicating the nature of the florets in the circumference and in the centre of the same capitulum; and in the last three divisions, the letters having reference to the nature of the separate capitula:—

	Jussieu.	De Candolle.
(<i>f. f. f.</i>)	<i>Cynarocephalæ</i>	{ <i>Tubulifloræ</i> 1. <i>Ligulifloræ</i> 3. <i>Labiatifloræ</i> 2.
(<i>l. f. l.</i>)	<i>Corymbiferæ</i>	
(<i>l. l. l.</i>)	<i>Cichoraceæ</i> ...	

	Heads of Flowers.	Linnean Orders.
(H. H. H.)	Homogamous.....	Polygamia æqualis.
(F. H. F.)	Heterogamous.....	{ ————— superflua.
(N. H. N.)		{ ————— frustranea.
(F. M. F.)	Monœcious	————— necessaria.
[(H.)]	Involucrate florets.....	————— segregata.
(M.)—(F.)	Diœcious	*
[(M.) (F.)]	Heterocephalous.....	*

915. The plants of this order are variously distributed over all quarters of the world. According to the calculations of Humboldt, they constitute $\frac{1}{7}$ of the phœnogamous plants of France, $\frac{1}{8}$ of Germany, $\frac{1}{15}$ of Lapland; in North America $\frac{1}{6}$, within the tropics of America $\frac{1}{2}$. Upon the authority of Brown, they only form $\frac{1}{16}$ of the Flora of the north of New Holland, and did not exceed $\frac{1}{23}$ in the collection of plants formed by Smith upon the western coast of Africa in Congo. In northern regions they are generally herbaceous, while in warm climates they sometimes become shrubby, or even arborescent. Cichoraceæ abound in cold regions, while Corymbiferae are common in hot climates. In the northern parts of the world Composites are universally herbaceous plants; they become gradually frutescent or even arborescent as we approach the Equator. Most of those of Chile are bushes, and the Composites of St. Helena are chiefly trees. The number of known genera amounts at present to upwards of 1000, comprehending 9500 species. They are considered as forming $\frac{1}{10}$ of the known species, and this seems to have been the proportion at different periods. Thus, Linnæus enumerates 785 Compositæ out of 8500 species of plants then known all over the world:—

	Compositæ.	Known Species in the World.
In 1809	2,800	27,000
1838	8,523	85,000
1846	9,500	95,000

Examples—Vernonia, Eupatorium, Aster, Bellis, Anthemis, Senecio, Centaurea, Carduus, Triptilion, Trixis, Cichorium, Hieracium, Sonchus.

916. The plants belonging to this vast order have all more or less bitterness, which is sometimes associated with astringent, acrid, and narcotic qualities.

Suborder *Cynarocephalæ*.—The plants of this suborder are usually tonic and stimulant. The bitterness of the plants of this section is often much lessened by cultivation, so that they become esculent. The root of *Arctium Lappa* (*majus* and *minus*), Burdock, is bitterish, and has been used in the form of infusion as a substitute for sarsaparilla. The root, leaves, and fruits (often called seeds), are diaphoretic, diuretic, and alterative. The leaves of *Carduus Benedictus*, Blessed Thistle, were formerly used in medicine as a stomachic and diaphoretic. The blanched stems and leaf-stalks of *Cynara Cardun-*

culus, Cardoon, are eaten, and so are the young succulent receptacles of *Cynara Scolymus*, the Artichoke. The dried flowers of *Carthamus tinctorius* constitute Safflower, which yields a pink dye. The genus *Carduus* includes the various species of Thistle. What is denominated by gardeners the Scotch Thistle, is *Onopordum Acanthium*, a doubtful native of Scotland, but not uncommon in England.

917. Suborder *Corymbiferae*.—The plants of this section have the general bitterness of the order, and some of them have an aromatic odour, from the presence of volatile oil. The flowers of *Anthemis nobilis*, Chamomile, are odoriferous, and yield a volatile oil, which is at first greenish or bluish, but afterwards yellowish brown. They are used as materials for fomentation, and an infusion of them acts as a diaphoretic and emetic. An extract is made from them, having bitter tonic qualities. The essential oil is an excellent carminative. *Anthemis tinctoria* supplies a yellow colour used for dyeing. *Pyrethrum Parthenium*, common Feverfew, is aromatic and stimulant. The root of *Anacyclus Pyrethrum*, Pellitory of Spain, or perhaps of *A. officinarum*, is an irritant and sialogogue; its properties depending on the presence of a volatile oil. *Tussilago Farfara*, Coltsfoot, has been used as a demulcent. The root of *Inula Helenium*, Elecampane, has stimulant and expectorant qualities. It contains a white amylaceous matter called Inulin. The species of *Artemisia* are remarkable for their strong odour and bitter taste. The heads of flowers of *Artemisia Absinthium* (*Absinthium officinale*, and *vulgare*), or Wormwood, and those of *Artemisia Santonica*, and of other species, under the name of Wormseed, are used as anthelmintics and tonics. Several of these species contain a crystalline bitter principle. *Artemisia mutellina*, and *spicata* are used in the preparation of a tincture or distilled spirit, called in France, Eau or Crème d'Absinthe, which is in request among those who are addicted to the pleasures of the table. The woolly leaves of *Artemisia Moxa* are used in China to form the inflammable cones or cylinders called Moxas, which are employed as counter-irritants. *Artemisia Dracunculus*, Tarragon, is used in pickles and salads, and in the medication of vinegar. *A. Abrotanum* is commonly called Southernwood, and is used on the continent in the preparation of beer. The leaves of *Tanacetum vulgare*, Tansy, have stimulant antispasmodic properties. They contain a bitter resin, and an aromatic volatile oil. *Arnica montana*, Mountain Tobacco, or Leopard's-bane, is an acrid stimulant. Its flowers, leaves, and root-stock, are administered in nervous diseases, as well as in gout and rheumatism. The seeds (properly fruits) of *Helianthus annuus*, common Sunflower, contain a bland oil, and when roasted they have been used as a substitute for Coffee. The name *Helianthus* (ἥλιος, the sun, and ἄνθος, a flower) is derived from the popular supposition, that its large heads of flowers follow the sun in its course (¶ 486). The roots of *Helianthus tuberosus*

us, Jerusalem, or more properly, Girasole Artichoke, are used as substitutes for potatoes. *Eupatorium Ayapana*, and *Mikania Guaco*, have been used to cure the bites of snakes. *Ceradia furcata* is a peculiar branching coral-like plant, which grows in dry sterile places the south and west of Africa, and yields a resinoid substance, called some African bdellium. *Madia sativa* has been cultivated on account of its bland oil. The species of *Lychnophora* give a peculiar nature to the mountains of Minas Gerães in Brazil. They grow like ellozias, and they are covered with a dense coat of long brownish-coloured wool, which is often used for beds and pillows.

918. Suborder *Cichoraceæ*.—Most of the plants of this section yield a milky juice, which is bitter, astringent, and sometimes narcotic. By cultivation, some of them are rendered esculent. *Cichorium Intybus*, Wild Succory, or Chicory, is cultivated for the sake of its root, which is used as a substitute for, and as an addition to Coffee. The blanched leaves of *Cichorium Endivia* constitute Endive. *Taraxacum Dens Leonis* (or *officinale*), Dandelion, yields a milky juice, which, in the form of extract, has been used medicinally as a diuretic and alterative. It contains a bitter crystalline principle called Taraxacine. Its root is mixed with Coffee in the same way as Chicory. The inspissated juice of *Lactuca sativa*, common Lettuce, and of *L. virosa*, wild or strong-scented Lettuce, receives the name of Lactucarium, or Lettuce-opium, and is used medicinally for allaying pain, and inducing sleep. It contains a neutral active principle called Lactucin. Other species of *Lactuca* yield an inspissated juice having similar qualities. *Scorzonera* is the esculent root of *Scorzonera hispanica*, while Salsafy is the root of *Tragopogon porrifolius*, which is called the Oyster-plant in America. Many of the plants of the Cichoraceous section, such as *Hieracium*, *Sonchus*, and *Tragopogon*, act as horological and meteorological flowers (§ 483, 484), their capitula opening and closing at certain periods of the day, and in different states of the weather.

919. Order 107.—**Brunoniaceæ**, the Brunonia Family. (*Monopet. Perigyn.*) Calyx persistent, 5-partite, with bracts at the base. Corolla inserted at the base of the calyx, monopetalous, nearly regular, withering; limb 5-parted, having central veins in its segments, which divide at the top into two recurrent marginal veins; æstivation valvate. Stamens 5, inserted with, but free from, the corolla, alternating with its segments; anthers articulated with the short filaments, dithecal, introrse, dehiscing longitudinally. Ovary free, unilocular; ovule solitary, erect, anatropal; style single; stigma enclosed in a 2-valved cup or indusium. Fruit a utricle, enclosed in the hardened calycine tube. Seed solitary, erect, exalbuminous; embryo straight; cotyledons fleshy, plano-convex; radicle minute, inferior.—Stemless herbaceous plants, with radical, exstipulate leaves, and capitate flowers, supported on scapes, and surrounded by an involucre of enlarged bracts. Natives of New Holland.

Their properties are unknown. The order contains as yet only 1 genus, and 9 species. *Example*—Brunonia.

920. Order 108.—**Goodeniaceæ**, the Goodenia Family. (*Monopet. Epigyn.* and *Perigyn.*) Calyx persistent, usually equal, with 3-5 divisions, sometimes obsolete. Corolla inserted into the calyx, monopetalous, more or less irregular, marcescent or deciduous; its tube split at the back, and sometimes separable into five pieces, when the calyx only coheres with the base of the ovary; its limb 5-partite, uni- or bilabiate, the thin part of the segments being at the edges, which are folded inwards in æstivation. Stamens 5, distinct, inserted with, but free from, the corolla, and alternate with its segments; anthers not articulated with the filaments, distinct or cohering, bilocular, with longitudinal dehiscence; pollen-grains either separate or united in fours. Ovary more or less united to the calycine tube, 1-2- or 4-celled, sometimes with a gland at its base; ovules definite or 00, attached to a central, often free, placenta; style 1, simple, rarely divided; stigma fleshy, undivided or 2-lobed, surrounded by a cup-like indusium. Fruit a 1-2- or 4-celled capsule, or drupaceous or nut-like. Seeds definite or indefinite, with a thickened, often hard testa; embryo straight, in fleshy albumen; cotyledons leafy; radicle inferior.—Herbs, rarely shrubs, not lactescent, with scattered, exstipulate, usually alternate leaves, and distinct, never capitate flowers. They are found chiefly in Australia, and in the South Sea Islands. The leaves of *Scævola Taccada* are eaten as potherbs. Some superstitious qualities are ascribed to its berries. The pith, which is soft and spongy, is fashioned by the Malays into artificial flowers. *Scævola Bela Modogam* appears to be emollient, and is used in India to bring tumours to a head. The order is divided into two suborders:—1. Goodeniæ, with dehiscent capsular fruit, and numerous seeds. 2. Scævoleæ, with indehiscent, drupaceous, or nut-like fruit, and seeds solitary, or two in each cell. There are 14 known genera, according to Lindley, and about 150 species. *Examples*—Goodenia, Velleia, Leschenaultia, Scævola, Dampiera.


921. Order 109.—**Stylidiaceæ**, the Stylidium or Stylewort Family. (*Monopet. Epigyn.*) Calyx adherent, persistent, with 2-6 divisions, bilabiate, or regular. Corolla gamopetalous, falling off late, limb usually irregular, 5-6-partite, segments with a central vein; æstivation imbricated. Stamens 2; filaments united with the style into a longitudinal column; anthers didymous, rarely simple, lying over the stigma; pollen simple, globose, or angular. Ovary cohering with the calyx, bilocular, or by contraction of the dissepiment unilocular, often surmounted by one gland in front, or by two opposite ones; ovules anatropal; style 1; stigma entire or bifid. Fruit a bivalvular, bilocular, or spuriously unilocular capsule, with septicidal dehiscence. Seeds 00, small, erect; embryo minute, enclosed in fleshy, somewhat

oily albumen.—Non-lactescent herbs or undershrubs, with alternate, scattered, or somewhat verticillate, entire, exstipulate leaves. They are well distinguished by their gynandrous structure. The column formed by the union of the filaments and style, possesses, in the species of the genus *Stylidium*, a peculiar irritability. It hangs down on one side of the flower, and when touched at the point of flexure, it springs over with considerable force from one side to the other. If not too far advanced to maturity, the column will recover its former position in the course of time. The flower may be cut off carefully without disturbing the column, and the irritability continues for a considerable length of time if the flower is put into water. The movement is said to be connected with the bursting of the anthers, and the discharge of the pollen on the stigma. The cause of this movement is very obscure, but it seems to depend on some changes in the cells (§ 497, 665). The plants are principally natives of marshy places in New Holland. Some are found at the southern point of South America. There are 5 known genera, and 121 species. *Examples*—*Stylidium*, *Forstera*.

922. Order 110.—**Campanulaceæ**, the Hare-bell Family. (*Monopet. Epigyn.*) (Fig. 635.) Calyx superior, usually 5-lobed (figs. 636, 637 c), sometimes 3-8-lobed, persistent. Corolla gamopetalous, inserted into the top of the calyx, usually 5-lobed (fig. 255), sometimes 3-8-lobed, regular, marcescent (fig. 461 e); æstivation valvate (figs. 636, 637 p). Stamens inserted into the calyx, alternating with the corolline lobes, and equal to them in number; anthers bilocular, free (fig. 637 e); pollen spherical. Ovary more or less completely inferior, composed of two or more carpels; ovules indefinite (fig. 638); style simple, covered with collecting hairs (fig. 637); stigma naked, simple, or with as many lobes as there are ovarian cells (figs. 293 s, 637 s). Fruit capsular, crowned with the withered calyx and corolla, dehiscing in a loculicidal manner by lateral apertures (figs. 461 t t, 639), or by valves at the apex. Seeds 00, attached to a central placenta; embryo straight, in the axis of fleshy albumen; radicle pointing to the hilum (figs. 640-642).—Lactescent herbs or undershrubs, with alternate, rarely opposite, exstipulate leaves. The hairs on the style are said to be retractile, and seem to be connected with the application of the pollen (§ 499). The flowers in most instances belong to the cyanic series. They are natives chiefly of northern and temperate regions. They abound in the alpine regions of Europe and Asia, and are also frequent in North America. Alphonse De Candolle states that the species whose capsule dehisces by lateral fissures are natives of the northern hemisphere, while those with apicilar dehiscence are principally found in the southern hemisphere. The milky juice found in the plants of this order has acrid properties. The roots and young shoots of *Campanula Rapunculus*, Rampion, are used as articles of diet. Lindley enumerates

Campanulaceæ.
 Superior Flower-
 Marig.
 Marcescent Corolla-
 Globular Pollen

28 genera, including 500 species. *Examples*—*Campanula*, *Phyteuma*, *Jasione*.



923. Order 111.—**Lobeliaceæ**, the Lobelia Family. (*Monopet. Epigyn.*) Calyx superior, 5-lobed or entire. Corolla gamopetalous, inserted on the calyx, irregular, more or less deeply 5-cleft. Stamens 5, attached to the calyx, alternate with the segments of the corolla; anthers cohering; pollen oval. Ovary inferior, 1-3-celled; ovules 00, attached either to central or parietal placentas; style glabrous, with a fringe of hairs below the stigma. Fruit a 1- or many-celled capsule, with apicilar dehiscence. Seeds numerous; embryo straight, in the axis of fleshy albumen; radicle pointing to the hilum.—Lactescent herbs or shrubs, with alternate, exstipulate leaves. They are found both in temperate and warm countries. There are 27 known genera, and 375 species. *Examples*—*Lobelia*, *Siphocampylus*, *Clintonia*.

Figs. 635-642.—Organs of fructification of *Campanula Rapunculus*, Rampion, to illustrate the natural order Campanulaceæ.

Fig. 635.—Diagram of the flower, showing five divisions of the calyx, five divisions of the corolla alternating with them, five alternating stamens, and five cells of the ovary.

Fig. 636.—Flower-bud. c, Calyx adherent to the ovary. p, Corolla, with valvate aestivation.

Fig. 637.—Vertical section of the flower. c, Calyx cohering with the ovary. o, p, Gamo-petalous corolla. s, Stamens with bilocular anthers. s, Lobed stigma at the apex of the style, which is covered with collecting hairs. o, Ovary containing numerous ovules attached to a central placenta.

Fig. 638.—Horizontal section of the ovary.

Fig. 639.—Fruit crowned by the limb of the calyx, dehiscing by openings at the base.

Fig. 640.—Seed in an entire state.

Fig. 641.—Seed cut vertically. p, Perisperm (albumen). e, Straight embryo in the axis of the albumen, with the radicle pointing to the hilum.

Fig. 642.—Embryo detached, showing its form, the cotyledons and radicle.

924. Acridity prevails more or less in the order. The milky juice of some, such as *Lobelia urens*, is said to be vesicant. *Lobelia inflata*, Indian Tobacco, a native of North America, is used medicinally as a sedative, expectorant, and antispasmodic. It is chiefly administered in cases of asthma. The whole plant is active, but the root and capsules are said to be most powerful. In large doses, the plant acts as a narcotico-acrid poison. It owes its properties to a bitter principle called Lobein. The root of *Lobelia syphilitica* is acrid and emetic. The milky juice of some of the plants of the order contains a considerable quantity of caoutchouc.

925. Order 112.—**Gesneraceæ**, the Gesnera Family. (*Monopet. Perigyn.*) Calyx partially adherent, 5-partite; æstivation valvate. Corolla monopetalous, tubular, more or less irregular, 5-lobed; æstivation imbricated. Stamens 4, didynamous, with the rudiment of a 5th, rarely 2; anthers dithecal, with a thick swollen connective. Ovary partly free, unilocular, formed by two carpels with parietal placentas, which are 2-lobed; ovules indefinite, anatropal; style continuous with the ovary; stigma capitate, concave, glandular or annular. Disk surrounding the base of the ovary. Fruit capsular or succulent, 1-celled, more or less adherent. Seeds 00, minute; testa thin, finely and obliquely veined; embryo erect in the axis of fleshy albumen; radicle pointing to the hilum.—Herbs or shrubs, often springing from scaly tubers, with opposite or whorled, rugose, exstipulate leaves and showy flowers. They are found principally in the warmer regions of America, and are interesting chiefly on account of their beauty, for they do not appear to possess any important qualities. There are 22 known genera, and upwards of 120 species. *Examples*—Gesnera, Columnea, Gloxinia, Achimenes.

926. Order 113.—**Ericaceæ**, the Heath Family. (*Monopet. Hypog.*) Calyx 4-5-cleft, nearly equal, persistent. Corolla inserted at the base of the calyx, or hypogynous, monopetalous (fig. 298), 4-5-cleft, sometimes tetra- or pentapetalous, regular or irregular, often marcescent; æstivation imbricated. Stamens definite, equal in number to the segments of the corolla, or twice as many, inserted with the corolla, and either free from it or attached to its base; anthers 2-celled, cells hard and dry, bifid (fig. 336), usually having appendages at the base (fig. 338 a) or apex, dehiscent by apicilar pores (fig. 340) or clefts. Ovary free, surrounded at the base by a disk or scales, multilocular; ovules 00, attached to a central placenta; style 1, straight; stigma 1, undivided (fig. 408), or toothed. Fruit capsular or baccate, many-celled, with loculicidal or septicidal dehiscence. Seeds 00, minute; embryo cylindrical, in the axis of fleshy albumen; radicle next the hilum.—Shrubs, undershrubs, or herbaceous plants, with evergreen, often rigid, entire, verticillate, or opposite, exstipulate leaves. The order contains many beautiful and showy plants, which abound at the

Many
Pores
dehiscent by
Pores
Ripet
Anthers
dehiscent
by
Pores
in
Ericaceæ
Sep. showy
Celled 000 in

Cape of Good Hope, and which are found also in Europe, North and South America, and Asia. The order has been divided into the following suborders:—1. *Ericææ*, with the testa closely adherent to the kernel, including the true Heaths with naked buds, and the *Rhododendron* tribe with scaly conical buds. 2. *Monotropeæ*, seeds having a loose winged testa, including the true *Monotropas* or Fir-rapes, scaly plants, with longitudinally or transversely dehiscent anthers, and *Pyroleæ*, or the Wintergreen tribe, leafy plants with porous anthers. There are 52 known genera, and nearly 880 species. *Examples*—*Erica*, *Calluna*, *Menziesia*, *Andromeda*, *Arbutus*, *Rhododendron*, *Azalea*, *Monotropas*, *Pyrola*.

927. The plants of the order are not distinguished for medicinal virtues. None of the species of *Erica* are put to any use. There are six species of the genus natives of Britain: two of which, *E. cineria* and *Tetralix*, are common; two are peculiar to Ireland, *E. Mackaiana* and *mediterranea*; one is restricted to England, *E. ciliaris*; and one is common to England and Ireland, *E. vagans*. *Calluna vulgaris*, is Ling, or the common Heather. It has astringent qualities, and has been used for dyeing. It is commonly made into brooms. The leaves of *Arbutus*, or *Arctostaphylos Uva-Ursi*, Bearberry, are used as astringents, especially in chronic mucous discharges. Many of the species of *Rhododendron*, *Azalea*, *Kalmia*, *Andromeda*, and *Ledum*, have poisonous narcotic qualities. These properties are well marked in *Rhododendron*, *Chrysanthum*, a Siberian species. It is said that *Azalea pontica* was the plant the flowers of which yielded the poisonous honey noticed by Xenophon in his account of the retreat of the 10,000. The fruits of many plants belonging to the order are eatable. *Gaultheria procumbens* and *Shallon* are American shrubs, which furnish succulent and grateful berries. They yield a volatile oil. *Arbutus Unedo* is called Strawberry-tree, from its fruit resembling a strawberry in aspect. It is, however, by no means agreeable as an article of food, and the specific name may possibly indicate, that to eat one is sufficient. The plant grows at the Lakes of Killarney in a native state. *Chimaphila (Pyrola) umbellata*, a North American plant, has been employed as a tonic and diuretic. The leaves have a bitter astringent taste, and the fresh plant is irritant.

928. Order 114.—**Vacciniacæ**, the Cranberry Family. (*Monopet. Epigyn.*) Calyx superior, entire, 4-6-lobed. Corolla monopetalous, 4-6-lobed; æstivation imbricated. Stamens distinct, 8-12, inserted into an epigynous disk; anthers bilocular, with two horn-like cells, dehiscing by pores (fig. 335). Ovary inferior, 4-5-celled; ovules 00; style simple; stigma simple. Fruit succulent, crowned by the persistent limb of the calyx. Seeds 1 or many in each cell, minute; embryo straight, in the axis of fleshy albumen; cotyledons very short; radicle long, inferior.—Shrubby plants, with alternate, undivided,

exstipulate leaves. They are closely allied to Ericaceæ, and differ from that order chiefly in their adherent (inferior) ovary. They are natives of temperate regions, and some of them are marsh plants. Some are astringent, others yield subacid edible fruits. Cranberries are produced by *Vaccinium Oxycoccus* (*Oxycoccus palustris* of some), and *V. Macrocarpum*. *V. Vitis-idea*, red Whortleberry, or Cowberry, yields a fruit which is often used as a substitute for Cranberries. The leaves of the plant are sometimes used to adulterate *Uva-Ursi*. *V. uliginosum*, found in alpine districts, produces the black whortleberry. *Vaccinium myrtillus*, yields the Bilberry or Blaeberry. There are 15 genera of the order, and 200 species. *Examples*—*Vaccinium*, *Oxycoccus*, *Thibaudia*.

929. Order 115.—**Epacridaceæ**, the Epacris Family. (*Monopet. Hypog.*) Calyx 5- rarely 4-parted, often coloured, persistent. Corolla inserted at the base of the calyx, or hypogynous, deciduous or marcescent, monopetalous, sometimes separable into 5 petals; limb with 5, rarely 4, equal divisions, sometimes by the cohesion of the segments, bursting transversely; æstivation imbricated or valvate. Stamens inserted with or on the corolla, equal in number to, and alternate with its segments, rarely fewer; anthers 1-celled (fig. 327), without appendages, opening longitudinally; pollen round, or formed of three united grains, attached to a single central receptacle. Ovary sessile, free, multilocular, rarely unilocular, surrounded by scales at the base; ovules solitary or 00; style 1; stigma simple, sometimes toothed. Fruit drupaceous, baccate, or capsular. Seeds albuminous; embryo slender, in the axis of fleshy albumen, and about half its length. —Shrubs, or small trees, with alternate, rarely opposite, exstipulate leaves, which are sometimes half-amplexicaul at the base. They are allied to Ericaceæ, and seem to occupy the place of heaths in Australia. They are distinguished from heaths by the structure of their anthers. They are cultivated for the beauty of their flowers. In some cases they yield edible fruits. One of the plants, called Native Currant in Australia, is *Leucopogon Richei*. The order has been divided into two sections:—1. Epacreæ, polyspermous. 2. Stypheliæ, monospermous. There are thirty known genera, and 320 species, according to Lindley. *Examples*—*Epacris*, *Sprengelia*, *Styphelia*, *Leucopogon*, *Lissanthe*.

930. Order 116.—**Columelliaceæ**, the Columellia Family. (*Monopet. Epigyn.*) Calyx superior, quinquepartite. Corolla rotate, inserted into the calyx, 5-8-parted; æstivation imbricate. Stamens 2, inserted in the throat of the corolla; anthers roundish, 3-lobed, extrorse, each consisting of six linear sinuous cells, arranged in pairs, dehiscing longitudinally, and attached to a 3-lobed fleshy connective. Disk fleshy, perigynous. Ovary adhering to the calycine tube, 2-celled; ovules 00; style simple, smooth; stigma capitate, 2-lobed. Fruit, a bilocular,

bivalvular capsule, with both septicidal and loculicidal dehiscence. Seeds 00; testa smooth and coriaceous; embryo straight, in the axis of fleshy albumen; cotyledons oval, obtuse; radicle long, pointing to the hilum.—Evergreen shrubs or trees, with opposite, entire, exstipulate leaves, and solitary yellow flowers. Natives of Mexico and Peru. Their properties unknown. There is 1 genus mentioned, including 3 species. *Example*—Columellia.

931. Order 117.—**Styracææ** (Symplocinæ of Don), the Storax Family. (*Monopet. Perigyn.*) Calyx persistent, with an entire or a 5- or 4-divided limb. Corolla gamopetalous, regular, inserted in the calyx; æstivation imbricated or valvate. Stamens definite or 00, attached to the corolline tube, of unequal length; filaments often slightly united at their base in one or more parcels; anthers innate, dithecal, introrse. Ovary either free or cohering more or less to the calycine tube, 2-5-celled, the septa occasionally deficient towards the centre; ovules 2-4 in each cell or 00, pendulous, sometimes the upper ones ascending; style simple; stigma simple. Fruit enclosed in the calyx, drupaceous, usually unilocular by abortion. Seeds usually solitary, erect or suspended; embryo slender, in the axis of fleshy albumen; cotyledons flat, foliaceous; radicle long, pointing to the hilum.—Trees or shrubs, with alternate, exstipulate leaves, and frequently stellate hairs. They are chiefly natives of warm countries. There are two sections:—1. *Styracææ*, with a more or less valvate æstivation of the corolla, and long anthers. 2. *Symplocææ*, with a quincuncial corolline æstivation, and roundish anthers. Lindley gives 6 genera, including 115 species. *Examples*—*Styrax*, *Halesia*, *Symplocos*.

932. The plants of the order have in general stimulant, aromatic, and fragrant properties. *Styrax officinale*, a tree inhabiting Syria, Arabia, and the southern parts of Europe, is supposed to be the source of the balsamic resinous substance called Storax. The resinous juice is procured after incisions or punctures by insects. Common Storax is imported into Britain from Trieste, in the form of little cakes, having a balsamic odour. Besides resin and a little volatile oil, it contains from 1-2½ per cent. of Benzoic acid. It has been employed medicinally as a pectoral remedy. *Styrax Benzoin* is a tree 70 or 80 feet high, a native of Sumatra and Borneo, which yields by incisions the concrete balsamic exudation called Benzoin. When fine, this substance contains about 80 per cent. of resin, and nearly 20 of benzoic acid. It is used medicinally as a stimulant expectorant, and is one of the ingredients in the celebrated Friar's balsam. It exists also in other empirical remedies, such as Riga balsam and Jesuits' drops. Benzoin is generally used for fumigation and incense. Pastilles are made by mixing it with balsam of tolu, sandal-wood, labdanum, charcoal, nitre, gum, and tragacanth. *Halesias* are the Snow-drop trees of

Carolina. Some of the species of *Symplocos* are used for dyeing; others are used as tea.

933. Order 118.—**Ebenaceæ**, the Ebony Family. (*Monopet. Hypog.*) Flowers hermaphrodite or unisexual. Calyx 3-7-divided, nearly equal, persistent. Corolla gamopetalous, regular, deciduous, somewhat coriaceous; limb with 3-7 divisions; æstivation imbricated. Stamens either attached to the corolla, or hypogynous, 2 or 4 times as many as the corolline segments, rarely equal to them in number, and then alternate with them; filaments usually in two rows, the inner row having smaller anthers; anthers erect, lanceolate, bilocular, with longitudinal dehiscence. Ovary free, sessile, multilocular; ovules 1-2 in each cell, pendulous; style divided, rarely simple; stigmas bifid or simple. Fruit fleshy, round or oval, the pericarp sometimes opening regularly. Seeds few; testa membranous; embryo straight, nearly in the axis of cartilaginous albumen; cotyledons leafy; radicle taper, next the hilum.—Trees or shrubs, not lactescent, with alternate, exstipulate, coriaceous leaves. They are chiefly found in tropical regions, and many species are met with in India. The plants are in general remarkable for the hardness and durability of their wood. Some yield edible fruit. *Diospyros Ebenus* and other African and Asiatic species, supply Ebony, which is the black duramen of the tree. Other species of *Diospyros* furnish Ironwood. *Diospyros virginiana*, the Persimmon, yields a fruit (sometimes called the Date-plum) which is austere when green, but becomes sweet and eatable when ripe, especially after being acted on by frost. Its bark has, by some practitioners in the United States, been employed with success as a febrifuge in cases of cholera infantum. *D. Kaki* is the Keg-fig of Japan, the fruit of which resembles a plum. Lindley notices 9 genera, including 160 species. *Examples*—*Diospyros*, Royena, Maba.

934. Order 119.—**Aquifoliaceæ** (Ilicineæ of some), the Holly Family. (*Monopet. Hypog.*) Sepals 4-6; æstivation imbricated. Corolla monopetalous, hypogynous, 4-6-parted; æstivation imbricate. Stamens inserted into the corolla, alternate with its segments, and equal to them in number; filaments straight; anthers adnate, bilocular, introrse. Disk 0. Ovary free, fleshy, somewhat truncate, 2-6-celled; ovules solitary, anatropal, pendulous from a cup-shaped funiculus; stigma nearly sessile, lobed. Fruit fleshy, indehiscent, with 2-6 monospermous nucules, and hence it is sometimes called a nuculanium (§ 553). Seed suspended; albumen large, fleshy; embryo small, lying next the hilum; cotyledons small; radicle superior.—Evergreen trees or shrubs, with alternate or opposite, coriaceous, simple, exstipulate leaves. They are found in various parts of the world, as in Europe, North and South America, and Africa. Lindley enumerates 11 genera, including 110 species. *Examples*—*Ilex*, *Prinos*.

935. Astringent and tonic properties seem to pervade the order. *Ilex Aquifolium*, the common Holly, is a native of Europe, and is one of the indigenous plants of Britain. It forms excellent fences and hedges. At Tynningham, in Scotland, there are 2952 yards of holly hedges, most of them upwards of 130 years old. These hedges vary in height from 10 to 23 feet, and they are 9 to 13 feet wide at the base. The leaves and bark of the Holly are said to possess tonic and febrifuge properties; while its succulent fruit (berries) are emetic and purgative. Haller recommends the juice of the leaves in icterus. Its wood is white and hard, and is much esteemed in turnery, joinery and cabinet work, while its bark furnishes bird-lime. *Ilex Paraguayensis* furnishes Yerba Maté or Paraguay Tea, which is used extensively in some districts of South America. The leaves of the plant yield the bitter principle called Theine, which has been mentioned as existing in Tea and Coffee. Other species of *Ilex* are employed in Brazil for a similar purpose. The black drink of the Creek Indians is prepared from the leaves of *Ilex vomitoria*.

936. Order 120.—**Sapotaceæ**, the Sappodilla Family. (*Monopet. Hypog.*) Flowers hermaphrodite. Calyx regular, with 5, sometimes 4-8 divisions, persistent; æstivation valvate or imbricate. Corolla monopetalous, hypogynous, deciduous, regular, its lobes equal to, rarely twice or thrice as many as those of the calyx. Stamens inserted on the corolla, definite, distinct; fertile ones as many as, rarely more than, the segments of the calyx, with which they alternate; sterile ones alternating with the fertile ones, rarely wanting. Disk 0. Ovary free, multilocular; ovules solitary, anatropal, ascending or pendulous; style 1; stigma simple, sometimes lobed. Fruit fleshy, multilocular, or by abortion unilocular. Seeds nut-like, solitary; testa bony and shining, with a long scar on its inner face; embryo large, erect, white; albumen usually fleshy, sometimes 0; cotyledons in the albuminous seeds, foliaceous, in the exalbuminous, fleshy; radicle straight or slightly curved, pointing to the hilum.—Lactescent trees or shrubs, with alternate, exstipulate, entire, coriaceous leaves. They are natives chiefly of the tropical parts of India, Africa, and America. A few are found at the Cape of Good Hope. The number of known genera noticed by Lindley is 21, species 212.

937. Many of the plants of this order yield edible fruits, while others supply oily matter. Some act as tonics, astringents, and febrifuges. *Achras Sapota*, and other species, furnish the Sappodilla Plum and Naseberry, well known West Indian fruits; while *Achras mammosa* yields the fruit called Marmalade. The bark of some of the species of *Achras* is tonic and astringent, and the seeds of several have laxative properties. The fruit of *Chrysophyllum Cainito* is the Star-apple. Another species of the genus supplies the Surinam Medlar of Europeans. Various species of *Bassia* yield oil. *B. Parkii* is said to be

See. 120 - Name said to be B. latifolia
of Long Island See p. 2005

the source of the Shea butter, and hence the tree is called the Butter-tree of Park. *B. butyracea*, the Madhuca-tree, gives a similar product, which is used as butter in Nipal. The milky juice of some of the plants contains elastic matter. *Isonandra Gutta* is the source of Gutta Percha, a kind of caoutchouc, which softens at a moderate temperature, and is now extensively used for the soles of shoes, ropes, straps, casts, and various articles for domestic use. The kernels of *Lucuma mammosa* abound with prussic acid. A small portion of the kernel, introduced into the stomach, produces sickness and coughing.

938. Order 121.—**Myrsinaceæ**, the Myrsine Family. (*Monopet. Hypog.*) Flowers hermaphrodite or occasionally unisexual. Calyx 4-5-cleft, persistent. Corolla monopetalous, hypogynous, 4-5-cleft, equal. Stamens 4-5, inserted into the corolla, and opposite to its segments; filaments distinct, rarely united, sometimes 0, occasionally 5 sterile petaloid alternating ones; anthers sagittate, erect, bilocular, with longitudinal dehiscence. Ovary free or slightly adherent, unilocular; ovules definite or indefinite, campylotropal, immersed in a free central placenta; style single; stigma simple or lobed. Fruit fleshy, 1- or many-seeded. Seeds angular or roundish, with a concave hilum, and a membranous spermoderm; albumen horny; embryo usually curved, often heterotropal; cotyledons short; radicle horizontal when the seed is solitary, inferior when there are several seeds.—Trees, shrubs, or undershrubs, with alternate or opposite, coriaceous, exstipulate leaves. They are much restricted as regards their geographical limits, and they are said to abound chiefly in islands with an equable temperature. They are found in Africa, Asia, and America. Little is known regarding their properties. *Theophrasta Jussiei* is a prickly-leaved shrub, which is called Coco in St. Domingo. Its seeds are eatable, and a kind of bread is made from them. The berries of *Myrsine bifaria* are said to possess cathartic properties. The *Ardisias* are prized for the beauty of their foliage. There are 31 known genera, and 325 species. *Examples*—Myrsine, Ardisia, Mæsa, Jacquinia.

939. Order 122.—**Jasminaceæ**, the Jasmine or Jessamine Family. (*Monopet. Hypog.*) Flowers ♂. Calyx with 5-8 divisions or teeth, persistent. Corolla monopetalous, hypogynous, regular, salver-shaped, with 5-8 divisions; æstivation twisted or valvate. Stamens 2, inserted on the corolla, included; anthers bilocular, with longitudinal dehiscence. Disk 0. Ovary free, 2-celled; ovules erect, anatropal, 1-4 in each cell; style 1; stigma 2-lobed. Fruit a double berry, or a pyxidium, or a 2-valved capsule. Seeds usually solitary, rarely in pairs, albuminous or exalbuminous; embryo straight; radicle inferior.—Shrubs often with twining stems, and opposite or alternate, pinnate leaves. They abound chiefly in the tropical parts of India. They have frequently fragrant flowers which yield oils, and their leaves and roots are sometimes bitter. The essential oil of Jasmine is procured from *Jasminum*

officinale, *grandiflorum*, *odoratissimum*, and *Sambac*. The bitter root of *Jasminum angustifolium*, ground small and mixed with powdered *Acorus Calamus* root, is considered in India as a valuable external application in cases of ringworm. In the East Indies the tube of the corolla of *Nyctanthes Arborescens* is used as an orange dye. There are 5 known genera, and 100 species. *Examples*—*Jasminum*, *Nyctanthes*, *Bolivaria*.

940. Order 123.—**Oleaceæ**, the Olive Family. (*Monopet. Hypog.*) (Fig. 254.) Flowers ♂, sometimes ♂ ♀. Calyx gamosepalous, divided, persistent. Corolla gamopetalous, hypogynous, 4-cleft, sometimes of 4 petals, which are connected in pairs by means of the filaments, sometimes 0, æstivation somewhat valvate. Stamens 2 (rarely 4), alternate with the corolline segments; anthers dithecal, with longitudinal dehiscence. Disk 0. Ovary free, 2-celled; ovules in pairs, collateral or pendulous; style 1 or 0; stigma entire or bifid. Fruit drupaceous, baccate or capsular, sometimes samaroid (fig. 437). Seeds often by abortion solitary; albumen dense, fleshy, abundant; embryo straight, about half the length of the albumen; cotyledons leafy; radicle superior.—Trees or shrubs, with opposite leaves (fig. 254), which are either simple or compound. Found chiefly in temperate regions. They occur in North America, Asia, Europe, and New Holland. There are two sections of the order:—1. *Oleæ*, with a drupaceous or berried fruit. 2. *Fraxineæ*, with a samaroid (winged) fruit. Lindley mentions 24 genera, including 130 species. *Examples*—*Olea*, *Ligustrum*, *Fraxinus*, *Syringa*.

941. The plants of the order are bitter, tonic, and astringent, and some yield fixed oil. *Olea europæa* is the Olive-tree, the *איל* of the Bible, the *ελαια* of the Greeks. It grows naturally on the coast of the Mediterranean, and is cultivated in many parts of the south of Europe. There are several varieties of the plant, two of which have been long distinguished—the wild and cultivated. The former is an evergreen shrub or low tree, with spiny branches and round twigs; the latter is a taller tree, without spines, and with four-angled twigs. The fruit is a drupe, about the size and colour of a damson. Its fleshy pericarp yields by expression olive oil, of which the finest comes from Provence and Florence. It consists of two oleaginous principles—Margarin and Elain. Olive oil has nutrient, emollient, and laxative properties. It is used in forming ointments, liniments, and plasters. The bark of the Olive-tree has been used as a tonic; and a resinous exudation from it, called Olivile, or Olive-gum, or Lecca-gum, is employed in the same way. Spanish or Castile soap is made by mixing olive oil and soda, while soft soap is made by mixing the oil with potash. The flowers of *Olea fragrans*, the Lan-hoa of the Chinese, are used to perfume teas. Several species of *Ornus*, more particularly *O. rotundifolia* and *O. europæa*, yield a sweet exudation called Manna, a substance completely different from the *איל* (manna) of the Bible, on

which the Israelites fed. The Manna or Flowering Ash is a native of southern Europe, and grows abundantly in the south of Italy and Sicily, whence the Manna of commerce is imported. The tree attains a height of 20 or 30 feet, and it has a fine appearance when its clusters of white flowers are produced.—There is a very fine Manna-ash in the Edinburgh Botanic Garden. Manna is the concrete juice of the tree, which flows out after incisions or insect punctures. It contains a peculiar sweet principle called Mannite. Manna is nutritive and laxative, and is sometimes administered to infants and young children, on account of the mildness of its action. *Syringa vulgaris*, common Lilac, has a febrifuge bark, which is extensively employed by the peasants in Brenne, for the cure of the endemic intermittent fever. According to Meillet this quality is owing to a principle which he calls Lilacine. *Fraxinus excelsior*, the common Ash, is one of the trees which comes late into leaf, and the leaves of which fall off early in autumn. Some specimens attain the height of 70, 90, or 100 feet, with a circumference of 20 or 30 feet. The wood of the tree is tough and elastic, and is used for oars, as well as by coachmakers, &c. The wood of its roots is beautifully veined. The pendulous variety, called Weeping-ash, is often engrafted on the common Ash, so as to produce a better effect. The leaves of *Ligustrum vulgare*, common Privet (fig. 254), are astringent. *L. lucidum* yields a kind of waxy excretion, which is used in China for economical purposes.

942. Order 124.—**Asclepiadaceæ**, the Asclepias Family. (*Monopet. Hypog.*) (Figs. 353, 643-651). Calyx 5-divided, persistent (fig. 646 c.) Corolla synpetalous (monopetalous), hypogynous, regular, 5-lobed (figs. 644, 645 p p), deciduous; æstivation imbricate, rarely valvate. Stamens 5, inserted into the base of the corolla, and alternate with its segments (fig. 646 e); filaments usually combined so as to form a tube; staminal tube rarely naked behind, generally furnished with a corona (crown) of variously-formed leaves, which are either distinct or connate. Anthers bilocular, each cell sometimes spuriously divided; pollen, when the anther dehisces, cohering in masses (pollinia), which are either as numerous as the cells, or are confluent in pairs, and adhere to the five stigmatic processes, either in sets of two or four, or singly (figs. 349, 353, 647). Ovaries 2 (fig. 646 o); ovules 00; styles 2, closely approaching each other (fig. 646 s), often very short; stigma common to both styles, dilated, quinquangular; the angles furnished with cartilaginous corpuscles which retain the pollinia, or with glands (figs. 645, 646 g). Fruit consisting of two follicles (sometimes only one by abortion) having a placenta on the ventral suture (fig. 649). Seeds 00, imbricate, pendulous, usually comose (hairy) at the hilum (fig. 650); albumen thin (fig. 651 p); embryo straight; cotyledons leafy; radicle superior (fig. 651 e).—Shrubs, or occasionally herbs, usually with milky juice,

and often twining. The leaves are usually opposite, sometimes alternate or verticillate, with interpetiolar cilia in place of stipules.



The gynostegium ($\gamma\upsilon\nu\eta$, pistil, and $\sigma\tau\acute{\epsilon}\gamma\omega$, I cover), staminal crown

Figs. 643-651.—Organs of fructification of *Asclepias nivea*, to illustrate the natural order Asclepiadaceæ.

Fig. 643.—Diagram of the flower, with five divisions of the calyx, five segments of the corolla, five stamens, and two ovaries.

Fig. 644.—The entire flower. *p*, Corolla, with five lobes. *a*, Appendages forming the staminal crown (corona).

Fig. 645.—The flower viewed from above. *p p*, Gamopetalous corolla with its five lobes. *a a*, Appendages forming the corona or crown. *g g*, Glandular bodies attached to the stigma, and bearing the pollen-masses (pollinia).

Fig. 646.—The flower cut vertically. *c*, Calyx. *p*, Corolla. *a a*, Coronal appendages. *e*, Stamens. *o*, Ovary. *s*, Styles which are united at the upper part by means of the large stigma, at the base of which, towards the points, *p p*, the pollen-tubes enter.

Fig. 647.—Two pollen-masses, *m*, attached by two prolongations, *g*, in the form of a candle or tail, to another body, *g*, formed by the union of two stigmatic glands. *p*, Pollen-grains with tubes beginning to escape from the masses.

Fig. 648.—One of the pollen-grains, with its tube separated and highly magnified.

Fig. 649.—Fruit at the period of dehiscence. *f f*, Two follicles. *p*, Placenta, which is detached. *g*, Comose seeds.

Fig. 650.—One of the comose seeds separated. *a*, The hairy appendage at the hilum.

Fig. 651.—Seed separated from the hairs and cut vertically. *t e*, External integument. *t i*, Internal integument. *p*, Perisperm or thin albumen. *e*, Embryo, with leafy cotyledons and superior radicle.

or peculiar-hooded (cucullate) appendages, prolonged from the tube of the filaments, which occur in many of the plants of this order, give a peculiar aspect to their flower (see fig. 353). They inhabit chiefly warm and tropical regions, but many species extend to northern climates. Many succulent species are found in the south of Africa. In tropical India and New Holland, and in all the equinoctial parts of America, they also abound. Lindley enumerates 141 genera, including 910 species. *Examples*—*Periploca*, *Asclepias*, *Calotropis*, *Cynanchum*, *Gonolobus*, *Stapelia*, *Hoya*, *Dischidia*.

943. The plants of the order have acrid, purgative, emetic, and diaphoretic properties. The milky juice is usually bitter and acrid, but occasionally it is bland, and is used as milk, as in the case of *Gymnema lactiferum*, the Cow-plant of Ceylon. *Asclepias tuberosa*, the Butterfly-weed, or Pleurisy-root, is used as a cathartic and diaphoretic in North America. The emetic properties of *Asclepias curassavica* have secured for it the name of Wild Ipecacuanha, in the West Indies. The leaves of *Solenostemma* (*Cynanchum*) *Argel*, are used to adulterate Alexandrian Senna. The fragrant roots of *Hemidesmus indicus* are used in Madras as a substitute for Sarsaparilla, under the name of Country Sarza. The bark of the root of several species of *Calotropis*, such as *C. procera*, *Hamiltonii*, and *gigantea*, furnish the substance called Mudar, which is used as a diaphoretic in India. It contains a principle called Mudarine, which gelatinizes on being heated, and becomes fluid on cooling. *Cynanchum monspeliacum* furnishes Montpellier Scammony, and *Periploca mauritiana* is the source of Bourbon Scammony. Both of these substances act as purgatives, and are used to adulterate true Scammony. *Marsdenia tinctoria* and *Gymnema tingens* are said to yield a dye similar to indigo. The milky juice of many of the plants contains caoutchouc in its composition. *Hoya carnosa* receives the name of wax-flower from the peculiar aspect of its blossoms. *Dischidia Rafflesiana*, an Indian climber, has remarkable ascidia (§ 164). The *Stapelias* are singular plants, resembling some of the Cactuses and Euphorbias. Their blossoms are often very fetid, and hence they are called Carrion flowers. Some of the species of *Asclepias* receive the name of Wild Cotton, on account of the hairs attached to their seeds. *Gomphocarpus fruticosus* is the silk-plant of Madeira.

944. Order 125.—**Apocynaceæ**, the Dogbane Family. (*Monopet. Hypog.*) Calyx usually 5-partite, persistent. Corolla hypogynous, gamopetalous, regular, usually 5-lobed, deciduous; æstivation contorted, twisting in some cases to the right, in others to the left. Stamens 5, inserted on the corolla, alternate with its segments; filaments distinct; anthers 2-celled, dehiscing longitudinally; pollen granular, globose, or 3-lobed, immediately applied to the stigma. Ovaries 2, and each unilocular, or uni-, and bilocular; ovules 00; styles 2 or 1; stigma 1, with a contraction in the middle. Fruit follicular or capsu-

lar, or drupaceous or baccate, double or single. Seeds 00, rarely definite, usually pendulous; albumen cartilaginous or fleshy, rarely 0; embryo foliaceous; radicle turned towards the hilum.—Trees or shrubs, usually lactescent, with entire, generally opposite, exstipulate leaves, with interpetiolar cilia or glands. They are chiefly found in tropical regions. They appear to be most abundant in the hot parts of Asia, are less common in the tropics of America, and still less abundant in Africa. Lindley enumerates 100 genera, including 566 species. *Examples*—*Apocynum*, *Echites*, *Nerium*, *Balfouria*, *Vinca*, *Tanghinia*, *Plumieria*, *Carissa*.

945. Many of the plants of this order are poisonous. Some are used medicinally, as cathartics, and there are a few which yield edible fruits. The order is in general to be regarded with suspicion. One of the most deadly plants of the order is *Tanghinia venenata* (*Cerbera Tanghin*), the seeds of which supply the famous Tanghin poison, used formerly in Madagascar as an ordeal in cases of criminals. *Nerium Oleander*, the common Oleander, is poisonous. The stomata of its leaves are formed by means of cellular hair-like processes (fig. 76), and the anthers are terminated by feathery appendages (fig. 334). Death has ensued from eating the flowers of this plant. Its branches, when divested of their bark and used as skewers, have rendered poisonous meat roasted on them, and so proved fatal to seven out of twelve of those who partook of it. The roots of *Apocynum cannabinum* and *androsæmifolium* are said to be emetic. The *Vincas*, Periwinkles, are astringent and acrid. *Allamanda cathartica*, a native of Ceylon and Java, is emetic and cathartic. Although the milky juice is generally acrid, still in some instances it is bland. Thus, the juice of *Tabernæmontana utilis*, Hya-hya, the Cow-tree or Milk-tree of Demerara, is used as milk. Many of the plants, such as *Urceola elastica* and *Vahea gummifera*, supply caoutchouc. *Wrightia tinctoria* yields a dye like indigo. *Aspidosperma excelsum* is a Guiana tree, remarkable for the sinuous arrangement of its wood, which gives the stem a deeply-fluted appearance.

946. Order 126.—**Loganiaceæ**, the Logania Family. (*Monopet. Hypog.*) Calyx 4-5-partite (fig. 287 c); æstivation valvate or imbricate. Corolla hypogynous, regular or irregular, 4-5- or 10-divided (fig. 287 t l); æstivation convolute or valvate. Stamens inserted on the corolla, 5 or 1, not always corresponding with the divisions of the corolla; pollen elliptical or triangular, simple, or marked with three bands. Ovary free, usually 2-celled; ovules 00 or solitary, peltate and amphitropal, or ascending and anatropal. Fruit a 2-celled capsule, with placentas finally becoming loose; or a nuculanium with 1 or 2-seeded nucules; or baccate, with seeds immersed in a pulp. Seeds usually peltate, sometimes winged; albumen fleshy or cartilaginous; embryo small; radicle turned towards the hilum, or parallel

with it.—Shrubs, herbs, or trees, with opposite entire leaves, and usually with stipules, which adhere to the footstalks, or form interpetiolar sheaths. They inhabit chiefly tropical and warm climates. The order is divided into three suborders:—1. *Loganiæ*, æstivation of corolla convolute, fruit a bilocular capsule or nuculanum, seeds peltate, sometimes winged. 2. *Strychnæ*, æstivation of corolla valvate, fruit a 2-3-celled berry or capsule, seeds peltate, embryo rather large. 3. *Spigeliæ*, æstivation of corolla valvate, fruit a didymous capsule, seeds apterous, embryo small, cotyledons inconspicuous. There are about 24 known genera, and nearly 170 species. *Examples*—*Logania*, *Potalia*, *Strychnos*, *Spigelia*.

947. The plants of this order are highly poisonous. They act energetically on the spinal marrow, causing tetanic spasms, or they produce narcotic symptoms by acting on the brain. Many are very bitter and a few are tonic. *Strychnos Nux-Vomica*, the Poison-nut or Koochla, a tree which abounds on the Malabar and Coromandel coasts, supplies the substance called Nux-Vomica. It yields fruit of the size and appearance of an orange, with a coriaceous reddish integument, enclosing a mucilaginous pulp. The seeds, which are embedded in the pulp, are the officinal part of the plant. They are circular and flat, umbilicated on one surface, and are thickly covered with brown silky hairs. All parts of the plant, especially the seeds and bark, are intensely bitter. The seeds contain two alkaloids, *Strychnia* and *Brucia*, to which they owe their poisonous properties. These alkaloids occur in combination with *Igasuric* or *Strychnic* acid. *Nux-Vomica* and *Strychnia*, in poisonous doses, cause death by producing tetanic spasms in the muscles of respiration. The bark of the *Nux-Vomica* tree is the false *Angustura* bark, and the wood is often called *Snake-wood*. *Strychnia* exists in other species of *Strychnos*, as *S. Ignatia* (*Ignatia amara*), St. Ignatius's Bean, *S. colubrina*, and *S. ligustrina*, *Snake-wood*, *S. Tieuté*, the source of a Java poison called *Upas Tieuté*. It is also said to exist in the *Woorali* or *Ourari* poison of *Guiana*, which some consider to be the produce of *S. toxicaria* or *guianensis*. The effects of this last mentioned poison, however, do not seem to agree with those of *Strychnia*. *Strychnia* stimulates the spinal cord without affecting the function of the brain. It causes convulsive twitches of the muscles of the arms and legs, and hence it has been recommended in cases of chronic palsy, unconnected with any signs of local irritation or determination of blood to the head. Its administration requires great caution, as $\frac{2}{3}$ of a grain have been known to produce alarming lock-jaw, and $\frac{1}{6}$ of a grain has killed a dog. Some species of *Strychnos* seem not to possess a poisonous principle in large quantity, for they are used as tonics and febrifuges. Among them may be noticed *Strychnos potatorum* and *pseudoquina*. The former is called *Clearing-nut*, and is used in *India* for purifying water. The

root of *Spigelia marilandica*, Carolina Pink-root (fig. 287), is used as an anthelmintic, more particularly in the United States. *S. Anthelmia*, Guiana Pink-root, is employed in Demerara for a similar purpose. These plants also possess narcotic qualities.

948. Order 127.—**Gentianaceæ**, the Gentian Family. (*Monopet. Hypog.*) (See fig. 248.) Calyx gamosepalous, usually in 5 divisions, sometimes 4-6-8 or 10 divisions, persistent. Corolla gamopetalous, hypogynous, usually regular and marcescent; limb sometimes fringed, divided into as many lobes as the calyx; æstivation plaited or imbricate-twisted. Stamens inserted upon the corolla, alternate with its segments, and equal to them in number, some of them occasionally abortive. Ovary composed of 2 carpels, unilocular or partially bilocular (fig. 389); ovules 00, anatropal; style 1, continuous; stigmas 1 or 2. Fruit capsular or baccate, 1-celled (fig. 389), usually bivalvular, with septicidal, or rarely loculicidal dehiscence. Seeds 00, small, embryo straight, minute, in the axis of soft fleshy albumen; radicle next the hilum.—Herbs, seldom shrubs, with opposite (fig. 248), rarely alternate, entire or divided, exstipulate leaves, which are often 3-5-ribbed. The plants of the order are distributed generally over the globe, inhabiting both cold and warm regions. They are rare in the arctic and antarctic islands. They exhibit great varieties of colours, and many are prized for their beauty. There are two suborders:—
1. Gentianeæ, æstivation of corolla imbricate-twisted, leaves opposite, simple, and entire. 2. Menyantheæ, æstivation of corolla plaited or induplicate, leaves usually alternate and compound, or divided. Lindley mentions 60 genera, including 450 species. *Examples*—*Gentiana*, *Chironia*, *Agathotes*, *Erythræa*, *Chlora*, *Menyanthes*, *Vilarsia*.

949. The general property of the plants of this order is bitterness, which pervades all their organs. Hence they are used as tonics. The medicinal gentian is the root of *Gentiana lutea*, a plant which grows abundantly on the Pyrenees, and on the Alps of Switzerland and Austria, usually at an elevation of 3000 to 5000 feet. It produces showy yellow flowers, and its root is yellow internally. It is administered in the form of extract, infusion, tincture, and wine, as a tonic. Its roots are often mixed with the roots of other species, such as *Gentiana punctata*, *purpurea*, and *pannonica*. *Gentiana Kurroo* of the Himalayas has similar properties. The British species, *Gentiana campestris* and *Amarella*, have also been used as bitter tonics. The officinal Chiretta is the herb and root of *Agathotes Chirayta* (*Ophelia Chirata*), a herbaceous plant found in the Himalayas. The whole plant is bitter, and has been long used in Bengal as a tonic and stomachic. The flowering cymes of *Erythræa Centaurium*, common Centaury (fig. 248), are used as a substitute for gentian, and so are the leaves of *Menyanthes trifoliata*, Buck-bean, Marsh-trefoil, or Bog-

Gentianaceæ
Cym-3-5
leaves
1-celled baccate

bean. The roots of *Frasera Walteri* sometimes receive the name of American Calumba.

950. Order 128—**Bignoniaceæ**, the Trumpet-Flower Family. (*Monopet. Hypog.*) Calyx divided or entire, sometimes spathaceous. Corolla monopetalous, hypogynous, usually irregular, 4-5-lobed. Stamens 5 and unequal, or 4 and didynamous, some of them occasionally sterile; anthers bilocular. Disk annular or glandular. Ovary superior, 1-2-celled, each cell being often spuriously divided; ovules indefinite; style 1; stigma bilamellar (fig. 406), or 2-4-cleft or entire. Fruit a 2-celled (sometimes spuriously 4-celled) and 2-valved capsule, occasionally succulent. Placentas parietal, sometimes extending to the centre, and forming a spurious dissepiment, which finally separates, bearing the seeds. Seeds winged or wingless, often flat and compressed, exalbuminous; embryo straight; radicle next the hilum.—Trees, shrubs, or herbs, with opposite, rarely alternate, exstipulate leaves. They abound generally in tropical regions, but some of them are widely distributed. The order has been divided into four sub-orders:—1. Bignoniæ, capsule 2-valved, 2-celled, sometimes spuriously 4-celled, with a dissepiment parallel or contrary to the valves, at length free, bearing the seeds, which are transverse, compressed, and winged. 2. Cyrtandræ (Didymocarpeæ), fruit succulent or capsular, or siliquose and 2-valved, seeds small, ovate, or cylindrical, suspended, apterous, sometimes comose. 3. Crescentiæ, fruit woody, and melon-shaped, enclosing large seeds, which are immersed in the pulp of the placentas. 4. Pedaliæ, fruit drupaceous, rarely capsular and 2-valved, spuriously many-celled; seeds few, large and apterous, pendulous, erect or transverse. These are reckoned separate orders by many. There are upwards of 100 known genera, and about 650 species. *Examples*—*Bignonia*, *Spathodea*, *Eccremocarpus*, *Cyrtandra*, *Didymocarpus*, *Crescentia*, *Petalium*, *Sesamum*.

951. There are many showy plants in this order. Their flowers are frequently large and trumpet-shaped. None of them are noted for marked medicinal properties. Some are timber trees, others furnish dyes and articles of diet, while a few have bitter and astringent qualities. The species of *Bignonia* are conspicuous objects in tropical forests. Their wood sometimes exhibits a crucial arrangement (fig. 107.) From *Bignonia Chica* the Indians extract a red ochreous colouring matter, with which they paint their bodies. *Crescentia Cujete*, the Calabash-tree, is found in the tropical regions of America, and pro-

with hooked processes. In the perfect fruit of *Pretrea* (*Martynia Zanguebarica*), there are 6 cells formed by the mode in which the placentas unite, and of these cells two are seedless.

952. Order 129.—**Polemoniaceæ**, the Phlox Family. (*Monopet. Hypog.*) Calyx inferior, in 5 divisions, persistent, sometimes irregular. Corolla regular, rarely irregular, 5-lobed. Stamens 5, inserted on the middle of the tube of the corolla, and alternate with its segments; pollen often blue. Disk lobed. Ovary free, 3-celled; ovules anatropal or amphitropal; style simple; stigma trifid. Fruit a 3-celled, 3-valved capsule, with septifragal dehiscence. Seeds angular or oval, or winged, often enveloped in mucus, containing spiral threads, ascending, in a single or a double row; embryo straight, in the axis of a fleshy or horny albumen; cotyledons foliaceous, elliptical or cordate; radicle inferior, next the hilum.—Herbaceous or climbing plants, with opposite or alternate, simple or compound leaves. They inhabit temperate countries chiefly, and they abound in the north-western part of America. There are 17 genera enumerated by Lindley, including 104 species. *Examples*—*Polemonium*, *Phlox*, *Cobæa* (fig. 318).

953. Many of the plants of this order have showy flowers, and are commonly cultivated in flower-borders. Connected with the episperm of various species of *Collomia* are numerous spiral cells, and when the seeds are moistened with water, the mucus surrounding the cells is dissolved, so that the spiral fibres are uncoiled. The movements of these fibres, when uncoiling, are beautifully seen under the microscope. The fibres carry with them a mucous envelope which has the appearance of a membrane. *Polemonium cæruleum*, Greek Valerian, or Jacob's ladder, is bitter. In Siberia, poultices are prepared from its leaves, and thought serviceable in syphilitic sores. The Russians fancy that a decoction of it is useful in hydrophobia.

954. Order 130.—**Hydrophyllaceæ**, the Hydrophyllum Family. (*Monopet. Hypog.*) Calyx 5-parted, persistent. Corolla monopetalous, hypogynous, regular, 5-cleft; æstivation plicate or imbricate. Stamens 5, inserted upon the corolla, and alternate with its segments; filaments sometimes petaloid; anthers deeply-lobed at the base, often versatile, 2-celled, dehiscing longitudinally or transversely. Disk annular or 0. Ovary free, 1-2-3-celled; ovules definite or indefinite; style 1 or 2; stigmas usually 2. Fruit capsular, 2-valved, 1-2-celled, with a parietal, or a large central placenta. Seeds with a brittle or reticulated testa; embryo in the midst of fleshy or cartilaginous albumen; radicle next the hilum.—Trees, shrubs, or herbs, with opposite, or alternate, exstipulate, often lobed leaves. They occur both in the northern and southern parts of America chiefly. They have no properties of importance. Many have showy flowers, and some have glandular or stinging hairs. The order has been divided into two suborders:—1. *Hydrophyllææ*, including *Hydrolleæ* of authors, with the anthers

dehiscing longitudinally, disk present, ovary 1-2-celled, styles 2.
2. Diapensiæ, with anthers dehiscing transversely, disk 0, ovary 3-celled, style single. There are 18 known genera, and 77 species.
Examples—Hydrophyllum, Hydrolea, Nemophila, Phacelia, Diapensia.

955.—Order 131.—**Convolvulaceæ**, the Convolvulus or Bindweed Family. (*Monopet. Hypog.*) Calyx in five divisions, persistent, imbricated, often bracteated (figs. 652-654). Corolla monopetalous, hypogynous, deciduous, regular; limb 5-lobed, with a plaited or imbricated æstivation (fig. 653 *p*); tube sometimes with scales, alternate



with the lobes of the limb. Stamens 5, inserted in the base of the corolla, and alternate with its lobes (fig. 654 *e*). Disk annular,

Figs. 652-659.—Organs of fructification of *Convolvulus* (*Calystegia*) *sepium*, to illustrate the natural order Convolvulaceæ.

Fig. 652.—Diagram of the flower, showing two bracts, five unequal divisions of the imbricated calyx, five lobes of the plicate corolla, five stamens alternating with the corolline lobes, and a quadrilocular ovary.

Fig. 653.—Flower bud. *b*, Large bracts. *c*, Calyx. *p*, Corolla.

Fig. 654.—Vertical section of the lower part of the flower. *b*, Bracts. *c*, Calyx. *p*, Tube of corolla, bearing the filaments of the stamens, *e*, *o*, Ovary. *s*, Style.

Fig. 655.—Summit of the style and stigmas.

Fig. 656.—Fruit, *f*, surrounded by the calyx, *c*, and the bracts, *b*, which are persistent.

Fig. 657.—Seed. *h*, Hilum.

Fig. 658.—Section of the seed showing the corrugated cotyledons.

Fig. 659.—Embryo separated.

Convolvulus
sepium habit—Plaited Corolla
 disc. 2-4 celled ovary

hypogynous. Ovary free, 2-4-celled, rarely by abortion 1-celled; ovules definite, erect, when more than one, collateral; style 1 (fig. 654 s), usually bifid, rarely 2; stigmas obtuse or acute (fig. 655). Fruit succulent or capsular (fig. 656), 1-4-celled, with septifragal and septicial, or circumscissile dehiscence. Seeds albuminous; embryo curved or spiral (figs. 659, 502); cotyledons corrugated (fig. 658) or inconspicuous; radicle inferior.—Herbs or shrubs, usually twining, sometimes parasitical, often with a milky juice, and with alternate, undivided or lobed, exstipulate leaves, rarely leafless. They occur chiefly in tropical and temperate regions. A few only are found in cold climates. The order has been divided into two suborders:—1. Convolvuleæ, true Bindweeds, leafy plants, with the corolline tube not scaly, embryo curved, cotyledons conspicuous. 2. Cuscutæ, Dodders, leafless parasites, having scales on the corolline tube, embryo spiral, and filiform (fig. 502), cotyledons inconspicuous. There are 45 genera, and upwards of 700 species. *Examples*—*Convolvulus*, *Ipomœa*, *Exogonium*, *Dichondra*, *Cuscuta*.

956. The order is characterized generally by the presence of an acrid juice in the roots, which has purgative properties. On this account several of the plants are used medicinally. The old genus *Convolvulus* has been split into various genera; such as *Ipomœa*, *Exogonium*, *Pharbitis*, *Batatas*, *Quamoclit*, *Calonyction*, and *Lepistemon*, according to the form of the corolla, the exertion or inclusion of the stamens, the form and nature of the stigma, and the structure of the ovary. *Exogonium Purga* (*Ipomœa Purga*, or *Convolvulus Jalapa*) is the Jalap plant, a native of Mexico, which grows well in this country, requiring only the protection of a frame during winter. The plant has flowered regularly for many years in a cold frame in the Edinburgh Botanical Garden. The root-stock is the officinal part. It has a roundish tuberous form, is black externally, white and milky within, and varies in size from that of a walnut to that of a moderate-sized turnip. It contains a resin, in which its active properties reside. It is used in the form of powder and tincture, as an active irritant cathartic. The root of *Convolvulus Scammonia* yields a gummy resinous exudation, which constitutes medicinal Scammony. The plant grows abundantly in Greece, the Grecian Islands, and various parts of the Levant. Scammony is procured by cutting the root across, and collecting the milky juice which soon concretes. The drug is imported into this country from Smyrna. Its active principle is a resin. It is used medicinally as a drastic purgative, in the form of powder, pill, and extract. A spurious kind of Scammony has been prepared from the root of *Convolvulus (Calystegia) sepium*; and several plants belonging to the natural order Asclepiadaceæ yield a purgative exudation, which has been used under the names of Montpellier and Bourbon Scammony (§ 124). The roots of some of the plants do not possess


purgative qualities, and have been used as articles of food. *Batatas perulidis* (*Convolvulus Batatas*) yields the sweet Potato, which contains much saccharine and amylaceous matter, and is used as food in tropical countries. The species of *Cuscuta*, or Dodder, generally reckoned a suborder of *Convolvulusæ*, have acrid purgative properties. Their seeds germinate in the soil, and the plants afterwards twine round others, and become attached to them by means of suckers. They then lose their connection with the soil, and are supported as true parasites. In this way they often destroy crops of Flax and Clover. M. Choisy objects to the separation of Didders from Bindweeds, but, as under the name of *Cuscuta* are included species with a very variable structure which might constitute a genera, adds that they should have a claim to be regarded as a peculiar order, if as many as 200 species were known instead of 50. Didders differ from Bindweeds in having a thread-shaped embryo composed almost exclusively of radicle, and twisted spirally in a mass of fleshy albumen. They also have generally, perhaps always, scale-like bodies at the base of the stamens, and apparently alternating with the lobes of the corolla. They have therefore by Lindley and others been classed as a separate order.

957. Order 132.—**Cordiaceæ**, the Cordia Family. (*Monopet. Hypog.*) Calyx 4-5-toothed, inferior. Corolla monopetalous, 4-5-cleft, regular. Stamens inserted on the corolla, alternate with its segments; anthers versatile. Ovary free, 4-8-celled; ovules solitary, pendulous, anatropal; style continuous; stigma 4-8-cleft. Fruit drupaceous, 4-8-celled. Seed exalbuminous, pendulous from the apex of the cell by a long funiculus, upon which it is turned back; radicle superior: cotyledons plaited longitudinally.—Trees, with alternate, rough, exstipulate leaves, and paniced flowers. They are chiefly natives of warm countries. Some yield edible fruits; their bark is occasionally bitter, tonic, and astringent, and their wood is used for various economical purposes. The succulent, mucilaginous fruits of *Cordia Myxa* and *Sebestena*, receive the name of Sebesten Plums. There are 11 genera enumerated by Lindley, including 180 species. *Examples*—*Cordia*, *Varronia*.

958. Order 133.—**Boraginaceæ**, the Borage Family. (*Monopet. Hypog.*) Calyx persistent, with 4-5-divisions (figs. 660, 661 c). Corolla gamopetalous, hypogynous, usually regular (figs. 296, 297), 5- rarely 4-cleft; æstivation imbricated (figs. 660, 661 pp). Stamens inserted on the corolla, equal in number to its segments, and alternate with them (fig. 661 e). Ovary usually 4-lobed, quadrilocular (fig. 661 o); ovules 4, each attached to the lowest point of the cavity, amphitropal; style simple, basilar (figs. 403, 661 s), (terminal in Ehretieæ and Heliotropieæ); stigma simple or bifid. Fruit (fig. 662) consisting of 2 to 4 distinct achænia (succulent and consolidated in Ehretieæ). Seed

Boraginaceae -
Alternat. opposit. leaves - Regular ^{vs} Pent. flowers
4 lobed corolla - Simple style -

exalbuminous, or with thin albumen; radicle superior; cotyledons plano-convex (fig. 662).—Herbs, shrubs, or trees, with terete stems,



alternate, rough, exstipulate leaves, and flowers generally in scorpioid (gyrate) cymes (fig. 253). On account of the asperities in the leaves, the plants have sometimes been called *Asperifoliæ*. The order is divided into three suborders:—1. *Boragineæ* (figs. 660-662), with a basilar style, 4-lobed ovary, achæmium-like fruit, and exalbuminous seeds; natives chiefly of temperate climates. 2. *Ehretiæ*, with a terminal style, a quadrilocular, concrete ovary, a succulent fruit, and usually albuminous seeds; natives of tropical countries. 3. *Heliotropiæ*, with a terminal style, an entire or 2-lobed ovary, a dry fruit separable into four achænia, and exalbuminous seeds; natives partly of temperate, and partly of warm climates. There are 67 known genera, and nearly 900 species. *Examples*—*Borago*, *Anchusa*, *Echium*, *Myosotis*, *Cynoglossum*, *Ehretia*, *Heliotropium*.

959. The plants of the order are generally mucilaginous and emollient. Some are astringent. Nitrate of potash exists in some, and imparts coolness to the water in which they are steeped. *Borago officinalis*, Borage, has been used for its mucilaginous emollient properties, as a remedy in pectoral affections; and with wine, water, lemon, and sugar, its leaves form an ingredient in what is called cool-tankard. Attached to the stamens in this plant, and others of the order, are scales, which may be considered as abortive stamens, formed by dilamination

Figs. 660-662.—Organs of fructification of *Anchusa Italica*, to illustrate the natural order Boraginaceæ.


Fig. 660.—Diagram of the flower, with five imbricated divisions of the calyx, five imbricated segments of the corolla, five stamens, and a 4-lobed ovary.

Fig. 661.—Vertical section of the flower. *c*, Hairy calyx. *p p*, Corolla. *e*, Stamens inserted into the corolla. *a a*, Staminal appendages or corolline scales. *o*, 4-lobed ovary, two of its divisions cut through vertically. *s*, Basilar style.

Fig. 662.—One of the carpels (achænia) cut vertically. *p*, Pericarp separable from the seed. *t*, Spermoderm or integuments of the seed. *e*, Embryo with superior radicle, and plano-convex cotyledons.

(fig. 312). *Anchusa tinctoria* supplies alkanet root, which is used as a reddish-brown dye. Some of the species of *Heliotropium* (as *H. peruvianum*) are distinguished by their fragrant odour. The leaves of *Steenhammera* (*Lithospermum*) *maritimum* have the taste of oysters, and hence it is called the Oyster-plant in Scotland. *Myosotis palustris* (fig. 279) is the true Forget-me-not. Miss Strickland remarks, that the banished and aspiring Henry of Lancaster appears to have been the person who gave to this plant its emblematical and poetical meaning, by uniting it in his exile with the initial letter of his watchword, 'Sou-veigne-vous-de-moy.'

960. Order 134.—**Solanaceæ**, the Nightshade Family. (*Monopet. Hypog.*) Calyx inferior, 5- rarely 4-partite, persistent (fig. 664 c). Corolla monopetalous, hypogynous, with the limb 5- rarely 4-cleft, regular, or somewhat unequal, deciduous; æstivation plicate or imbricated (fig. 663). Stamens inserted on the corolla (fig. 664 e), equal



in number to the corolline segments, and alternate with them (fig. 663); anthers with longitudinal or porous dehiscence (fig. 664 e). Ovary usually 2-celled (fig. 664 o), sometimes 4-5- or many celled; ovules indefinite; style continuous; stigma simple (fig. 664 s). Fruit with 2, 4, or more cells, rarely unilocular; either a capsule dehiscing in a

Figs. 663-668.—Organs of fructification of *Solanum tuberosum*, the Potato, to illustrate the natural order Solanaceæ.

Fig. 663.—Diagram of the flower, with five divisions of the calyx, five plicate segments of the corolla, five stamens, and a 2-celled ovary with polyspermous placentas. *a*, Axis.

Fig. 664.—Vertical section of the flower. *c*, Calyx. *p p*, Lower part of the corolla. *e*, Stamens, with porous dehiscence of the anthers. *o*, Bilocular ovary, polyspermous. *s*, Style and stigma.

Fig. 665.—Fruit, baccate.

Fig. 666.—Horizontal section of the fruit, showing the seeds and placenta.

Fig. 667.—The seed.

Fig. 668.—Vertical section of the seed. *t*, Integument (spermoderm) of the seed. *p*, Fleshy perisperm (albumen). *e*, Embryo, which is curved and excentric, with the radicle next the hilum.

Solanaceæ - Potandrom - Flowers - Plicate corolla - bilocular ovary

septicidal or circumscissile manner, and having a double dissepiment parallel to the valves, or a berry (figs. 665, 666) with the placentas adhering to the dissepiment, or a nuculanum with 5 or more nucules. Seeds 00; embryo straight or curved (fig. 668), often excentric, lying in fleshy albumen; radicle next the hilum.—Herbs or shrubs, with alternate leaves. Natives of most parts of the world, but abundant in the tropics, in which the mass of the order exists, in the form of the genera *Solanum* and *Physalis*. The order has been divided into two sections, which are not, however, well defined:—1. *Rectembryæ*, with the embryo nearly straight. 2. *Curvembryæ*, with the embryo curved in a semicircular, annular, or spiral manner. These are subdivided into tribes, chiefly according to the nature of the fruit. There are 66 known genera, and 935 species enumerated; but there is considerable difficulty in the allocation of the genera. *Examples*—*Cestrum*, *Habrothamnus*, *Nicotiana*, *Petunia*, *Datura*, *Hyoscyamus*, *Solanum*, *Physalis*, *Nolana*.

961. The plants of this order have in general narcotic qualities. These are sometimes developed in a great degree, so as to render the plants very poisonous; at other times they are obscured by the prevalence of nutritious and starchy matter. In many instances, certain parts of the plant have poisonous narcotic properties, while other parts are harmless, and are used as articles of diet. These facts will be illustrated by a consideration of different genera and species. *Solanum Dulcamara*, Bitter-sweet or woody Nightshade, has slightly narcotic properties, which appear to be owing to the presence of an alkaloid called Solania. The twigs of the plant are used medicinally, in the form of decoction, in various cutaneous diseases. They are supposed to possess sudorific and alterative properties. The scarlet berries of the plant are reckoned poisonous. *Solanum nigrum* has more active narcotic qualities. *Solanum tuberosum*, the Potato (fig. 92), has slight narcotic qualities in its leaves and fruit, but in the tubers there is an abundance of starch, and when cooked they are wholesome and nutritious. In the genus *Solanum* the anthers open by pores. *Atropa Belladonna*, Deadly Nightshade or Dwale, is a highly poisonous plant belonging to this order. All parts of the plant are narcotic. The fruit is a dark brownish-black shining berry, which often proves attractive to children. The leaves are the part used in medicine, and from these an extract is prepared. The watery extract is best made in vacuo, but the alcoholic extract is probably the best. *Belladonna* is one of our most active indigenous poisons. It owes its properties to the presence of an alkaloid called Atropia, which exists in the plant in combination with malic acid. *Belladonna* is used medicinally to allay pain and spasmodic action, to cause dilatation of the pupil, and as a prophylactic against scarlatina. Similar properties exist in the species of *Hyoscyamus*, more especially in *H. niger*, Henbane, a biennial plant,

with dingy-yellow flowers, exhibiting beautiful purple reticulations, hairy viscous leaves, and a bilocular operculate capsule (fig. 459). The leaves yield by expression a large quantity of juice, whence an extract is prepared. A tincture of Henbane is often used in place of laudanum, on account of not causing constipation. It is employed in medicine to procure sleep and allay pain, and it acts also in dilating the pupil. The narcotic properties seem to be owing to an easily decomposed alkaloid called Hyoscyamia. An empyreumatic oil is obtained from the plant, which is an energetic narcotic poison. The roots of the plant have sometimes caused poisoning by being mistaken for parsnips. Many species of *Datura* are powerfully narcotic. *D. Stramonium* is the Thorn-apple, so called on account of its prickly capsule. Its leaves and seeds are used medicinally as narcotics, their qualities being due to an alkaloid called Daturia. They are prescribed as anodynes and antispasmodics, in the form of powder, extract, and tincture, and the leaves are smoked in cases of asthma. *Datura Tatula* and *Metel sanguinea*, *ferox* and *fastuosa* have similar properties. Several species of *Nicotiana* furnish Tobacco. That chiefly used in Europe is procured from *N. Tabacum*, a plant naturally inhabiting the hotter parts of North and South America. It is an annual plant, attaining a height of six feet, having dingy-red infundibuliform flowers (fig. 294) and viscid leaves. The leaves are the official part, and their active properties depend on a peculiar oily-like alkaloid called Nicotina. They are employed in the form of infusion, tincture, and wine. Tobacco is an energetic narcotic poison. Its oil, which is inhaled and swallowed in the process of smoking, is one of the most deadly known poisons. The Hottentots are said to kill snakes by putting a drop of it on their tongues: the death of these reptiles is said to take place instantaneously. It is employed medicinally as a sedative, and its depressing action is useful in cases of hernia. Its depressing action is indicated by its effect on the cerebral functions and on the heart. The flavour and strength of tobacco depend on climate, cultivation, and the mode of manufacture. That most esteemed by the smoker is Havannah tobacco; but the Virginian is the strongest. It is said that small Havannah cigars are prepared from the leaves of *Nicotiana repanda*; Syrian and Turkish tobacco from *N. rustica*, and fine Shiraz tobacco from *N. persica*. The fruit of different species and varieties of *Capsicum* supply Cayenne-pepper, and what are called Chillies. Chilli is the Mexican name for all the varieties of *Capsicum*. They are natives of the East and West Indies, and of other hot climates. *Capsicum annuum* is the species commonly noticed, but of it there seems to be numerous varieties, which by many are reckoned species. Thus, *C. frutescens* is a shrubby plant which, along with *C. minimum*, supplies the variety called Bird-pepper; *C. baccatum* has a globular fruit, and furnishes Cherry or berry-capsicum. In *Capsicum irritant*

properties prevail, so as to obscure the narcotic action. Their acridity is owing to an oleaginous substance called Capsicin. Cayenne-pepper is used chiefly in the form of tincture as a rubefacient and stimulant, especially in cases of ulcerated sore throat. It acts on the stomach as an aromatic condiment, and, when preserved in acetic acid, it forms Chilli vinegar. *Mandragora officinalis* (*Atropa Mandragora*), Mandrake, acts as a stimulant on the nervous system, and its forked root was long celebrated for its properties in this respect. It is the דַּמְסָא, from דָּם of the Bible. Its root is easily made to assume the human form, and hence has arisen the stories of the plant shrieking when torn out of the ground. By the Arabs the plant is called Tufah-al-Sheitan, or Devil's apple. The species of *Physalis* are remarkable for their accrescent calyx (fig. 280). The fruit of some, such as *P. peruviana*, Peruvian Winter Cherry, is eaten. The fruit of *Lycopersicum esculentum* is the edible Tomato, or Love-apple; that of *Solanum Melongena* and *ovigerum* is the Egg-apple.

A division of the genera of this order has been made by M. Fee into—1. Those which contain species, all of which are more or less poisonous, as *Datura*, *Hyoscyamus*, *Atropa*, *Nicotiana*, &c. 2. Those which contain some species truly poisonous, and others innocuous or stimulant, as *Solanum* and *Capsicum*. 3. Those having species which are all innocuous, as *Lycopersicum*. Miers has revised the genera included in the orders Solanaceæ and Scrophulariaceæ, and he proposes to institute an intermediate order, Atropaceæ. The general definition of these orders is:—

1. Solanaceæ; Isomerous flowers with a valvate or induplicato-valvate æstivation; inflorescence extra-axillary.

2. Atropaceæ; Isomerous flowers or nearly so, with an imbricate or a peculiar æstivation; inflorescence extra-axillary.

3. Scrophulariaceæ; Anisomerous flowers with imbricate æstivation; organs of inflorescence truly-axillary.

962. Order 135.—**Orobanchaceæ**, the Broom-rape Family. (*Monopet. Hypog.*) Calyx divided, persistent, inferior. Corolla monopetalous, hypogynous, irregular, usually bilabiate, persistent; æstivation imbricated. Stamens 4, didynamous. Disk fleshy. Ovary free, 1-celled, composed of two carpels which stand fore and aft, with 2 or more parietal placentas; ovules 00; style 1; stigma 2-lobed, each of the lobes belong half to each carpel. Fruit capsular, enclosed within the withered corolla, 1-celled, 2-valved. Seeds 00, minute; embryo very minute, at one end of fleshy albumen.—Herbaceous parasitical plants, having scales in place of leaves. They are natives of Europe, more especially the southern parts, and of Asia, North America, and the Cape of Good Hope. Lindley gives 12 genera, and 116 species. *Examples*—Orobanche, Lathræa.

963. The properties of the plants of the order are, in general, astrin-

gency and bitterness. Some have been used as tonics, and as applications to indolent ulcers. The species of *Orobanche* are called Broom-rapes, on account of the ravages they are supposed to commit on the Broom tribe. They attach themselves to the roots of various plants, and are hence called Root-parasites. Different species infest and injure different tribes of plants. Thus, *Orobanche rapum* is parasitical upon Broom and Furze; *O. ramosa*, upon Hemp; *O. rubra*, upon common Thyme; *O. minor*, upon red Clover; *O. barbata*, upon the Ivy; *O. elatior* and *arenaria*, upon different species of Compositæ, as Centaury and Milfoil. The stems of Orobanches have a large central cellular portion, surrounded by numerous fibro-vascular bundles, which are arranged in a circle without any medullary rays. Tubers exist at the lower part, whence subterranean buds are developed. Sometimes the fibro-vascular bundles of the plants, to which the Broom-rapes are attached, are found ramifying in the substance of the parasite. As Broom-rapes possess tubers and ordinary roots, Henfrey is disposed to think that they also derive nourishment from the soil in the usual way. *Lathræa squamaria*, Tooth-wort, is parasitical upon the roots of Hazels, Cherry-laurels, and other trees. *Epiphegus virginiana*, Beech-drops, has been used in powder as an application to cancerous sores. In conjunction with white oxide of Arsenic, it is supposed to have constituted the specific known in North America under the name of Martin's Cancer Powder.

964. Order 136.—**Scrophulariaceæ**, the Figwort Family. (*Monopet. Hypog.*) Calyx divided into 4 or 5 parts, unequal, persistent, inferior (fig. 288 c). Corolla monopetalous, more or less irregular and bilabiate (fig. 288 p), or personate (fig. 300), sometimes spurred or saccate at the base; æstivation imbricate. In the bud, the flowers are regular (fig. 305). Stamens usually 4, didynamous (figs. 344, 346), rarely 5, sometimes 2; anthers bilocular, or unilocular by abortion or adhesion. Ovary free, 2-celled; ovules usually 00; style simple; stigma 2-lobed, rarely entire. Fruit capsular, rarely fleshy, dicarpellary, 2-celled (fig. 445), 2-4-valved, opening by septicidal or loculicidal dehiscence, rarely by pores (fig. 462) or lids, the dissepiments becoming finally loose in the centre (fig. 446). Placentas attached to the dissepiment, and sometimes in the mature fruit becoming central. Seeds definite or 00; embryo straight or slightly curved, included within fleshy albumen.—Herbs, undershrubs, or shrubs, with opposite, whorled or alternate leaves. They are found generally distributed over the globe, both in cold and warm regions. The order has been divided by Bentham into three sections:—1. Salpiglossidæ. 2. Antirrhinidæ. 3. Rhinanthidæ. The characters of these divisions are founded on the nature of the inflorescence, whether centrifugal, centripetal, or compound, and the æstivation of the corolla. There are 176 known genera, and 1,814 species. *Examples*—Schizanthus, Salpiglossis, Cal-


Scrophulariaceae. tree
 Inflorescence corolla -
 Standard or didynamous
 Stamens
 bicarp many -

ceolaria, Verbascum, Antirrhinum, Scrophularia, Pentstemon, Mimulus, Digitalis, Veronica, Rhinanthus, Melampyrum.

965. The plants of the order are usually scentless, or at all events not aromatic. They are acrid and slightly bitter, and some of them are sedative and poisonous. Some of the plants of the order belong to the tribe of Root-parasites. This is particularly the case with species of *Euphrasia*, *Rhinanthus*, *Bartsia*, *Melampyrum*, and *Pedicularis*. These parasites differ from Broom-rapes in having green leaves, and they seem to be apparently independent after they have acquired a certain degree of development. The species of *Mimulus* have a bilamellate stigma, the two lamellæ of which are irritable, and close when irritated. The movements of the stigma are probably in some way connected with fertilization. One of the species, *Mimulus luteus*, has become naturalized in many parts of Britain, as in the neighbourhood of Edinburgh, on the shores of the Clyde, the Isle of Skye, Perthshire, &c. *Mimulus moschatus* is cultivated on account of its musk-like odour. The most important medicinal plant of the order is *Digitalis purpurea*, Foxglove, the leaves and seeds of which are employed in the form of powder, tincture, and infusion. The leaves have a bitter taste, which they retain when carefully dried. In large doses they act as a narcotico-irritant poison, and in small doses they are used as sedative of the circulation, and diuretic. Their continued use causes great slowness of the pulse, and hence their employment in diseases of the heart and in hæmorrhages, such as hæmoptysis. In dropsical cases, especially those connected with diseased heart, *Digitalis* is extensively used. Its active properties are due to the presence of a crystalline principle called Digitalin. Several other species of *Digitalis*, such as *D. lævigata*, *grandiflora*, *lutea*, and *tomentosa*, have similar properties. The leaves of *Scrophularia nodosa*, knotted Figwort, have irritant qualities, and have been used as emetic and cathartic remedies. In the form of ointment and fomentation, they have been applied to diseases of the skin and tumours. The woolly leaves of *Verbascum Thapsus*, Great Mullein, are emollient and slightly narcotic. They have been used in some pectoral affections. The species of *Melampyrum* are called Cow-wheat, in consequence of being relished by cows. *Euphrasia officinalis*, Eye-bright, or Euphrasy, was formerly used in cases of ophthalmia. Some of the species of *Linaria* and *Calceolaria* are used for dyeing. *Linaria vulgaris* exhibits what Linnæus called Peloria (§ 654, 655), by the flowers being 5-spurred in place of 1-spurred, and thus becoming regular. *Gratiola officinalis*, Hedge-hyssop, is bitter and acrid, and is said to enter into the composition of the Eau médicinale, so much vaunted as a remedy for gout. This was formerly called Gratia Dei, on account of its efficiency as a medicine. In over doses it acts as a poison. According to Haller, it renders by its abundance some of the Swiss meadows useless as pastures. The

leaves of *Veronica officinalis* are bitter and astringent, and are sometimes used as tea.

966. Order 137.—**Labiata** (Lamiaceæ of Lindley), the Labiate Family. (*Monopet. Hypog.*) Calyx tubular, inferior, regular or bilabiate, persistent (fig. 670, 672 c). Corolla monopetalous, hypogynous, bilabiate; upper lip entire or bifid, lower 3-lobed (figs. 299, 670, 671). Stamens 4 (fig. 669), didynamous (fig. 671 e), sometimes 2 by abortion, inserted into the corolla, and alternate with the lobes of the lower lip; anthers 2-celled, or 1-celled by abortion, or



by absorption of the septum (fig. 333); connective sometimes large and distractile (fig. 333 c). Disk fleshy. Ovary free, deeply 4-lobed

Figs. 669-673.—Organs of fructification of *Lanium album*, to illustrate the natural order Labiata.

Fig. 669.—Diagram of the flower, with the pentamerous calyx; pentamerous corolla, having two lips, the upper lip being formed of two united petals, the lower of three; four stamens, in consequence of one being undeveloped, and four divisions of the ovary.

Fig. 670.—Entire flower viewed laterally. *c*, Five-cleft calyx. *t*, Tube of the corolla. *ls*, Upper lip of two petals. *li*, Lower lip of three. *s*, Style.

Fig. 671.—The flower cut vertically. *c*, Calyx. *p*, Corolla. *e*, Didynamous stamens. *s*, Style and bifid stigma. *o*, Ovary.

Fig. 672.—Fruit (a tetraechænum) cut vertically, showing the carpels, two of which have been removed. *c*, Persistent calyx. *g*, Fleshy disk or gland. *r*, Gynobasic receptacle bearing the style, *s*, which is basilar, *i. e.* arises from the lower part of the carpels. *o*, Two carpels, which form achænia when ripe.

Fig. 673.—A carpel cut vertically. *p*, Pericarp. *t*, Integument of the seed. *e*, Embryo erect with inferior radicle.

Labiatic Orphos ^{neous} - Irregular dichotomous flowers - 4 lobed ovary -
Simple style

(figs. 402, 669); ovules 4; style 1, basilar (figs. 402, 671 s); stigma bifid (fig. 671 s), usually acute. Fruit consisting of 1-4 achænia, enclosed within the persistent calyx (figs. 402, 672). Seeds erect (fig. 673); albumen either 0, or in small quantity; embryo erect (fig. 673 e); cotyledons flat; radicle inferior.—Herbs or undershrubs, with tetragonal stems, opposite exstipulate leaves, and cymose inflorescence, the flowers being often in verticillasters. Linnæus looked upon the fruit as naked seeds, and hence included many of the plants in the order Gymnospermia of his Didynamous class. They are natives chiefly of temperate regions. Lindley mentions 125 genera, including 2,350 species. *Examples*—*Mentha*, *Salvia*, *Melissa*, *Lamium*, *Teucrium*, *Scutellaria*.

967. The plants of this order are in general fragrant and aromatic, and none of them are poisonous or injurious. Scarcely any are used for ordinary food, although many form grateful condiments. Their leaves contain receptacles of volatile oil, and many of them furnish a stearoptin resembling camphor. Medicinally, many of them are used as carminatives. The species of *Mentha* yield volatile oils. *M. Piperita*, Peppermint, is used as a powerful diffusible stimulant in cases of colic and gastrodynia. The oil is procured by distillation with water, and, when dissolved in rectified spirit, it forms the essence of Peppermint. *Mentha viridis*, Spearmint, is used in the same way as Peppermint; while *M. Pulegium*, Penny-royal, is employed as a pectoral and antispasmodic. *Lavandula vera* (*L. spica*, *officinalis*, and *angustifolia* of authors) yields the best oil of Lavender; while *L. latifolia* furnishes Spike-oil. Like the other volatile oils of the Labiatae, oil of Lavender consists of a fluid oil, or Elæoptin, and a solid crystalline substance, or Stearoptin, analogous to camphor. Lavender is a tonic, stimulant, and carminative. The flowering tops of *Rosmarinus officinalis*, Rosemary, furnish an oil which has similar properties. It is used much in perfumery, and enters into the composition of Eau de Cologne. It is reputed as possessing considerable efficacy in encouraging the growth of hair and in curing baldness. The admired flavour of Narbonne honey is ascribed to the bees feeding on the flowers of this plant. Oils of the same nature are procured from *Origanum vulgare*, Wild Marjoram, *O. Majorana*, Sweet Marjoram, *Melissa officinalis*, common Balm, and *Marrubium vulgare*, white Horehound. Some consider the Hyssop of Scripture, *היסק*, as being *Hyssopus orientalis* (*H. officinalis*, var. *angustifolius*); but Royle looks upon it as one of the Caper plants (*Capparis ægyptiaca*). *Plectranthus graveolens* of some, *Pogostemon suavis* or *P. Patchouly* of others, is the Patchouli plant of the East Indies, which is used as a perfume. It is called in India puchá pát. It yields a volatile oil of a yellowish-green colour. *Lycopus virginicus*, Bugle-weed, and *L. europæus*, Gipsy-wort, are used as astringents and sedatives. Many Labiates, such as Thyme (*Thymus*), Mint (*Men-*

tha), Sage (*Salvia*), Basil (*Ocimum*) Savoury (*Satureia*), &c., are used as culinary vegetables, more particularly to flavour sauces and dishes. The species of *Salvia* are distinguished by having only two stamens in consequence of the abortion of the rest, and by their distractile connective, which separates the anther lobes (fig. 333). In the episperm of the seeds of the species of *Salvia* there are beautiful spiral cells, the fibres of which, like those of *Collomia*, uncoil when moistened with water, and form an interesting microscopic object. *Salvia officinalis*, common sage, has been used in the form of tea as a stomachic. What are called Sage-apples, are galls produced by the puncture of insects on *Salvia pomifera*. The roots of *Ocimum tuberosum* are said to be esculent. *Hyptis membranacea*, one of the Brazilian Labiates, attains the height of twenty or thirty feet.

968. Order 138.—**Verbenaceæ**, the Vervain Family. (*Monopet. Hypog.*) Calyx tubular, persistent, inferior. Corolla monopetalous, tubular, hypogynous, deciduous, limb usually irregular; æstivation imbricated. Stamens usually 4, didynamous, rarely equal, sometimes 2. Ovary free, 2-4-celled; ovules usually 4, erect or pendulous, anatropal or amphitropal; style 1, terminal; stigma bifid or entire. Fruit nucamentaceous or baccate, composed of 2 or 4 achænia united. Seeds 1-4; albumen 0 or fleshy; embryo straight; radicle either inferior or superior.—Trees or shrubs, rarely herbs, with opposite or alternate, exstipulate leaves. The order has been divided into three suborders:—1. Myoporineæ, anthers 2-celled, seed pendulous, radicle superior; natives of the southern parts of America and Africa, and of Australia. 2. Verbeneæ (fig. 237), anthers 2-celled, seed erect, radicle inferior; natives both of the tropical and temperate regions of America, and found also in Asia and in Europe. 3. Selagineæ, anthers 1-celled, seed pendulous, radicle superior; natives chiefly of the Cape of Good Hope, but some are European. There are 75 known genera, and upwards of 770 species. *Examples*—*Myoporum*, *Avicennia*, *Verbena*, *Vitex*, *Tectona*, *Selago*, *Globularia*.

969. Many of the plants of the order are fragrant and aromatic, some are bitter, tonic, and astringent, others are acrid. None of them occur in the British Pharmacopœias. *Aloysia citriodora*, Sweet-scented Verbena, is commonly cultivated for its fragrance. In the leaves, Mr. Murchison has noticed peculiar glands containing oily matter. The species of *Avicennia* have adventitious roots like the Mangrove. The bark of *Avicennia tomentosa* is used in Brazil for tanning. *Tectona grandis* is the Teak-tree of India, the wood of which is very hard and durable, and is used for ship-building. A fine specimen exists in the Edinburgh Botanic Garden. The trunk of the tree in Eastern forests sometimes attains a height of two hundred feet, and its leaves are twenty inches long by sixteen broad. Roxburgh asserts that the wood is the only useful part of it, but Endlicher states that its flowers are diuretic,

that its foliage yields a red dye, and that a decoction of it is employed by the Malays in cholera. The fruit of several species of *Vitex* is acrid and aromatic. Some species of *Lantana* and *Stachytarpheta* are used for tea. The Vervain was a sacred plant among the Greeks, and it was looked upon by the Druids with superstitious reverence.

970. Order 139.—**Acanthaceæ**, the Acanthus Family. (*Monopet. Hypog.*) Calyx with 4-5 divisions, equal or unequal, occasionally multifid, or entire and obsolete, persistent. Corolla monopetalous, hypogynous, usually irregular, with the limb ringent or bilabiate, or rarely unilabiate, sometimes nearly equal, deciduous. Stamens inserted on the corolla, usually 2, sometimes 4, didynamous, the shorter ones being occasionally sterile; anthers 1-2-celled, with longitudinal dehiscence. Disk glandular. Ovary free, 2-celled; placentas adhering to the axis; ovules 2 or more in each cell, curved; style 1; stigma 2-lobed, rarely entire. Fruit a 2-celled capsule, dehiscing by 2 elastic valves, in a loculicidal manner. Seeds 2 or many in each cell, sometimes solitary, roundish, attached to hard, persistent, hooked or subulate ascending processes of the placenta; testa loose; albumen 0; embryo curved or straight; cotyledons large, leafy; radicle cylindrical, next the hilum.—Herbaceous plants or shrubs, with opposite, exstipulate, simple leaves, and bracteated flowers; 2 or 3 large leafy bracts accompanying each flower. They abound in tropical regions. The order has been divided into three tribes:—1. Thunbergiæ, with the placental processes in the form of a hard cup, supporting the seed. 2. Nelsoniæ, with the placental processes contracted into a papilla, bearing the seed, which is small and pitted. 3. Echmatacanthi, with the placental processes hooked. There are 105 genera, according to Lindley, and about 750 species. *Examples*—Thunbergia, Nelsonia, Acanthus, Justicia, Ruellia.

971. The plants of the order have mucilaginous and bitter properties in general, but they are not put to important uses. The leaves of *Acanthus*, with their sinuated lobes, gave origin to the capital of the Corinthian pillar. *Acanthus mollis* has emollient qualities. The seeds of *Acanthodium spicatum* have beautiful spiral cells in their episperm (§ 11). The style of *Ruellia* (*Goldfussia*) *anisophylla* exhibits a peculiar irritability; its curved stigmatic apex becoming gradually straightened so as to come into contact with the hairs of the corolla upon which the pollen is scattered. A valuable dye of a deep-blue colour, called Room, is obtained in Assam from a species of *Ruellia*. Many of the species of *Justicia*, *Ruellia*, and *Aphelandra*, are cultivated in hot-houses on account of their showy flowers.

972. Order 140.—**Lentibulariaceæ**, the Butterwort Family. (*Monopet. Hypog.*) Calyx inferior, divided, persistent. Corolla monopetalous, hypogynous, irregular, bilabiate, usually spurred. Stamens 2, inserted into the base of the corolla, and included; anthers monothecal,


sometimes contracted in the middle. Ovary free, composed of 2 carpellary leaves, unilocular; ovules 00, anatropal; placenta free, central; style 1, very short; stigma bilamellar. Fruit a 1-celled capsule, dehiscing transversely, or by an apicular cleft. Seeds numerous, minute, exalbuminous; embryo sometimes undivided; radicle next the hilum.—Aquatic or marsh herbaceous plants, with radical leaves, which are sometimes compound, and bear little bladders or ampullæ. Flowers often on scapes. They are found in all parts of the world, and abound in the tropics. Lindley enumerates 4 genera, including 173 species. *Examples*—*Utricularia*, *Pinguicula*.

973. The plants of the order have no properties of importance. The name of Butterwort, given to the species of *Pinguicula*, may be derived from the property of giving consistence to milk. Others say that it has reference to the greasy appearance of their foliage. Linnæus declares that the solid milk of the Laplanders is prepared by pouring it warm and fresh from the cow over a strainer on which fresh leaves of *Pinguicula* have been laid. Of the four British species, one (*P. grandiflora*) is peculiar to Ireland, and another (*P. alpina*) is peculiar to Scotland. *Utricularias*, Bladderworts, are so called on account of the utricles or bladders connected with the leaves (§ 164). In the interior of these vesicles a mucous fluid is found along with cellular projections in the form of hairs. *Utricularia nelumbifolia* grows in the water which collects in the bottom of the leaves of a large *Tillandsia* in Brazil. It sends out runners and shoots, and often in this way unites several plants of *Tillandsia*. The leaves are peltate, and more than three inches across, while the flowering stem is two feet long.

974. Order 141.—**Primulaceæ**, the Primrose Family. (*Monopet. Hypog.*) Calyx 5- rarely 4-cleft (fig. 272), inferior or half superior, regular, persistent (figs. 675 c, 677). Corolla monopetalous (fig. 295 p), hypogynous (fig. 675), rarely perigynous, with the limb 5- rarely 4-cleft, sometimes 0 (fig. 543). Stamens inserted on the corolla, equal in number and opposite to its segments (figs. 674, 675). Ovary free, (figs. 675, 676 o), rarely adherent to the base of the calyx, 1-celled; ovules 00, usually amphitropal; style 1 (fig. 675 s); stigma capitate (fig. 675). Fruit a capsule, opening with valves (fig. 677), or with a lid (fig. 458). Seeds numerous, peltate (fig. 678), attached to a free central placenta (fig. 677); embryo straight (fig. 680), enclosed within fleshy albumen, and lying across the hilum (fig. 679).—Herbaceous plants, with leaves usually opposite, and frequently radical, exstipulate; flowers on simple or umbellate scapes. They are natives chiefly of temperate and cold regions in the northern hemisphere; some occur in elevated situations in warm countries. Lindley notices 29 genera, including 215 species. *Examples*—*Primula*, *Androsace*, *Glaux*, *Trientalis*, *Anagallis*, *Samolus*.

Primulaceæ - Stamens opposite - Free Central Placenta - Segments of Corolla

975. None of the plants of this order occur in the British Pharmacopœias, and few of them have any important medicinal properties. Acridity prevails more or less in the order. They are cultivated as showy garden



annuals and perennials. All the fine forms of *Auricula* are derived from the yellow *Primula Auricula*, a native of the Swiss Alps. The British species of *Primula*, are *P. veris*, the Cowslip, the flowers of which are said to be narcotic; *P. elatior*, the Oxlip; *P. vulgaris*, the Primrose; *P. farinosa*, the Bird's-eye Primrose; and *P. scotica*, the Scottish Primrose. The species of *Cyclamen*, or Sowbread, have large tuberous-like partially subterranean stems, with acrid properties, and their English name is derived from the circumstance of their being eaten as food by wild boars. In them, as well as in the species of *Dodecatheon*, the petals are reflexed. The flowers of the species of *Anagallis* are meteoric (§ 484), and their seed-vessel is a pyxidium (fig. 458). They are said to be acrid, and to cause inflammation of the mucous membrane. *Trientalis europæa* is the only British plant belonging to the Linnæan class Heptandria. It is slightly acrid. In *Samolus Valerandi*, Brook-weed, the calyx is partially adherent to the ovary, and in *Glaux maritima* the corolla is abortive, and the calyx becomes coloured (fig. 543).

976. Order 142.—**Plumbaginaceæ**, the Sea-pink Family. (*Monopet. Hypog.*) Calyx tubular, persistent, sometimes coloured; æstivation plaited. Corolla monopetalous, or pentapetalous, regular. Stamens 5,

Figs. 674-680.—Organs of fructification of *Primula elatior*, illustrating the natural order Primulaceæ.

Fig. 674.—Diagram of the flower, with five imbricate divisions of the calyx, five segments of the corolla, five stamens opposite the corolline segments, and five carpellary leaves, surrounding a free central placenta.

Fig. 675.—Vertical section of the flower. *c*, Inferior calyx. *p*, Monopetalous corolla. *e*, Stamens attached to the corolla. *o*, Superior ovary. *s*, Style with capitate stigma.

Fig. 676.—Ovary cut vertically, to show the free central placenta covered with ovules. *s*, Base of the style.

Fig. 677.—Vertical section of the fruit. *f*, Pericarp. *p*, Placenta with numerous seeds, some of which have been detached.

Fig. 678.—Peltate amphitropal seed separated. *h*, Hilum.

Fig. 679.—Seed cut vertically. *t*, Integuments (spermoderm). *h*, Hilum. *p*, Fleshy perisperm (albumen). *e*, Transverse embryo lying across the hilum.

Fig. 680.—Embryo with cotyledons and radicle.

hypogynous when the corolla is gamopetalous, attached to the base of the petals when they are separate. Ovary free, 1-celled; ovule solitary, pendulous from a funiculus which arises from the bottom of the cell; styles 5, seldom 3 or 4, each bearing a subulate stigma. Fruit a utricle. Seed pendulous; spermoderm simple; embryo straight, in the axis of mealy albumen; radicle superior.—Herbs or undershrubs, with alternate or fasciculate exstipulate leaves, somewhat sheathing at the base; flowers panicle or capitate. They inhabit the sea-shores and salt marshes chiefly in temperate regions. There are two sections of this order:—1. Plumbagineæ, with a synpetalous corolla and connate styles. 2. Staticeæ, with a pentapetalous corolla and distinct styles. Lindley mentions 8 genera, and 160 species. *Examples*—Plumbago, Statice, Armeria.

977. Some of the plants are acrid, others have tonic qualities. *Armeria maritima*, Thrift, or common Sea-pink, grows both on the sea-shores and on the top of the highest Scottish mountains. Its inorganic chemical ingredients are said to vary in these positions (§ 228). In *Armeria* the funiculus curves over the foramen of the ovule in a young state, but slips off at the period of fecundation, and allows an ovular process to proceed from the exostome towards the placenta. In this genus also, the scaly bracts unite so as to form an inverted cylindrical sheath below the heads or shortened panicles of flowers. The root of *Statice caroliniana* is one of the most powerful astringents in the vegetable materia medica. *Plumbago europæa* has been employed for the relief of toothache, and has hence been called Toothwort. Its root is so acrid that it is used in Roumelia for causing issues. Administered internally in small doses it is said to be as effectual an emetic as Ipecacuanha. Some of the species of this genus act as vesicants.

978. Order 143.—**Plantaginaceæ**, the Ribwort Family. (*Monopet. Hypog.*) Calyx 4-parted, persistent, æstivation imbricate. Corolla monopetalous, hypogynous, scarious, persistent, with a 4-parted limb. Stamens 4, inserted into the corolla, and alternate with its segments; filaments long, filiform, folded inwards in the bud; anthers dithecal, versatile. Disk inconspicuous. Ovary free, 2-4-celled; ovules solitary, or in pairs, or 00; style simple, capillary; stigma hispid, simple, rarely bifid. Fruit an operculate capsule enclosed within the persistent corolla. Seeds sessile, peltate, or erect; spermoderm mucilaginous; embryo in the axis of fleshy albumen, transverse; radicle inferior.—Herbs, which are often stemless, with radical ribbed leaves, and spiked hermaphrodite flowers, or solitary unisexual ones. The species are chiefly found in temperate and cool regions. There are 3 genera noticed by Lindley, including 120 species. *Examples*—Plantago, Littorella.

979. The plants of this order are frequently bitter and astringent.

Their mucilaginous seeds are sometimes used as demulcents. *Plantago maritima* is found both on the sea-shores, and on the top of the highest mountains in Scotland. Its inorganic constituents are said to differ in these localities (§ 228). *Plantago major*, Way-bred, is said to follow the footsteps of man in his migrations. Its spikes are used for feeding birds. Sometimes the bractlets become large, and at other times they assume a verticillate appearance.

Subclass IV.—MONOCHLAMYDEÆ.*

980. Corolla wanting; a calyx, or what is called a simple perianth, present; flowers sometimes Achlamydeous. This subclass includes the Apetalous orders of Jussieu, and many of his Diclinous irregular orders. It corresponds to the Apetalæ and Gymnospermæ of Endlicher.


Section A.—ANGIOSPERMÆ.†

a. SPERMOGENEÆ.

Having true seeds with a Dicotyledonous embryo.

981. Monochlamydeous or Achlamydeous plants, having their seeds contained in an ovary, and fertilized by the action of the pollen on a stigma. It is the Apetalous division of Endlicher's Acramphibrya.

982. Order 144.—**Nyctaginaceæ**, the Marvel of Peru Family. (*Apet. Hypog.*) Perianth tubular, coloured, contracted in the middle, be-



* *Mónes*, one, and *χλαμύς*, a cloak or covering.

† *Άγγος*, a vessel, and *στιγμη*, a seed.

Figs. 681-687.—Organs of fructification of *Mirabilis Jalapa*, illustrating the natural order Nyctaginaceæ.


Fig. 681.—Diagram of the flower, with an imbricated involucre, five divisions of the perianth, five alternate stamens, and a unilocular ovary.

Fig. 682.—Lower part of the flower cut vertically. *i*, Involucre. *c*, Base of the perianth, green and swollen around the ovary. *t*, Part of its coloured tube. *e*, Lower part of the filaments. *s*, Part of the style. *o*, Ovary with its erect ovule.

Fig. 683.—Stamens with convex swelling at the base of the filaments.

Fig. 684.—Style and stigma.

coming indurated at the base (fig. 682); limb entire, or toothed and deciduous; æstivation plicate (fig. 681). Stamens definite, hypogynous (fig. 682 *e*); anthers dithecal (fig. 683). Ovary superior, 1-celled; ovule solitary, erect; style 1; stigma 1 (figs. 682, 684). Fruit a caryopsis, enclosed within the enlarged persistent tube of the



perianth (figs. 441, 685, 686). Embryo peripheral (figs. 522, 686 *e*); albumen farinaceous; cotyledons foliaceous; radicle inferior (figs. 686, 687).—Herbs, shrubs, or trees, with opposite, often unequal, sometimes alternate leaves, and involucrate flowers (figs. 681, 682). They are natives principally of warm regions. Lindley notices 14 genera, including 100 species. *Examples*—*Mirabilis* (Nyctago), *Boerhaavia*, *Pisonia*.

983. The plants of the order have in general purgative qualities. *Mirabilis Jalapa* was at one time considered the Jalap-plant, in place of *Exogonium Purga*, one of the Convolvulaceæ. *M. dichotoma* is the Marvel of Peru, which is commonly cultivated in gardens. It is called in the West Indies “four o’clock flower,” on account of opening its blossoms at that hour in the afternoon. Some of the species of *Pisonia* present a peculiar arrangement of the vascular bundles of the woody stem, which resembles in appearance that of *Endogens*.

984. Order 145.—**Amaranthaceæ**, the Amaranth Family. (*Apet. Hypog.*) Perianth 3-5-partite, hypogynous, scarious, persistent, usually with two bractlets at the base. Stamens hypogynous, either 5 and opposite the segments of the perianth, or double that number, distinct, or united, sometimes partly abortive; anthers either dithecal or monothecal. Ovary superior, single, 1-celled; ovules solitary or several, amphitropal, hanging from a free central funiculus; style 1, or 0; stigma simple or compound. Fruit a utricle or a caryopsis, rarely baccate. Seeds lentiform, pendulous; testa crustaceous; embryo peripheral; albumen farinaceous; radicle next the hilum.—Herbs and shrubs, with simple, opposite, or alternate exstipulate leaves; flowers

Fig. 685.—Fruit enclosed by the persistent and indurated base of the perianth.

Fig. 686.—The same cut vertically. *i*, Involucre. *c*, Perianth. *p*, Pericarp. *e*, Curved peripheral embryo, surrounding mealy albumen.

Fig. 687.—Horizontal section of the fruit. *c*, Perianth. *t*, Integument of the seed with the pericarp. *p*, Perisperm. *r*, Radicle. *co*, Cotyledons.

in heads or spikes, usually hermaphrodite. They are natives of tropical and temperate regions. There are 38 known genera, and 282 species. *Examples*—*Amaranthus*, *Achyranthes*, *Celosia*, *Deeringia*, *Gomphrena*.

985. The plants are principally mucilaginous and demulcent. Many of them are known in cultivation, such as *Amaranthus hypochondriacus*, Prince's-feather; *A. caudatus*, Love-lies-bleeding; *Celosia cristata*, Cockscomb; *Gomphrena globosa*, Globe amaranth. *Amaranthus Blitum*, *A. oleraceus*, or Chusan Han-tsi, and other species, are used as potherbs. In the Cockscomb, the flowers form at the apex a peculiar crest of flattened or fasciated peduncles (fig. 230).

986. Order 146.—**Chenopodiaceæ**, the Goosefoot Family. (*Apet. Perigyn.* and *Hypogyn.*) Perianth deeply divided, sometimes tubular at the base, persistent, without bracts; æstivation imbricate. Stamens inserted into the base of the perianth or hypogynous, opposite to its segments, and equal to them in number, or fewer (fig. 544). Ovary single, superior, or sometimes cohering to the tube of the perianth, 1-celled; ovule solitary, attached to the base of the cell; style 2-4-parted; stigmas simple. Fruit membranous, indehiscent, enclosed in the calyx, sometimes fleshy. Seed erect or resupinate; embryo curved around farinaceous albumen, often like a horse-shoe, or spiral or doubled together without albumen; radicle next the hilum.—Herbs or undershrubs, with alternate, sometimes opposite, exstipulate leaves, and hermaphrodite or unisexual flowers. They are found in almost all parts of the world, but do not abound in the tropics. Most of the plants are inconspicuous weeds. There are 67 known genera, and 372 species. *Examples*—*Chenopodium*, *Salicornia*, *Salsola*, *Atriplex*, *Beta*, *Basella*.


987. Many of the plants of this order are used as esculent potherbs, such as *Spinacia oleracea*, Spinage, *Beta vulgaris*, Beet, and var. *campestris*, Field Beet or Mangold Wurzel, *Atriplex hortensis*, Garden Orach, *Chenopodium Bonus Henricus*, English Mercury. The seeds of *Chenopodium Quinoa* are used as food in Peru, under the name of petty rice. The plant grows at a great elevation. Its leaves are used for spinage. They contain much starch and oil, combined with a bitter substance which appears to reside in the integuments. Many of the plants grow in salt marshes, and are called Halophytes (ἅλς, salt, and φυτόν, a plant). They yield a quantity of soda. Among them may be enumerated species of *Salicornia*, *Salsola*, *Halimocnemis*, and *Kochia*. Beet-root yields a quantity of sugar. *Ambrina anthelmintica* yields a volatile oil, which is used in the cure of worms. Some of the *Chenopodiums* have a very fetid odour. The genus *Atriplex* has polygamous flowers, and was placed by Linnæus in his class polygamia. *Salvadora persica*, the true Mustard-tree, σίγηπι or σίναπι of Scripture, has been referred to this order. Lindley considers it the type of a new order in his Echial alliance. The tree grows in Syria,

and it has been found in Ceylon. Its root is acrid. Its succulent fruit has the taste of cresses, and its seeds are very small.

988. Order 147.—**Phytolaccaceæ**, the Phytolacca Family. (*Apet. Perigyn.*) Perianth 4-5-partite. Stamens usually perigynous, indefinite, or equal to the segments of the perianth, and alternate with them. Ovary of 1 or several carpels, distinct or combined; ovule 1 in each carpel, ascending or erect; styles equal to the carpels in number, terminal or lateral; stigmas simple or divided. Fruit fleshy and dry, indehiscent, sometimes samaroid. Seeds solitary, erect or ascending; embryo straight or curved; albumen mealy or 0; radicle next the hilum.—Undershubs or herbs, with alternate, entire leaves, which are often dotted. They are natives both of tropical and warm countries, and are found in America, Asia, and Africa. The order has been divided into two suborders;—1. Phytolaccæ, with ascending seeds, embryo curved round mealy albumen, and exstipulate leaves. 2. Petiveriæ, with an erect seed, exalbuminous straight embryo, and stipulate leaves. There are 12 known genera, including about 70 species. *Examples*—Phytolacca, Rivina, Petiveria.

989. There is frequently much acidity in the plants of this order, and some of them act as irritant emetics and purgatives. The succulent fruit of *Phytolacca decandra*, common Poke, yields a red juice. It has been used as a remedy in cases of chronic syphilitic pains, and it possesses also emetic and purgative qualities. The plant is said to yield much potash. *Petiveria alliacea* is the Guinea-henweed, so called on account of these animals being fond of it.

990. Order 148.—**Polygonaceæ**, the Buckwheat Family. (*Apet. Hypog. and Perigyn.*) Perianth inferior (fig. 688 *c c*), divided, often coloured; æstivation imbricate (fig. 689). Stamens definite, inserted



Figs. 688-690.—Organs of fructification of *Fagopyrum esculentum* (*Polygonum Fagopyrum*), to illustrate the natural order Polygonaceæ.

Fig. 688.—Vertical section of the flower. *c c*, Perianth. *e e*, Outer stamens, which are introrse. *e i*, Inner stamens, which are extrorse. *a*, Glandular appendages. *o*, Ovary with its erect ovule. *s*, Styles and stigmas.

Fig. 689.—Diagram of the flower, showing five divisions of the imbricate perianth, stamens opposite the divisions, with glands and triangular unilocular ovary.

Fig. 690.—Seed cut vertically, showing the embryo with its superior radicle curved at one side of mealy albumen.

into the bottom of the perianth (fig. 688 *ee, ei*); anthers with longitudinal dehiscence. Ovary free (fig. 688 *o*), usually formed by 3 carpels, unilocular; ovule solitary, orthotropal (fig. 418); styles and stigmas equal to the carpels in number (fig. 688 *s*). Fruit a nut, usually triangular, naked or covered by the persistent perianth (fig. 271). Seed erect; albumen farinaceous; embryo antitropal, generally on one side (fig. 690), sometimes in the axis of the albumen; radicle superior (fig. 690).—Herbaceous, rarely shrubby plants, with alternate, stipulate, or exstipulate leaves, and often unisexual flowers. They are found in almost all parts of the world, more especially in the temperate regions of the northern hemisphere. They grow in fields, waste-grounds, ditches, mountains, &c. The order has been divided into two tribes:—1. *Polygonæ*, with loose flowers, embryo usually abaxial (fig. 521), ochreate stipules (fig. 132). 2. *Eriogonæ*, with involucrate flowers, embryo axial, leaves generally exstipulate. Lindley enumerates 29 genera, including 490 species. *Examples*—*Polygonum*, *Rumex*, *Rheum*, *Eriogonum*.

991. The plants of this order have astringent and acid properties; some of them are purgative, and a few are acrid. Their astringency depends on the presence of tannin, and their acidity chiefly on oxalic acid. The root of *Polygonum Bistorta*, Bistort, so called on account of its double twist, contains much tannin, some gallic acid and starch, and is a powerful astringent. The leaves of *P. Hydropiper*, Water-pepper, are acrid and vesicant. *P. tinctorium* yields a blue dye. The fruit of *P. aviculare* is emetic and purgative. The fruit of *Fagopyrum esculentum*, and other species of Buckwheat, is used as food. The plant is cultivated in some northern countries. The leaves of *Rumex Acetosa*, Common Sorrel, and of *R. Acetosella*, Field Sorrel, are acid and astringent. The roots of *Rumex aquaticus*, Water Dock, *R. Hydrolapathum*, Great Water Dock, and of other species, are used as astringents, and alteratives, while those of *R. alpinus*, under the name of Monk's-rhubarb, were formerly employed as purgatives. One of the most important plants of the order is the Rhubarb-plant. The officinal Rhubarb is the root of an undetermined species of *Rheum*. The extent of country from which rhubarb of one kind or another is actually collected, according to Christison, stretches from Ludak, in $77\frac{1}{2}^{\circ}$ east longitude, to the Chinese province of Shen-si, 29° farther east, and from the Sue-chan mountains, in north latitude 26° , nearly to the frontiers of Siberia, 24° northward. The best rhubarb is said to come from the very heart of Thibet, within 95° east longitude, and 35° north latitude, five or six hundred miles north of Assam. The following are the species *Rheum* said to yield rhubarb:—

1. *Rheum Palmatum*, L. This has perhaps the best title to be considered the true rhubarb-plant.
2. *Rheum undulatum*, L., which yields much of the French rhubarb.

3. *Rheum Compactum*, L. Another species yielding French rhubarb, and often cultivated in Britain for its acid petioles.
4. *Rheum Emodi*, Wall. This species yields a kind of Himalayan rhubarb. Its petioles are used for their acid properties.
5. *Rheum rhaponticum*, L. Used in France and Britain in the same way as the third species.
6. *Rheum hybridum*, Murr. Much cultivated in Germany for its root, and in Britain for its stalks.
7. *Rheum Webbianum*, Royle. A Himalayan species.
8. *Rheum spiciforme*, Royle. Another Himalayan species.
9. *Rheum Moorcroftianum*, Royle. Another Himalayan species.
10. *Rheum crassinervium*, Fisch. A Russian species.
11. *Rheum leucorhizum*, Pall. A Siberian and Altai species, said to yield imperial or White Rhubarb.
12. *Rheum Capsicum*, Fisch. A Caspian and Altai species.
13. *Rheum Ribes*, L. An Affghanistan and Persian species.

All these species grow in the cold parts of the world, as on the Altai mountains, in Siberia, Thibet, North of China, and on the Himalayan range. The rhubarb procured from one or more of these species, is known in commerce under the names of Russian or Turkey, Chinese or East Indian, and English rhubarb. Rhubarb contains raphides of oxalate of lime (§ 18), along with tannin, gallic acid, resin, and a peculiar yellow-coloured principle called rhabarberin, which seems to be identical with chrysophanic acid. Raphides form from 35 to 40 per cent. of Turkey rhubarb, and give rise to its grittiness. These crystals are less abundant in the other varieties of rhubarb. Rhubarb is employed medicinally as a cathartic, astringent, and tonic, in the form of powder, pill, extract, tincture, wine, and infusion. *Coccoloba uvifera*, Sea-side-grape, so called from the appearance of its fruit, yields an astringent substance, called Jamaica Kino.

992. Order 149.—**Begoniaceæ**, the Begonia Family. (*Apet. Diclin.*) Flowers unisexual. Perianth coloured, having usually 4 divisions in the male flowers, and 5 or 8 in the female, some being smaller than others; æstivation imbricate. Stamens 00, distinct, or united into a solid column; anthers collected in a head, dithecal, with a thick connective, and longitudinal dehiscence. Ovary adherent to the tube of the perianth, winged, 3-celled, with three placentas meeting in the axis; ovules 00, anatropal; stigmas 3, sessile, 2-lobed, somewhat spirally twisted. Fruit a membranous, triangular, winged capsule, dehiscing below in a loculicidal manner. Seeds 00, minute; testa thin and reticulated; albumen 0; embryo oblong; radicle next the hilum.—Semi-succulent herbaceous plants and undershrubs, with alternate oblique leaves, having large scarious stipules. They are sometimes called Elephant's-ear, from the form of the leaves. They are natives of warm countries, as the East and West Indies, and South America. The stomata on the lower side of the leaves of many of the species of Begonia are arranged in clusters, and exhibit a beautiful

appearance under the microscope. Their leaves and young stems are acid, and have been used for tarts. Their roots are astringent and slightly bitter. *Begonia obliqua* is said to have purgative roots, and it is sometimes called wild rhubarb. There are 3 genera, and 159 known species. *Example*—*Begonia*.

993. Order 150.—**Lauraceæ**, the Laurel Family. (*Apet. Perigyn.*) Perianth with 4 or 6 divisions, which are usually in 2 rows (figs. 691, 692), the limb sometimes obsolete; æstivation imbricate (fig. 692). Stamens perigynous, definite, often twice as many as the divisions of



the perianth, and arranged usually in two rows; those of the inner row (often three) being frequently sterile (staminodia), (fig. 693 *es*), while those of the outer (often six in number) are fertile (figs. 692, 693 *ef*); if the inner stamens are fertile, they are extrorse, while the outer are introrse; filaments of the inner row often with glands at their base (figs. 325, 694 *g*); anthers 2-4-celled, cells opening by

Figs. 691-697.—Organs of fructification of *Cinnamomum zeylanicum* (*Lauris Cinnamomum*), to illustrate the natural order Lauraceæ.

Fig. 691.—Flower entire, with 6-divided perianth.

Fig. 692.—Diagram of the flower, with six imbricate divisions of the perianth; stamens in two rows, the outer six introrse, the inner three extrorse; glandular disk, and unilocular ovary.

Fig. 693.—The flower cut vertically. *c*, The perianth. *ef*, Fertile outer stamens with valvular introrse dehiscence. *es*, Sterile inner stamens with glandular bodies. *o*, Monotheal ovary with pendulous ovule. *s*, Style, and obtuse stigma.

Fig. 694.—Stamens separated. *f*, Filament with two glandular bodies, *gg*, at its base. *a*, Anther with valves.

Fig. 695.—Anther viewed separately, showing its mode of dehiscence from below upwards by four longitudinal valves.

Fig. 696.—Fruit, which is succulent and partially enclosed in the persistent perianth.

Fig. 697.—The fruit deprived of the perianth, and cut vertically. *p*, Pericarp. *t*, Integument of the seed. *e*, Embryo.

longitudinal valves (figs. 325, 695). Ovary superior, unilocular (fig. 693 o); ovule solitary, pendulous (fig. 693); style simple; stigma obtuse (fig. 693 s). Fruit baccate or drupaceous, naked, or covered by the enlarged perianth (fig. 696); peduncle of the fruit sometimes becoming fleshy. Seed solitary, pendulous; albumen 0; embryo inverted (fig. 697 e); cotyledons large, plano-convex, peltate near the base; radicle very short, superior; plumule conspicuous. Trees, with exstipulate, alternate, rarely opposite leaves; sometimes twining, parasitic, and leafless herbs or undershrubs. They are natives chiefly of the tropical regions of Asia and America. Few are found in Africa. The order has been divided into two suborders:—1. Laureæ, true Laurels, trees with leaves. 2. Cassythæ, Dodder-laurels, climbing parasitic plants without leaves. There are 47 known genera, and 459 species. *Examples*—*Laurus*, *Cinnamomum*, *Persea*, *Nectandra*, *Tetranthera*, *Cassynia*.

994. The plants of this order are in general aromatic and fragrant. Many of them yield volatile and fixed oils, others furnish camphor, and others have bitter and tonic barks. Some supply useful timber. *Laurus nobilis* is the Victor's-laurel, the leaves of which were used to crown the conquerors in battle, and in the Olympic games. It is probably the *אֶרְוֶה** of the Bible. It is often called Sweet-bay, and is quite distinct from the common Bay, or Cherry-laurel (*Cerasus Lauro-cerasus*), both as regards structure and properties. It does not yield any hydrocyanic acid. The leaves and fruit are used medicinally as aromatic stimulants. The leaves contain a volatile oil, and the dark-coloured fruit yields, by expression, an odoriferous concrete oil of a green colour, called Oil of Bays. It is the only species found in Europe in a wild state. *Camphora officinarum* (*Laurus Camphora*), a native of China, Japan, and Cochin-China, is the Camphor-tree. Many plants supply a kind of Camphor, but the common camphor of the shops is the produce chiefly of this tree. All parts of the tree supply it, but it is obtained principally from the wood by distillation and subsequent sublimation. It is used in medicine as a sedative antispasmodic, in the form of mixture and tincture. The Borneo camphor has been noticed under the natural order Dipterocarpaceæ (¶ 789). *Sassafras officinale* (*Laurus Sassafras*), is an American tree, the root, wood, and flowers of which have been used in medicine. The root is prescribed in Britain as a warm aromatic stimulant and diaphoretic. It contains a volatile oil. *Cinnamomum zeylanicum* (*Laurus Cinnamomum*), is the true Cinnamon-tree, which is extensively cultivated in Ceylon. The tree attains the height of 30 feet. The bark of the tree constitutes the cinnamon of commerce, the *קַיְמֹן* of the Bible. The young twigs furnish the best cinnamon. The bark yields by distillation an oil,

* *אֶרְוֶה*, virida, according to Gesenius, from *אור*, lux. It is supposed because green is a bright colour.

which is at first of a yellow colour, but soon assumes a reddish hue. The ripe fruit yields a concrete oil called cinnamon-suet. The root yields camphor. Cinnamon is administered as a tonic, stomachic, and carminative. The average importation of cinnamon into London is estimated at 500,000 pounds. The leaves of the cinnamon-tree are more or less acuminate; they have three principal ribs, which come into contact at its base, but do not unite; its young twigs are not downy, and its leaves have the taste of cloves. *Cinnamomum Cassia* or *aromaticum* (*Laurus Cassia*), seems to be the chief source of the Cassia lignea, or Cassia-bark of commerce, the $\pi\pi$ of the Bible. It differs from the true cinnamon in many particulars. Its leaves are oblong-lanceolate; they have three ribs, which coalesce into one at the base; its young twigs are downy, and its leaves have the taste of cinnamon. There is a fine specimen of the plant about twenty feet high in the Edinburgh Botanical Garden, which bears flowers regularly every year, and occasionally has produced fruit. Cassia-bark is imported from Canton through Singapore. It yields a yellow volatile oil called Oil of Cassia. Both the bark and oil are administered as aromatic stimulants. It is probable that Cassia buds, which consist of the flower-bud (perianth and ovary), are the produce of the Cassia-bark tree. They are used chiefly in confectionery, and they have the flavour and pungency of Cassia. Malabar Cassia appears to be the produce of another species of *Cinnamomum*, perhaps *C. eucalyptoides*. *Nectandra Rodicea*, a large tree found in British Guiana, yields a bark known as the Bebeeru-bark. The wood of the tree is imported for ship-building, under the name of Green-heart. The bark was used by Dr. Rodie, who detected the existence of an alkaloid called Bebeerina. Dr. Douglas MacLagan obtained it pure, and found along with it another alkaloid called Sipeerina. Sulphate of Bebeerina is used as an antiperiodic. The cotyledons of the seed contain much starch, and are used for food. *Persea gratissima* (*Laurus Persea*) yields a pear-shaped succulent fruit called Avocado or Alligator-pear, or Subaltern's-butter. It contains a fixed oil. The clove nutmegs of Madagascar are produced by *Agathophyllum aromaticum*, and Brazilian nutmegs are the produce of *Cryptocarya moschata*. *Benzoin odoriferum* is the Spice-wood or Fever-bush of North America.

995. Order 151.—**Myristicaceæ**, the Nutmeg Family. (*Apet. Di-clin.*) Flowers unisexual. Perianth trifid, rarely quadrifid, in the female deciduous; aestivation valvate. Stamens 3-12; filaments combined into a cylinder; anthers united or distinct, dithecal, extrorse, dehiscing longitudinally. Ovary free, composed of one or more carpels, unilocular; ovule solitary, erect, anatropal; style very short; stigma somewhat lobed. Fruit succulent, 1-celled, 2-valved. Seed solitary, usually covered by a lacinated arillus; embryo small, orthotropal, at the base of ruminant albumen; cotyledons foliaceous;

radicle inferior.—Trees with alternate, exstipulate, entire, not dotted leaves. Natives of the tropical regions of Asia and America. There are 5 known genera, and between 30 and 40 species. *Example*—*Myristica*.

996. Acridity and aromatic fragrance are the properties of the order. The most important plant is *Myristica officinalis* (*M. moschata* or *aromatica*), a tree attaining a height of thirty feet, found in the Moluccas, and cultivated in many tropical countries. The fruit is drupaceous, and opens by two valves when ripe, displaying the beautiful reticulated scarlet arillus which constitutes mace. Within this is a thin, hard, dark-brown, glossy shell, covering the kernel, which is the nutmeg of the shops. It is said that a single tree will yield on an average about six pounds of nutmegs. By expression they are made to yield a concrete oil called *Adeps Myristicæ*, or sometimes erroneously oil of mace. A volatile oil is also procured by distillation. Mace is an arillode or additional covering of the seed commencing at the exostome (§ 581). It has a fine crimson hue, and yields a fatty matter and volatile oil, resembling those of the nutmeg. Nutmeg and mace are used medicinally as aromatic stimulants and condiments. In large doses they have a narcotic effect. The fleshy part of the fruit is used as a preserve. The kernels of *Myristica tomentosa* are also used as aromatics, under the name of wild or male nutmegs. The bark of many plants of the order yields an acrid juice, which is sometimes of a crimson colour. A red pigment is furnished by *Pyrrhosa tingens*.

997. Order 152.—**Proteaceæ**, the Protea Family. (*Apet. Perigyn.*) Perianth more or less deeply 4-divided; æstivation valvate. Stamens perigynous, 4 (1 sometimes sterile), opposite the segments of the perianth; anthers dithecal, with longitudinal dehiscence. Ovary single, superior, unilocular; ovules single or in pairs, anatropal or amphitropal; style simple; stigma undivided, discoid. Fruit dehiscent or indehiscent. Seed exalbuminous, sometimes winged; embryo straight; cotyledons 2 or more; radicle inferior, next the hilum.—Shrubs or small trees, with hard, dry, opposite or alternate, exstipulate leaves. They are natives principally of Australia and the Cape of Good Hope. In general they occur in land unfit for cultivation, and seldom attain to a considerable size. The order has been divided into two sections:—1. Nucumentaceæ, with nucumentaceous indehiscent fruit. 2. Folliculares, with follicular dehiscent fruit. Lindley mentions 44 genera, including 650 species. *Examples*—*Protea*, *Persoonia*, *Grevillea*, *Hakea*, *Banksia*, *Dryandra*.

998. The plants of this order have no medicinal properties of importance. They present great diversity of appearance, hence the name of the order, and they are cultivated for their handsome habit, and the peculiarity of their flowers. The clustered cone-like heads of the

flowers of *Banksias* have a remarkable appearance. In *Grevillea* the style is at first bent downwards, and the discoid stigma is enclosed within the upper part of the perianth, where the anthers are placed; but after the pollen has been scattered, the stigma is emancipated, and the style rises upwards. The fruit and seeds of a few plants of the order are eaten, and the wood is used for economical purposes. *Guevina Avellana* yields nuts, which are sold in Chili under the name Avellano. *Protea mellifera* is called Sugar-bush, on account of the honey furnished by its flowers. *Leucadendron argenteum* is the Witteboom of the Cape. The bark of *Protea grandiflora* is used by the Cape settlers in diarrhœa. It is reported by Endlicher that the root of *Banksia marcescens* is emetic. This statement has not been confirmed.

999. Order 153.—**Elæagnaceæ**, the Oleaster Family. (*Apet. Diclin.* and *Perigyn.*) Flowers usually unisexual, rarely hermaphrodite. Male flowers amentaceous, with 2-4 leaves forming the perianth; stamens 3, 4, or 8; anthers nearly sessile, dithecal, introrse, and dehiscing longitudinally. In the female and hermaphrodite flowers, perianth tubular, persistent, with an entire or 2-4-toothed limb. Disk fleshy. Ovary superior, 1-celled; ovule solitary, ascending, on a short funiculus, anatropal; style short; stigma simple, subulate, glandular. Fruit a crustaceous achæmium, enclosed within the enlarged succulent perianth. Seed ascending; embryo straight, surrounded by thin fleshy albumen; cotyledons fleshy; radicle inferior.—Trees or shrubs, with alternate or opposite, entire, exstipulate leaves, which are often covered with scurfy scales (fig. 83). They are found in all parts of the northern hemisphere. They have no marked medicinal properties. The fruit of some is eaten. *Hippophaë rhamnoides*, Sea Buckthorn, is furnished with sharp spines, and forms a good hedge near the sea. Its fruit is eaten, although it is said by some to have narcotic qualities. The plant yields a yellow dye. The fruit of *Elæagnus parvifolia* is eaten. Its flowers are highly fragrant, and abound in honey, which is esteemed as a remedy for malignant fevers in some parts of Europe. There are 4 known genera, and 30 species. *Examples*—*Elæagnus*, *Hippophaë*.

1000. Order 154.—**Penæaceæ**, the Sarcocol Family. (*Apet. Perigyn.*) Perianth coloured, salver-shaped, with a 4-lobed limb, and with two or more bracts at its base, persistent. Stamens perigynous, either 4 or 8, alternate with the lobes of the perianth; anthers dithecal, introrse. Ovary superior, 4-celled; ovules usually in pairs, collateral, anatropal, ascending or suspended; style simple; stigmas 4. Fruit a 4-celled, 4-valved capsule. Seed erect or pendulous; testa brittle; hilum with a fungus-like aril; nucleus a fleshy mass, without distinction of albumen or embryo.—Shrubs, with opposite, entire, exstipulate leaves. They are found at the Cape of Good Hope. They have no known

properties of importance. The gum-resin called Sarcocol is said to be produced on the perianth of *Penæa Sarcocolla*, and other species. There are two sections of this order:—1. *Penææ*, æstivation valvate, stamens 4, connective fleshy, ovules ascending. 2. *Geissolomeæ*, æstivation imbricate, stamens 8, connective not fleshy, ovules suspended. There are 3 known genera, and 21 species. *Examples*—*Penæa*, *Geissoloma*.

1001. Order 155.—**Thymelæaceæ**, the *Daphne* Family. (*Apet. Perigyn.*) Perianth tubular, coloured, 4- rarely 5-cleft, inferior; occasionally with scales in its orifice; æstivation imbricate. Stamens perigynous, definite, often 8, sometimes 4 or 2 and then opposite the segments of the perianth; anthers dithecal, with longitudinal dehiscence. Ovary free, 1-celled; ovule suspended, anatropal (fig. 426); style 1; stigma undivided. Fruit either nut-like or drupaceous. Seed solitary, pendulous; albumen 0, or thin and fleshy; embryo straight; cotyledons plano-convex, or somewhat lobed and shrivelled; radicle superior.—Shrubby, rarely herbaceous plants, with alternate, or opposite, entire, exstipulate leaves. Natives of various parts of the world, both in warm and temperate regions. There are two sections of the order:—1. *Daphnææ*, with hermaphrodite or rarely unisexual flowers, and plano-convex cotyledons. 2. *Hernandiææ*, with polygamous flowers, and lobed and shrivelled cotyledons. Lindley enumerates 38 genera, including 300 species. *Examples*—*Daphne* (*Thymelæa*), *Passerina*, *Pimelea*, *Gnidia*, *Lagetta*, *Exocarpus*, *Hernandia*, *Inocarpus*.

1002. The bark of many of the plants is acrid and irritant, the fruit is often narcotic. The bark of the root, as well as that of the branches of *Daphne Mezereum*, *Mezereon*, is used in decoction as a diaphoretic in cutaneous and syphilitic affections. In large doses it acts as an irritant poison, causing hypercatharsis; and, when applied externally, it acts as a vesicant. It contains a neutral crystalline principle called Daphnein. The succulent fruit is also poisonous. The barks of *Daphne Gnidium*, *D. alpina*, *D. Cneorum*, *D. pontica*, and *D. Laureola*, Spurge-laurel, have similar properties. The berries of *Daphne Laureola* are poisonous to all animals except birds. The fruit of *Dirca palustris*, Leather-wood, is said to be narcotic. The bark of many of the plants is made into ropes and paper (fig. 101). The inner bark of *Lagetta lintearia* (*Daphne Lagetta*,) when cut into thin pieces after maceration, assumes a beautiful net-like appearance, whence' it has received the name of Lace-bark. The bark, young leaves, and seed of *Hernandia*, are slightly purgative. The seeds of *Inocarpus edulis* have the taste of chestnuts, and are eaten when roasted.

1003. Order 156.—**Aquilariaceæ**, the *Aquilaria* Family. (*Apet. Perigyn.*) Perianth coriaceous, imbricate or tubular, limb 4-5-lobed; æstivation imbricate. Stamens usually 10 fertile, alternating with 10

sterile, in the form of petaloid scales, sometimes 8 or 5; filaments inserted into the orifice of the perianth, often united; anthers dithecal, with longitudinal dehiscence. Ovary free, ovate compressed, 2-celled; ovules 2, suspended, anatropal; stigma usually sessile, large and simple. Fruit a pyriform, sessile, or stipitate 2-valved capsule, or drupaceous and indehiscent. Seeds 2, one on each placenta, pendulous; albumen 0; cotyledons fleshy, hemispherical; radicle straight, superior.—Trees, with alternate or opposite, entire, stalked, and exstipulate leaves. They are natives of the tropical regions of Asia. They have no known medical properties. *Aquilaria ovata* and *Agallochum* furnish a fragrant wood called Eagle-wood, or Aloes-wood. It is probably the דגל, the trees of Aloes or Lign-Aloes, of the Bible, yielding an aromatic perfume. It has been considered a cordial by some Asiatic nations, and has been prescribed in Europe in gout and rheumatism. There are 6 genera noticed, including 10 species. *Examples*—*Aquilaria*, *Gyrinopsis*.

1004. Order 157.—**Chailletiacæ**, the Chailletia Family. (*Apet. Perigyn.*) Perianth 5-parted, with an incurved valvate æstivation. Stamens inserted into the base of the perianth, 5 inner fertile opposite the segments of the perianth, 5 outer sterile, petaloid, usually with glands at their base; anthers ovate, versatile, dithecal. Ovary free, 2-3-celled; ovules twin, pendulous; styles 2-3, distinct or combined; stigmas capitate or obscurely 2-lobed. Fruit dry, 1-2- or 3-celled. Seeds solitary, pendulous, exalbuminous; embryo thick; cotyledons fleshy; radicle superior.—Trees or shrubs, with alternate, stipulate leaves, and axillary peduncles, often cohering to the petiole. They are natives of the warm parts of Africa and South America. The fruit of *Chailletia toxicaria* is said to be poisonous; it is called Rats-bone in Sierra Leone. There are 4 genera, and 10 species known. *Examples*—*Chailletia*, *Tapura*.

1005. Order 158.—**Samydacæ**, the Samyda Family. (*Apet. Perigyn.*) Perianth 4-5-divided, usually coloured inside; æstivation somewhat imbricate. Stamens inserted into the tube of the perianth, 2, 3, or 4 times as many as its divisions, either all fertile, or the alternate ones sterile, shorter and fringed; filaments monadelphous at the base; anthers erect, ovate, 2-celled. Ovary free, 1-celled; ovules 00, attached to parietal placentas, semi-anatropal; style 1, filiform; stigma capitate or slightly lobed. Fruit a coriaceous, unilocular, 3-5-valved capsule, partially dehiscent. Seeds 00, fixed irregularly on the pulpy inner surface of the valves, with a fleshy arillus, and a hollowed hilum; embryo large, in the midst of oily or fleshy albumen; cotyledons ovate, foliaceous; radicle pointing to the extremity remote from the hilum.—Trees or shrubs, with alternate, simple, stipulate leaves, usually having pellucid, round, or linear markings. Natives of tropical regions, chiefly in America. Some of the species of *Casearia* are

bitter and astringent. A decoction of the leaves of *Casearia lingua*, called by the Brazilians *Cha de Frade* and *Lingua de Fiu*, is also used internally in inflammatory disorders and malignant fevers. There are 5 known genera, and 80 species. *Examples*—*Samyda*, *Casearia*.

1006. Order 159.—**Homaliaceæ**, the *Homalia* Family. (*Apet. Perigyn.*) Perianth funnel-shaped, with 5 to 15 divisions, and having usually alternating petaloid segments, and glands or scales in front of the outer divisions. Stamens perigynous, either single or in parcels of 3 or 6, alternating with the outer divisions of the perianth; anthers dithecal, with longitudinal dehiscence. Ovary partly adherent to the tube of the perianth, 1-celled; ovules numerous, anatropal, pendulous, attached to 2, 3, or 5 parietal placentas; styles 3-5, simple, filiform, or subulate. Fruit either baccate or capsular. Seeds small, ovate; embryo in the axis of fleshy albumen; cotyledons leafy; radicle superior.—Trees or shrubs, with alternate leaves, having deciduous stipules. Many look upon the petaloid division of the perianth as true petals. Lindley puts this order in his Cactal alliance, and considers it as allied to Loasaceæ. It contains tropical plants, which do not possess any important properties. Lindley mentions 8 genera, including 30 species. *Examples*—*Homalium*, *Nisa*.

1007. Order 160.—**Santalaceæ**, the Sandal-wood Family. (*Apet. Epigyn.*) Perianth superior, 4-5-cleft; æstivation valvate. Stamens 4-5, opposite the segments of the perianth, and inserted into their bases. Ovary coherent, 1-celled; ovules 1-4, pendulous from the apex of a central placenta; style 1; stigma often lobed. Fruit nut-like or drupaceous. Seed solitary; embryo minute, in the axis of fleshy albumen; radicle superior.—Trees, shrubs, or herbs, with alternate or nearly opposite exstipulate leaves. Found in various parts of the world, as Europe, Asia, America, and New Holland. Lindley gives 18 genera, including 110 species. *Examples*—*Santalum*, *Osyris*, *Thesium*.

1008. Some are astringent, others yield fragrant wood. *Santalum album*, and other Indian and Polynesian species, yield Sandal-wood, which is used both medicinally and as a perfume. The seeds of some of the plants of the order are eaten. The species of *Thesium* seem to be root-parasites. The large seeds of *Pyrolaria oleifera*, Buffalo-tree, or Oil-nut, yield a fixed oil.

1009. Order 161.—**Aristolochiaceæ**, the Birthwort Family. (*Apet. Epigyn.*) Perianth adherent, tubular, 3-cleft (fig. 699), regular or sometimes very irregular (fig. 698); æstivation valvate or induplicate. Stamens 6-12, epigynous, distinct or gynandrous (fig. 701). Ovary inferior, 3-6-celled (figs. 700, 702); ovules 00 (fig. 700), anatropal, horizontal; style simple, short; stigmas radiating, 3-6 (fig. 701 s). Fruit dry or succulent, 3-6-celled (fig. 703). Seeds (fig. 704) numerous; embryo very minute, at the base of fleshy albumen (fig. 705);

cotyledons inconspicuous; radicle next the hilum (fig. 706).—Herbs or shrubs, often climbing, with alternate, simple, often stipulate leaves,



and solitary axillary flowers. Found in abundance in the warm regions of South America, and growing also in the temperate and cold regions of Europe, Asia, and America. There are 8 known genera, and 130 species. *Examples*—*Asarum*, *Aristolochia*.

1010. The plants of the order are generally bitter, tonic, and stimulant. Some are acrid, and act as emetics. The leaves of *Asarum europeum* are used as an acrid emetic under the name of *Asarabacca*.

Figs. 698-706.—Organs of fructification of *Aristolochia Clematitis*, to illustrate the natural order Aristolochiaceæ.

Fig. 698.—Flower entire, consisting of an inferior ovary, and a superior, irregular, funnel-shaped perianth. *o*, Part of the perianth adherent to the ovary. *l*, Part of the tube of the perianth, with a swollen portion at the base, enclosing the anthers and stigma. *l*, Limb of the perianth prolonged laterally in a tongue-like form.

Fig. 699.—Diagram of the flower, showing three divisions of the perianth, six anthers, and six cells of the ovary.

Fig. 700.—Lower part of the flower cut vertically. *o*, Ovary with numerous ovules. *s*, Radiating stigma. *a*, Anthers. *c*, Swollen part of the tube of the perianth.

Fig. 701.—*s*, Stigma with the anthers adhering to the column in pairs. *o*, Summit of the ovary. *c*, Swollen part of the tube of the perianth.

Fig. 702.—Horizontal section of the six-celled ovary.

Fig. 703.—Ripe fruit. Fig. 704.—Angular seed.

Fig. 705.—Seed cut vertically. *l*, Integument thickened near the chalaza. *p*, Fleshy perisperm. *e*, Minute embryo.

Fig. 706.—Embryo separated, with cotyledons and radicle.

The roots appear to have greater activity than the leaves. The powdered root and leaves enter into the composition of cephalic snuffs, which cause sneezing by their irritation, and are used in cases of headache and ophthalmia. An active crystalline substance, called Asarin, exists in the plant. *Asarum canadense*, Wild Ginger, or Canada Snake-root, is used as a spice in Canada. The shrubby species of *Aristolochia* have a peculiar arrangement of vascular bundles in their wood. There are no concentric zones, but a number of separable wedges (§ 90). The name of Birthwort, given to *Aristolochias*, depends on their supposed action on the uterus. Some of them are used as emmenagogues. The root of *Aristolochia serpentaria*, Virginian Snake-root, is a stimulant tonic. The plant is a native of the United States. It was formerly used as an antidote to snake-poison. It is now employed occasionally as a tonic diaphoretic. *Aristolochia longa*, *rotunda*, and *Clematitis*, were celebrated in ancient times as uterine remedies. The roots of many of the species have a strong aromatic taste. Those of *Aristolochia anguicida* are said to stupify snakes.

1011. Order 162.—**Nepenthaceæ**, the Pitcher-plant Family. (*Apet. Diclin.*) Flowers diœcious. Perianth 4-parted, inferior; æstivation imbricated. Male flowers: stamens united in a solid central column; anthers about 16, forming a spherical head, extrorse, and with longitudinal dehiscence. Female flowers: ovary free, four-cornered, 4-celled; ovules 00; stigma sessile. Fruit a 4-celled, 4-valved capsule, with loculicidal dehiscence. Seeds 00, ascending, very minute, fusiform, with a loose testa; nucleus less than the seed, suspended by the chalaza; embryo in the midst of fleshy albumen; cotyledons plano-convex; radicle pointing to the hilum.—Herbs, or half-shrubby plants, with alternate leaves, slightly sheathing at the base, having a foliaceous petiole, which forms an ascidium at its extremity, and the lamina in the form of a lid (fig. 184). Natives of swampy ground in the East Indies and China. They have no known properties. The pitchers have been found to contain a solution of binoxalate of potash. Some chemists have detected muriate of soda, malic, and other acids in them. Spiral vessels abound in all parts of the pitcher plants; and the woody bundles are without concentric zones. Lindley gives 1 genus, and 6 species. *Example*—*Nepenthes*.

1012. Order 163.—**Datisceæ**, the Datisca Family. (*Apet. Diclin.*) Flowers unisexual. Male flowers: perianth 3-4-divided. Stamens 3-7; anthers linear, membranous, dithecal, with longitudinal dehiscence. Female flowers: perianth adherent, 3-4-toothed. Ovary inferior, unilocular; ovules 00, anatropal, attached to 3 or 4 parietal placentas; styles as many as the placentas. Fruit a 1-celled capsule, opening at the apex. Seeds 00, strophiolate, with a reticulated spermoderm; albumen 0; embryo straight; cotyledons very short; radicle pointing to the hilum.—Herbaceous branched plants or trees, with alternate, ex-


stipulate leaves. They are scattered over North America, various parts of Asia, and the south-eastern part of Europe. Some of the plants are said to be bitter, and others of them have purgative qualities. It has been stated that female plants of *Datisca cannabina* have produced seed without the application of pollen. Facts are still wanting to prove this. Lindley mentions 3 genera, and 4 species. *Examples*—*Datisca*, *Tetrameles*.

1013. Order 164.—**Empetraceæ**, the Crowberry Family. (*Apet. Diclin.*) Flowers unisexual. Perianth bud-like, consisting of persistent imbricated scales, in 2 or 4 alternating rows, the inner row often petaloid. Male flowers: stamens 2-3, equal in number to the scales in each row, and alternating with the innermost, hypogynous; anthers roundish, dithecal, with longitudinal dehiscence. Female flowers: ovary free, seated on a fleshy disk, 3-6- or 9-celled; ovules solitary, anatropal, ascending; style 1; stigma with as many radii as there are ovarian cells. Fruit a nuculanum, seated within the persistent perianth. Seeds solitary in each nucule, ascending; embryo in the axis of fleshy albumen; radicle inferior.—Heath-like shrubs, with alternate or somewhat verticillate, evergreen exstipulate leaves. They inhabit chiefly Europe and North America. The fruit of some is slightly acid. *Empetrum nigrum*, the black Crowberry, is common on the mountainous and northern parts of Europe. The fruit is watery, and very slightly acid and astringent. Lindley notices 4 genera, and 4 species. *Examples*—*Empetrum*, *Corema*.

1014. Order 165.—**Euphorbiaceæ**, the Spurge Family. (*Diclin.*) Flowers unisexual, sometimes enclosed within an involucre (fig. 707). Perianth lobed, inferior (figs. 314 c, 317 c), with various glandular or petaloid, scaly, internal appendages (figs. 314 p a, 405); sometimes the flowers are naked (fig. 708). Male flowers (fig. 707 fm fm); stamens definite or 00, distinct (fig. 708) or monadelphous (fig. 314, 1), or polyadelphous (fig. 317); anthers bilocular, sometimes with porous dehiscence (fig. 323). Female flowers (figs. 405, 707 ff): ovary free, sessile or stalked, 1-2-3- or many-celled (fig. 709); ovules solitary or twin, suspended; styles equal in number to the cells (figs. 314, 2, 709 s), distinct or combined, sometimes 0; stigmas several, or 1 with several lobes. Fruit usually tricocous (figs. 447, 453), with the cocci separating in an elastic manner, and opening by 2 valves (figs. 710, 711), or indehiscent and fleshy. Seed solitary (fig. 712) or in pairs, suspended, often arillate (fig. 453 g g); embryo enclosed in fleshy albumen (fig. 483); cotyledons flat (fig. 509); radicle superior (fig. 713).—Trees, shrubs, and herbs, often abounding in acrid milk, with opposite or alternate, often stipulate leaves, sometimes none. Some look on this order as apetalous, with a tendency to develop a corolla, while others consider it polypetalous, with a tendency to have the corolla suppressed. In European plants of the order there are usually no

Euphorbiaceæ - Unisexual Flowers - Involucres - Sticky Juice - Fruit

petals present, but in those of tropical countries the corolla is frequently well marked. In the Euphorbias of Britain, there is an evident involucre, surrounding a number of achlamydeous male and



female flowers, which by Linnæus were looked upon as merely stamens and pistils, and hence the plants were put by him in Dodecandria in place of Monœcia (see ¶ 401). The flowers in Euphorbiaceæ vary much in the number of their parts, as may be seen in figs. 545-550. Sometimes the general peduncle or rachis becomes flattened and leaf-like (fig. 229). The inflorescence is occasionally amentaceous, as in the division *Scapaceæ*, which is separated, as a distinct but not fully defined order, by Lindley. The plants of the order abound in warm regions, especially in Equinoctial America, where they occur as trees or bushes, or lactescent herbs, and often present the appearance of Cactuses, from which their milky juice at once distinguishes them. They are also found in North America and in Europe. In Britain there are 18 species. There are about 192 known genera, and upwards of 2,500 species. *Examples*—Euphorbia, Hippomane, Hura, Acalypha, Croton, Jatropha, Ricinus, Phyllanthus, Buxus.

1015. The plants of this order are acrid and poisonous, in some instances furnished with stinging hairs. These properties reside especially in their milky juices, which are contained in laticiferous

Figs. 707-713.—Organs of fructification of *Euphorbia palustris*, to illustrate the natural order Euphorbiaceæ.

Fig. 707.—Inflorescence, with the involucre, *ii*, opened and spread out, to show the position of the male and female flowers which it encloses. *gg*, Glands (glandular lobes) alternating with the divisions of the involucre. *bb*, Membranous laminae, or bracts, at the base of the flowers. *fm, fm*, Achlamydeous male flowers, consisting of a single stamen, supported on a pedicel, to which it is attached by an articulation. *ff*, Achlamydeous female flower in the centre; the ovary and styles supported on a long pedicel.

Fig. 708.—Achlamydeous male flower separated. *b*, Bract. *p*, Pedicel. *f*, Filament articulated with the pedicel. *a*, Anther.

Fig. 709.—Female flower. *p*, Summit of the pedicel which supports it. *c*, A flattened portion of the pedicel, which some call a perianth. *o*, Tricocous ovary. *s*, Styles and stigmas.

Fig. 710.—One of the coeci (carpels), *c*, separated, and seen on its inner surface. *g*, The seed seen across the opening by which the nourishing vessels enter.

Fig. 711.—A coccus separated, after dehiscence and expulsion of the seed.

Fig. 712.—Seed separated.

Fig. 713.—Seed cut vertically. *t*, Integument (spermoderm). *p*, Perisperm (fleshy albumen). *e*, Embryo with flat cotyledons, and a superior radicle.

vessels (fig. 66), in which the movements of Cyclosis were observed by Schultz (§ 262). In many cases the elaborated sap contains caoutchouc and resin. The acrid properties of the order are also found in the seeds, many of which yield oils, both of a bland and of an irritating nature.

1016. The milky juice of many species of *Euphorbia* is caustic, and has been used for destroying warts, and causing vesication. At other times the juice has been used for its purgative and emetic properties. The root of *Euphorbia Ipecacuanha* has been employed as a substitute for Ipecacuanha. The resinous substance called Euphorbium is procured, in all probability, from two African species, *Euphorbia officinarum* and *antiquorum*, as well as from *E. canariensis*, a native of the Canary islands. The resin is a powerful irritant, and has been used as a vesicant. It causes great irritation of the mucous membrane when applied to the nostrils and eyes, and it acts as a cathartic when taken internally. Many species of *Euphorbia* yield resins of a similar nature. The juice of *Hippomane Mancinella*, Manchineel, is very acrid and poisonous. When applied to the skin, it excites violent inflammation, followed by ulceration. The juice of *Hura crepitans*, Sand-box-tree, or Monkey's dinner bell, is also very acrid. The fruit of this tree is composed of numerous 1-seeded cocci, which, when dry, separate from each other with great force. *Mercurialis perennis*, and *annua*, produce vomiting and purging.


1017. Many important medicinal oils are furnished by the plants of this order. Castor-oil is expressed from the seeds of *Ricinus communis* (*Palma Christi*), a plant with peltate-palmate leaves (fig. 146), which is found native in Greece, Africa, and the East Indies, and is cultivated in the West Indies, as well as in North and South America. In the temperate and more northern parts of Europe, the plant is a herbaceous annual of from three to eight feet high; in the more southern parts it becomes shrubby, and even attains a height of twenty feet; while in India it is often a tree thirty or forty feet high. The best oil is got by expression from the seeds without heat, and is called cold-drawn Castor-oil. It is entirely soluble in alcohol, and, by the action of hyponitrous acid, it is converted into a solid yellow substance called Palmin. The oil acts as a mild laxative. Besides this comparatively bland oil, there exists in the seed a powerfully cathartic constituent, which remains behind when the oil is expressed, and which is destroyed or evaporated under the process of ebullition. Croton-oil is obtained by expression from the seeds of *Croton Tiglium*, an Indian and Asiatic shrub. It acts as an irritant purgative in the dose of one drop. In large doses it is a dangerous poison. When applied externally it produces pustules. Other species of *Croton*, as *C. Parana* and *Roxburghii*, yield a purgative oil. The oil procured from the seeds of *Euphorbia Lathyris*, Caper-spurge, has cathartic

properties, and so has that procured from the seeds of *Jatropha Curcas* (*Curcas purgans*), Physic or purging-nut, *Jatropha multifida*, and *Hura crepitans*. The fatty matter obtained from the seeds of *Stilingia sebifera*, the Tallow-tree of China, is used for making candles; the plant also yields a bland oil. The roots of *Euphorbia pilosa* and *palustris* are used as purgatives, and are said to have been useful in hydrophobia.

1018. Cascarilla is the bark of *Croton Eleuteria*, and of other species of *Croton*. It acts as a tonic and stimulant. When burned it gives out a musky odour, and is often used in pastilles. The bark of another species of *Croton* (*C. suberosus*), is known by the name of Copalchi bark, and used as a tonic. The bark of *Buxus sempervirens*, Box-tree, is said to be alterative, and its leaves have bitter and purgative qualities. Its wood is much used for wood-engraving. It is conjectured that the hard wood called African Teak, is the produce of a Euphorbiaceous tree. In the root of *Janipha Manihot* (*Manihot utilissima*), a shrub about six feet high, extensively cultivated in tropical countries, there is much starchy matter deposited, usually along with a poisonous narcotic substance, which is said to be hydrocyanic acid. The latter can be removed by washing, or it can be driven off by roasting, and then the starch is used in the form of Cassava bread. There are two varieties of the Cassava or Manioc plant; one (called sometimes *Janipha Læslingii*) having a spindle-shaped root, brown externally, about six ounces in weight, which contains amylaceous matter, without any bitterness, and is used as food under the name of Sweet Cassava; another, called Bitter Cassava, having a knotty root, black externally, and sometimes 30 lbs. in weight, which is bitter and poisonous, and requires to be rasped and washed thoroughly before the amylaceous matter can be used. From the starch of the bitter Cassava, Tapioca is prepared by elutriation and granulating on hot plates. Manihot starch is sometimes imported into Europe under the name of Brazilian Arrow-root. The milky sap of *Euphorbia phosphorea* is said to emit a peculiar phosphorescent light. That of *Siphonia elastica* contains much caoutchouc, and supplies the bottle India-rubber. *Aleurites laccifera* furnishes gum-lac in Ceylon. *Crotophora tinctoria* supplies a purple dye called Turnsole, which becomes blue on the addition of ammonia. The seeds of a few species of *Aleurites*, *Anda*, and *Omphalea* are edible. It is said that, in some instances, the elaborated sap has poisonous properties, while the ascending sap is innocuous (§ 263). A species of *Cœlebo-gyne* is stated to have produced perfect seeds without the application of pollen (§ 501).

1019. Order 166.—*Urticaceæ*, the Nettle Family. (*Apet. Diclin.*) Flowers unisexual (fig. 716), hermaphrodite or polygamous, scattered, or collected into catkins or heads. Perianth usually divided (fig. 716). Stamens definite, inserted into the perianth; filaments some-

times curved in æstivation (fig. 715). Ovary free (figs. 717, 718), rarely coherent, 1-2-celled; ovule solitary, erect (fig. 718), or suspended; stigmas 1 or 2, simple or bifid (fig. 718). Fruit an inde-



hiscent nut, surrounded by the persistent pericarp, or a samara, or a syconus, or a sorosis (figs. 245, 246, 475). Seed solitary, erect, suspended, or pendulous, albuminous, or exalbuminous; embryo straight, or curved, or spiral; radicle superior (figs. 719, 720).—Herbs, shrubs, or trees, with alternate, stipulate leaves, which are usually hispid or scabrous. This order has been divided into the following suborders:—

Figs. 714-720.—Organs of fructification of *Urtica urens*, to illustrate the natural order Urticaceæ, section Urticæ.

Fig. 714.—Bud of the male flower, viewed from above.

Fig. 715.—Stamen taken from the bud of the male flower, with the elastic incurved filament, and the anther bent down before dehiscence.

Fig. 716.—Male flower expanded. *c*, Perianth with four divisions. *eeee*, Four hypogynous stamens, thrown back by the elasticity of the filaments, with the anthers burst. *pr*, Abortive rudiment of the central pistil.

Fig. 717.—Female flower. *c*, Perianth with four unequal segments, the two exterior ones being very small. *o*, Unilocular ovary. *s*, Sessile stigma.

Fig. 718.—Pistil cut vertically, to show the direction of the erect ovule, *o*. *p*, Parietes of the ovary. *s*, Stigma.

Fig. 719.—Seed cut vertically, parallel to the cotyledons. *t*, Integument (spermoderm). *h*, Hilum. *p*, Perisperm. *e*, Embryo straight, with the radicle superior.

Fig. 720.—Seed cut perpendicularly to the cotyledons. *t*, Integument. *h*, Hilum. *p*, Perisperm. *e*, Embryo cut.

1. Urticeæ, True Nettles (figs. 714-720): rough-leaved plants, often with stinging hairs, filaments elastic, and curved in æstivation (fig. 715); fruit an indehiscent nut; seed erect, albuminous; embryo straight; juice watery. They are widely scattered over the globe, and many of them follow the footsteps of man in his migrations.
2. Cannabineæ, Hemp tribe: scabrous plants with erect filaments; fruit indehiscent; seed suspended, exalbuminous; embryo hooked or spiral; juice watery. They occur chiefly in temperate regions.
3. Ulmaceæ, Elm tribe: rough-leaved trees or shrubs, filaments erect in æstivation; fruit 1-2-celled, samaroid or drupaceous; seed pendulous, usually exalbuminous; embryo straight or curved; juice watery. Natives of the northern and mountainous parts of Europe, Asia, and America.
4. Moreæ, Mulberry tribe: usually rough-leaved trees or shrubs; filaments erect in æstivation; fruit a sorosis or syconus (figs. 246, 475); seed solitary, pendulous, albuminous; embryo hooked; juice milky. Natives of temperate and tropical regions.
5. Artocarpeæ, Bread-fruit tribe: trees or shrubs, with leaves often rough; filaments generally erect in æstivation; fruit often a sorosis; seed erect or pendulous, albuminous; embryo straight; juice milky. Natives of tropical regions.

There are between 60 and 70 known genera, and about 600 species. *Examples*—*Urtica*; *Cannabis*, *Humulus*; *Ulmus*, *Celtis*; *Morus*, *Ficus*, *Dorstenia*; *Artocarpus*, *Antiaris*.

1020. The properties of the order are various. Many yield valuable fibres, others supply important edible fruits, others furnish caoutchouc, and others form important forest trees. Various species of *Urtica*, Nettle, such as *U. dioica*, *urens*, *pilulifera*, *stimulans*, *crenulata*, and *urentissima*, have stinging hairs (fig. 86). The young shoots of the common nettle are sometimes used like spinach or greens. *Urtica cannabina* and *tenacissima* furnish fibres fit for cordage. *Bahmeria nivea* supplies fibre for the Chinese grass-cloth, and *Bahmeria Puya* gives the Pooah or Puya fibre of Nepal and Sikkim. In Nettles and Pellitories, the elastic filaments turn the anthers back with elasticity, and cause the scattering of the pollen (§ 497). Specimens of Tree-nettle, *Urtica gigantea*, were measured by Backhouse in Australia, and found to be 18, 20, and 21 feet in circumference. Their sting is very severe, causing violent inflammation. *Cannabis sativa* is the source of the valuable fibre called Hemp. A variety called *Cannabis indica* is used in India for producing intoxication. It is also employed medicinally in the form of extract, as an antispasmodic and anodyne, in cases of tetanus and neuralgia. The properties of the hemp plant appear to be much modified by climate. The Indian variety has a marked resinous varnish, called Churrus, on its leaves. What is called Bhang in India, consists of the dried larger leaves and fruit, while Gunjah is the whole plant dried after flowering, and the Haschisch of the Arabs is composed of the tops and tender parts of the plant dried. The strobili of the female plants of *Humulus Lupulus* constitute hops, the bitterness of which resides in the resinous glandular scales surrounding the fruit, and to which the name of Lupulinic

glands, or Lupulin, has been applied. The latter name is also given to the bitter principle of the hops. Hops are employed as a tonic and narcotic, in the form of extract, infusion, and tincture. Their tonic properties depend on their bitterness. A pillow stuffed with hops is a popular means of procuring sleep. The twigs of hops have been used to adulterate Sarsaparilla. Several species of Elm are cultivated for timber. *Ulmus campestris*, English or small-leaved Elm, rarely produces fruit in this country. It often attains a height of 70 to 90 feet, with a diameter of 4 to 5 feet. Its wood is compact and durable under water, and it has been used for sleepers on railways, and for wooden pavements. Its inner bark is bitter, mucilaginous, and astringent. *Ulmus montana*, the Mountain, Wych, or Scotch Elm, produces fruit freely in this country, but its wood is inferior to that of the English Elm. *Celtis*, the Nettle-tree, or Sugar-berry, has a sweet drupaceous fruit.

1021. The common Fig is the fruit of *Ficus Carica*. It consists of a succulent hollow receptacle, enclosing numerous single-seeded carpels (fig. 246), and is called a syconus (§ 558). The fruit is demulcent and laxative, and is used for cataplasms. Many other species of *Ficus* yield edible fruits. The plants belonging to the Fig tribe are generally remarkable for the adventitious roots which they send out from the stems. One of the most celebrated in this respect is *Ficus indica*, the Banyan (§ 121, 635). Many of the species can live suspended in the air for a long time. A specimen of *Ficus australis* has grown in this way in the Botanic Garden of Edinburgh for upwards of twenty years (§ 217). *Ficus* (*Urostigma*) *religiosa* is the Pippul-tree, or Sacred Fig of India. *F. elastica* is an Indian tree which supplies a large quantity of caoutchouc; so also do *Ficus Radula*, *elliptica*, and *prinoides*. Peculiar clusters of raphides are found in the cellular tissue of some of them (fig. 39). The milky juice is not in all instances bland and innocuous; it occasionally has acrid qualities. *Ficus Sycomorus* (*Sycomorus antiquorum*), is probably the Sycamore of the Bible, the שִׁטָּה אֲבִיחָה of the Old Testament, and the συκομορέα of the New. The wood of the tree is said to be very durable. *Morus nigra* supplies the common black Mulberry, which is an anthocarpous fruit, composed of numerous succulent flowers, forming a sorosis (fig. 475). The Mulberry is the συράμινος, or Sycamine-tree of the New Testament. Some have considered it as the שִׁטָּה* of the Old Testament. The white Mulberry, a less esteemed fruit, is the produce of *Morus alba*. Both of these mulberries are subacid. Their leaves are the favourite food of silk-worms. The root of the white Mulberry is anthelmintic. *Dorstenias* have a slightly concave broad receptacle, bearing numerous flowers (fig. 245). *D. Contrayerva*, *D. Houstoni*, and *D. brasiliensis*,

* The שִׁטָּה of the Bible is the name of some tree or shrub, and it is doubtfully translated Mulberry-tree.

furnish the Contrayerva-root of commerce. The officinal part is the root-stock, which is used as a stimulant, tonic, and diaphoretic. *Broussonetia papyrifera* is the Paper-mulberry, so called on account of being used in China and Japan in the manufacture of a kind of paper. It is called Crape-paper, and is prepared by pounding the bark, steeping it in water, then mixing it with glue, and taking it up with a mould of Bamboo-screen of the size required. The dye-wood called Fustic is the produce of *Maclura (Broussonetia) tinctoria*.

1022. The Artocarpus section is important as regards its uses. *Artocarpus incisa*, the Bread-fruit tree, supplies an amylaceous fruit, which furnishes an abundant supply of food in tropical countries. The properties of this tree are thus enumerated by Hooker:—The fruit serves for food; clothes are made from the fibres of the inner bark; the wood is used for building houses and making boats; the male catkins are employed as tinder; the leaves for table-cloths and for wrapping provisions in; and the viscid milky juice affords birdlime. *A. integrifolia* is the Jack or Jaca, the fruit of which attains a large size, sometimes weighing thirty pounds, and is inferior in quality to the Bread-fruit. In both instances the fruit is a sorosis, consisting of numerous flowers on a common axis, which becomes succulent. The milky juice of many of the Artocarpus tribe supplies caoutchouc, and in some instances it is used as a substitute for milk. This is the case with the juice of *Galactodendron utile* (perhaps a species of *Brosimum*,) which is called Palo de Vaca, or the Cow-tree, in Demerara. While the juice of some is nutritive, that of others is highly poisonous. Thus *Antiaris toxicaria* is the source of the famous poison called Bohun-Upas, or Upas-Antiar, by the Javanese, and which is said to owe its properties to the presence of Strychnia. Another Upas poison, called Upas-Tieuté, has already been noticed under the order Loganiaceæ, as being the produce of a species of Strychnos. The wood of *Piratinera guanensis* (probably a species of *Brosimum*) is called Snake-wood, or Letter-wood in Demerara, and is used for articles of furniture. Specimens sent by Dr. Campbell from Demerara have been beautifully manufactured in Scotland. The bark of *Lepurandra saccidora* (probably a species of *Antiaris*,) is used for forming sacks. Mr. N. B. Ward, in London, has one of these bags in his Museum, which has been formed by separating the bark entire from the wood throughout the whole extent, with the exception of a small portion at one end. The wood has then been removed from the interior, a part being left with the bark attached to form the bottom of the sack. The seeds of many of the Artocarpus tribe are eaten. *Brosimum Alicastrum* yields Bread-nuts, which, when boiled or roasted, are nutritious and agreeable articles of food. *Cecropia peltata* is the Trumpet-wood, so called on account of the hollowness of its stem and branches, which are used for wind instruments. The fibrous bark of the tree is used for cordage.

The two first of these suborders only are by recent writers included under the head. The third, or Ulmaceæ, constitute an independent order, which has been subdivided into two suborders:—1. Celtideæ, Nettle-trees, with a 1-celled ovary and amphitropal ovules. 2. Ulmeæ, true Elms, with a 2-celled ovary and anatropal ovules. The plants of the two last divisions, the Mulberry and Bread-fruit tribe, have also been constituted into a separate order, under the title of Artocarpaceæ, which differs from the true Urticaceæ chiefly in being lactescent, and in their fruit being a sorosis or syconus.

1023. Order 167.—**Ceratophyllaceæ**, the Hornwort Family. (*Apet. Diclin.*) Flowers unisexual. Perianth inferior, 10-12-parted. Male flowers: stamens 12-20; anthers sessile, bilocular. Female flowers: ovary free, 1-celled; ovule solitary, pendulous, orthotropal; style filiform, oblique; stigma simple. Fruit a 1-celled indehiscent nut, terminated by the hardened style. Seed solitary, pendulous, exalbuminous; cotyledons 2, but apparently 4; radicle inferior.—Aquatic submersed herbs, with verticillate leaves cut into filiform lobes. They are found in ditches in various parts of Europe, Asia, and America. The affinities of the order are still obscure. Some authors consider it as allied to Lythraceæ, others to Chenopodiaceæ, while Lindley puts it in his Urticle alliance. Its properties are unimportant. There is only 1 genus, and about 6 species. *Example*—Ceratophyllum.

1024. Order 168.—**Podostemaceæ**, the Podostemon Family. (*Apet. Monoclin.*) Flowers naked, or with a more or less perfect perianth, bursting through an irregularly lacerated spatha. Stamens hypogynous, definite or indefinite, distinct or monadelphous; anthers dithecal, with longitudinal dehiscence. Ovary free, 2-3-celled; ovules numerous, anatropal, attached to a fleshy central placenta; styles or stigmas 2 or 3. Fruit slightly pedicellate, capsular, 2-3-valved. Seeds 00; embryo exalbuminous, orthotropal.—Herbaceous, branched, floating plants, with capillary, or linear, or lacerated, or minute and imbricated leaves. Natives chiefly of South America, and of the islands to the east of Africa. Little is known in regard to their properties. The affinities of the order are not well determined. Some authors put it among the Monocotyledons. There are 9 known genera, and 25 species, according to Lindley. *Examples*—Podostemon, Lacis.

1025. Order 169.—**Stilaginaceæ**, the Stilago Family. (*Apet. Diclin.*) Flowers unisexual. Perianth 2-3- or 5-partite. Male flowers: stamens 2 or more, arising from a swollen receptacle; filaments capillary; anthers innate, 2-lobed, with a fleshy connective, and vertical cells opening transversely. Female flowers: ovary free, 1-2-celled; ovules 2, anatropal; stigma sessile, 3-5-toothed. Fruit drupaceous. Seed solitary, suspended; embryo in fleshy albumen; cotyledons leafy; radicle superior.—Trees or shrubs, with alternate, stipulate leaves. Natives chiefly of the East Indies. Some yield edible fruits, others

are used as potherbs. The position of this order in the natural system is obscure. Lindley places it in the Urtical alliance, others consider it as allied to Amentaceæ. There are 3 known genera, and about 20 species. *Examples*—*Stilago*, *Antidesma*.

1026. Order 170.—**Monimiaceæ**, the Monimia Family. (*Apet. Diclin.*) Flowers unisexual. Perianth somewhat globose, in one or more rows, divided at the border. Male flowers: stamens indefinite, covering the whole interior of the perianth; filaments often with 2 scales at the base; anthers dithecal, with longitudinal dehiscence. Female flowers: ovaries several, superior, enclosed within the tube of the perianth, each with 1 style and 1 stigma; ovule solitary, pendulous, anatropal. Fruit consisting of several achænia, enclosed within the enlarged perianth. Seeds pendulous; embryo at the end of copious fleshy albumen; radicle superior.—Trees or shrubs, with opposite exstipulate leaves. They are natives chiefly of South America and Australia. The bark and leaves are aromatic and fragrant. The succulent fruit of some is eaten. There are 8 known genera, and about 40 species. *Examples*—*Monimia*, *Boldoa*.

1027. Order 171.—**Atherospermaceæ**, the Plume-Nutmeg Family. (*Apet. Diclin.*) Flowers unisexual. Perianth tubular, divided at the top into several segments in 2 rows, the inner often petaloid, and accompanied in the female flowers with a few scales. Male flowers: stamens 00, inserted in the bottom of the perianth; filaments with scales at the base; anthers 2-celled, with valvular dehiscence. Female flowers: ovaries usually 00; ovule solitary, erect; style simple, lateral or basilar; stigmas simple. In some flowers, though rarely, stamens and pistils are found, and in that case the stamens are fewer, and arise from the orifice of the perianth. Fruit consisting of achænia, with persistent, ultimately feathery styles, enclosed within the tube of the perianth. Seed solitary, erect; embryo small, at the base of soft fleshy albumen; radicle inferior.—Trees, with opposite, exstipulate leaves, found in Australia, and in some parts of South America. They are generally fragrant. There are 3 known genera, and 4 species, according to Lindley. *Examples*—*Atherosperma*, *Laurelia*.

1028. Order 172.—**Lacistemaceæ**, the Lacistema Family. (*Apet. Diclin.*) Flowers polygamous. Perianth in several narrow divisions, covered by an enlarged bract. Stamen 1, hypogynous; anther having 2 cells, which are separated by a thick 2-lobed connective, and which dehisce transversely. Disk fleshy. Ovary superior, 1-celled; ovules several, anatropal, attached to 2-3 parietal placentas; stigmas 2-3, nearly sessile. Fruit a unilocular, 2-3-valved capsule, with loculicidal dehiscence. Seed usually, by abortion, solitary, suspended, with a fleshy arillus; spermoderm crustaceous; embryo in fleshy albumen; cotyledons flat; radicle cylindrical, superior.—Small trees or shrubs, with simple, alternate, exstipulate leaves, and amentaceous flowers.

They are natives of the warm parts of America. Their properties are unknown. There are 2 genera, and 6 species. *Example*—*Lacistema*.

1029. Order 173.—**Chloranthaceæ**, the Chloranthus Family. (*Achlamyd.* *Monoclin.* or *Diclin.*) Flowers bisexual or unisexual, with a supporting scale. Perianth 0. Stamens definite, lateral, and if more than 1, connate; anthers monothechal, with longitudinal dehiscence, each adnate to a fleshy connective. Ovary unilocular; ovule solitary, pendulous, orthotropical; stigma sessile, simple. Fruit drupaceous, indehiscent. Seed pendulous; embryo minute, at the apex of fleshy albumen; cotyledons divaricate; radicle inferior, remote from the hilum.—Herbs or undershrubs, with jointed stems, opposite, simple, stipulate leaves, sheathing petioles, and spiked flowers. Natives of the warm regions of India and America. Some of them, as *Chloranthus officinalis*, are aromatic and fragrant, and have been used as stimulants and tonics. There are 3 known genera, and 15 species. *Example*—Chloranthus.

1030. Order 174.—**Saururaceæ**, the Lizard's-tail Family. (*Achlamyd.*) Flowers bisexual. Perianth 0, a scale or bract supporting the flowers. Stamens 3-6, clavate, hypogynous, persistent; filaments slender; anthers 2-celled, continuous with the filament, with a thick connective separating the lobes, dehiscence longitudinal. Ovaries 3-4, distinct, with 1 ascending orthotropical ovule, and a sessile recurved stigma, or united so as to form a 3-4-celled pistil, with several ovules and 3-4 stigmas. Fruit either consisting of 4 fleshy indehiscent nuts, or a 1-3-4-celled capsule, dehiscing at the apex, and containing a few ascending seeds. Seeds with a membranous spermoderm; embryo minute, lying in a fleshy vitellus, outside of hard mealy albumen at the apex of the seed.—Herbs growing in marshy places, with alternate, stipulate leaves, and spiked flowers. Natives of North America, India, and China. Their properties are said to be acrid. There are 4 known genera, according to Lindley, and 7 species. *Examples*—*Saururus*, *Houttuynia*.

1031. Order 175.—**Piperaceæ**, the Pepper Family. (*Achlamyd.*) Flowers ♂. Perianth 0, flowers supported on a bract. Stamens 2-3-6, arranged on one side or all round the ovary; anthers 1- or 2-celled, with or without a fleshy connective; pollen roundish, smooth. Ovary solitary, free, 1-celled; ovule solitary, erect, orthotropical; stigma simple, sessile, rather oblique. Fruit somewhat fleshy, indehiscent, unilocular. Seed erect; embryo in a fleshy vitellus outside the albumen, and at the apex of the seed.—Shrubs or herbs, with articulated stems, leaves opposite (sometimes alternate by abortion of one of the pair of leaves), or verticillate, exstipulate or stipulate, and spiked or racemose flowers. Natives of the hottest quarters of the globe. Common in South America and India. The wood is often arranged in wedges, with medullary rays, but without concentric zones. There

are 21 known genera, and upwards of 600 species. *Examples*—Piper, Artanthe, Peperomia.

1032. The plants of the order have pungent, acrid, and aromatic properties. Most of them contain an acrid resin, and a crystalline principle called Piperin, in which their active qualities reside. Some are narcotic and astringent. The dried fruiting spikes of *Piper longum*, an Indian creeper, constitute Long-pepper. Of late, however, the genus Piper has been subdivided, and this kind of pepper has been referred to various species of a new genus, *Chavica*, viz. *C. peepuloides*, *Roxburghii*, and *officinarum*. The dried unripe fruit (drupes) of *Piper nigrum* constitute Black-pepper, a climbing plant common in the East Indies. The ripe fruit, when deprived of its outer fleshy covering by washing, forms the White-pepper of the shops. These peppers are hot aromatic condiments, and they are used medicinally as tonic, stimulant, febrifuge, and stomachic. The fruit of *Piper Cubeba* (*Cubeba officinalis*), a climbing plant of Java and other Indian islands, is the medicinal Cubeb-pepper, which is used extensively in arresting discharges from mucous membranes. It contains a resin, a volatile oil which is very active, and a peculiar principle called Cubebin. The substance called Matico or Matica, consists of the leaves and unripe fruit of *Piper angustifolium* (*Artanthe elongata*.) It possesses aromatic, fragrant, and astringent qualities. It has been particularly recommended for checking hæmorrhage, a property which seems, in part, to be a mechanical one, depending on the structure of the leaf, which also abounds in tannin. The root of *Piper methysticum* is the Kava of the South Sea Islanders, which is used by them for preparing a stimulating beverage. The leaf of Betel-pepper (*Piper Belle*) is chewed with the Areca nut in the East, as a means of intoxication.

1033. Order 176.—**Amentaceæ**, the Catkin-bearing Family. (*Diclin.*) Flowers unisexual. Male flowers (fig. 721) capitate or in catkins



Figs. 721-727.—Organs of fructification of *Corylus Avellana*, the Hazel, to illustrate the natural order Amentaceæ, section Cupuliferæ.

Fig. 721.—Male flower separated from the catkin (amentum). *e*, Scale or bract bearing the stamens, *a*, with their dithecal anthers.

Fig. 722.—Female flower, *ff*, in a very young state, with its involucre, *i*.

Fig. 723.—Female flower more advanced. *i*, Involucre opened to show the ovary, *o*, covered by the perianth, *c*. *s*, Two styles.

(amenta), (fig. 238,) sometimes with a membranous perianth. Female flowers clustered, solitary (fig. 722), or in catkins. Stamens varying



from 1 to 20, distinct (fig. 721) or monadelphous; anthers dithecal (fig. 721). Ovary usually simple (fig. 723 o); stigmas 1 or more (fig. 723 s). Fruit membranous or bony, or drupaceous, indehiscent (fig. 726), or dehiscent. Seeds solitary or numerous, erect or pendulous (fig. 724), usually exalbuminous; embryo straight or curved; radicle mostly superior (fig. 727).—Trees or shrubs, with alternate, stipulate, or exstipulate leaves. Natives chiefly of temperate climates. The order has been divided into the following suborders:—

1. Salicinæ, the Willow tribe: flowers naked, or with a membranous cup-like calyx; ovules 00, erect, anatropal; fruit naked, coriaceous, unilocular, 2-valved; seeds comose; embryo erect; radicle inferior; leaves stipulate. Found in woods in temperate and cold regions.
2. Myricæ, the Gale tribe: achlamydeous flowers; stamens 2-8 in the axil of a scale; ovary 1-celled, with hypogynous scales; ovule solitary, erect, orthotropal; fruit drupaceous, often with a waxy secretion, and with fleshy adherent scales; radicle superior. Natives both of temperate and tropical regions, and found in North and South America, in India, and at the Cape of Good Hope.
3. Casuarinæ, the Beef-wood tribe: flowers with bracts; stamen 1; ovary 1-celled; ovules 1-2; fruit consisting of winged achænia, collected into a cone; seed erect; radicle superior. Australian trees or shrubs, with filiform branches, bearing membranous toothed sheaths in place of leaves.
4. Betulinæ, the Birch tribe: flowers with bracts, which are sometimes verticillate; ovary 2-celled; ovules solitary, pendulous, anatropal; fruit membranous, indehiscent, forming a sort of cone; seeds pendulous; radicle superior; leaves with deciduous stipules. Natives of temperate and cold regions in Europe, Asia, and America, and extending to arctic and antarctic regions.
5. Balsamaceæ (Altingiaceæ), the Liquidambar tribe: flowers with verticillate bracts or minute scales; anthers numerous; ovary 2-celled; ovules 00, amphitropal; fruit consisting of 2-celled capsules, united together so as to form a hard cone; seeds usually numerous, winged, albuminous; radicle superior; leaves stipulate. Balsamic trees, natives of tropical and warm regions.
6. Platanæ, the Plane tribe: flowers in globose catkins; stamen 1, with scales; ovary 1-celled; style thick and subulate; ovules solitary or in pairs; suspended, ortho-

Fig. 724.—Female flower cut longitudinally, to show the two loculaments with a pendulous ovule in each.

Fig. 725.—Female flower more advanced. c, Perianth. s, Styles.

Fig. 726.—Ripe fruit, f, enveloped in its involucre or bracts, i.

Fig. 727.—Seed separated. i, Integument, half of which is removed to show the exalbuminous embryo, e. r, Superior radicle.

tropical; fruit consisting of compressed clavate nuts, terminated by a recurved style; seeds 1-2, pendulous, albuminous; radicle inferior; leaves palmate or toothed, and stipulate. Natives chiefly of temperate regions.

7. Cupuliferæ (Corylaceæ), the Nut tribe: flowers amentaceous (fig. 238) or aggregate; stamens 5-20, attached to scales (fig. 721) or a perianth; ovary surrounded by a coriaceous involucre (cupula), crowned by the remains of a persistent perianth (figs. 723, 725), multilocular; ovules 2 or 1, pendulous; fruit a glans (figs. 257, 726); seed usually solitary; radicle superior (fig. 727); leaves stipulate, often feather-veined (fig. 134). Natives of temperate regions chiefly. Some extend to warm countries.

This extensive Amental alliance, with which perhaps the next order should be incorporated, embraces 18 known genera, and 600 species. The suborders are by some considered as separate orders, and placed in various parts of their system of alliances. *Examples*—*Salix*, *Populus*; *Myrica*; *Casuarina*; *Betula*, *Alnus*; *Liquidambar*; *Platanus*; *Corylus*, *Fagus*, *Castanea*, *Quercus*.

1034. The Amentaceous order contains trees which yield valuable timber, and are extensively cultivated. Many have bitter, tonic, and astringent qualities. Some yield resinous and balsamic fluids; and the seeds of some are used as articles of food. The bark of many species of Willow, such as *Salix Capræa*, *alba*, *Russelliana*, *fragilis*, *pentandra*, *vitellina*, *purpurea*, and *Helix*, yields a neutral crystalline bitter substance called Salacin, which is employed as a febrifuge and tonic. The bark also possesses astringent qualities from the presence of tannin. Salicin assumes a carmine-red tint when moistened with a few drops of concentrated sulphuric acid. Various species of Willows (osiers) are used for basket-making, while others are employed in forming charcoal. *Salix babylonica* is the Weeping-willow. Its specific name is founded on the supposition, that it was the species on which the Israelites hanged their harps by the waters of Babylon. The word *אֲרֵז* in the Bible is doubtfully translated Willow. *Populus alba* is commonly called the Abele, while *P. tremula* is the Aspen, and *P. fastigiata* and *dilatata* the Lombardy Poplar. The buds of *Populus nigra* and *balsamifera* are covered with a resinous exudation, to which the name of Tacamahac has been given; it is said to be diuretic and antiscorbutic. The leaves and bark of some Poplars secrete a saccharine substance, which has been termed Populine. *Salix arctica* and *polaris* extend to the arctic regions, and form the most northern woody plants. *Salix herbacea*, a small creeping Willow, occurs abundantly on the Scotch mountains. The downy matter surrounding the seeds of Poplars and Willows, is used for stuffing pillows and cushions, as well as for the manufacture of a kind of paper.

1035. The species of *Myrica* are aromatic, and yield resinous and oily matter. *Myrica Gale* is the Gale or Scotch Myrtle, which is common in marshy grounds and damp heaths in Britain. The fruit of *Myrica cerifera*, called Wax Myrtle, or Bay Myrtle, or Candleberry,

466
 466
 466

 yields a greenish-coloured wax, which is used for candles. The drupaceous fruit of some *Myricas* is eaten. The leaves of *Comptonia asplenifolia*, Sweet Fern, are found by Mr. Murchison to contain peculiar glands. Resinous matter is also procured from several species of *Liquidambar*, and is known by the name of Liquid Storax. *Liquidambar styraciflua*, *orientale*, and *Altingia*, are said to be the chief sources of this resin or balsam, containing Benzoic acid. The species of *Casuarina* (Cassowary-tree) yield excellent timber, called Beefwood, from its having some resemblance to raw beef. What is called the She-oak in Australia, is *C. quadrivalvis*. In the integument of the seeds of *Casuarinas* there are numerous spiral cells.

1036. The species of *Betula*, Birch, have astringent and resinous barks. The oil from the bark of the common Birch (*Betula alba* and *glutinosa*) gives the peculiar odour to Russia leather. In North America the bark of the Canoe Birch (*Betula papyracea*) is used for making boats. A saccharine matter exists in the sap of the Birch. *Betula lenta* is the Black Birch of America, and is called Mountain Mahogany on account of the beauty and hardness of its timber. *Alnus glutinosa*, common Alder, grows well on the muddy ground on the banks of rivers. *Platanus orientalis*, the Oriental Plane, has broad palmate leaves, resembling the Sycamore, which is often erroneously called the Plane in Scotland. Some say that this is the Sycamore of the ancients. *Platanus occidentalis*, another species of Plane, is also cultivated as a showy tree.

1037. The Hazel-nut, with its involucre appendage, is the produce of *Corylus Avellana*. The bark of *Quercus pedunculata* (*Robur*), the common Oak, contains much tannin, and is used as an astringent. Another British species, *Q. sessiliflora*, having sessile fruit, is said to yield the best timber. It is said that, in the wood of *Q. pedunculata*, there are numerous medullary rays (silver-grain), (fig. 99), while in that of *Q. sessiliflora*, Durmast, there are scarcely any visible. There is some doubt as to the existence of more than one species in Britain, and no permanent characters have been established. Babington mentions three species. It has been stated that 2000 well-grown oaks, equal to 3000 loads of timber, are required to build a seventy-four gun-ship. The acorns of *Quercus Ægilops*, Valonia or Balonia Oak, under the name of Velonia, are used by dyers. Oaks are liable to the attacks of insects, whose punctures give rise to the formation of galls. These excrescences occur on the buds, bark, and leaves. The medicinal galls are the produce of *Quercus infectoria*, a native of Asia Minor, and the best are imported from Aleppo. They are caused by punctures from the ovipositor of the *Diplolepis* (*Cynips*) *Gallæ-tinctorum*. Blue galls are those which still contain the young insect in their interior, while white galls are those from which it has escaped. In medicine they are employed as powerful astringents, and in the arts they are

used for dyeing, tanning, and forming ink. The bark (Epiphloeum) of *Quercus Suber* constitutes cork (fig. 100). The bark of *Quercus tinctoria* is called Quercitron, and yields a yellow dye. *Quercus Ilex*, Evergreen Oak, is commonly cultivated in gardens. The Oak, אֵילָנָה of the Bible, is said by some to be *Quercus Grammuntia*. The Beech-tree (*Fagus sylvatica*), the Horn-beam (*Carpinus Betulus*), and the Spanish Chestnut (*Castanea vesca*), belong to the Cupuliferous section of this order. *Fagus Forsteri* is the Evergreen Beech of South America, found at Terra del Fuego. A species of Beech (*F. antarctica*) is found in the antarctic regions.

1038. Order 177.—**Juglandaceæ**, the Walnut Family. (*Apet. Diclin.*) Flowers unisexual. Male flowers amentaceous: perianth membranous, oblique, irregularly-lobed, with a scaly bract. Stamens definite or 00; filaments short, free; anthers dithecal, erect. Female flowers in terminal clusters, or in loose racemes, with separate or united bracts: perianth single or double, the outer 3-5-parted, inner, when present, in minute separate pieces. Ovary adherent to the perianth, 1-celled; ovule solitary, erect, orthotropal; styles 1-2, very short; stigmas 2-4, fringed or sessile, discoid and 4-lobed. Fruit a drupe, sometimes with an adherent involucre; endocarp bony, 2-valved or valveless, 2-4-celled at the base, and 1-celled at the apex, with partial dissepiments. Seed exalbuminous, 2-4-lobed, with a membranaceous testa; embryo large; cotyledons fleshy, oily and sinuous; radicle superior.—Trees with alternate, pinnated leaves, having neither dots nor stipules. They are chiefly natives of North America. There are 4 genera, according to Lindley, and 27 species. *Examples*—*Juglans*, *Carya*.

1039. While the plants belonging to this order yield edible oily nuts, their bark is often acrid, and there is frequently bitterness and astringency in the coverings of their fruit and seed. The seeds of *Juglans regia*, common Walnut, yield a bland oil, which may be used as a substitute for olive-oil. *Carya alba* yields the American Hickory-nut. Purgative and resinous properties prevail in some of the plants. The timber of many of the trees is valuable. That of the Black Walnut (*Juglans nigra*) has a fine dark-brown colour when polished.

1040. Order 178.—**Garryaceæ**, the Garrya Family. (*Apet. Diclin.*) Flowers unisexual, amentaceous. Male flowers: perianth of 4 parts. Stamens 4, alternate with the segments of the perianth. Female flowers: perianth superior, 2-toothed. Ovary unilocular; ovules 2, pendulous on short funiculi; styles 2. Fruit baccate, indehiscent. Seeds 2; embryo minute, at the base of fleshy albumen.—North American shrubs, with opposite, exstipulate leaves. The male plants of *Garrya elliptica* are commonly cultivated in shrubberies, and are prized for their peculiar silky catkins. Lindley associates with this order the Helwingiaceæ, which agree in their unisexual flowers, adhe-

rent fruit, pendulous ovules, minute embryo, at the base of solid albumen. There are 2 known genera, and 6 species. *Example*—Garrya.

b. SPOROGENÆ OR RHIZANTHÆÆ.

Having spore-like seeds with an Acotyledonous embryo.

1041. Order 179.—**Rhizanthææ**, the Rhizanth or Rhizogen Family. (*Apet. Diclin.*) Flowers usually monœcious or diœcious, sometimes ♂. Perianth more or less perfect, superior, trimerous, tetramerous or pentamerous, sometimes obsolete or 0. Stamens united, often in a fleshy column, to which the anthers cohere, dithecal, extrorse, opening longitudinally or by pores. Ovary inferior, 1-2-celled; ovules definite or 00. Fruit indehiscent, pulpy, usually unilocular. Seeds sometimes solitary and pendulous, at other times 00, and attached to parietal placentas; embryo albuminous or exalbuminous.—Leafless, scaly, parasitic plants, having a fungus-like appearance. They are never green, but assume a brown, yellow, or purple colour. They are composed chiefly of cellular tissue, with a few scalariform and spiral vessels. They are often stemless, and sometimes are furnished with a creeping rhizome. In their mode of decay they resemble Fungi. Their seeds present a peculiar appearance, resembling spores rather than true seeds. The nature of their embryo is undetermined, and their place in the natural system is still disputed. Lindley has placed them in a separate class, intermediate between Thallogens and Endogens. They have been divided by him into three distinct orders:—1. Balanophoracæ, male flowers pedicellate; stamens 1-3; filaments and anthers both united; ovule solitary, pendulous; fruit monospermous. 2. Cytinacæ, flowers in spikes; perianth 3-6-lobed; anthers sessile on a column, dehiscing by slits; ovules 00, attached to parietal placentas; fruit polyspermous. 3. Rafflesiacæ, flowers sessile, solitary; perianth 5-lobed, with calli in its throat; anthers attached to a column, dehiscing by pores; ovules 00, attached to parietal placentas; fruit polyspermous. They are natives chiefly of tropical countries, but some extend into temperate climates. They are found in the East Indies, South America, Cape of Good Hope, and the south of Europe. Lindley enumerates 21 genera, and 53 species. *Examples*—Balanophora, Cynomorium, Cytinus, Rafflesia.

1042. Some of the plants are astringent, and have been employed as styptics. *Cynomorium coccineum*, commonly known as Fungus melitensis, grows in Malta, and was long celebrated for arresting hæmorrhage. *Cytinus* is parasitic on the roots of Cistuses in the south of Europe. *Cytinus hypocistis* is said to contain gallic acid. The species of *Rafflesia* are gigantic parasites, the perianth being sometimes three feet in diameter, and capable of holding twelve pints of fluid. They

are parasitic on the stems of a species of *Cissus*, as well as on some leguminous plants. *R. Patma* is employed in Java as an astringent and styptic.

Section B.—GYMNOSPERMÆ.*

1043. Monochlamydeous or Achlamydeous plants, with an Exogenous structure as regards their stems and organs of vegetation, but differing from Exogens in having naked ovules, which are fertilized by the direct application of the pollen to the foramen, without the intervention of stigma, style, and ovary. Their woody tissue is marked by the presence of disks (figs. 47, 48, 808). They are included in Lindley's class of Gymnogens, and Endlicher's Gymnospermous division of Acramphibrya.

1044. Order 180.—**Coniferæ**, the Cone-bearing Family. (*Achlamydl. Diclin.*) It includes the orders Pinaceæ, Taxaceæ, and Gnetaceæ of Lindley. Flowers unisexual. Male flowers monandrous or monadelphous: stamens collected in a deciduous amentum, about a common rachis (fig. 728); anthers 1-2 or many-lobed, with longitudinal dehiscence, often terminated by a scaly crest (fig. 729). Female flowers in cones (fig. 730), sometimes solitary: ovary none, its place being supplied by the flat scales of the cones, arising from the axil of membranous



bracts (fig. 731); ovules naked, usually in pairs on the face of the scales (figs. 731, 732 oo), inverted or erect; style 0; stigma 0. Fruit a cone (figs. 201, 476, 477), or a solitary naked seed (fig. 442.) Seed

* *Γυμνός*, naked, and *σπίγυα*, seed.

Figs. 728-734.—Organs of fructification of *Pinus sylvestris*, Scotch Fir, to illustrate the natural order Coniferæ.

Fig. 728.—Collection of male catkins, c, clustered round a common axis. f, Leaves. b, Terminal buds, with young leaves and scaly sheaths.

Fig. 729.—Male flower, or the two-lobed anther, separated.

Fig. 730.—Three collections of female flowers, or young cones, c, at the extremity of a branch.

Fig. 731.—A scale detached from one of these young cones, and seen on the exterior. b, Bract. e, Scale. oo, Summit of the naked ovules.

with a hard crustaceous integument, sometimes winged (fig. 733); embryo in the midst of fleshy oily albumen (fig. 734); sometimes more than one embryo; cotyledons 2, or many and verticillate (fig. 734);



radicle next the apex of the seed, organically connected with the albumen.—Trees or shrubs, with branched, usually resinous trunks, the wood marked with circular disks (figs. 47, 48, 808), the leaves usually narrow, rigid or acerose, entire (fig. 147), sometimes fascicled, and with a scaly sheath at their base (fig. 728 *b*). They are found in various parts of the world, both in cold and hot regions. They abound in the temperate regions of Europe and America, and many occur in Australasia. Four genera of Coniferæ, *Araucaria*, *Phyllocladus*, *Microcachrys*, and *Arthrotaxis*, are peculiar to the southern hemisphere. The following attain their maximum to the south of the tropics,—*Calitris*, *Podocarpus*, and *Dacridium*. *Dammara* has one species in each hemisphere.

1045. The order is a very extensive one, and has been divided into the following suborders:—

1. Abietineæ, the Fir and Spruce tribe: fertile flowers in cones, with 1 or 2 inverted ovules at the base of each scale; embryo in the axis of fleshy and oily albumen, di- or poly-cotyledonous. Under this tribe are included the following sections and sub-sections:—

A. Scales 2-seeded, seeds adnate to the scale, and at length separating from it; anthers bilocular.

a. Scales with a thickened apophysis, which is either entire or dimidiate.

Pinus.—Leaves in twos, threes, fours, or fives.

b. Scales without an apophysis.

Fig. 732.—Scale of a young cone seen on the inside. *c*, The scale. *i*, The point by which it is attached to the axis of the cone. *o o*, The two naked inverted ovules. *m*, Their upper opening or foramen to which the pollen is applied. The foramen was formerly described erroneously as a stigma.

Fig. 733.—A scale from a mature cone. *c*, The scale. *i*, Point of insertion. *g*, One of the winged seeds; the other having been removed.

Fig. 734.—The seed cut longitudinally. *a*, Base of the wing. *t*, Integument. *p*, Perisperm (albumen). *e*, Polycotyledonous embryo. Near the radicle are the remains of two other abortive embryos.

* Leaves solitary.

Abies.—Scales deciduous, leaves flat.

Picea.—Scales persistent, leaves tetragonous.

Tsuga.—Scales persistent, leaves flat.

* * Leaves fasciculated.

Larix.—Leaves flat, annual.

Cedrus.—Leaves tetragonous, perennial.

B. Scales 1-seeded, seed adnate to the scale, and not separating from it, anthers multilocular.

Araucaria (*Eutassa* and *Alingia*).

C. Scales 1- or many-seeded, seeds free, anthers by- tri- or multi-locular.

Dammara.—Anthers multilocular.

Cunninghamia.—Anthers trilocular.

Arthrotaxis.—Anthers bilocular.

2. Cupressineæ, the Cypress tribe : ovules erect ; fruit an indurated cone (fig. 477), or fleshy, with the scales connected, forming a galbulus (fig. 478) ; embryo di- or poly-cotyledonous. *Examples*—Cupressus, Juniperus, Thuya, Taxodium, Cryptomeria, Callitris.
3. Taxineæ, the Yew tribe : anthers usually bilocular, with longitudinal dehiscence : fertile flowers, solitary, terminal ; ovule solitary, sessile in the centre of a fleshy disk, when in fruit forming a sort of drupe (fig. 442) ; embryo dicotyledonous. *Examples*—Taxus, Podocarpus, Dacrydium, Phyllocladus Salisburia.
4. Gnetaceæ, the Joint-fir tribe : male flowers with a perianth, anthers uni- or quadrilocular, opening by a short cleft ; ovules with a projecting process formed from the intimate covering of the nucleus ; seed solitary ; embryo with a long spirally-twisted funiculus ; stems jointed ; zones of wood, often separated by marked cellular circles (fig. 104). *Examples*—Gnetum, Ephedra.

The order embraces 31 known genera, and 165 species, according to Lindley. Zuccarini enumerates 208 species of Coniferæ,—in the northern hemisphere, 165, and in the southern, 51 ; some species being common to both hemispheres.

1046. The plants of this order furnish valuable timber, and yield various important products, such as turpentine, pitch, and resin. The various kinds of Pine, Fir, Spruce, and Cedar, belong to this family. *Eutassa* (*Araucaria*) *excelsa* is the Norfolk-island Pine, famed for its size and for its wood. *Cedrus Libani* is the Cedar of Lebanon, the צֶדֶר of the Bible, while *Cedrus Deodara* is the Deodar or Himalayan Cedar. By exudation, and partly by the aid of heat, the plants of this order yield various kinds of turpentine, resin, tar, and pitch. Common turpentine is procured from *Pinus sylvestris*, the Scotch Fir, *Pinus Pinaster*, the Cluster-Pine, and var. *maritima*, Bourdeaux Pine, *Pinus palustris*, Swamp Pine, and *Pinus Tæda*, Frankincense Pine. Oil of turpentine is obtained from it by distillation. Venice turpentine and Strasburg turpentine are the produce of *Abies Picea* (*Abies* or *Picea pectinata*), the Silver Fir, and *Larix europæa*, the Larch ; while

Canada Balsam is collected from *Abies* or *Picea balsamea*, Balm of Gilead Fir, and *A. canadensis*, Hemlock Spruce. *Dammara australis*, the Kawrie-tree of New Zealand, yields a hard resin, and so does *D. orientalis*, the Amboyna Pitch-tree. *Callitris quadrivalvis* (*Thuja articulata*), the Arar-tree, supplies a solid resin called Sandarach or Pounce, which is used to strew over manuscripts. Thus, or Common Frankincense, and Burgundy-pitch, are yielded by *Abies excelsa*, Norway Spruce Fir. *Pinus Pumilio* gives Hungarian balsam, *Pinus pinea*, the Stone Pine, is the source of Carpathian balsam. Essence of Spruce, used in making Spruce-beer, is got by boiling in water the leaves of the Scotch Fir (*Pinus sylvestris*), the Black Spruce (*Abies nigra*), and other species. A kind of Manna is procured from the Larch and from the Cedar of Lebanon. The Bark-bread of the Norwegians is prepared from the inner bark of *Pinus sylvestris*. These various kinds of resin and pitch are used for stimulating and healing plasters, while the oil of turpentine (*oleum terebinthine*) is employed medicinally as a stimulant, diuretic, cathartic, and anthelmintic. The vapour of tar has been recommended in affections of the chest. The succulent cones (fig. 478) (commonly called berries) of *Juniperus communis*, Common Juniper, and the oil procured from them, are used medicinally as diuretics. The oil enters into the composition of the spirituous liquor called Hollands. The young branches and leaves of *Juniperus Sabina*, Savin, contain an active volatile oil, which is used as an anthelmintic and emmenagogue. In large doses it acts as a violent irritant poison. The wood of *Juniperus Bermudiana* furnishes Pencil Cedar. *J. Virginiana*, the Red Cedar, yields a rubefacient oil. *Thuja occidentalis* is the common Arbor-vitæ of gardens. *Thuja orientalis* is also in cultivation. *Cupressus sempervirens*, common Cypress, yields a durable wood, which is supposed to be the Gopher-wood, גפרית, of the Bible. *Podocarpus Totarra* and *Dacrydium taxifolium*, both supply good timber in New Zealand. *Taxus baccata*, the Yew, is a valuable timber tree. It yields resin, and its leaves and seeds are said to be narcotico-acrid. *Salisburia* has remarkable cuneate leaves, and the fruit of *S. adiantifolia*, the Ginko, is said to be eatable. *Gnetum urens* has singular stinging hairs within the epispERM or outer integument of the seed.

1047. Order 181.—*Cycadaceæ*, the Cycas Family. (*Achlamydeclin.*) Flowers unisexual. Males collected into cones, the scales bearing on their lower surface 1-celled anthers, which are united often in sets of two, three, or four. Females consisting of naked ovules, placed at the base of flat scales, or beneath peltate ones, or seated on the margins of altered leaves. Seeds hard and nut-like, sometimes with an external spongy coat; embryos 1 or 2, suspended in a central cavity; albumen fleshy or mealy; cotyledons unequal; radicle superior, having a long cord-like prolongation by which the

embryo is suspended.—Trees or shrubs, with cylindrical trunks, usually simple, sometimes dichotomous, marked with the scars of the leaves, and in many respects having the aspect of Palms (fig. 817). The internal structure is more or less distinctly that of dicotyledons. Pitted tissue and spiral vessels occur. The leaves are pinnate, and their venation is circinate, thus resembling ferns. The plants of this order are found in the temperate and warm regions of America and Asia, as well as at the Cape of Good Hope. There are 6 genera, according to Lindley, and 45 species. *Examples*—*Cycas* (fig. 817), *Zamia* (fig. 818).

1048. The Cycadaceous family yields much starchy matter, along with mucilage. From the stems of *Cycas revoluta* and *circinalis*, a kind of Sago is made. A clear insipid mucilage also exudes from them, which hardens into a transparent gum resembling tragacanth. This gum when dried in the air, coagulates into a mass, which when applied to malignant ulcers quickly excites suppuration. *Zamia pumila*, and other species in the West Indies, supply an amylaceous matter which has been sold as Arrow-root. The Bread-tree is a name applied by the Hottentots to various species of *Encephalartos*.

CLASS II.—MONOCOTYLEDONES, *Juss.* ENDOGENÆ, *DC.* ENDOGENS AND DICOTYGENS, *Lindl.* AMPHIBRYA, *Endl.*

1049. In this great class the plants have a *cellular* and *vascular* system, the latter consisting partly of elastic spiral vessels (fig. 51). The woody *stem* (as in Palms, fig. 115, 1) is usually more or less cylindrical, simple, and unbranched. There is no true separable bark, no concentric zones, and no true pith (figs. 112, 113). The *wood* is endogenous, *i.e.* increases by additions, which first tend towards the centre, and then curve outwards in an interlacing manner (fig. 114, 2) towards the circumference, where much hard ligneous matter is deposited, so as to make the exterior the hardest part (figs. 826, 827). The development of the stem usually takes place by a single central and terminal bud; occasionally lateral buds are produced (fig. 115, 2), and at times the stem is hollow. The *leaves* are parallel-veined (figs. 135, 173, 194), except in the subclass Dictyogens, where a kind of reticulation is visible. The parts of the *flower* are arranged in a ternary manner (fig. 538), and they are often petaloid (fig. 260), sometimes scaly or glumaceous. The ovules are contained in an ovary, and are fertilized by the application of the pollen to the stigma. The *embryo* has one cotyledon (fig. 504), and the germination is endorhizal (fig. 527).

Subclass 1.—DICTYOGENÆ.*

1050. Leaves reticulated, often articulated with the stem, branches

* Δίκτυον, a net, and γένναι, to produce.

having the usual structure of Endogens, rhizomes or underground stems having the woody matter disposed in a compact circle, or in wedges containing central cellular tissue, and often showing medullary processes.

1051. Order 182.—**Dioscoreaceæ**, the Yam tribe. (*Mono-epigyn.*) Flowers unisexual. Perianth in 6 divisions, adherent. ♂. Stamens 6, inserted into the base of the perianth; anthers introrse, with longitudinal dehiscence. ♀. Ovary inferior, 3-celled; ovules 1-2, anatropal; style bifid; stigmas undivided. Fruit a compressed trilocular capsule, with 2 cells, sometimes abortive, occasionally fleshy and indehiscent. Seeds compressed, winged or wingless, in the succulent fruit, ovate; embryo small, near the hilum, lying in a large cavity of cartilaginous albumen.—Twining shrubs, with large epigeal or hypogeal tubers, alternate, sometimes opposite, reticulated leaves, and small, spiked, bracteated flowers. Natives chiefly of tropical countries; a few only found in temperate regions. There are 6 genera, according to Lindley, and 110 species. *Examples*—Dioscorea, Tamus.

1052. Although farinaceous matter exists in the tubers of many species, yet there is a prevalent acidity throughout the order. Various species of *Dioscorea*, as *D. alata*, *sativa*, and *aculeata* produce the tubers called Yams, which are used in warm countries as a substitute for the potato. *Testudinaria Elephantipes* is the Tortoise plant of the Cape, or Elephant's-foot, so called on account of its peculiar shortened and thickened stem (§ 92). *Tamus communis*, Black Bryony, is common in hedges in England. It produces red succulent fruit, and has a large root, which is acrid. This acidity does not extend to the young suckers, which may be eaten with impunity. The acidity of the order sometimes manifests itself in purgative qualities.

1053. Order 183.—**Smilacææ**, the Sarsaparilla Family. (*Mono-perigyn.*) Flowers bisexual or polygamous. Perianth petaloid, 6-parted. Stamens 6, inserted into the base of the perianth, rarely hypogynous. Ovary free, 3-celled; cells uni- or multi-ovulate; ovules orthotropal; styles usually 3-cleft; stigmas 3. Fruit globular and succulent. Seeds with fleshy, cartilaginous albumen; embryo very small; usually distant from the hilum.—Herbs or undershrubs, often climbing, with netted-veined leaves. Natives of the temperate and tropical regions of Asia and America. There are 4 or 5 known genera, and upwards of 120 species. *Examples*—Smilax, Philesia.

1054. Mucilaginous and demulcent properties prevail throughout the order. The root of various species of *Smilax* constitutes the Sarsaparilla or Sarza of the pharmacopœias. Linnæus considered *Smilax Sarsaparilla*, a native of the United States, as the plant which furnished Sarsaparilla, but recent observers state that this is not the case. The following are enumerated as sources whence Sarsaparilla of various kinds is derived.

1. *Smilax officinalis*, found in woods near the Rio Magdalena in Columbia. It furnishes Jamaica Sarza, which is the best in the market.
2. *Smilax medica*, native of the Mexican Andes. It is thought to furnish Vera Cruz Sarza.
3. *Smilax syphilitica*, found in Brazilian Guiana. It in part supplies Brazil and Lisbon Sarza.
4. *Smilax cordato-ovata*, a Brazilian species. Brazil Sarza in part.
5. *Smilax papyracea*, a Brazilian species. Brazil and Lisbon Sarza in part.
6. *Smilax Brasiliensis*, a Brazilian species. Brazil Sarza.
7. *Smilax*, sp. ? from Angostura, supplies Rio Negro Sarza.
8. *Smilax*, sp. ? from Honduras, supplies Honduras Sarza.

Besides these species, which supply in a greater or less degree the officinal drug, there are others which furnish substitutes in different countries, viz:—

9. *Smilax Purhampuy*, a Peruvian species.
10. *Smilax excelsa* and *aspera*, are used as substitutes in Europe.
11. *Smilax Zeylanica*, *glabra*, *perfoliata*, and *leucophylla*, in Asia.
12. *Smilax China*, a Chinese plant yielding China-root.
13. *Smilax Pseudo-China*, from the United States.

The officinal part is the roots which come off from the rhizomes. The roots are mucilaginous, bitterish, and slightly acrid. They contain mucilage, starch, oil, resin, and a crystalline principle called Pariglin or Smilacin. Sarsaparilla is used in decoction and infusion, as a tonic and alterative, in cachectic and syphilitic cases.

1055. Order 184.—**Trilliaceæ**, the Trillium Family. (*Mono-perigyn.*) Flowers usually bisexual. Perianth in 6, sometimes 8 divisions, coloured or herbaceous. Stamens 6, 8, or 10; filaments subulate; anthers linear, with a prolonged connective. Ovary free, 3- 4- or 5-celled; styles as many, distinct; ovules 00, anatropal. Fruit succulent, 3- 4- or 5-celled. Seeds 00; embryo minute, in fleshy albumen.—Natives of the temperate parts of Europe, Asia, and America. Some of them are more or less acrid, others are narcotic. The rhizome of *Trillium cernuum* is used as an emetic. The juice of the berries mixed with alum gives a blue colouring matter. *Paris quadrifolia*, Herb Paris, is narcotic. There are about 10 known genera, and upwards of 60 species. *Examples*—Trillium, Paris, Triuris?

Subclass 2.—PETALOIDEÆ.

1056. Flowers having usually a perianth consisting either of verticillate leaves, which may sometimes be separated into calyx and corolla, and are often coloured (petaloid), or of a few whorled scales. Occasionally the perianth is abortive.

a. Perianth adherent, Ovary inferior, Flowers usually hermaphrodite.

1057. Order 185.—**Hydrocharidaceæ**, the Frog-bit Family. (*Mono-epigyn.*) Flowers spathaceous, unisexual, rarely ♂. Perianth with a 6-partite limb, the 3 outer segments herbaceous, and equivalent to the


calyx, the 3 inner petaloid, and equivalent to the corolla. Stamens definite or indefinite, epigynous. Ovary adherent, 1 or many-celled; ovules 00, anatropal, frequently attached to parietal placentas; stigmas 3-6. Fruit dry or succulent, indehiscent, uni- or multilocular. Seeds numerous, exalbuminous; embryo straight, radicle remote from the hilum.—Floating or aquatic plants, with parallel-veined leaves, sometimes spiny. Chiefly found in Europe, Asia, and North America. The plants of this order are not remarkable for their properties. Some are mucilaginous and astringent. *Vallisneria spiralis* (fig. 228) is a dioecious plant, the male flowers of which, at the time of flowering, are said to be detached from the mud of the water in which they grow, and to float on the surface. At the same time the female flower develops a long spiral peduncle, by means of which it reaches the surface of the water, so as to allow the application of the pollen (¶ 496). The order has been divided into two sections:—1. *Vallisneriæ*, ovary 1-celled. 2. *Stratioteæ*, ovary many-celled. There are 12 known genera, according to Lindley, including 20 species. *Examples*—*Vallisneria*, *Udora*, *Anacharis*, *Stratiotes*, *Hydrocharis*.

1058. Order 186.—**Orchidaceæ**, the Orchis Family. (*Mono-epigyn.*) Flowers bisexual. Perianth adherent, herbaceous, or coloured, with a 6-partite limb (fig. 736 *pe*, *pi*), the segments being arranged in 2 rows; exterior row (fig. 735 *ce*), called the calyx (although Lindley says it is more properly the corolla, the true calyx or calyculus being usually abortive), consisting of 3 segments (rarely 2 by adhesion), the odd one of which is often next the axis by a twisting of the ovary; interior row (fig. 735 *ci*), called the corolla (regarded by Lindley as petaloid stamens), consisting usually of 3 segments (very rarely 1), the odd one of which is called the labellum or lip. This labellum (figs. 735, 736, 737 *l*) frequently differs from the other divisions of the perianth, assuming remarkable forms, being lobed, spurred at the base, or furnished with peculiar appendages, which are sometimes derived from the stigma. It is sometimes divided by contraction, so as to exhibit three distinct portions, the lowest being the hypochilium (ὕπῳ, under, and χεῖλος, lip); the middle, mesochilium (μέσος, middle); and the upper, the epichilium (ἐπί, upon or above). Stamens 3, epigynous, united in a central column along with the style; the two lateral stamens are usually abortive (figs 736 *ss*), the central one opposite the odd exterior segment being fertile (fig. 736 *e*); but at times the two lateral are fertile, and the central one is abortive; anthers 1-2-4-celled (fig. 738); pollen powdery or cohering in definite (fig. 744) or indefinite waxy masses (pollinia) (figs. 739, 743), which often adhere by a caudicle (fig. 743 *c*) to a gland connected with the beak (rostellum) of the stigma. This gland is sometimes naked, at other times in a sac or pouch (bursicula). Ovary adherent, 1-celled (fig. 740), composed of 6 carpels, of which 3 only are placentiferous (Lindley); style incor-


porated with the column (gynostemium, γυνή, pistil, and στήμιν, stamen); stigmas a viscid hollow space in front of the column (fig. 737 s),



735




737



736



738



739



740



741



742



744

743

Figs. 735-741.—Flower of *Spiranthes autumnalis*, to illustrate the natural order Orchidaceæ.

Fig. 735.—Flower after the ovary has twisted on itself, seen laterally. *o*, Ovary with the adherent perianth. *ce*, *ce*, Outer divisions of the perianth, called by some calyx, and by Lindley corolla. *ci*, *ci*, Inner divisions of the perianth, called by some the corolla, and considered by Lindley as petaloid stamens. *l*, The labellum or lip, being the lower of the three inner segments.

Fig. 736.—Diagram of the flower in the young state, before the twisting of the ovary has taken place. *a*, The axis of the spike of flowers. *pe*, *pe*, *pe*, Outer perianth. *pi*, *pi*, Two divisions of the inner perianth. *l*, Third division of the inner perianth, in this state placed next the axis. *e*, Fertile anther. *ss*, Two abortive anthers or staminodia. *o*, Ovary.

Fig. 737.—Summit of the flower cut vertically. *o*, Adherent ovary with parietal ovules, *g*, *l*, Labellum or lip, *s*, Stigma, *a*, Anther.

Fig. 738.—Anther separated. Its inner surface shown with its two cells.

Fig. 739.—Granular pollen-masses taken from the anther.

Fig. 740.—Horizontal section of the ovary, with three parietal placentas bearing numerous ovules.

Fig. 741.—A seed separated, with its external reticulated integument, *t*.

Fig. 742.—Embryo of *Aceras anthropophora* deprived of its integuments.

Fig. 743.—Pollen-masses (Pollinia) of *Orchis maculata*, with the grains united in little conical masses. *c*, Caudicle terminated by the retinaculum and glands.

Fig. 744.—The conical masses which the pollen-grains form by their cohesion.

Superior mycelium Spiranthes autumnalis - Pollen
1 bellid ovary - 3 parietal Placenta

communicating directly with the ovary by an open canal. The upper part of the united stigmas is often extended into a beak-like process (rostellum). Placentas 3, parietal (figs. 457, 740). Fruit a capsule, opening by 3 or 6 valves, rarely fleshy, and indehiscent. Seeds 00, very minute, with a loose reticulated spermoderm (fig. 741), exalbuminous; embryo solid, fleshy (fig. 742); large radicle next the hilum.—Perennial herbs or shrubs, with fibrous or tubercular roots (fig. 121), either no stem or a pseudo-bulb, entire parallel-veined often sheathing leaves, and generally showy, attractive flowers. Sometimes buds are produced on the margins of the leaves (fig. 214). They are natives of almost all parts of the world, but they abound in moist tropical regions. They are not found in the arctic regions, nor in very dry climates. Some are terrestrial, and others are epiphytic. The former are commonly seen in temperate climates, the latter in warm regions. *Disa grandiflora* is found on Table Mountain at an elevation of 3,582 feet. The only known locality for it is in a marshy hollow, near the eastern extremity of the summit, where it is abundant among rushes on the margins of small pools and streamlets in a black boggy soil. Two rare species of *Disa* are also seen there, *D. ferruginea* and *tenuifolia*. *Oncidium nubigenum* is found on the Andes, near Quito, at an elevation of 14,000 feet above the level of the sea. *Epidendrum frigidum* occurs in Columbia at an elevation of 12,000 to 13,000 feet (mean temperature 46°), and is covered with a sort of varnish. Lindley, who is our best authority on Orchids, has divided them into various tribes. He enumerates 396 genera, including about 3000 species; of these, 16 genera and 40 species are British.

1059. The plants of this order are well distinguished by the peculiar forms of their flowers, their remarkable lip, gynandrous stamens, and pollen-masses. Their flowers often resemble insects, as butterflies, moths, bees, flies, and spiders; or birds, as doves and eagles; or reptiles, as snakes, lizards, and frogs. The colours and spots on the perianth sometimes give the appearance of the skins of quadrupeds, as the leopard and tiger. These resemblances are often indicated in the generic and specific names. The labellum, in some instances, displays peculiar irritability (¶ 665).

1060. Mucilaginous properties occur in many of the plants of this order. Some are aromatic and fragrant; others are antispasmodic and tonic. The tuberous roots of some yield a nutritious substance called Salep, which consists chiefly of bassorin, some soluble gum, and a little starch. The orchid yielding it is not well known; some say that it is *Orchis mascula*, others *O. papilionacea*, *Morio* and *militaris*, others *Eulophia vera* and *campestris*. Salep forms an article of diet fitted for convalescents, when boiled with water or milk. The roots of *Aplectrum hyemale* contain a very glutinous matter, and hence the plant is called in America Putty-wort. The fleshy pod-like fruit of

*
in
Chakish
Diction
p. 358

Salep considered in the East - and called
شکر صندل

Vanilla planifolia, *claviculata*, and other species, constitutes the substance called Vanilla, which is used in confectionery, and in flavouring chocolate. It contains an oil and much benzoic acid. A blue colouring matter has been found in some of the orchids. The odour of many of them is very fragrant; sometimes it is oppressive; at other times, as in *Malachadenia clavata*, it is very fetid, resembling carrion.

1061. Order 187.—**Zingiberaceæ**, or **Scitamineæ**, the Ginger Family. (*Mono-epigyn.*) Perianth superior, in 2 whorls; outer (calyx) tubular, 3-lobed, short; inner (corolla) tubular, elongated, 3-parted, segments nearly equal. Stamens in 2 whorls; outer sterile, petaloid, having the appearance of a 3-parted corolline whorl, with the intermediate segment (labellum) larger than the rest, and often 3-lobed, sometimes the lateral segments are inconspicuous or nearly abortive; inner stamens 3, the two lateral being abortive, the intermediate one opposite the labellum, fertile; filament not petaloid, often prolonged beyond the anther; anther 2-celled, dehiscing longitudinally. Ovary 3-celled, or imperfectly so; ovules several, anatropal, attached to a placenta in the axis; style filiform; stigma dilated, hollow. Fruit usually a 3-celled capsule, sometimes baccate. Seeds roundish or angular, sometimes with an arillus; embryo enclosed in a vitellus (the remains of the embryo-sac), surrounded by farinaceous albumen, which is deficient near the hilum.—Herbs with a creeping rhizome, and simple sheathing leaves, having parallel veins proceeding from the midrib to the margin. The flowers arise from membranous spathaceous bracts. Natives of tropical countries. By far the greater number inhabit various parts of the East Indies; some are found in Africa, and a few in America. They form a part of the singular Flora of Japan. Lindley mentions 29 genera, and 247 species. *Examples*—Zingiber, Curcuma, Amomum, Hedychium, Renealmia.

1062. Plants often with showy flowers, having aromatic stimulant properties, which reside chiefly in their rhizome or root, and in their seeds. Some yield starchy matter. The rhizome of *Zingiber officinale* (*Amomum Zingiber*), constitutes the Ginger of commerce, which is imported from the East and West Indies. In the young state the rhizomes are fleshy and slightly aromatic, and they are then used as preserves; while in a more advanced state, the aroma is fully developed, their texture is more woody, and they are then fit for ordinary ginger. When dried, after immersion in hot water, they form Black ginger; when simply dried in the sun, after being cleaned, they receive the name of White ginger. The rhizome contains an acrid resin and volatile oil, starch and gum. It is used as a tonic and carminative, in the form of powder, syrup, and tincture. *Curcuma longa*, a native of Eastern Asia, furnishes Turmeric. This consists of the branches of the rhizome, or root-stock. Its powder is lemon-yellow,

and it is used as a dyestuff. It contains starch, an acrid volatile oil, and a yellow colouring matter called Curcumin. It is employed medicinally as an aromatic carminative, and, as a condiment, it enters into the composition of curry-powder. The root-stocks of *Alpinia racemosa*, and *A. Galanga*, Galangale, and many other plants of the order, have the same properties as ginger. They have been used by the Indian doctors in cases of dyspepsia and in the treatment of coughs. Various species of *Amomum*, *Elettaria*, and *Renealmia*, appear to furnish the Cardamoms of the shops, which consist of the oval trivalvular capsules containing the seeds. The following are some of the sources whence Cardamoms are procured :—

1. *Amomum Cardamomum*, a Java plant, supplies the round Cardamoms.
2. *A. angustifolium* furnishes the large Madagascar Cardamoms, and also supplies some of the seeds called Grains of Paradise.
3. *A. maximum* produces the Java Cardamoms of the London market.
4. *A. repens* (*Renealmia* or *Elettaria Cardamomum*) is the source of Malabar Cardamoms.
5. *Elettaria major* is said to yield Ceylon Cardamoms.

The seeds of these plants are used as aromatic tonics and carminatives. Their active ingredient is a pungent volatile oil. Grains of Paradise are the seeds of *Amomum Melegueta*, Melegueta Pepper, and have the same properties as Cardamoms. East Indian Arrow-root is procured in part from *Curcuma angustifolia*, and a similar kind of starch is yielded by *Curcuma Zerumbet*, *C. leucorrhiza*, and *Alpinia Galanga*.

1063. Order 188.—**Marantaceæ** or **Cannaceæ**, the Arrow-root Family. (*Mono-epigyn.*) Perianth superior, in 2 whorls; outer (calyx) 3-lobed, short; inner (corolla) tubular, elongated, 3-parted, segments nearly equal. Stamens in 2 whorls; outer sterile, petaloid, irregular, resembling a tubular trifold corolla, with one of the lateral segments different from the others; inner petaloid, 2 sterile, and 1 lateral fertile; filament of the latter petaloid, entire, or 2-lobed; anther on the margin of the filament, 1-celled, dehiscing longitudinally. Ovary 3-celled, rarely 1-celled; ovules solitary and erect, or numerous and attached to the axis; style petaloid or swollen; stigma either the naked apex of the style, or hollow, hooded, and incurved. Fruit a 3-celled capsule, or baccate, 1-celled and 1-seeded. Seeds round, without arillus; embryo straight, in hard, somewhat floury albumen, without a vitellus; radicle lying against the hilum (fig. 527).—Herbaceous plants, with tuberous rhizomes, and leaves and flowers, similar to those of the Ginger Family. They are natives of tropical America, and Africa; several are found in India; none are known in a wild state beyond the tropics. Lindley enumerates 6 genera, including 160 species. *Examples*—*Maranta*, *Canna*, *Phrynium*.

1064. The plants of the order contain much starch in the rhizomes and roots. They are destitute of aroma. Arrow-root is the produce

of the tuberous rhizomata of *Maranta arundinacea* and *M. indica*. The best West Indian arrow-root comes from Bermuda. Its globules are much smaller and less glistening than those of *tous-les-mois* or potato starch. Amylaceous matter of a similar kind is produced from other species of *Maranta*, as well as from species of *Canna*. *Tous-les-mois* is the produce of *Canna coccinea*, *C. Achiras*, *C. edulis*, &c. The seeds of Cannas are round and black, and are commonly known under the name of Indian shot. They have been used as a substitute for coffee. *Calathea zebrina*, Zebra plant, is so called from the peculiar variegation of its leaves, which have a velvety aspect. Barnéoud states that the two outer verticils of the flowers in *Cannas* are always developed, one after the other, precisely like the calyx and corolla; while the verticil, sometimes called petals, is really metamorphosed stamens, and hence its irregular aspect.

1065. Order 189.—**Musaceæ**, the Banana Family. (*Mono-epigyn*.) Perianth 6-cleft, adherent, petaloid, in 2 whorls, more or less irregular. Stamens 6, inserted on the middle of the segments of the perianth, some usually abortive; anthers linear, dithecal, introrse, often with a membranous petaloid crest. Ovary inferior, 3-celled; ovules numerous, anatropal; style simple; stigma usually 3-lobed. Fruit either a 3-celled capsule, with loculicidal dehiscence, or succulent and indehiscent. Seeds sometimes surrounded by hairs; testa usually crustaceous; embryo erect, in the axis of mealy albumen; radicle touching the hilum.—Plants without true aerial stems, or nearly so, having shoots proceeding from subterranean root-stocks, which form spurious stems, composed of the sheathing leaf-stalks. Veins in the limb of the leaf parallel, and proceeding in a curved manner from the midrib to the margin (fig. 135). Flowers bursting through spathas. Natives of warm and tropical regions. There are 5 known genera, and 21 species. *Examples*—*Musa*, *Strelitzia*, *Ravenala*.

1066. The plants of this order furnish a large supply of nutritious fruit, and their leaves afford valuable fibres. Spiral vessels abound in them. *Musa sapientum* and *Cavendishii* furnish different kinds of Banana, while *M. paradisiaca* yields the Plantain. These fruits in their ripe state contain much starchy matter. From their spurious stems, the fibres of the spiral vessels may be pulled out in such quantity as to be used for tinder. The ribbon-like fibre in these vessels is composed of several threads united together (pleiotrachæ) (fig. 51). The produce of the Banana is of great value to the inhabitants of warm countries. The same extent of ground which in wheat would only maintain two persons, will yield sustenance under the Banana to fifty. *Musa textilis* yields a kind of woody fibre, which is used in India in the manufacture of fine muslins. Manilla Hemp is the produce of *Musa textilis*. The woody tissues of many species of *Musa* is used for manufacture in warm climates. The young shoots of the

Banana are used as a culinary vegetable. The juice of the fruit and the lymph of the stem of *Musa* are slightly astringent and diaphoretic. *Urania speciosa* or *Ravenala* is the Water-tree of the Dutch, so called on account of the great quantity of water which flows from its stem or leaf-stalk when cut across. The juice of the fruit of *Urania* is used for dyeing.

1067. Order 190.—**Iridaceæ**, the Iris Family. (*Mono-epigyn.*) Perianth adherent, 6-parted, coloured, in 2, often unequal whorls (figs. 745, 746). Stamens 3, epigynous, opposite the outer segments of the perianth (figs. 745, 746 *ee*); filaments distinct or monadelphous; anthers 2-celled, extrorse. Ovary inferior (fig. 746 *o*), 3-celled; ovules numerous (fig. 746 *g*), anatropal; style 1; stigmas 3, often petaloid (fig. 746 *s*), sometimes bilabiate. Fruit a 3-celled, 3-valved capsule,



with loculicidal dehiscence (fig. 448). Seeds numerous; embryo enclosed in horny or fleshy albumen; radicle next the hilum (fig. 747). —Herbs, rarely undershrubs, with rhizomes or underground corms, having their leaves often equitant or distichous, and their flowers spathaceous. Natives chiefly of warm and temperate regions. They

Figs. 745-747.—Organs of fructification of *Iris Germanica*, to illustrate the natural order Iridaceae.

Fig. 745.—Diagram of the flower, showing six divisions of the perianth in two verticils, three exserted stamens, and the 3-celled capsule with numerous ovules. *a*, Position of the axis of inflorescence.

Fig. 746.—Vertical section of the flower. *ce*, Outer divisions of the coloured perianth. *ci*, Inner divisions of the perianth. *t*, Tube of the perianth, above the part which is adherent to the ovary. *o*, Inferior 3-celled ovary. *g*, Numerous anatropal ovules. *ee*, Stamens. *ss*, Petaloid stigmas.

Fig. 747.—Seed separated and cut longitudinally. *t*, Integuments (spermoderm). *p*, Perisperm. *e*, Embryo enclosed in the perisperm. *m*, Micropyle (foramen).

abound at the Cape of Good Hope. The Crocus occurs only in Europe and Asia. There are 53 known genera, and 550 species. *Examples*—*Iris*, *Sisyrinchium*, *Witsenia*, *Gladiolus*, *Ixia*, *Crocus*.

1068. Some of the plants have fragrant and stimulant, and some acrid, rhizomes and corms ; others yield dyes. The root-stock of *Iris Florentina* yields orris-root, which has a pleasant odour like violets, and an acrid taste, depending on the presence of a volatile oil. It is imported from Florence and Leghorn. Orris-root is used chiefly for giving a pleasant odour to the breath, and in perfumery and tooth-powder. Orris-root starch is used for hair-powder. *Crocus sativus*, the זרע of the Old Testament, furnishes the colouring material called Saffron. It consists of the stigmata, which have a fine deep-orange colour. These stigmata are either dried in the loose state, forming Hay Saffron, or compressed into masses, constituting Cake Saffron. The yellow colouring ingredient is Polychroite, which possesses the property of being totally destroyed by the action of the solar rays, and of forming blue tints when treated with sulphuric and nitric acid. Saffron contains an active volatile oil, and it has been used in the form of tincture and syrup, as an emmenagogue and antispasmodic. The stigmata of *Crocus autumnalis* and *C. odoratus*, also supply saffron. The roasted seeds of *Iris pseudacorus* have been used as a substitute for coffee.

1069. Order 191.—**Burmanniaceæ**, the Burmannia Family. (*Monopigyn.*) Perianth coloured, tubular, 6-cleft, the three outer segments (calyx) often keeled at the back, the three inner (petals) minute. Stamens 3, inserted in the tube of the perianth, opposite its inner segments, sometimes with 3 alternating sterile filaments ; anthers dithecal, opening transversely, with a fleshy connective. Ovary inferior, either 1- or 3-celled, in the latter case the cells opposite the outer segments of the perianth ; ovules 00 ; style simple ; stigmas 3. Fruit a 1-3-celled, 3-valved capsule, crowned by the persistent perianth. Seeds 00, minute, striated.—Herbs, with radical leaves and bisexual flowers. Natives of moist grassy places in tropical regions. They have no properties of importance. There are about 10 known genera, and 35 species. *Examples*—*Burmanna*, *Apteria*, *Apostasia* ?

1070. Order 192.—**Hæmodoraceæ**, the Blood-root Family. (*Monopigyn.*) Perianth petaloid, more or less woolly, 6-cleft. Stamens inserted on the perianth, either 3, and opposite the inner segments of the perianth, or 6 ; anthers introrse. Ovary adherent to the tube of the perianth, usually 3-celled, rarely 1-celled ; ovules 1-2 or numerous ; style simple ; stigma undivided. Fruit a 3-valved capsule, sometimes indehiscent. Seeds either definite or 00, sometimes peltate ; embryo in cartilaginous albumen.—Herbs with fibrous roots, equitant distichous leaves, and bisexual flowers. They are found in various parts of the world, more especially in the warm parts of South America, at the

Cape of Good Hope, as well as in North America and Australia. Lindley mentions 13 genera, and 50 species. *Examples*—*Hæmodorum*, *Anigosanthus*, *Velozia*, *Barbacenia*.

1071. The plants receive the name of Blood-root, from the red colour of their roots, which are used for dyeing. *Vellozias*, Tree Lilies, give a decided feature to the vegetation of the mountains of Minas Geraes in Brazil. Their trunks are covered by the withered remains of the leaves, and their branches are dichotomous, and bear tufts of leaves at the extremities. The outer surface of their stems is covered thickly with numerous adpressed rootlets.

1072. Order 193.—**Amaryllidaceæ**, the Amaryllis Family. (*Monoepigym.*) (Fig. 251). Perianth petaloid, regular, 6-cleft, the outer segments overlapping the inner. Stamens 6, inserted in the perianth, sometimes cohering by the dilated bases, and forming a kind of cup; occasionally there are additional sterile stamens, which sometimes form a corona above the tube of the perianth; anthers introrse. Ovary inferior, 3-celled; ovules 00, anatropal; style 1; stigma 3-lobed. Fruit either a 3-celled, 3-valved capsule, with loculicidal dehiscence, or baccate. Seed with a thin or thick, or black and brittle spermoderm; albumen fleshy; embryo nearly straight; radicle next the hilum.—Usually bulbous plants, sometimes with fibrous roots; leaves ensiform, with parallel veins; flowers spathaceous; stem sometimes woody and tall. Natives chiefly of the Cape of Good Hope, but species are found in Europe, East and West Indies, South America, and Australia. Lindley enumerates 68 genera, and 400 species, and he divides them into 4 tribes:—1. *Amaryllææ*, bulbs, flowers without a corona. 2. *Narcisseæ*, bulbs, flowers with a corona. 3. *Alströmeriææ*, fibrous roots, outer segments of the perianth different in form from the inner. 4. *Agavææ*, fibrous roots, both segments of the perianth alike. *Examples*—*Amaryllis*, *Galanthus*, *Crinum*, *Narcissus*, *Alströmeria*, *Agave*.

1073. The bulbs of many plants of this order have narcotic poisonous qualities. Some of them act as emetics, others are used in the preparation of a kind of intoxicating spirit. The tough fibres of some are used for flax. The root of *Hæmanthus toxicarius* is poisonous. The flowers of the Daffodil (*Narcissus pseudo-narcissus*) are also said to be poisonous. The fibres of *Agave Americana*, American Aloe, yield Pita flax. This plant does not flower often, but when flowering begins, it proceeds with great rapidity and vigour. Its roots are sometimes used to adulterate Sarza. Its juice is fermented so as to form an intoxicating beverage. *Agave Saponaria* is used in Mexico for washing. The bulbs of *Narcissus poeticus*, *N. Jonquilla*, *N. odoratus*, *N. pseudo-narcissus*, *N. Tazetta*, and of some species of *Pancratium* are emetic. The Guernsey Lily is also reputed poisonous. Some *Alströmerias* are diuretic. In *Alströmerias* (fig. 251), the leaves are twisted, so that what

Amaryllidaceæ - Superior Hemodorum - Narcissus - Tree Lilies - Flowers -

should be the upper surface becomes the lower. In *Narcissus* the corona or crown of abortive filaments projects beyond the flower; while in *Pancratium* the dilated filaments of the fertile stamens unite together, and are included within the perianth. Many ornamental garden plants belong to the order.

1074. Order 194.—**Hypoxidaceæ**, the Hypoxis Family. (*Monoepigyn.*) Perianth petaloid, superior, usually 6-parted, regular. Stamens 6, inserted into the base of the segments of the perianth, filaments distinct; anthers introrse. Ovary inferior, 3-celled; ovules numerous, amphitropal; style simple; stigma 3-lobed. Fruit indehiscent, sometimes succulent, 1-2-3-celled. Seeds 00, with a lateral hilum and a beaked caruncle; testa black, and crustaceous; embryo straight, in the axis of fleshy albumen; radicle remote from the hilum.—Herbaceous and usually stemless plants, with tuberous and fibrous roots, radical plaited leaves, and simple or branched scapes. Natives of warm countries. Some have bitter roots, others have edible tubers. There are 4 known genera, including 60 species. *Examples*—Hypoxis, Curculigo.

1075. Order 195.—**Bromeliaceæ**, the Pine-apple Family. (*Mono-perigyn.*) Perianth tubular, 6-divided, in 2 verticils; outer whorl (calyx) persistent, more or less adherent to the ovary; inner petaloid, marcescent or deciduous, with imbricated æstivation. Stamens 6, inserted into the base of the segments of the perianth; anthers introrse. Ovary either free or partially adherent, 3-celled; ovules 00, anatropal; style single; stigma 3-lobed or entire, often twisted. Fruit capsular or succulent (figs. 256, 474), 3-celled. Seeds 00; embryo minute, curved or straight, lying in the base of mealy albumen; radicle next the hilum.—Stemless or short-stemmed plants, with rigid, channelled leaves, which are often spiny at the margin, and are covered with scurfy matter. Natives of the warm parts of America chiefly. There are 23 genera, according to Lindley, and 170 species. *Examples*—Bromelia, Ananassa, Tillandsia, Bonaparteia.

1076. The plants of this order are all more or less epiphytic, being able to grow without any direct attachment to the soil. In hothouses they are frequently kept suspended in moistened moss. Some of the *Tillandsias* are hung from balconies in South America as air-plants. *Tillandsia usneoides* has the appearance of the Beard-moss (a lichen commonly seen on trees in Britain), and it is used for stuffing cushions, &c. The plant has been called Tree-beard or Black Moss. The leaves of *Tillandsias* frequently contain much water in their hollowed-out bases. The fruit of *Ananassa sativa* is well known as the Pine-apple or Ananas (fig. 256). It is an anthocarpous fruit, consisting of numerous flowers and bracts united together, and becoming succulent. The fruit is more or less acid in the wild state, but when cultivated it becomes sweet and highly aromatic. The fibres of the

leaves are used in the preparation of fine muslins. The woody fibres of many *Bromelias* are used in manufactures. *Bromelia Pinguin* is used as a vermifuge in the West Indies. Its ovaries do not combine into one mass, and therefore illustrate well the formation of the Pine-apple. The crown of the Pine-apple consists of the leaves arising from the prolonged axis (fig. 474 f).

b. Perianth free, Ovary superior, Flowers usually hermaphrodite.

1077. Order 196.—**Liliaceæ**, the Lily Family. (*Mono-perig.* and *Mono-hypog.*) Flowers usually bisexual. Perianth coloured, in 2 rows, regular, with 6 divisions (figs. 259, 260, 748, 749). Stamens 6 (fig. 538), perigynous, inserted into the segments of the perianth (figs. 259, 748, 749); anthers introrse (fig. 750 e). Ovary free, 3-celled




748



749



750



751

(fig. 749); ovules 00; style 1; stigma simple or 3-lobed (figs. 259, 260, 750 s). Fruit 3-celled, either succulent or dry and capsular. Seeds numerous, packed one above the other in 1 or 2 rows (fig. 750); embryo in the axis of fleshy albumen (fig. 751).—Herbs, shrubs, or trees, with bulbs (figs. 207–209), or tubers, or arborescent stems, or rhizomes (fig. 90); leaves not articulated, usually narrow, with parallel veins. They are found both in temperate and tropical climates. In warm regions some of them are arborescent, as in the case of *Dracenas*; others are very succulent, as species of *Aloe*. The order has not been sufficiently defined, and there are still many differences of opinion as to its limits. Under it are included by some the following suborders:—

1. Tulipeæ, Tulip tribe: bulbous plants, segments of perianth scarcely adherent in a tube, testa pale and soft.
2. Hermerocallidæ, Day-lily tribe: bulbous plants, with a tubular perianth, testa pale and soft.

Figs. 748–751.—Organs of fructification of *Scilla autumnalis*, to illustrate the natural order Liliaceæ.

Fig. 748.—Flowers seen from above. ce, Outer verticil of the perianth (calyx). ci, Inner verticil of the perianth (corolla).

Fig. 749.—Diagram of the flower, showing three outer and three inner leaves of the perianth, six alternating stamens in two rows, and three carpels of the ovary with the ovules.

Fig. 750.—Vertical section of the flower. c c, Perianth. e, Stamens. o, Ovary. s, Style and stigma. g, Ovules attached to a placenta in the axis.

Fig. 751.—Seed separated and cut lengthwise. t, Integument. p, Perisperm. e, Embryo.

3. Scilleæ or Allicæ, the Squill or Onion tribe: bulbous (figs. 207-209), with the testa black and brittle.
4. Anthericæ or Asphodelæ, Asphodel tribe: not bulbous, roots fascicled (fig. 120) or fibrous, leaves not coriaceous nor permanent.
5. Convallariæ, Lily of the Valley tribe: stem developed as a rhizome or tuber (fig. 90).
6. Asparagæ, Asparagus tribe: stem usually fully developed, arborescent, in some cases branched, leaves often coriaceous and permanent.
7. Aloinæ, Aloes tribe: stem usually developed, arborescent, leaves succulent.
8. Aphyllanthæ, Grass-tree tribe: having a rush-like habit and membranous imbricated bracts.

Lindley adds to these the tribes Conantheræ, Wachendorfeæ, Aspidistreæ, and Ophiopogonæ. He enumerates 133 genera, including 1,200 species. *Examples*—*Tulipa* *Lilium*; *Hemerocallis*, *Phormium*; *Anthericum*, *Asphodelus*; *Convallaria*; *Asparagus*, *Dracæna*; *Aloe*; *Aphyllanthes*, *Xanthorrhæa*; *Conanthera*; *Wachendorfia*; *Aspidistra*; *Ophiopogon*.

1078. Many of the plants of this order are showy garden flowers, such as Tulips, Lilies, Fritillaries, Day-lilies, Tuberoses (*Polianthes*), and Dog-tooth-violets (*Erythronium*). Some of them are used medicinally as purgatives, stimulants, emetics, and diaphoretics. Some yield valuable fibres, others supply resinous matter. The bulb of *Scilla* or *Squilla* (*Urginea*) *maritima* supplies the officinal squill. The plant grows on the sandy coasts of the Mediterranean. Its bulbs vary in weight from half a pound to four or five pounds. In their fresh state they are very acrid. They contain a bitter crystalline principle called Scillitina. Squill is used medicinally in the form of powder, vinegar, syrup, and tincture, as an emetic, diaphoretic, expectorant, and diuretic. The drug called Aloes is the inspissated juice of the leaves of various species of *Aloe*, as *A. spicata*, *vulgaris*, *socotrina*, *indica*, *rube-scens*, *arabica*, *linguæformis*, and *Commelini*. It is imported under the names of Socotrine, East Indian or Hepatic, Barbadoes, Cape and Caballine Aloes. It contains a substance called Aloesin, which some regard as its active principle. Aloes is used medicinally as a cathartic, acting chiefly on the large intestines and on the rectum. The bulb of *Allium sativum*, Garlic, is used as an irritant, stimulant, and diuretic. It is the שום of the Bible, the σκόροδον of the Greeks. The bulb of *Allium Cepa*, the Onion, the נֶזֶק of the Bible, is used in the same way as garlic, and so is the bulb of *Allium Porrum*, the Leek, the קֶרֶן of the Bible (figs. 207, 208). Besides the Onion and Leek, several species of *Allium*, under the name of Chive (*A. Schænoprasum*), Shallot (*A. ascalonicum*), and Rocambole (*A. Scorodoprasum*), are used as articles of diet. These plants contain free phosphoric acid, and a sulphuretted oil, which is in a great measure dissipated by boiling or roasting. In the Oregon and Missouri districts of North America, the bulbs of *Gamassia esculenta*, Gamass or Squamash, are also employed in a

similar manner. It is called by the Indians Biscuit-root. The turios or young shoots sent up from the underground stem of *Asparagus officinalis* (fig. 110), are the parts employed in cooking. The bulbs of species of *Lilium*, found in the east of Siberia, are eaten like potatoes. Fibres are procured from *Phormium tenax*, New Zealand Flax, and from the species of *Yucca*, Adam's Needle (fig. 234). *Dracæna Draco*, and other species, yield an astringent resin called Dragon's-blood. The *Dracænas* often branch in a dichotomous manner, and attain a large size. The Grass-tree of New South Wales, *Xanthorrhæa hastilis*, gives a peculiar feature to the vegetation of that country. It yields a yellow gum-like substance. The base of the inner leaves of some Grass-trees is used as food. Some of the Lilies have bulbils or bulb-lets in the axils of their leaves (fig. 213). In the Crown-Imperial there is a nectariferous depression in the base of the segments of the perianth (fig. 302). *Lilium chalcedonicum* is said to be the Lilies of the field, *τα κρίνα του ἀγρου*, mentioned in Scripture. *Hyacinthus orientalis* is the common cultivated Hyacinth, of which the Haarlem florists had at one time upwards of 2,000 varieties. The mania for Tulip bulbs was at one time carried to a great extent, and the price given for approved kinds was enormous. Many hundred varieties of tulips are known.

1079. Order 197.—**Melanthaceæ**, the Colchicum Family. (*Mono-perigyn.*) Perianth petaloid, in 6 pieces, which are sometimes slightly coherent, usually involute in æstivation. Stamens 6; anthers extrorse. Ovary 3-celled; ovules numerous; style 3-parted; stigmas 3, undivided. Fruit a 3-celled capsule, with septicidal or loculicidal dehiscence. Seeds with a membranous spermoderm; albumen dense, fleshy; embryo very minute.—Plants with bulbs, tubers (fig. 93), or fibrous roots, having parallel-veined leaves, sheathing at the base. The flowers are sometimes polygamous. They are natives of various parts of the globe, but are most abundant in northern countries. The order has been divided into three suborders:—1. Veratreæ or Melantheæ, rhizome fibrous, dehiscence of capsule septicidal, flowers frequently unisexual. 2. Colchicææ, rhizome bulbous, dehiscence septicidal. 3. Uvulariææ, rhizome bulbous or fibrous, dehiscence loculicidal. There are 30 known genera, and 130 species. *Examples*—*Melanthium*, *Veratrum*, *Tofieldia*; *Colchicum*; *Uvularia*.

1080. The plants of the order have in general poisonous properties. Many are acrid, purgative, and emetic, and some are narcotic. Among the medicinal plants of the order, the most important is *Colchicum autumnale*, Meadow Saffron, or Autumn Crocus. The bulb or corm (fig. 93) and the seeds are the officinal parts. They contain a peculiar alkaloid called Colchicia, which seems to be analogous to Veratria. Colchicum in large doses acts as a narcotico-acrid poison. In medicinal doses, in the form of extract, vinegar, tincture, and wine, it is used in the cure of gout and rheumatism. It is sedative, cathartic,

*Separate the roots from Lilium which is from
Lilium Chalcidicum*

and diuretic. The rhizome of *Veratrum album*, the White Hellebore of the Greeks, is an irritant narcotic poison, its properties being due to the presence of an alkaloid called Veratria. It has been used as an emetic and purgative, especially in mania, and it has been administered as a remedy for gout. Cevadilla is the fruit of *Helonias* or *Asagrea officinalis*, as well as of *Veratrum Sabadilla*. The fruit and seeds contain the alkaloid veratria in combination with a peculiar fatty acid called cevadic or sabadillic acid. Cevadilla is used in cases of neuralgia and rheumatism.

1081. Order 198.—**Gilliesiaceæ**, the Gilliesia Family. (*Mono-perigyn.*) Perianth 6-parted, sometimes 5-parted by cohesion of two of the pieces, in a double row; the outer, petaloid and herbaceous; the inner, smaller, and more coloured; æstivation twisted. Stamens in a double series; outer whorl sterile, in the form of a 6-toothed urceolate body, or of scale-like bodies, one of which forms a sort of labellum; inner whorl of six stamens, of which three are sometimes sterile. Ovary superior, 3-celled; style 1; stigma simple. Fruit a 3-celled, 3-valved capsule, with loculicidal dehiscence. Seeds numerous, attached to the axis; spermoderm black and brittle; embryo curved in the midst of fleshy albumen.—Herbs with tunicated bulbs, grass-like leaves, and umbellate spathaceous flowers. Natives of Chili. Their properties unknown. The description of the flower is in accordance with Arnott's view, and differs from that of Lindley, who considers the perianth as bracts, and the outer verticil of stamens as the perianth. There are 2 genera and 5 species. *Examples*—Gilliesia, Miersia.

1082. Order 199.—**Pontederiaceæ**, the Pontederia Family. (*Mono-perigyn.*) Perianth tubular, coloured, 6-parted, more or less irregular; æstivation circinate. Stamens 3-6, perigynous; anthers introrse. Ovary free, or slightly adherent, 3-celled; ovules numerous, anatropal; style 1; stigma simple. Fruit a 3-celled, 3-valved capsule, with loculicidal dehiscence. Seeds 00, attached to a central axis; testa membranous; hilum small; embryo straight, in the axis of somewhat mealy albumen; radicle next the hilum. Aquatic or marsh plants with sheathing, parallel-veined leaves, which are sometimes cordate or sagittate, and have inflated petioles. The flowers are spathaceous. They are natives of North and South America, East Indies, and Africa. Their properties are unimportant. There are 6 genera, according to Lindley, and 30 species. *Examples*—Pontederia, Leptanthus.

1082. Order 200.—**Xyridaceæ**, the Xyris Family. (*Mono-perigyn.*) Perianth 6-parted, in two verticils; the outer glumaceous, the inner petaloid. Stamens 6, 3 fertile, inserted into the claws of the inner perianth; anthers extrorse. Ovary single, 1-celled; ovules 00, orthotropal, attached to parietal placentas; style trifid; stigmas obtuse, multifid or undivided. Fruit a 1-celled, 3-valved capsule. Seeds numerous; embryo on the outside of mealy albumen, remote from

the hilum.—Herbs having a sedge-like aspect, with radical leaves, equitant and sheathing at the base, and scaly heads of flowers. Natives chiefly of tropical regions, having no important properties. There are about 6 genera, and 70 species. *Examples*—*Xyris*, *Abolboda*, *Philydrum*?

1084. Order 201.—**Juncaceæ**, the Rush Family. (*Mono-hypoperigyn.*) Perianth 6-parted, more or less glumaceous. Stamens 6, inserted into the base of the segments, sometimes 3, and opposite the outer segments; anthers 2-celled, introrse. Ovary 1-3-celled; ovules 1, 3, or many in each cell, anatropal; style 1; stigmas generally 3, sometimes 1. Fruit a 3-valved capsule, with loculicidal dehiscence, sometimes indehiscent. Seeds with the testa neither black nor crustaceous; embryo very minute, near the hilum, within fleshy or cartilaginous albumen.—Herbs with fasciculated or fibrous roots, hollow, grooved, or flat leaves, with parallel veins. They are natives chiefly of the colder regions of the globe. Many species of *Juncus* are used for making the bottoms of chairs, mats, &c., and the central cellular tissue forms the wicks of rushlights. There are 14 known genera, and upwards of 200 species. *Examples*—*Juncus*, *Luzula*, *Narthecium*, *Astelia*.

1085. Order 202.—**Palmæ**, the Palm Tribe. (*Mono-perigyn.*) Flowers bisexual or unisexual, or polygamous. Perianth 6-parted in a double row (fig. 757); 3 outer (calyx) fleshy, or leathery and persistent (figs. 752 *c e*, 756 *c*), 3 inner (corolla) often larger (fig. 752 *c i*), and sometimes deeply connate. Stamens 6 (figs. 753, 757), rarely 3, sometimes 00, inserted into the base of the perianth. Ovary free, 1-3-celled, usually composed of 3 carpels, which are more or less completely united (fig. 755); ovules 1-3. Fruit drupaceous, or nut-like (fig. 756), or baccate, often with a fibrous covering. Seed with cartilaginous or horny albumen (fig. 520), which is often ruminant (fig. 497), or furnished with a central or lateral cavity; embryo small, cylindrical, or flat, in a cavity of the albumen, remote from the hilum (figs. 497, 520).—Arborescent plants (fig. 115, 1), with simple, rarely branched trunks, marked with the scars of the leaves, which are terminal, pinnate, or fan-shaped, with plicate veneration, and parallel simple veins, and often spiny petioles. Flowers on a terminal, often branched spadix, enclosed in a 1- or many-valved spathe. Natives of tropical regions chiefly, and imparting to them much of their botanical physiognomy. Most of them have unbranched stems, attaining sometimes a height of 190 feet, and sending out clusters of large leaves, from the axil of which bunches of flowers proceed. Although the flowers are small, still the inflorescence, taken collectively, has often a most imposing aspect. Humboldt describes their effect on the landscape in glowing colours, and Martius has illustrated the order by splendid delineations. Linnæus called them the Princes of the Vegetable

Kingdom. Lindley states that there are 73 known genera, and 400 species; but this estimate probably falls short of the total amount, for much still remains to be done in the elucidation of the species.



Martius estimates the known species at 582, of which 91 have fan-shaped leaves. They have been divided by him into various tribes, depending chiefly on the nature of the ovary, ovules, and fruit; and sections are formed according as the leaves are pinnate or flabelliform, and the stems are spiny or not. The following are the tribes:—1. Arecinæ, the Betel-nut tribe. 2. Lepidocaryinæ, the Sago tribe. 3. Borassinæ, the Palmyra Palm tribe. 4. Coryphinæ, the Talipot and Date tribe. 5. Cocoinæ, the Coco-nut tribe. *Examples*—*Areca*, *Euterpe*, *Caryota*; *Lepidocaryum*, *Calamus*, *Sagus*; *Borassus*, *Lodoicea*; *Corypha*, *Livistona*, *Phœnix*; *Cocos*, *Elais*, *Acrocomia*; *Phytelphas*.

1086. The properties of the plants of this order are very various. In the countries in which they grow, they are used for supplying food,

Figs. 752-756.—Organs of fructification of *Areca Catechu*, Betel-nut Palm, to illustrate the natural order Palmæ.

Fig. 752.—Unexpanded flower-bud. *ce*, Outer division of the perianth. *ci*, Inner perianth.

Fig. 753.—Diagram of the flower, showing the two verticils of the perianth, the six stamens, and three abortive carpels. *a*, The position of the axis of inflorescence in reference to the flower.

Fig. 754.—A flower deprived of its perianth, in which the stamens, *e*, are partly abortive, while the ovary, *o*, is developed. *s*, Stigma.

Fig. 755.—Diagram of the last flower, showing the double perianth, the partially abortive stamens, and the 3-celled ovary. *a*, Axis of inflorescence.

Fig. 756.—Fruit, *f*, surrounded at its base by the persistent perianth, *c*.

Fig. 757.—Flower of *Charærops humilis*, European Fan-Palm, seen from above. There are six divisions of the perianth, six stamens, and the ovary.

and for forming habitations. The fruit of some is eatable, while that of others is extremely hard. Many supply oil, wax, starchy matter, and sugar, which is fermented so as to form an intoxicating beverage. Their fibres are employed for ropes, and the reticulum surrounding their leaves is sometimes manufactured into brushes.

The Palm of the Bible, *תאנה*, seems to be *Phoenix dactylifera*, the Date, the drupaceous fruit of which supplies food to many of the inhabitants of Arabia and Africa. *Cocos nucifera* (fig. 115, 1), the Coco-nut Palm, is one of the most useful, supplying food, clothing, materials for houses, and utensils of various kinds, ropes and oil. The sugar procured from it is called Jagery, and is fermented so as to form arrack. The fibrous part of its fruit is manufactured into Coir-rope. The wood of the Coco-nut Palm is known by the name of porcupine wood. The terminal bud of the Coco-nut Palm, as well as that of *Euterpe montana*, the Cabbage Palm, are used as culinary vegetables. The Double Coco-nut of the Seychelles islands is produced by *Lodoicea Seychellarum*. The palm-oil imported from the west coast of Africa is obtained by bruising the fruit of *Elais guineensis* and *melanococca*. The oil-bearing palms are in the tribe Coccoinæ. The Betel-nut is the produce of *Areca Catechu*, and from it an extract is prepared of an astringent nature resembling Catechu. The seeds or nuts form an ingredient in the Eastern masticatory called Pan or Betle, and which seems to owe its stimulating properties to the leaves of the Piper Betel. Sago, and starchy matter allied to it, is obtained from many Palms. It is contained in the cellular tissue of the stem, and is separated by bruising and elutriation. Fine Sago is said to be procured from *Sagus lævis* or *inermis*, a native of Borneo and Sumatra; *S. Rumphii* or *farinifera*, a native of Malacca; and *Saguerus Rumphii* or *saccharifer*, which is found in the eastern islands of the Indian Ocean. After the starchy matter is washed out of the stems of these Palms, it is then granulated so as to form sago. A single tree, it is said, will yield 500 to 600 pounds. The last-mentioned Palm also furnishes a large supply of sugar. Sago, as well as sugar and a kind of Palm-wine, are procured from *Cayota urens*. The date sugar of Bengal is the produce of *Phoenix sylvestris*. *Ceroxylon* or *Iriarteia Andicola* yields wax, which forms a coating over its trunk. *Corypha cerifera*, Carnahuba Palm, is another wax palm. Its trunks are imported into Britain, and have been used for veneering. It is much used in the northern parts of Brazil, as at Aracaty, for thatch, hats, packsaddles, stakes, and palisades. The wax is procured by shaking the leaves, which have a glaucous bloom. Each leaf will yield fifty grains. A reddish resinous matter is yielded by *Calamus Draco*. It is one of the substances called Dragon's-blood, and is used for colouring. The whalebone-like bristles which surround the base of the leaves of some species of *Sagus* and *Saguerus* are used for brushes. The thinner stems of Palms, as of

Calamus Scipionum and *Rotang*, are used as canes under the name of Rattans. *Calamus Rudentum*, the Cable Cane, a native of the East Indies, Cochin-China, and the Moluccas, grows sometimes to the length of 500 feet. The fruit of *Attalea funifera* is known by the name of Coquilla-nuts, and its hard pericarp is used for making umbrella-handles, &c. It seems to supply a fibre used in manufacture under the name of Piassaba. The hard albumen of *Phytelephas macrocarpa* is used in the same way as ivory. Hence the plant is called the Ivory Palm. The spathe of *Manicaria saccifera* comes off in the form of a conical cap, and is used as a covering for the head in the West Indies. *Chamærops humilis* is the only European species of Palm. It is able to stand the climate of this country with slight protection during winter. A specimen in the Edinburgh Botanical Garden has lived in the open air for upwards of thirty years. It is covered with matting during winter. The Doom-palm of Egypt (*Hyphcne thebaica*) has a trunk which divides in a dichotomous manner. Its pericarp is used as food, and has the taste of gingerbread. *Acrocomia sclerocarpa* is the Macahuba-palm of Brazil; *Mauritia vinifera* is the Buritipalm, the stem of which, when perforated, yields a reddish juice, having the taste of sweet wine. Among the Palms which have flowered in the Edinburgh Botanical Garden may be mentioned—various species of *Chamædorea*, *Euterpe montana*, *Chamærops humilis*, *Livistona chinensis*, *Harina caryotoides*, *Areca*, *triandra*, *Corypha umbraculifera*? *Phoenix sylvestris*.

1087. Order 203.—**Commelynaceæ**, the Spider-wort Family. (*Mono-hypogyn.*) Perianth in 2 verticils; outer (calyx) herbaceous and tripartite; inner (corolla) petaloid, tripartite or trifold. Stamens 6 or fewer, hypogynous, some of them occasionally abortive or deformed; anthers introrse. Ovary 3-celled; ovules few in each cell; style 1; stigma 1. Fruit a 2-3-celled, 2-3-valved capsule, with loculicidal dehiscence. Seeds often in pairs, with a lateral and linear hilum; embryo pulley-shaped, antitropal, in a cavity of fleshy albumen, remote from the hilum.—Herbs with flat narrow leaves, which are usually sheathing at the base. Natives chiefly of warm climates. Some have fleshy rhizomes, which are used for food. *Tradescantias*, Spider-worts, have moniliform staminal hairs, in which a movement of granules is distinctly seen under the microscope (fig. 225). There are 17 genera, and 264 species. *Examples*—*Commelyna*, *Tradescantia*, *Mayaca*.

1088. Order 204.—**Alismaceæ**, the Water-plantain Family. (*Mono-hypog.*) Perianth in 6 divisions and 2 verticils; outer whorl usually herbaceous; inner usually petaloid; sometimes the perianth is wanting. Stamens definite or 00, hypogynous; anthers introrse or extrorse; ovaries 3, 6, or more, distinct or united; ovules erect or ascending, solitary or in pairs. Styles and stigmas equal to the number of car-

pels. Fruit of several dry, indehiscent carpidia. Seeds 1-2 in each carpel, exalbuminous (fig. 525); embryo straight, or curved like a horse-shoe; radicle next the hilum.—Plants growing in flowing or stagnant water, usually with a creeping rhizome, parallel-veined leaves, and hermaphrodite or unisexual flowers. Natives both of tropical and temperate regions. The limits of the order are not well defined. It has been divided into two suborders:—1. *Alismæ*, inner perianth petaloid, anthers introrse, embryo curved or hooked. 2. *Juncagineæ*, inner perianth herbaceous, sometimes perianth 0, anthers extrorse, embryo straight, plumule coming through a slit in the embryo (fig. 504). They have few important properties. Some are acrid, others have eatable rhizomes. Various Brazilian *Sagittarias* are very astringent; their expressed juice being employed in the preparation of ink. There are 5 known genera, and about 70 species. *Examples*—*Alisma*, *Sagittaria*; *Triglochin*, *Scheuchzeria*.

1089. Order 205.—*Butomaceæ*, the Flowering-rush Family. (*Mono-hypog.*) Perianth of 6 parts, in 2 verticils (fig. 381, 2); outer usually herbaceous; inner petaloid. Stamens definite (fig. 381, 2, *eo*, *ei*), or 00, hypogynous. Ovaries 3, 6, or more, distinct or united, 1-celled (fig. 381); ovules 00; stigmas simple, as many as the carpels. Fruit consisting of several follicles, which are either distinct (fig. 393) and beaked, or combined. Seeds 00, minute, attached to the whole inner surface of the pericarp (fig. 394), exalbuminous; embryo often curved like a horse-shoe; radicle next the hilum.—Aquatic plants, often lactescent, with parallel-veined leaves, and frequently umbellate flowers. They are chiefly found in northern countries, and some of them have acrid and bitter properties. *Butomus umbellatus*, Flowering-rush, is the only British plant in the class Enneandria of Linnæus. Lindley gives 4 genera, including 7 species. *Examples*—*Butomus*, *Limncharis*.

c. Flowers incomplete, often unisexual, without a proper perianth, or with a few verticillate scales.

† 1090. Order 206.—*Pandanaceæ*, the Screw-pine Family. (*Mono-hypog.*) Flowers unisexual or polygamous, covering the whole of the spadix. Perianth 0, or a few scales. Male flowers: stamens numerous; filaments with single anthers, which are 2-4-celled. Female flowers: ovaries 1-celled, united in parcels; ovules solitary or numerous, anatropal; stigmas sessile, equal to the carpels in number. Fruit either fibrous drupes collected into parcels, or berries. Seeds solitary in the drupes, numerous in the berries; embryo at the base of fleshy albumen; radicle next the hilum.—Trees or bushes, sometimes with adventitious roots (fig. 115, 2), long, imbricated, amplexicaul leaves, usually with spiny margins and backs. Natives of tropical regions.

See N. p. 167/1 كيطور = *Pandanus Obvatissimus*.

The order is subdivided into two sections:—1. Pandaneæ, undivided leaves and no perianth. 2. Cyclantheæ, fan-shaped or pinnate leaves, flowers with a few scales. There are 7 genera, according to Lindley, and 75 species. *Examples*—*Pandanus*, *Freycinetia*; *Cyclanthus*, *Carludovica*, *Nipa*.

1091. The flowers of some of the plants are fragrant, and their seeds are sometimes used as food. The juice has in some instances astringent properties. The species of *Pandanus* are remarkable for their aerial roots, with large cup-like spongioles. These roots are sent out regularly from all parts of their stems, and appear like artificial props (fig. 115, 2). Their spermoderm has numerous raphides. Their leaves are arranged in a spiral manner in three rows, and in their aspect they have some resemblance to those of the pine-apple—hence the name Screw-pine. *Pandanus Candelabra* is the Chandelier-tree of Guinea, and is so called on account of its mode of branching.

1092. Order 207.—**Araceæ**, the Arum Family. (*Mono-hypog.*) Flowers generally unisexual, rarely bisexual, enclosed within a spatha, and usually on a spadix (fig. 239), having male flowers at its upper part, female below, and abortive flowers between them (fig. 239, 2). Perianth either 0, or in the ♂ flowers rudimentary and scaly. Stamens definite or 00, hypogynous; anthers extrorse. Ovary free, 1- 3- or more celled; ovules solitary or numerous; style short or 0; stigma simple. Fruit succulent or dry, indehiscent, one, very rarely three-celled; seeds one or several; embryo in the axis of fleshy or mealy albumen, sometimes with a lateral cleft for the plumule; radicle usually next the hilum.—Herbaceous or shrubby plants, often with tubers or creeping rhizomes, leaves sheathing at the base, and having parallel or branching veins (fig. 239, 1). They occur in dry and marshy places, and in lakes in various parts of the world, abounding in the tropics. The order has been divided into four suborders:—1. Arineæ, Cuckoo-pint tribe (fig. 239); naked flowers with a spadix and spatha, ♂ ♀, anthers sessile, ovules several, fruit succulent, seeds pulpy. 2. Typhineæ, Bulrush-tribe; marsh or ditch plants, with nodeless stems, flowers ♂ ♀, with a scaly or hairy perianth, arranged on a spadix without a spatha, anthers wedge-shaped on long filaments, ovule solitary, fruit dry, seed with adherent pericarp. 3. Acoreæ, Sweet-flag tribe; flowers ♂, having usually a scaly perianth, arranged on a spathaceous spadix, ovules 1 or more, fruit a berry. 4. Pistiæ, Duckweed tribe; flowers ♂ ♀, naked, enclosed in a spatha without a spadix, ovary 1-celled, ovules 2 or more, fruit membranous or capsular. The order includes 47 genera, and 273 species. *Examples*—*Arum*, *Caladium*, *Colocasia*, *Calla*; *Typha*, *Sparganium*; *Acorus*, *Orontium*, *Pothos*; *Pistia*, *Lemna*.

1093. The general property of the order is acidity. Sometimes the plants are dangerous irritant poisons. In some instances the

Spatheaceous flowers
 Spadiceous
 Spadix
 Unisexual flowers
 Araceæ

rhizomes yield much starchy matter, and when boiled or roasted, are used as substitutes for yams, under the name of Coco. The starch may be separated and used as Arrow-root. Thus, Portland Sago is prepared from the rhizome of *Arum maculatum*, common Cuckoo-pink, or Wake-Robin. *Dieffenbachia seguina* (*Caladium seguinum*), is called Dumb-cane, on account of the swelling of the tongue caused by chewing the plant. Many of the plants of this order give out heat in a marked degree during flowering (§ 475). Some send out aerial roots, by means of which they climb upon trees. *Dracontium pertusum* has perforated leaves (§ 137). *Symplocarpus fœtidus*, Skunk-cabbage, has a very disagreeable odour. Its rhizome and seeds have been used as antispasmodics. *Richardia africana*, with its white spatha, is commonly cultivated under the name of Ethiopian Lily. The root-stock of *Acorus Calamus*, Sweet-flag, has an aromatic odour, combined with a bitterish acrid taste. It has been used as a stimulant and tonic. In *Typha latifolia*, Great Reed Mace, the pollen is abundant and easily collected, and from its inflammable nature has been used as a substitute for the Lycopodium spores. The young shoots of *T. latifolia* and *angustifolia* are eaten by the Cossacks like asparagus. The large, fleshy, amylaceous rhizomes are eaten by the Kalmucks. *Lemnas*, Duckweeds, are common in ditches in temperate regions. Their flowers are very simple, one male, and the other female, without a perianth, enclosed in a membranous bag; their roots are simple fibres, covered with a sheath. *Pistia Stratiotes* floats in lakes in tropical countries.

1094. Order 208.—**Naiadaceæ** or **Potameæ**, the Naias or Pondweed Family. (*Mono-hypog.*) Flowers hermaphrodite or unisexual. Perianth of two or four herbaceous or scaly pieces, often deciduous, sometimes 0. Stamens definite, hypogynous. Ovary free, of one or more carpels; ovule solitary; style 1 or 0; stigma entire, rarely 2-3 parted. Fruit dry, 1-celled, usually indehiscent. Seed solitary, erect, or pendulous, exalbuminous; embryo straight or curved, usually with a lateral slit for the plumule (fig. 758); radicle large (figs. 499, 758).—Plants living in



fresh and in salt water, having cellular leaves with parallel veins and inconspicuous flowers. They are found in various parts of the world. They have no properties of importance. *Zostera marina* is used in the dried state for stuffing mattresses, and has been recommended for hospitals. *Ouvirandra* (*Hydrogeton*) *fenestralis* has peculiar skeleton-like leaves (fig. 131). *Aponogeton distachyum*, a Cape aquatic, has grown well for many years in the open pond of the Edinburgh Botanical Garden. *Caulinia fragilis* is one of the plants in which Rotation has been observed. There are 19 known genera, and upwards of 70

Fig. 758.—Embryo of *Zostera*, in the natural order Naiadaceæ. *c*, Cotyledon. *r*, Radicle. *b*, Lateral swelling connected with the radicle. *f*, Slit for the plumule, which lies in a cavity of the very large radicle.

species. *Examples*—*Naias*, *Zannichellia* (fig. 505), *Potamogeton* (fig. 130), *Ruppia*, *Zostera*.

1095. Order 209.—**Restiaceæ**, the Restia or Cord-rush Family. (*Mono-perigyn.*) Flowers frequently unisexual. Perianth glumaceous, sometimes 0. Stamens definite, perigynous, when two or three in number opposite the inner glumes; anthers usually 1-celled. Ovary 1 or more celled, sometimes composed of several carpels; ovules solitary, pendulous; styles and stigmas 2 or more. Fruit capsular or nucumentaceous. Seeds pendulous; embryo lenticular, outside mealy albumen, remote from the hilum.—Herbs or undershrubs, with narrow simple leaves or none, naked or sheathed culms, and spiked or capitate, bracteated flowers. They are found chiefly in America and New Holland. They have few properties of importance. The tough wiry stems of *Willdenovia teres* and some *Restias*, are used for making baskets and brooms. *Eriocaulon septangulare* is a native of Britain, being found in the Isle of Skye, as well as in the west of Ireland. In Brazil there exist branched *Eriocaulons* six feet high. In 1764, Linnæus described only 5 species of *Eriocaulon* in all the world, while Gardner collected in Brazil 100 species. The Diamond districts of Brazil are great centres of *Eriocaulons*. There are according to Lindley, 36 genera, and 286 species. *Examples*—*Restio*, *Centrolepis*, *Eriocaulon*.

Subclass 3.—GLUMACEÆ.

1096. Flowers glumaceous, consisting of bracts or scales, which are imbricated, and not arranged in true verticils. Leaves with parallel veins.

1097. Order 210.—**Cyperaceæ**, the Sedge Family. (*Mono-hypog.*) Flowers hermaphrodite or unisexual, generally without a perianth. Each flower furnished with a solitary bract (glume or scale). These bracts are imbricated upon a common axis, and the lowermost are often empty. Occasionally they enclose two or three opposite membranous bracts or glumes. (In the female flower of *Carex*, the two inner bracts receive the name of Perigynium.) Stamens hypogynous, definite, 1-12; anthers dithecal, innate. Ovary 1-celled, often surrounded by hypogynous bristles (setæ) which are probably abortive filaments; ovule erect, anatropal; style single, 2-3-cleft; stigmas undivided, sometimes bifid. Fruit a crustaceous or bony achæmium or nut (fig. 519); embryo lenticular, enclosed within the base of fleshy or farinaceous albumen (fig. 519 e); plumule inconspicuous (fig. 759).—Grass-like herbs with fibrous roots. Their stems are solid, often without joints, sometimes creeping



Fig. 759.—Embryo of *Carex depauperata*, separated to show the structure of that body in the natural order Cyperaceæ. *r*, Radicle. *c a*, Cotyledon. *f*, Slit for the plumule.

Cyperaceæ - jointless stem - Embryo lenticular - Plumule in the center - Albumen -


(fig. 91), frequently angular. The leaves are narrow, and their sheaths are entire, not slit. They are found in all quarters of the globe, and in various localities, from the sand on the shore, to the tops of the mountains. Many of them occur in marshy ground. Lindley enumerates 112 genera, including 2,000 species. *Examples*—*Cyperus*, *Eriophorum*, *Scirpus*, *Fuirena*, *Cladium*, *Schœnus*, *Scleria*, *Elyna*, *Carex*.

1098. None of the plants of the order possess important medicinal qualities. The creeping stems of *Carex arenaria*, *disticha*, and *hirta*, are diaphoretic and demulcent, and have been used in medicine under the name of German Sarsaparilla. *Papyrus antiquorum* is the Papyrus of the Nile, the cellular tissue of which was used in the manufacture of paper. Some say that the word *קנה* in the Bible, translated Bulrush, is either the *Papyrus* or a species of *Cyperus*. The word *קנה* has been translated Paper-Reeds. The species of *Eriophorum* are called Cotton-grass, on account of the woolly-like substance which is attached to the base of the ovary. Some species of *Cyperus* have tubers at the lower part of their stems, which are used as food. The roots of *Cyperus longus* have been used as bitter and tonic remedies, while those of *C. odoratus* are aromatic. Some species of *Scirpus* are used for making chair bottoms. Some of the *Carices*, with their creeping stems, tend to bind together the loose sand on the sea-shore.

1099. Order 211.—**Gramineæ**, the Grass Family. (*Mono-hypogyn.*) Flowers usually ♂, sometimes unisexual or polygamous; 1, 2, or more (some occasionally abortive), are attached to a common axis, and enclosed within bracts, the whole together forming a locusta or spikelet (figs. 760–762). The outer imbricated bracts are called glumes; they are usually 2 (figs. 760, 761 *g e g i*), sometimes 1, rarely wanting, and often unequal. They are either awned (aristate) or awnless (muticous). The bracts enclosed within the glumes are called paleæ or glumellæ; they immediately enclose the stamens, are usually 2, the lower being simple, and the upper being formed of 2 united by their margins (fig. 761 *p e p i*). The innermost set of bracts consists of two or three hypogynous scales (squamulæ, glumellulæ, or lodiculæ), which are either distinct or combined (fig. 763 *p*), and are sometimes wanting. Stamens hypogynous, 1–6 or more; anthers dithecal, versatile (figs. 337, 763 *e*). Ovary simple (fig. 763 *o*); ovule ascending, anatropal; styles 2 (fig. 763) or 3, sometimes united; stigmas feathery or hairy (fig. 410, 763 *s s*). Fruit a caryopsis (fig. 467). Seed incorporated with the pericarp; embryo lenticular, lying on one side of farinaceous albumen (fig. 495), near its base (figs. 764, 765); endorhizal in germination (fig. 124).—Herbaceous plants, with cylindrical, hollow (fig. 111), and jointed stems, called culms; alternate leaves, with a split sheath and a membranous expansion at the junction of the petiole and blade, called a ligule (fig. 194), the

collections of flowers (locustæ) being arranged in spikes, racemes, or panicles.

Discussions have arisen as to the true nature of the palcæ in



grasses. Brown thinks that the upper palea is composed of two parts united, while the inferior palea is the third part. The arrange-

Figs. 760-765.—Organs of fructification of *Avena sativa*, common cultivated Oat, to illustrate the natural order Gramineæ.

Fig. 760.—Spikelet of the Oat. *a*, Axis of inflorescence or rachis. *ge*, Exterior or lower glume. *gi*, Inner or upper glume. *ff*, Inferior fertile flower, *fa*, Two upper abortive flowers.

Fig. 761.—The same spikelet with the envelopes separated to show the internal parts. *aa*, Axis of inflorescence. *ge*, Outer glume. *gi*, Inner glume. *pe*, Outer palea of the fertile flower, with its awn (arista). *pi*, Inner palea, cleft at the apex, and apparently formed by two united. *e*, Three stamens. *o*, Pistil consisting of the ovary, and two styles. *fa*, Two abortive flowers.

Fig. 762.—Diagram of the spikelet. *ge*, Outer glume. *gi*, Inner glume. *pe*, Outer palea with awn; the inner palea being opposite. *e*, Stamens. *o*, Pistil. *ll*, Scales or lodiculæ. *fs*, *fs*, Barren flowers.

Fig. 763.—Fertile flower deprived of glumes and palcæ. *e*, Three stamens with versatile cleft anthers. *p*, Scales (squamæ or lodiculæ) partially united. *o*, Ovary ultimately forming the grain, which consists of pericarp and seed combined. *ss*, Two styles with feathery stigmas.

Fig. 764.—Vertical section of the Caryopsis (fruit or grain), with the upper portion cut off. *tt*, Integuments of the caryopsis and of the seed united. *pp*, Perisperm. *e*, Embryo. *r*, Radicle. *ca*, Cotyledon. *f*, Slit corresponding to the plumule.

Fig. 765.—Embryo separated. *r*, Radicle. *ca*, Cotyledon. *f*, Slit corresponding to the plumule.

Rolled Inflorescence - Ligule
 Embryo outside - Althous
 Gramineæ - Abortive Stamens (Culms)
 Abortive anthers -

ment is thus trimerous, and bears a relation to the scales or lodiculæ. Mohl, on the other hand, states, that the inferior palea is not on a level with the other, and is in fact a bract from which the other is developed. From their alternate position, the parts of the flowers of grasses are in general looked upon as bracts, rather than as parts of a true perianth. The following may be given as a general view of the parts of the flower.

1. Outer envelope: One or two glumes, the calyx of some authors, containing one or more flowers, with distinct insertions on a common axis. When one glume is suppressed it is the exterior or lower.
2. Inner envelope: One or two paleæ, corolla of some authors, calyx of Jussieu. Inner or upper palea sometimes suppressed. This palea (valve) consists usually of two confluent valves, as shown by two ribs equidistant from the axis. Hence this envelope is, according to some, a ternary perianth.
3. Squamulæ (scales, lodiculæ, or glumellulæ), occur within the last envelope, and at the base of the ovary.

Grasses are found in all quarters of the globe, and are said to form about $\frac{1}{2}$ part of known plants. In tropical regions they sometimes assume the appearance of trees. They generally grow in great quantity together, so as to receive the name of social plants. The order has been divided into numerous sections, founded on the number of flowers in a spikelet, their hermaphrodite, unisexual, or polygamous nature, the number and form of the different sets of bracts, and the nature of the fruit. Lindley enumerates 291 genera, including about 3,800 species. *Examples*—*Oryza*, *Zea*, *Phalaris*, *Panicum*, *Stipa*, *Agrostis*, *Arundo*, *Echinaria*, *Chloris*, *Avena*, *Bromus*, *Festuca*, *Bambusa*, *Lolium*, *Triticum*, *Hordeum*, *Nardus*, *Rottboellia*, *Andropogon*, *Saccharum*.

1100. This is one of the most important orders in the vegetable kingdom, whether we regard it as supplying food for man, or herbage for animals. To the former division belong the nutritious cereal grains, as Wheat (*Triticum*), Oats (*Avena*), Barley (*Hordeum*), Rye (*Secale*), Rice (*Oryza*), Maize (*Zea*), Guinea-corn and Millet (*Sorghum* and *Panicum*); to the latter, the various pasture grasses, as Rye-grass (*Lolium*), Timothy-grass (*Phleum*), Meadow-grass (*Poa*), Cock's-foot-grass (*Dactylis*), Sweet-Vernal-grass (*Anthoxanthum*), Fescue (*Festuca*), Dog's-tail-grass (*Cynosurus*), &c. The grains of many other grasses are used for food. *Zizania aquatica* supplies a kind of rice in Canada, *Setaria germanica* yields German millet, *Panicum miliaceum* gives a kind of millet in India, and *Andropogon Sorghum* has been sent to this country under the name of Durra, an Indian grain. *Phalaris canariensis* is the source of the common Canary seed. The cereal grains have been so extensively distributed by man, that all traces of their native country are lost. They seem to be in many instances examples

of permanent varieties or races kept up by cultivation. Their grain or caryopsis contains a large amount of starch (figs. 35, 36) and gluten. The grasses used for fodder in some parts of the world attain a large size, such as *Anthistiria australis*, the Kangaroo-grass of New Holland, *Tripsacum dactyloides*, the Gama-grass of Mexico, and *Dactylis cæspitosa*, the Tussac-grass of the Falkland islands. Some of these are five or six feet in height, and are, nevertheless, sufficiently delicate to be used as food for animals. The Tussac has been introduced into this country, and it thrives well in peaty soils within the influence of the sea spray. It promises to be a valuable grass in the Hebrides of Scotland.

1101. Sugar is a valuable product obtained from many grasses. It has been procured in Italy from *Sorghum saccharatum*, sweet Sorgho; in China, from *Saccharum sinense*; in Brazil, from *Gynerium saccharoides*; in the West Indies, from *Saccharum violaceum*; and in many other parts of the world, from *S. officinarum*. The last two are commonly known as Sugar-cane, and they are generally considered as varieties of a single species, *Saccharum officinarum*, which is now widely spread over various parts of the world. Six or eight pounds of the saccharine juice of the plant yield one pound of raw sugar. The following were the imports of sugar into Great Britain in 1848:—

West Indies.....	121,600 tons.
Mauritius.....	43,600 —
East Indies.....	65,200 —
Java and Manilla.....	11,000 —
Havana, Porto-Rico, and Brazil.....	76,900 —
Total.....	318,300 tons.

1102. Some grasses have a very agreeable fragrance. This has been remarked in *Anthoxanthum odoratum*, which is hence called sweet-scented vernal grass, and is said to impart the odour to new-made hay in this country. Other grasses have the same property, which has been referred to the presence of benzoic acid. A fragrant oil is procured from some species of *Andropogon*, as *A. Shenanthus*, Lemon-grass, and *A. Calamus aromaticus*, which seems to be the הקנה or הקנה נשם the Sweet-cane of the Bible. Grasses contain a large quantity of siliceous matter in their stalks. This is deposited so as to form part of their structure, and in some cases it accumulates in the joints. The tabasheer in the joints of *Bambusa arundinacea*, the Bamboo, is composed of silica. This is one of the tree-like branching grasses, which sometimes attains a height of fifty or sixty feet. It shoots up with great rapidity. In the Edinburgh Botanic Garden the young shoots attain a height of thirty or forty feet in a few months; and the late superintendent (Mr. W. M'Nab) measured, during a long summer day, a growth of the young stem to the extent of seven or eight inches.

Most of the species of Bamboo have hollow stems, which often attain a diameter of many inches. Gardner mentions a large species of Bamboo (*B. Tagoara*) having a stem 18 inches in circumference, and attaining a height of 50 to 100 feet. The touch-paper of the Chinese is made from a variety of Bamboo, by beating the young shoots flat, steeping them in a lime pit for a month, and then washing and drying. A kind of paper is made from the Bamboo in India. Its young shoots are used as pickles. The hollow stems of some reeds in warm climates supply refreshing water to travellers.

1103. The stems of some grasses run under ground, and form a sort of net-work, which is useful in consolidating the sand of the sea shore. *Elymus arenarius* and *Ammophila* (*Psamma*) *arenaria* constitute the Bent and Marram of the British shores. This tendency to creep under ground renders some grasses, such as *Triticum repens*, Couch-grass, difficult of extirpation. The grains of some grasses are used for ornaments. Beads are made from those of *Coix Lachryma*, commonly called Job's-tears, from their form and hardness. A few grasses, as *Bromus purgans* and *catharticus*, have purgative properties; and one, *Lolium temulentum* (*infelix lolium*), Darnel-grass, is poisonous. Some suppose that it is the ζιζάνια, tares, of Scripture. The grains of Rye, and other grains, are liable to a disease called Ergot, depending on the attack of a fungus which alters the texture of the ovary, and makes it assume an elongated spurred form. The Ergot of Rye, or spurred Rye, has a peculiar effect in promoting the contraction of the uterus, and is on this account used in medicine. Ergotted rye, when regularly used for food, has the effect of causing what has been called convulsive and gangrenous ergotism, the former disease being distinguished by insensibility and convulsions, ending in death; the latter by dry gangrene, which attacks the fingers and toes, causing sloughing of these parts, and sometimes proving fatal by exhaustion. The poisonous effects of Ergot are attributed to the presence of a fixed oil.

SECTION II.—CRYPTOGAMOUS PLANTS.

CLASS III.—ACOTYLEDONES, *Juss.* CELLULARES AND MONO-CRYPTOGAMEÆ, *DC.*
THALLOPHYTA AND ACROBRYA, *Endlich.* THALLOGENS AND ACOGENS, *Lindl.*

1104. The plants belonging to this class are in some instances composed entirely of *cellular tissue*; in other instances both cells and vessels are present. The *vascular tissue* in the higher orders consists partly of closed spiral and scalariform (fig. 62) vessels. Many of them have no true stem nor leaves. The *woody stem*, when present,

consists of simultaneous vascular bundles, which increase in an acrogenous manner (§ 103). The stem of Tree-ferns (which illustrates this class) is unbranched, more or less uniformly cylindrical, hollow in the interior, and marked by the scars of the leaves (fig. 116). *Stomata* occur in the epidermis of the higher divisions. *Leaves*, when present, have frequently no true venation, at other times the venation is forked. There are no *flowers*, and no distinct stamens nor pistils. *Reproduction* takes place in some cases apparently by the union of cells of different kinds (antheridia and pistillidia) (§ 492), by means of which germinating bodies called *spores* are formed (fig. 498). In other cases it is difficult to trace this process of fertilization. The *spore* may be considered as a cellular embryo which has no cotyledons, and germinates from any part of its surface, being heterorhizal (§ 591, fig. 530).

Subclass 1.—ACROGENÆ OR CORMOGENÆ.*

1105. Acotyledons, having usually distinct stems and leaves, stomata, a certain amount of vascular tissue, and thecæ or cases containing spores. This subclass corresponds in a great measure with the division of Cormophyta, called Acrobrya by Endlicher, and with the Foliosæ or Ætheogamæ of De Candolle.

1106. Order 212.—**Equisetaceæ**, the Horse-tail Family. Stem striated, hollow, usually branched, containing much silica in its composition, articulated, the joints being separable, and surrounded by a membranous toothed sheath. There are no true leaves, green-coloured branches having a straight veneration occupying their place. The cuticle exhibits a longitudinal series of stomata. A spiral structure is observed in some of the vessels. Reproductive organs collected into



cones; spore-cases (thecæ or sporangia) attached to the lower surface of peltate polygonal scales (fig. 766), and opening by an internal

* 'Ακρον, summit, πορὸς, a stalk or stem, and γεννάειν, to produce.

Figs. 766-769.—Reproductive organs of Equisetum, to illustrate the natural order Equisetaceæ.
Fig. 766.—A peltate or polygonal scale, *e*, taken from the terminal cone-like fructification of an Equisetum. *c*, Thecæ or spore-cases arranged in a verticil on the under surface of the scale. *p*, Stalk by which the scale is attached to the axis.

Fig. 767.—*c*, Spore-case seen on its inner surface, with the slit or opening by which the spores are discharged.

Fig. 768.—A spore, *s*, with the two clavate filaments rolled up in a spiral manner around it.

Fig. 769.—Spore, *s*, with the two filaments, which are clavate at each extremity, unrolled. These filaments or elaters are hygrometric, and move about under the influence of moisture.

longitudinal fissure (fig. 767); spores in the form of rounded cells, surrounded by 2 elastic, club-shaped, hygrometric filaments or elaters (figs. 768, 769).—Plants with simple or branched stems, the branches being jointed and placed in whorls at the articulations of the stem, each whorl consisting of as many branches as there are teeth in the sheath. Found in ditches, lakes, and rivers, in various parts of the world. In South America, Gardner measured an *Equisetum* fifteen feet high, and three inches in circumference at the lower part of the stem. There is only 1 known genus, comprehending about 12 species. *Example*—*Equisetum*.

1107. From the quantity of silicic acid contained in them, some of the species of *Equisetum* are used in polishing mahogany. An analysis of them is given at ¶ 226. The spiral filaments which surround their spores are interesting objects under the microscope, exhibiting marked movements according to the moisture or dryness of the atmosphere around them. The stomata are arranged in lines on the cuticle. In *Equisetum hyemale*, often called Dutch Rushes, the siliceous stomatic apparatus is well seen after the action of nitric acid on the stem. There are regular rows of tubercles of a siliceous nature, in each of which is a transverse fissure, and at the bottom of the fissure a stoma is placed, with its opening at right angles to that of the tubercle. Each portion of the stoma has a pectinated (comb-like) appearance. The distinctions between the species of *Equisetum* are founded on the nature of the fertile and barren stems, the number of striæ or furrows, and the number of teeth at the articulations.

1108. Order 213.—**Filices**, the Fern Family. Stem, a rhizome (fig. 770), which creeps along or under the surface of the ground, emitting descending roots and ascending fronds (leaves), or which rises into the air so as to form an acrogenous trunk (fig. 116). This trunk (stipe) is of nearly uniform diameter, is hollow in the interior, marked on the hard outer rind by the scars (cicatrices) of the leaves, and contains vascular bundles of woody, dotted, and scalariform vessels, which are enclosed in hard plates, and are arranged in an irregular manner (fig. 117). Sometimes the trunk is dichotomous (fig. 118). The outer fibrous covering is formed by the bases of the leaves, and is thicker at the lower than at the upper part of the stem. The leaves (fronds) have a circinate (gyrate) vernation (fig. 770 *f' f''*); their veins are generally of equal thickness, and either simple or dividing in a forked manner (fig. 771), or somewhat reticulated, and occasionally stomata occur. Reproductive organs, consisting of spores cases (thecæ, sporangia), which arise from the veins on the under surface of the fronds (figs. 770 *f'''*, 771 *s*, 772), or from their margin. Spore-cases, either stalked, with the pedicel passing round them in the form of an elastic ring (fig. 773), or sessile and destitute of a ring. The thecæ sometimes arise from the surface of the frond, while at other times they

spring from below, having a cuticular covering in the form of an indusium or involucre (fig. 771). The clusters of thecæ are called sori (fig. 772). The margin of the frond sometimes is folded so as to cover the thecæ, and at times the whole frond is converted into clusters of thecæ. The subject of embryonic growth in Ferns has latterly attracted much attention. It is now believed that the prothallium simultaneously bears antheridia and archegonia. The antheridia occur in the under surface of the prothallium, and consist of cellular papillæ, having a central cavity. This cavity contains free cellules, within each of which a ciliated spiral filament or spermatozoid rests. This cavity discharges those cellules by a rupture at its apex. The spiral filaments then burst the cellules, and being set free reach the ovule containing the embryo germ, which ovule, embedded in the substance of the prothallus, is constituted and surrounded by the archegonial cells. Those cells are larger than the antheridia, and present a canal leading to the ovule, which canal remains closed till the period when the spermatozoid is matured. Mercklen asserts that the spermatozoid enters the canal of the archegonium. Suminski believes that the end of a spermatozoid is developed into the embryo cell, which after a time becomes converted into an embryo, from which the future plant results.—Ferns are elegant leafy plants, occurring chiefly in moist insular climates, and abounding in the tropical islands. In mild and warm climates they occur in the form of large Tree-ferns, fifty to sixty feet high, which give a char-

Fig. 770.—Rhizome of *Scolopendrium officinale*, with several fronds (leaves), *f*, *f'*, *f''*, *f'''*, in different degrees of development. In *f'* and *f''*, the circinate or gyrate vernation is seen. In *f'''*, the linear transverse sori or clusters of thecæ are seen, having the appearance of dark lines on the lower surface of the frond.

Figs. 771-773.—Frond and fructification of *Polystichum angulare*, to illustrate the natural order Filices.

Fig. 771.—Part of the frond seen on the lower surface. *p*, Two pinnæ covered with sori, *s*, having an indusium. *r*, Rachis or central stalk of the frond.

Fig. 772.—One of the sori or clusters of thecæ cut vertically. *n*, The vein bearing it. *i*, Indusium or fold of the frond covering it. *c*, Thecæ or sporangia (spore-cases).

Fig. 773.—One of the thecæ separated at the period of dehiscence. *a*, Incomplete annulus or ring, which is elastic, and causes transverse dehiscence of the theca. *p*, Stalk of the theca. *s*, Spores discharged.



acter to the landscape. The theca of ferns has been looked upon as a modified leaf, having the same gyrate or circinate development as the frond. Leaves have occasionally been produced in place of thecæ. Ferns having the thecæ on the back of the frond, and furnished with an elastic ring or band, are called dorsiferous and annulate; while those having no thecal ring are exannulate.

1109. The order has been divided into several suborders:—

1. Polypodiæ, thecæ on the back of the frond, pedicellate or sessile, distinct, annulate, ring vertical, usually incomplete, bursting irregularly and transversely.
2. Hymenophyllæ, thecæ marginal or dorsal, nearly sessile, distinct, annulate, ring horizontal, complete, occasionally oblique, bursting lengthwise.
3. Osmundæ, thecæ dorsal, or forming a separate stalked mass (an altered frond), distinct, with a terminal or dorsal ring more or less incomplete, bursting lengthwise.
4. Ophioglosseæ, thecæ collected into a spike, formed at the edges of an altered frond, distinct, exannulate, two-valved.
5. Danaæ, thecæ united in masses, exannulate, opening irregularly by a central cleft.

The generic characters of Ferns are founded on the position and direction, covered or uncovered nature of the sori, as well as on the venation. Lindley notices 192 genera in his Filical Alliance, including upwards of 2,000 species. Kunze estimates the species at 3,000. There are not above 700 species cultivated in Europe. In 1846 there were 387 species grown in the Royal Garden at Kew. *Examples*—Polypodium, Aspidium, Lastrea, Grammitis, Adiantum, Pteris, Davallia, Woodsia, Cyathea; Hymenophyllum, Trichomanes, Aneimia, Lygodium; Osmunda; Ophioglossum, Botrychium; Danaea.

1110. Few of the Ferns are used medicinally. They are in general demulcent and astringent. Some yield food. The rhizome of *Lastrea Filix-mas*, Male-shield fern, has been used as a vermifuge, especially in cases of tape-worm. It contains starch, gum, saccharine matter, tannin, green fixed oil, and resin. Its properties are ascribed to the fixed oil. The rhizome has been used for tanning, and its ashes contain much carbonate of potash. The syrup called capillare, and certain pectoral mixtures are prepared from *Adiantum pedatum* and *A. Capillus Veneris*. The rhizome of *Pteris esculenta* is used as food in Australia, and that of *Marattia alata* in the Sandwich islands. Many other species of Ferns are esculent. The stems and leaf-stalks of Ferns are often covered with scales, and with woolly matter. One (*Davallia canariensis*) is called Hare's-foot Fern on this account, and another (*Aspidium Barometz*) receives the name of Scythian or Tartarian-lamb, because, when prepared in a particular way, it resembles that animal.

1111. Order 214.—*Marsileaceæ*, or *Rhizocarpeæ*, the Pepperwort Family. Stem wanting. Leaves often stalked, with the lamina divided into three or more wedge-shaped pieces, sometimes the lamina


is abortive; vernation circinate. Reproductive organs near the root or along the petiole, enclosed in an involucre; these organs are of two kinds:—1. Stalked or sessile clustered membranous sacs, containing minute granules, which some consider as pollen; hence the bodies are called anthers. 2. Membranous sacs, containing cells which divide into four, one of which only is developed as a germinating body; the sacs have been called ovule-sacs, and the single developed cell is considered by some as an ovule which is impregnated by the so-called pollen. The thecæ are the bodies from which germination proceeds.—Creeping or floating plants, found in ditches and pools in various parts of the world, more especially in temperate climates. They are not put to any important use. There are 4 genera, and upwards of 20 species. *Examples*—Marsilea, Pilularia, Salvinia.

1112. Order 215.—**Lycopodiaceæ**, the Club-moss Family. Stems creeping or corms; annular vessels in the axis. Leaves imbricated, more or less setaceous, sometimes subulate. Thecæ axillary and sessile, 1-3-celled, opening by valves or indehiscent; often of two kinds. One, round, reniform, or crescentic, containing minute powdery matter, and called by some antheridia, though, perhaps, erroneously; the other of a roundish tetrahedral form, enclosing a cell which produces four spores capable of germinating; the spores are considered by some as equivalent to ovules, and the mother-cell as an ovary or oophoridium (ὠόν, an egg or ovule, and φορέω, I bear). In *Isoetes* the two kinds of reproductive bodies are embedded in the substance of the base of the leaf.—They are moss-like plants, intermediate between ferns and mosses, and in some respects allied to cone-bearing plants. They abound in warm, moist, insular climates. There are 6 genera, and about 200 species. *Examples*—*Lycopodium*, *Selaginella*, *Isoetes*.

1113. Some of the *Lycopodiums* are emetic and cathartic. The powdery matter in the thecæ is inflammable, and has been used as a substitute for sulphur, under the name of Lycopode or vegetable brimstone. It is also employed to cover pills, so as to prevent their being acted upon by moisture. *Lycopodium squamatum*, a Brazilian species, coils up into a ball during the dry season, and unrolls during the wet season.

1114. Order 216.—**Musci**, the Moss Family. Plants having a distinct axis of growth, often giving off branches or innovations; no vascular system. Leaves minute and imbricated (fig. 774 *f*), entire, or serrated; sometimes with condensed cells in the form of ribs or nerves. Reproductive organs of two kinds:—1. Antheridia (fig. 368), cylindrical or fusiform stalked bags, containing powdery matter and phytozoa (fig. 368, 3), and mixed with empty jointed filaments or paraphyses (παράφυσις, an offset). 2. Urn-shaped pistillidia (fig. 777), enclosed at first within a calyptra (καλύπτρα, a covering), which is

ultimately carried up with them (fig. 775 c), leaving often a sheath (vaginula) round the bottom of the fruit-stalk. These pistillidia finally become the thecæ (fig. 774) or spore-cases, supported on a stalk or seta which has leaves at its base, called perichætil (περί, around, and χαίτη, a bristle) leaves; on removal of the calyptra, the theca is



found to consist of a case with an operculum or lid (fig. 776), which, when it falls off, shows the mouth of the urn either naked or crowned with a peristome (περί, around, and στόμα, mouth), consisting of one or more rows of teeth (in number, four, or a multiple of four), distinct, or united in various ways (fig. 776 p). In the centre of the theca is a columella (fig. 777 c), and the bag formed between it and the parietes of the theca, contains spherical cells called spores, each of which divides into four small spores or sporules, the germinating bodies (fig. 777).

Figs 774-777.—Figures to illustrate the natural order Musci.

Fig. 774.—*Funaria hygrometrica* slightly magnified. *f*, Leaves, those connected with the seta being called perichaetial. *u*, Urn-like theca, sporangium or spore-case supported on a long twisted stalk or seta. *p*, *c*, Calyptra, which exists on one of the thecæ, and has fallen from the other. *o*, Operculum or lid.

Fig. 775.—Theca of *Encalypta vulgaris*. *u*, Theca or spore-case. *c*, Mitridiform entire calyptra. *o*, Operculum or lid. *s*, Top of the seta. The calyptra is transparent, and the operculum and theca are seen through it.

Fig. 776.—The same theca, *u*, with the calyptra removed. *o*, Operculum detached, showing the peristome, *p*, with its sixteen cilia or teeth.

Fig. 777.—Very young theca of *Splachnum* cut longitudinally. *a*, Apophysis or swelling of the seta at the base of the theca. *c*, Central columella. *s*, Cavity or bag between the columella and the walls of the theca, containing spores. The integument of the theca is formed of different cellular layers; the first, *e*, forms the epidermis, and is thickened at the summit to form the operculum; there are then two intermediate layers, which ultimately form the teeth of the peristome; and lastly, an inner integument, *s*, which forms the parietes of the spore-bearing cavity.

In some cases the operculum remains persistent and the theca opens by four valves. At the base of the theca there is occasionally a fleshy protuberance at one side called a struma, or a swelling of the seta called an apophysis (ἀποφύσις, excrescence), (fig. 777 *a*). The calyptra is sometimes split on one side (dimidiate), at other times it is entire (fig. 775 *c*) or split into short clefts all round its base (mitriform). Between the teeth of the peristome and the edge of the theca, an elastic ring or annulus is formed; and occasionally a horizontal septum or epiphragmum (φραγμα, a partition) extends across the mouth of the thecæ. The setæ are sometimes twisted, and so are the teeth of the peristome.—Mosses are either erect or creeping, terrestrial or aquatic plants found in all moist countries, extending from the arctic to the antarctic regions. They abound most in temperate climates. They are among the first plants which appear on newly-formed islands. In speaking of the morphology of mosses, Lindley states that the calyptra may be considered as a convolute leaf, the operculum another, the peristome one or more whorls of minute flat leaves, and the theca itself as the excavated distended apex of the seta. I have a specimen of *Tortula fallax*, which I received from the late Mr. E. Quekett, in which leaves appear at the top of the seta in place of the spore-case.

1115. Mosses have been divided according as their seta is terminal (*acrocarpi*, ἀκροα, summit, and καρπός, fruit), or lateral (*pleurocarpi*, πλευροα, side), according as the operculum is adherent or not, and according as the mouth of the theca is naked, or has a single or double peristome. Divisions have also been adopted, founded on the position of the antheridia and pistillidia, &c. Lindley separates *Andræa* from true Mosses, on account of its spore-case opening by four valves. According to him there are 46 known genera, and upwards of 1,100 species. *Examples*—*Andræa*, *Phascum*, *Gymnostomum*, *Splachnum*, *Orthotrichum*, *Dicranum*, *Bryum*, *Funaria*, *Polytrichum*, *Hypnum*, *Sphagnum*.

1116. Order 217.—**Hepaticæ**, the Liverwort Family. Plants having an axis which either bears cellular leaves (fig. 778), or is leafless and is bordered by a membranous expansion or thallus. Stomata are found in the epidermis of some. The reproductive organs are—1. Antheridia, which are either embedded in the frond, or situated on rounded sessile and stalked receptacles. 2. Pistillidia, either enclosed in involucre and solitary (fig. 778 *ii*), or occurring at the edge of the frond, or on the lower side of stalked peltate expansions (fig. 412). Thecæ or developed pistillidia, having no operculum, opening irregularly, or by four valves (fig. 778). Spores (fig. 498) often mixed with spiral filaments called Elaters (fig. 779). Heterorhizal in germination (fig. 530).—Terrestrial plants found in damp places, or inhabiting water; some having a moss-like appearance. They are natives both of cold and warm climates, and are generally distributed over the globe.

1117. The order has been divided into three sections:—1. *Marchantieæ*: thecæ collected in heads, bursting irregularly, no operculum, spores with elaters. 2. *Jungermanniæ*: thecæ solitary, opening by four valves, no operculum, spores with elaters. 3. *Ricciæ*: thecæ solitary, decaying so as to allow the spores to escape, no operculum, no elaters (fig. 411). Many of the *Hepaticæ* produce gemmæ or buds, which are developed on the frond in the form of cup-shaped receptacles, and ultimately fall off so as to become distinct plants. *Marchantia hemispherica* has been recommended in dropsical

cases. There are, according to Lindley, 65 genera, and about 700 species. *Examples*—*Marchantia*, *Jungermannia*, *Riccia*.

Subclass 2.—THALLOGENÆ* OR CELLULARES.

1118. Acotyledons composed entirely of cellular tissue, having no distinct axis, nor leaves, nor stomata, propagated by means of spores, which are often enclosed in asci. It corresponds to Endlicher's division of *Thallophyta*, and includes the *Amphigamæ* of De Candolle.

1119. Order 218.—*Lichenes*, the Lichen Family. Plants forming a thallus, which is either foliaceous, crustaceous (fig. 780), or pulverulent; these different forms depending on the mode in which the cells are developed and combined. The reproductive organs appear on the frond in the form of protuberances of various kinds, consisting of an outer layer of thick-walled roundish cells, more dense than the tissue of the thallus, and of a different colour (fig. 782 *cc*), and of an internal medullary layer of paraphyses and sporangia, lying perpendicularly to the outer layer (figs. 782 *cm*, 781 *tp*). The fructification gradually projects more from the surface, and either remains covered with the outer layer, or bursts through it. When it remains closed, there is a nucleus in the centre. When the fructification bursts through the cortical or outer layer, it expands in the form of shield-like discs, called


* Θαλλός, a green leaf, γένεσις, I produce.

Figs. 778, 779.—Organs of fructification of *Jungermannia Tamarisci*, to illustrate the natural order *Hepaticæ*.

Fig. 778.—*f*, Branches covered with imbricated leaves, arranged in a distichous manner. Two of the branches bearing thecæ, supported on stalks which arise from an involucre at the base. *i* *i*, Involucres. *c*, Theca closed in the young state. *c'*, Theca opening by four valves to discharge the spores and elaters.

Fig. 779.—*r*, Receptacle bearing elaters, *e*, or spiral filaments, one of which shows the double spiral fibre. *s*, Free spores.

apothecia (ἀποθήκη, a repository), or patellæ (figs. 780 s, 781 a) (*patella*, a hollow disk), or linear expansions called lirellæ (*lira*, a furrow). Sometimes the cortical matter forms a border round the fructification,



at other times it grows up in the form of a stalk, so as to give rise to a podetium (ποῦς, a foot). The young thecæ (asci) contain spores, varying from 4 to 8 (fig. 413), or from 12 to 16. Occasionally, the spores are in sets of two (fig. 413, 2). Separated cells of the medullary layer, of a green colour, called gonidia (γόνι, generation, and εἶδος, resemblance), or gongyli, are considered as another kind of reproductive organ. There is much uncertainty as to the real character of the spherical or subspherical green bodies called gonidia, which are characteristic of true lichens. When separated from the parent structure, they are capable of forming new plants. Thwaites says that they are not gemmæ, but that they are analogous to the vesicles of *Nostoc*, one of the Algæ.—Lichens are found in all quarters of the globe, adhering to stones, rocks, trees, &c. During part of their growth, they appear to be capable of deriving most of their nourishment from the atmosphere. They have the power of acting on hard rocks, so as to disintegrate them in process of time, and many of them contain much inorganic matter in their composition. They all grow in the air; none are found submersed.

Figs. 780-782.—Organs of fructification of *Parmelia Acetabulum*, to illustrate the natural order Lichenes, section Hymenothalameæ.

Fig. 780.—*t*, Thallus of the Lichen. *s*, Apothecia in the form of shields in different degrees of development.

Fig. 781.—Apothecium, *a*, cut vertically and magnified in order to show the layer, *t p*, formed by the union of thecæ and paraphyses.

Fig. 782.—A small portion of the apothecium much more magnified, showing, *cm*, the central medullary layer. *cc*, The cortical layer. *tt*, Thecæ in different degrees of development. *p*, Paraphyses.

1120. The order has been divided into four sections:—

1. Hymenothalamæ (ἵμις, a membrane, θάλαμος, a receptacle): shields open, discoid permanent nucleus bearing the sporangia on its surface (fig. 780).
2. Gasterothalamæ (γαστήρ, a belly): shields either closed always, or opening by bursting through the cortical layer of the thallus, the nucleus containing the deliquescent or shrivelled sporangia.
3. Idiothalamæ (ἴδιος, peculiar): shields closed at first, opening afterwards, containing free spores in a nucleus composed of the gelatinous remains of the paraphyses and sporangia.
4. Coniothalamæ (κίνη, powder), pulverulent lichens; shields open, without a nucleus, cavity filled with free spores.

1121. Lichens furnish articles of food and important dyes. *Cetraria islandica*, commonly called Iceland Moss, contains a nutritious matter called Lichenin, or Lichen-starch. There exists in it a bitter principle also, to which the name Cetrarin has been given. The plant is used as a demulcent and tonic, in the form of decoction or jelly. *Cladonia rangiferina* is a Lichen upon which the Reindeer feeds. Several species of *Gyrophora* constitute the Tripe de Roche, on which Franklin and his companions subsisted for some time. Many other Lichens, such as *Sticta pulmonaria* and species of *Lecanora*, furnish articles of food. *Rocella tinctoria* from the Canaries, and *R. fuciformis*, furnish valuable dyes, under the name of Orchil or Archil. The dye procured from them, and from other Lichens, is called Litmus. *Lecanora tartarea* supplies the dye called Cudbear. *Parmelia parietina* contains a yellow colouring matter called Parietin or Chrysophanic acid. Some species of *Variolaria* contain a large quantity of oxalate of lime. Some plants of the order are aromatic.

1122. Order 219.—**Fungi**, the Mushroom Family. The plants belonging to this order consist of cells, sometimes round, sometimes elongated in the form of filaments, either placed closely together or separated. They are variable in their consistence, being soft or hard, fibrous or gelatinous, fleshy or leathery. They never contain green gonidia like Lichens, and they rarely grow in water. There exists a vegetative system called spawn or mycelium (μύκης, fungus), formed of elongated, simple, or articulated filaments, concealed within the matrix, or expanded over its surface, from which varied forms of fructification proceed. The mycelium occurs either in a filamentous, a membranous, a tubercular, or a pulpy form. The reproductive organs consist of spores or spherical cells (usually 4 or some multiple of 4), which are either attached to the cellular tissue, and supported often on simple or branched filamentous processes (fig. 785, 787 b) called sporophores (σπορά, a spore, and φορέω, I bear) or basidia (βάσις, a base); or are contained in thecæ (*theca*, a sac), cystidia (κύστις, a bladder), or asci (άσκος, a bag), (fig. 785 c), accompanied with bodies called antheridia or paraphyses; in the latter case, the term sporidia

is sometimes applied to the spores. The sporophores sometimes end in delicate cells bearing the spores, and called sterigmata (στερήγμα, a support). In the Agarics or Mushrooms, which are among the best

784


786



783



785



787



known fungi, there is observed first a roundish protuberance on the mycelium. This swelling is called the volva or wrapper, and it gradually enlarges, containing in its interior what appears afterwards as the agaric with its reproductive bodies. When the volva is ruptured, the fully-formed agaric is seen, consisting of an upper rounded portion called the pileus or cap (fig. 783 *c c*), supported on a stalk or stipes (fig. 783 *p*). On its under surface is situated the hymenium (ὑμὴν, membrane), or the part where the spores are produced (fig. 783 *h*), covered at first by a thin membrane called a veil (indusium or velum), which is ultimately ruptured; and when the rupture takes place at the edge of the pileus, an annulus or ring is left on the stipes (fig. 783 *a a*). The hymenium, or the part on which the organs of reproduction are placed, consists in the agaric of cellular plates, lamellæ, or gills, radiating from the centre (fig. 783 *h*). In other genera of Fungi (fig. 786) it

Figs. 783-787.—Figures to illustrate the natural order Fungi.

Fig. 783.—A cluster of plants of *Agaricus campestris*, Mushroom, in different stages of development. *p*, Stipe or stalk. *c c c*, Pileus, hat or cap. *v*, Velum or indusium which unites the pileus and stipe, and when ruptured forms the annulus or ring, *a a*. *h*, Lamellæ or gills radiating from the centre, on the under surface of the pileus, and bearing the hymenium or receptacle of the spores.

Fig. 784.—Hymenium seen from above, the spores, *b*, being scattered over it in sets of four (quaternary).

Fig. 785.—A small portion of the Hymenium much magnified and viewed laterally. *h*, Its tissue composed of cells. *b*, Basidia or sporophores bearing the spores; one of these is figured separately, bearing a large number of spores. *c*, Cystidia or thecæ.

Fig. 786.—A small portion of the pileus of *Clathrus cancellatus*, in the form of a sort of network. The Hymenium covers its inner surface, and is seen following the contour of the lacunæ, *l l*, of the net-work.

Fig. 787.—Hymenium much more highly magnified to show the particular form of the basidia, *b*. *s*, Spores.

consists of tubes or solid columns, or fleshy or gelatinous matter. Sometimes the hymenium is on the upper surface of the fungus.—Cellular plants, often growing on decaying organic matter, generally very fugacious, and presenting various colours. They are found in all parts of the world.

1123. The following are the divisions usually recognized, as defined by Berkeley:—

1. Hymenomycetes (ἡμῆν, a membrane, and μύκης, a fungus): Hymenium naked, spores in sets of four (fig. 784 *b*), and borne on distinct sporophores (figs. 783, 785).
2. Gasteromycetes (γαστήρ, a belly): Hymenium enclosed in a membrane (peridium), spores as in section 1 (figs. 786, 787).
3. Coniomycetes (κόνις, powder): Flocci of the fruit obsolete or mere peduncles, spores single, often partitioned, and on more or less distinct sporophores.
4. Hyphomycetes (ὑφάω, I weave): Thallus floccose, spores naked, often septate.
5. Ascomycetes (ἀσκή, a bag): Sporidia (spores) contained often in sets of eight in asci or tubes.
6. Physomycetes (φύσα, a bladder): Thallus floccose, spores surrounded by a vesicular veil or sporangium.

Under these sections Berkeley enumerates 598 genera, including about 4,000 species. *Examples*—*Agaricus*, *Polyporus*, *Hydnum*, *Clavaria*; *Phalus*, *Geaster*, *Bovista*, *Craterium*, *Nidularia*; *Bactridium*, *Torula*, *Uredo*, *Æcidium*; *Ceratium*, *Tubercularia*, *Botrytis*, *Penicillium*; *Helvella*, *Peziza*, *Tuber*, *Erysiphe*, *Onygena*; *Phycomyces*, *Mucor*.

1124. The plants of this order deserve attention, whether we regard their esculent or their poisonous qualities, or the destruction which they cause by their parasitic growth. In this country the chief species eaten are *Agaricus campestris*, the common Mushroom, *Agaricus Georgii*, *Morchella esculenta* and other species of Morel, *Tuber cibarium*, Truffle. In foreign countries, as in France, Italy, Germany, and Russia, numerous Fungi are used as food, which have acted as poisons in this country. The process of cooking, as well as the climate, may have some effect in modifying their qualities. *Agaricus procerus* is eaten abroad; but a case of poisoning from it has been known to occur in Edinburgh. In Rome it is stated that the yearly average of taxed mushrooms, from 1837 to 1847, was between 60 and 80,000 pounds weight. The finest mushroom is said to be the *Agaricus Prunulus*. *Amanita muscaria* is a poisonous species, which is used as a means of intoxication in Kamtschatka. It is said to give this property to the urine of those who eat it. It is not easy to distinguish between edible and poisonous Fungi. It has been said that the latter are often highly coloured, have scales or spots on their surface, tough watery flesh, and grow in clusters on wet ground, and often in the shade; while the former are seldom highly coloured, generally white or brownish, rarely show scales or spots, have brittle flesh, and grow



solitary in dry pastures, not in the shade. In some cases Fungi form a staple article of food. Darwich states that the inhabitants of Terra del Fuego live upon a globular fungus of a bright yellow colour (*Cyttaria Darwinii*), found on the bark of the beech. Many species of *Boletus* are used as food in Western Australia, according to Drummond. *Mytila australis* is known in Australia as Native Bread.

1125. Some Fungi are limited to certain kinds of decaying matter. Many species of *Onygena* are found only on the dung, feathers, and hoofs of particular animals. Peculiar species of *Mycoderma* are developed in vinegar, in yeast, and in flour. The rapidity with which Fungi sometimes grow is remarkable. Ward noticed *Phallus impudicus* shoot up three inches in the course of twenty-five minutes, and attain its full elevation of four inches in an hour and a-half. *Bovista gigantea*, in a single night, has increased from the size of a pea to that of a melon. The force also with which they expand has been shown by their raising pavements under which they had been developed. Some Fungi, as *Agaricus oreades*, *coccineus*, and *personatus*, are developed in a centrifugal manner, forming fairy rings. Certain species of *Agaricus* give out a sort of phosphorescent light. This has been remarked in *Agaricus olearius*, *Agaricus Gardneri*, and some species of *Agaric* from the Swan River. A similar kind of light is produced by species of *Rhizomorpha*, which occur in coal mines. *Polyporus fomentarius* forms amadou, and it, as well as *P. betulinus*, have been made into razor-straps.

1126. The diseases caused by Fungi are numerous (§ 693-696). Blight, mildew, rust, and smut, are diseases of grain due to the attacks of Fungi. Dry-rot is owing to the presence of *Merulius lacrymans* and *vastator*, and *Polyporus destructor*, the mode of preventing which has been already alluded to (§ 699). The disease called ergot, which attacks Rye and other grasses, is produced by the *Spermoedia Clavus* (§ 1103). The various moulds which occur on bread, cheese, preserves, and fruits, are plants of this extensive order. *Penicillium glaucum* is one of the most common moulds, occurring on organic substances, on books, &c. A species of *Racodium* is found in low cellars, as at the London docks. Some Fungi are produced on living animals. Thus, the disease called muscardine in the silkworm is produced by *Botrytis Bassiana*. Certain wasps in the West Indies are affected by a similar disease. *Sphæria sinensis*, a celebrated Chinese drug, grows from a caterpillar; *Sphæria Robertsii* is developed on the larva of *Hepialus virescens* in New Zealand; and *Sphæria Taylora* on an Australian caterpillar. So are also *Sphæria sobolifera*, *entomorrhiza*, *militaris*, and others. Particular kinds of mould sometimes grow on the mucous membrane of birds. Some mycodermatous Fungi are connected with certain cutaneous and other diseases in the human species. Thus, cellular filaments called Porrigophytes are found in the crusts of

Porrigo fuvosa, Mentagrophytes in those of *Mentagra* or *Sycosis menti*, and Aphthaphytes in *Aphthæ*.

1127. Order 220.—**Algæ**, the Sea-weed Family. Cellular plants found both in salt and in fresh water. Fronds composed of variously formed, often elongated cells, which are either simple or branched filaments, continuous or articulated, separate, or combined in different ways (fig. 29) so as to constitute fronds of various kinds (fig. 788). Growth takes place by the division of cells, or by cellular prolongations, in the form of lateral branches. Reproductive organs consist of spores, which are contained in mother-cells or perispores (περι, around,



and σπορά, seed), or sporocarps (καρπός, fruit). These are sometimes congregated together in receptacles of different sorts (figs. 788 *c c*, 789). The spores occasionally divide into 3 or 4 cells, constituting tetraspores (τετρας, four). In addition to spores or sporocarps (fig. 791 *s p*), there are sometimes round, or clavate, or filamentous cellular bodies present, to which some give the name of antheridia (fig. 791 *f*). In some of the simplest Algæ, the whole plant is concerned in producing new individuals by division of the parent cells into 2 or 4. In others there is a union of 2 filaments, and a passage of certain granular particles (endochrome) from the one to the other, ending in the formation of the spore. This process is termed *conjugation*, and is one of great

Figs. 788-792.—Frond and organs of reproduction of *Fucus serratus*, to illustrate the natural order Algæ.

Fig. 788.—The entire plant much diminished in size. *f*, Frond composed of cells, so united as to form a flat expansion. *c c*, Conceptacles at the extremities of the frond, containing the organs of reproduction.

Fig. 789.—Extremity of the frond covered with conceptacles.

Fig. 790.—Vertical section of a conceptacle, *c*, with its inner surface covered with spores (sporocarps), and paraphyses, or antheridia. *t*, The superficial cellular tissue of the frond, in which the conceptacle is buried. *o*, Foramen by which the conceptacle opens externally.

Fig. 791.—Spore, *s p*, covered with its perispore or sporocarp, *p*. *f*, Filaments or paraphyses, by some called antheridia.

Fig. 792.—Spore, *s*, separated and deprived of its perispore or outer covering.

interest. It has been observed in some of the *Conservaceæ* and *Diatomaceæ*. In certain cases, the terminal cell of the filament is that in which a spore is formed without any conjugation, and in these cases the spore is frequently provided with ciliary processes, which exhibit for a time spontaneous movements (figs. 431-434); hence called zoospores. In the higher Algæ, the sporocarps containing 2, 4, or more reproductive cellules, are united together in conceptacles along with filaments containing phytozoa, and called antheridia (figs. 790, 791). In *Characeæ* there are two distinct organs of reproduction.

1128. This extensive order has been divided into the following sub-orders:—

1. *Characeæ*: water plants formed of parallel tubes, which are sometimes encrusted with carbonate of lime; reproductive organs are of two kinds—*a*, a round red globule consisting of eight valves which enclose cells of different kinds, containing granular matter and peculiar spiral filaments or phytozoa (fig. 369); *b*, an oval nucule formed by a large central cell or spore, with five elongated cells wound spirally round it, surmounted by five teeth. Some consider the globule as an antheridium, and as equivalent to an anther. The nucule contains the germinating body. The rotation of granules takes place in the plants of the tribe (§ 275, 276).
2. *Fucaceæ*, the Sea-wrack tribe: usually growing in salt water; frond consisting of cells which are often united by gelatinous matter (fig. 29), and which sometimes form a broad expansion (a membranous thallus), supported on a stalk; organs of reproduction consist of sporocarps and antheridia, contained in conceptacles opening externally (fig. 790), which are united in club-shaped expansions or receptacles, situated at the end or margins of the fronds (figs. 788, 789). In germinating, the nucleus bursts the epispore or outer covering of the spore, and sends out filamentous processes.
3. *Floridæ* or *Ceramiaceæ*: rose or purple-coloured sea-weeds, with fronds formed of a single row of articulated cells, or of several rows of cells combined into a flat expansion; organs of reproduction consist of sporocarps or perispores, intermingled with clavate filaments called antheridia. The sporocarps contain cells or spores often divided into four (tetraspores), and enclosed in conceptacles of various kinds.
4. *Conservaceæ*: aquatic plants, often of a green colour, consisting of one or more cells of a rounded or cylindrical form, united together so as to form an articulated or flat frond. They increase by the merismatic division of cells. Reproduction effected by spores which are formed in the interior of the cells by a change in the arrangement of the granular matter, or by the union of filaments of different plants, a process of conjugation, by which granular matter passes from one to the other. The spores are discharged by the opening of the cells, and they often have moving cilia (figs. 431-434).
5. *Diatomaceæ*: inhabiting still waters and moist places; fronds consisting of frustula or fragments, which are either angular or cylindrical, often siliceous and brittle (non-siliceous in *Desmidiæ*), united by a gelatinous sort of substance; propagated by the division of parent cells into two halves, which become more or less completely detached, and form new individuals. Conjugation also takes place in some instances, in the same way as in the *Conservaceæ*.

The order Algæ, exclusive of Characeæ, has been divided by Harvey as follows:—

1. Melanospermæ: plants of an olive-green or olive-brown colour, and cellular or filamentous structure; growing in the sea. Fructification contained in definite capsules or receptacles, or in distinct sori. Seeds dark-coloured.
2. Rhodospermæ: plants usually marine, of a rose-red, purple, or red-brown colour, leafy, cylindrical, or filamentous. Fructification usually double; the primary contained in capsules, receptacles, or immersed in the frond; secondary, when present, minute granules forming sori, or embedded in distinct receptacles. Seeds red or red-brown.
3. Chlorospermæ: plants growing either in the sea, in fresh water, or in damp situations; either filamentous, membranaceous, or shapeless; either colourless, or (owing to the presence of an internal, granular, sporular mass) of a grass-green, very rarely purple or red colour. Fructification, green or purple sporules, either filling the frond or collected into sporidia, rarely contained in external capsules.
4. Diatomaceæ: as already defined above.

Lindley enumerates 283 genera, including about 2,000 species. *Examples*—Chara, Nitella; Fucus, Sargassum, Laminaria, Padina, Ectocarpus, Bryopsis; Ceramium, Delesseria, Rhodymenia, Chondrus; Conferva, Ulva, Oscillatoria, Palmella, Protococcus; Diatoma, Homœocladia, Desmidiium.

1129. The plants of the order are widely distributed over the globe, being found in salt and fresh water, in moist places, as on damp rocks and stones, and the glass and pots of hothouses, and even in hot springs. Sometimes they present collectively the appearance of green slime. They derive nourishment chiefly from the medium in which they grow; and the root-like processes with which some of them are provided, seem to be merely for the purpose of fixing them. Some of the species are very gigantic, others very minute, requiring the aid of the microscope for their detection. The lowest members of the order approach very nearly to the lowest tribes of animals, and it is difficult to draw a line of demarcation. Many species now considered vegetable, such as *Corallina officinalis* and many *Diatomaceæ*, are figured as animals by Ehrenberg. There are interesting movements connected with the cells of many Algæ, such as *Oscillatoria* and *Nostoc*. Some of the species found in the ocean have conspicuous stems, which sometimes present the appearance of zones in their interior (§ 106). Among the large-stemmed species may be noticed *Durvillea utilis* and *Lessonia fuscescens*. *Scytosiphon* (*Chorda*) *Filum* attains in the British seas a length of 30 or 40 feet, while *Macrocystis pyrifera* in the Pacific ocean reaches the length of 500 to 1,500 feet. Some of the *Laminarias* of Britain have stalks of considerable size. *Sargassum bacciferum*, the Gulf-weed, is found floating in great quantities on each side of the equator in the Atlantic, Pacific, and Indian oceans. *Protococcus nivalis* and *viridis* are said to occur in red and

green snow. The red and green colours of certain lakes and seas are attributed to species of *Trichodesmium* and *Sphærozyga*. According to Dr. Joseph Hooker, *Diatomaceæ* are found in countless numbers between the parallels of 60° and 80° S., where they give a colour to the sea, and also to the icebergs floating on it. The death of these bodies in the South Arctic ocean is producing a submarine deposit, consisting entirely of the siliceous particles which entered into the composition of these plants. *Conferva crispa*, called Water-flannel, forms beds of entangled filaments on the surface of water. Species of *Tyndaridea* also occur in thick green patches. *Hydrodictyon utriculatum*, Water-net, has the appearance of a green net, composed of filaments which enclose pentagonal and hexagonal spaces. *Achlya prolifera*, and other *Confervæ*, are developed occasionally on living animals, such as on the gills of the gold-fish and of trout. Certain organisms have been detected in the human stomach, which appear to belong to this order. One of these is called *Sarcinula ventriculi* by Goodsir, and was ejected by vomiting in a case of pyrosis. It consists of square cells united together in sets of four, and propagating by division.

1130. The plants of this order supply a quantity of gelatinous matter, and many of them are used for food. Kelp is obtained by the burning of Sea-weeds, and iodine is procured from them. *Sphærococcus* (*Chondrus*) *crispus*, Carrageen or Irish Moss, supplies a nutritious article of diet. *Rhodymenia palmata*, Dulse, *Alaria esculenta*, *Iridaea edulis*, young plants of *Laminaria digitata* and *saccharina*, Tangle, as well as various species of *Porphyra*, Laver, and *Ulva*, Green Laver, are esculent. The edible swallows'-nests of the East are said to be formed of a species of *Galidium*. *Sphærococcus cartilagineus*, var. *setaceus*, is used in China as a substitute for these nests. Agar-agar is a sea-weed of a similar kind. *Nostoc edule* is used in China as an article of food. The use of burnt sea-weed, in cases of scrofulous swellings, has been superseded by the discovery of iodine, the active ingredient. *Plocaria* (*Gigartina*) *Helminthocorton*, under the name of Corsican Moss, was formerly used as a vermifuge. The *Charas* have frequently a peculiarly fetid odour, and their presence is said to give rise to malaria. Occasionally they communicate their odour to the water of reservoirs, and render it unpleasant. It is of importance for Water Companies to see that *Charas* do not exist in the streams which supply the water for their reservoirs.

For full details on the subject of Classification, the arrangement and characters of natural orders, and the properties of plants, the student should consult Lindley's Vegetable Kingdom.

PART III.

GEOGRAPHICAL BOTANY, OR THE DISTRIBUTION OF PLANTS OVER THE GLOBE.

1131. THIS department of Botany treats of the manner in which plants are affected by climate and station, and endeavours to investigate the conditions under which particular families, or species of plants, are confined to certain zones of latitude and altitude. It is a subject of great interest, and one which cannot be prosecuted with success until the vegetation of the globe is more fully known. So long as there are vast tracts of continents unexplored by botanical travellers, the facts upon which Botanical Geography is founded must be imperfect.

I.—EPIRRHEOLOGY, OR THE INFLUENCE OF VARIOUS EXTERNAL AGENTS ON PLANTS.

1132. It is a matter of common observation, that the localities and soils in which plants grow vary much. Thus, some species grow in the shade, while others thrive best in full exposure to light; some grow in mountainous or alpine districts, while others prefer the plains; some are found in dry, others in marshy places; some are submersed in lakes or in the sea; while others live on muddy banks, or on sandy shores. The plants growing on a granitic or micaceous soil differ frequently from those found on trap, limestone, or sandstone. It is equally well known that climate exercises a powerful influence on vegetation, modifying the Floras in different regions of the globe. Some plants are fitted to bear the rigour and duration of an arctic winter, with a moderate summer heat, others require the heat and light of the torrid zone; and between these two extremes, there are all varieties of gradation. Thus vegetation extends over the whole globe, from

one pole to the other—from the summit of the highest mountains to the bosom of the ocean. There are, notwithstanding this general diffusion of vegetable structures, a few spots in which their traces have not been detected. Among such districts may be mentioned the hot sands of Africa, and some of the Antarctic Islands. In other districts their number and species are limited to a few. Each zone may be said to have its own peculiar vegetable features, the number of species being found to increase as we approach the equator, and to decrease as we retire from it. Palms, Bananas, Tree Ferns, and Orchideous Epiphytes, are chiefly confined to the tropics; Cruciferous and Umbelliferous plants are found in temperate regions; some Coniferous and Amentaceous plants flourish in more northern countries, while Saxifrages and Lichens extend to the arctic regions. In warm regions are found those fruits which are so necessary for the well-being of the inhabitants; in temperate climates chiefly, occur the cereal grains for the food of man, and the green pastures for the nourishment of cattle; and in the arctic regions, the Lichen, on which the reindeer feeds, grows luxuriantly.

1133. The number of known species of plants amounts to about 100,000. The following is an estimate of the known species of plants on the globe at different dates:—

Linnaeus.....	1753	5,323	Phanerog.	615	Cryptog.	5,938
Persoon.....	1807	19,949	—	6,000	—	25,949
Steudel.....	1824	39,684	—	10,965	—	50,649
Steudel.....	1841	78,000	—	13,000	—	91,000
Steudel.....	1844	80,000	—	15,000	—	95,000

In 1846, Lindley gave the following estimate of known genera and species:—

Thallogens.....	939	Genera,	8,394	Species.
Acrogens.....	310	—	4,086	—
Rhizogens.....	21	—	53	—
Endogens.....	1,420	—	13,684	—
Dictyogens.....	17	—	268	—
Gymnogens.....	37	—	210	—
Exogens.....	6,191	—	66,225	—
Total.....	8,935		92,920	

Much yet remains to be done in regard to the Floras of India, China, Africa, New Holland, and South America. Meyen conjectures that the total vegetation of the globe may be about 200,000 species.

1134. The distribution of species over different quarters of the globe is regulated by various external agents, the study of which is termed Epirrheology (*ἐπιρρῆσις*, I influence). These agents are chiefly temperature and moisture, and the nature of the soil. The effects produced on plants by increase or decrease of light, and by changes in the state of the atmosphere, have not been sufficiently determined.

1.—EFFECTS OF TEMPERATURE.

1135. The effects of this agent must be considered both as regards its latitudinal and its altitudinal ranges. In proceeding from the equator to the poles, or in ascending from the surface of the ocean to the summit of a lofty mountain, there is a gradual decrease of temperature, and, at the same time, marked changes in the nature of the vegetation. The scale of atmospherical temperature serves as a scale for the progress of vegetation. As regards the latitudinal distribution of heat, the globe has been divided into eight regions, four northern and four southern—viz., a tropical region, from the equator to the limits of the tropics in each hemisphere; subtropical, between this and 40° of latitude; temperate, between 40° and 60° of latitude; arctic and antarctic, beyond 60° of latitude.

1136. Each species of plant is adapted to thrive best between certain limits of temperature. These limits do not necessarily coincide with any definite parallels of latitude; for it is well known that the climate of different places in the same latitude is very different. It is of importance, therefore, to ascertain the mean temperature of the year, or rather of different seasons. By drawing lines through different places where the mean annual temperature is the same, Humboldt has established a series of isothermal (*ἴσος*, equal, and *θερμῆν*, heat,) lines intersecting the parallels of latitude. These lines run in curves, which rise in their course from the eastern coast of America towards western Europe, and sink towards the south in the interior of the continent,—and that so quickly, that Scotland lies in the same isothermal line as Poland, and England as Hungary. It is clear, therefore, that the isothermal lines in the higher latitudes do not, by any means, correspond with the parallels of latitude. At the equator, however, these lines coincide more nearly. Much depends upon the temperature of the different seasons. Thus, a place which has a very cold winter, and a very warm summer, may be in the same isothermal line with one in which the temperature of both these seasons is moderate, and plants which succeed well in the one may not grow in the other. Cherry-laurels and other Evergreens, which grow well in the open air in England, will not stand the winter of places on the continent in the same isothermal line. It is necessary, in determining the geographical distribution of plants, to take into account the mean summer and the mean winter heat, and, better still, the mean monthly temperature. The distribution of temperature among the different months of the year is of importance, especially in reference to the heat and duration of the summer months; for many plants protected by a covering of snow, are enabled to brave rigorous winters, provided the summer be hot enough, and of sufficient duration. Lines passing through places having the same mean summer temperature are called isothermal (*ἴσος*, equal, and *θερος*, summer); those passing through places

with an equal mean winter temperature, are isotheimonal (*χρειμών*, winter). The isotheimonal lines in the interior of continents bend considerably towards the south. The isothermal lines bend to the north, but in the interior of continents they approach the parallels of latitude. Many circumstances conspire to influence the temperature of countries. Insular and coast climates are more equable, from the effect of the sea in preventing the atmosphere from being much heated during the day, and much cooled during night. In the interior of vast continents the extremes of temperature are often great. Winds have a powerful effect on climate. In China, the north-east monsoon brings great cold in February. The state of a country as regards forests, has a decided effect on the temperature. In different quarters, the nature of the exposure, also, whether to the east or west, north or south, and the intervention of elevated ranges of mountains, materially affect the temperature.

1137. Temperature, in its hypsometrical (*ὑψος*, altitude, and *μέτρον*, measure,) relation, or as regards its altitudinal range, requires to be considered. In ascending into the atmosphere, a decrease of temperature is observed, which varies in its amount at different stages of ascent. The following table shows the temperature at different heights in the equatorial and temperate zones:—

Height in feet.	Equat. Zone. Lat. 0°—10°	Mean.	Temp. Zone. Lat. 45°—47°
0	81°·50 F.		53°·60 F.
3,197	71 ·24	41 ·00
6,394	64 ·40	31 ·64
9,591	57 ·54	23 ·36
12,789	44 ·60		
15,985	37 ·70		

Taking an average, it may be said that there is a fall of 1° in the thermometer for every 340 feet of ascent. Prof. Forbes states that 549·5 feet of ascent give a difference of 1° of the thermometer in the boiling point of water. The elevation at which constant frost takes place is called the snow-line or line of perpetual congelation. Its limit does not exactly correspond with the height at which the temperature is equal to 32° F. The following table gives the height of the snow-line (in feet) at different latitudes:—

Lat.	Height of Snow Line.	Lat.	Height of Snow Line.
0	15,207	50	6,334
5	15,095	55	5,034
10	14,764	60	3,818
15	14,220	65	2,722
20	13,278	70	1,778
25	12,557	75	1,016
30	11,484	80	457
35	10,287	85	117
40	9,001	90	0
45	7,671		

The decrease of temperature on lofty mountains regulates, in a great measure, the nature of the plants which grow at different heights. The same changes take place as have been shown to occur in proceeding from the equator to the poles. The following observations made on the growth of certain trees on the Grimsel, show the relation between height and latitude :—

	Lat.	On the Grimsel.
Beech, which extends to	60°	grows about the height of 3000 feet.
Oak.....	61° 2600 —
Fruit Trees.....	63° 3400 —
Hazel.....	64° 5000 —
Norway Spruce.....	67° 40' 6000 —
Scotch Fir.....	70° 6400 —
Birch.....	70° 40'

2.—EFFECTS OF MOISTURE.

1138. The quantity of moisture in the air has a decided effect on the distribution of plants. Nothing checks vegetation more than extreme dryness. Hence the barrenness of those hot sandy deserts, which exhibit only an arid waste, without a single blade of grass to relieve the eye of the weary traveller. In warm and dry climates, succulent plants occur, with hard epidermal coverings, capable of resisting the effects of evaporation and transpiration. Among these may be noticed Cactaceæ, Mesembryaceæ, Euphorbias, and some of the Aloe tribe. In the districts of Australia, where a dry climate prevails, many plants, such as Proteas, Banksias, and leafless Acacias, have hard and dry foliage, capable of enduring much drought without injury. In warm climates, the effect of the dry season on vegetation is very remarkable. This season may be said to correspond with our winters. In some parts of South America, where no rain falls for eight months of the year, the leaves during the dry season fall, buds are developed in their axils, and it is only when the wet season arrives that the trees become clothed with verdure, and the herbage appears. Forests appear to keep up the humidity of the atmosphere in a country, and thus have a powerful influence on the climate.

3.—EFFECTS OF SOIL, LIGHT, AND OTHER AGENTS.

1139. The physical localities in which plants grow vary considerably. These variations are connected with the dryness and moisture of the soil, as well as with its mechanical and chemical composition. Some plants are fitted to grow in water, others in marshes; some grow in peaty soil, others in sandy soil. Thurmann has endeavoured to show that the nature of the soil, whether siliceous, clayey, calcareous, or saline, has an effect in modifying the vegetation. Prof. E. Forbes states, that, in Syria, he could easily distinguish the serpentine from

the limestone, not merely by their geological characters, but also by the disposition of the arborescent vegetation. On the serpentine, usually pines only grew, and never in thick forest masses, but scattered; whereas the limestone bore thick clustered oaks and a luxuriant underwood, with now and then clumps of lofty pines. In the low countries near the sea, the serpentine was marked by *Senecio squalidus*, a little *Erophila*, and *Cheilanthes odora*; while on the limestone, *Acrostichum lanuginosum* was a conspicuous fern.

The following is a division of plants according to the botanical stations or physical localities in which they grow, whether placed there by nature or by art. :—

A.—Plants growing in Water, whether Salt or Fresh.

1. Marine Plants, such as Sea-weeds, Lavers, &c., which are either buried in the ocean, or float on its surface: also, such plants as *Ruppia* and *Zostera*. In the Sargasso Sea there are floating meadows of *Sargassum bacciferum*, gulf-weed. This sea extends from 22° to 36° north lat., and from 25° to 45° west long. from Greenwich, and extends over 40,000 square miles.

2. Maritime or saline plants. These are plants which grow on the border of the sea, or of salt lakes, and require salt for nourishment, as *Salicornia*, glasswort, *Salsola*, saltwort, *Anabasis*. Such plants are often called Halophytes (ἅλα, salt, and φυτὸν, a plant). Under this head may be included littoral and shore plants, such as *Armeria*, sea-pink, *Glaux*, and *Samolus*.

3. Aquatic plants, growing in fresh water, either stagnant or running; as *Sagittaria*, arrowhead, *Nymphaea*, water-lily, *Potamogeton*, pondweed, *Subularia*, awlwort, *Utricularia*, bladderwort, *Stratiotes*, water-soldier, *Lemna*, duckweed, *Pistia*, *Conservea*, *Oscillatoria*, and *Ranunculus fluitans*. Some of these root in the soil, and appear above the surface of the water; others root in the soil and remain submersed; while a few swim freely on the surface without rooting below.

4. Amphibious plants, living in ground which is generally submerged, but occasionally dry, as *Ranunculus aquatilis* and *sceleratus*, *Polygonum amphibium*, *Nasturtium amphibium*. The form of the plants varies according to the degree of moisture. Some of these, as *Limosella aquatica*, grow in places which are inundated at certain periods of the year; others, such as *Rhizophoras* (mangroves), and *Avicennias*, form forests at the mouths of muddy rivers in tropical countries.

B.—Land Plants which root in the Earth and Grow in the Atmosphere.

5. Sand plants; as *Carex arenaria*, *Ammophila arenaria*, *Elymus arenarius*, and *Calamagrostis arenaria*, which tend to fix the loose sand, *Plantago arenaria*, *Herniaria, glabra*, *Sedum acre*.

6. Chalk plants; plants growing in calcareous and cretaceous soils, as some species of *Ophrys*, *Orchis*, and *Cypripedium*.

7. Meadow and pasture plants; as some species of *Lotus*, bird's-foot trefoil, a great number of grasses and trefoils, the daisy, dandelion, and buttercups.

8. Plants found in cultivated ground. In this division are included many plants which have been introduced by man along with grain; as *Centaurea Cyanus*, corn blue-bottle, *Sinapis arvensis*, common wild mustard, *Agrostemma*, corn-cockle, several species of *Veronica* and *Euphorbia*, *Lolium temulentum*, *Convolvulus arvensis*, *Cichorium Intybus*; also plants growing in fallow ground, as *Rumex Acetosella*, *Carduus*

nutans, *Echium vulgare*, *Artemisia campestris*, and *Androsace septentrionalis*. In this division, garden weeds are included; such as Groundsel, Chickweed, *Lamium amplexicaule*, *Chenopodium album*, and *urbicum*.

9. Rock or wall plants; Saxifrages, Wall-flower, *Linaria Cymbalaria*, *Draba muralis*, species of *Hieracium* and *Sedum*, *Asplenium Ruta muraria*, and some lichens and mosses.

10. Plants found on rubbish heaps, especially connected with old buildings. Some of these seem to select the habitations of man and animals, on account of certain nitrogenous and inorganic matters, which enter into their composition. Among them may be noticed, Nettles, Docks, Borage, Henbane, *Xanthium*. Here, also, have been placed some plants immediately connected with the habitation of man, such as *Racodium cellare*, a fungus found on wine casks, *Conferva fenestralis*, an alga produced on window-panes, and *Conferva dendrita*, one developed on paper. Some plants, as *Sempervivum tectorum*, select the roofs of houses.

11. Plants growing in vegetable mould; such as bog-plants, or those growing on wet soil, so soft that it yields to the foot, but rises again; and marsh plants, growing in wet soil, which sinks under the foot and does not rise. To the former class belong such plants as *Pinguicula alpina* and *Primula farinosa*; to the latter, such as *Menyanthes*, *Comarum*, *Bidens cernua*.

12. Forest plants, including trees which live in society, as the Oak, the Beech, Firs, &c., and the plants which grow under their shelter, as the greater part of the European Orchises, some species of *Carex* and *Orobanche*. Some plants especially grow in pine and fir-woods, as *Linnæa borealis*, and some *Pyrolas*.

13. Plants of sterile places, found in barren tracts, by roadsides. This is a heterogeneous class, and contains many plants of uncertain characters. Under it are included the plants of uncultivated grounds, as those found in moors, where *Calluna vulgaris*, common heather and various heaths, Juniper, *Andromeda*, and some species of *Polytrichum* occur.

14. Plants of the thickets or hedges, comprehending the small shrubs which constitute the hedge or thicket, as the Hawthorn and Sweet-briar; and the herbaceous plants which grow at the foot of these shrubs, as *Adoxa*, Wood-sorrel, Violets; and those which climb among their numerous branches, as Bryony, Black Bryony, Honeysuckle, Traveller's Joy, and some species of *Lathyrus*.

15. Plants of the mountains, which De Candolle proposes to divide into two sections: 1. Those which grow in alpine mountains, the summits of which are covered with perpetual snow, and where, during the heat of summer, there is a continued and abundant flow of moisture, as numerous Saxifrages, Gentians, Primroses, and Rhododendrons. 2. Those inhabiting mountains on which the snow disappears during summer, as several species of Snap-dragon, among others the Alpine Snap-dragon, Umbelliferous plants, chiefly belonging to the genus *Seseli*, meadow Saxifrage, Labiate plants, &c.

C.—Plants Growing in Special Localities.

16. Parasitic plants, which derive their nourishment from other vegetables, and which, consequently, may be found in all the preceding situations; as the Mistletoe, species of *Orobanche*, *Cuscuta*, (Dodder), *Loranthus*, *Rafflesia*, and numerous Fungi.

17. Pseudo-parasitic plants, or Epiphytes, which live upon dead vegetables, as Lichens, Mosses, &c. or upon the bark of living vegetables, but do not derive much nourishment from them, as *Epidendrum*, *Aerides*, and other Orchids, as well as *Tillandsia*, *Bromelia*, *Pothos*, and other air-plants.

18. Subterranean plants, or those which live under ground, or in mines and caves, almost entirely excluded from the light, as *Byssus*, Truffles, and some other Cryptogamic plants.

19. Plants which vegetate in hot springs, the temperature of which ranges from 80° to 150° of Fahrenheit's thermometer ; as *Vitex Agnus-castus*, and several Cryptogamous plants, as *Ulea thermalis*, the hot-spring Laver.

20. Plants which are developed in artificial infusions or liquors, as various kinds of *Mucor*, causing mouldiness.

21. Plants growing on living animals ; as species of *Sphaeria* and *Sarcinula* and various other Fungi and Algæ.

22. Plants growing on certain kinds of decaying animal matter ; such as species of *Onygena*, found on the hoofs of horses, feathers of birds, &c., some species of Fungi, which grow only on the dung of animals, and certain species of *Splachnum*.

1140. Light is an agent that has a powerful influence on plants, as regards their vigour, irritability, secretions, and colour. Hence, in those regions where the light is intense, the vegetation presents certain peculiarities. The luxuriance and greenness of the leaves, the nature of the woody matter deposited, of the fruit produced, and of the secretions formed, are all influenced in some degree by the intensity of the sun's rays. Little is known in regard to the effects of increased or diminished atmospheric pressure on plants. Humboldt believed that vegetation was influenced by the amount of atmospheric pressure. Further research has not corroborated his suppositions. The effects of the atmosphere have been studied chiefly as regards dryness and moisture, and the mixture of certain gases with it, especially in the vicinity of manufacturing towns (§ 297).

1141. The effects of climate and season on the leafing, flowering, and fruiting of plants, may be seen in the case of some species which are found distributed over various countries in Europe. Berghaus has made an extensive series of observations on the subject. The Lilac (*Syringa vulgaris*), according to him, unfolds its leaves at Naples, in latitude 41°, during the first half of the month of January ; near Paris, in latitude 49°, on the 12th March. The Elder unfolds its leaves

At Naples.....	January 1-15.
At Paris.....	February 14.
In England.....	March 8.
At Upsal.....	March 1-8.

The Beech unfolds its leaves

At Naples.....	End of March.
In England.....	1st May.
At Upsal.....	Beginning of May.

In regard to flowering, Berghaus states, that in the middle latitudes of Europe and North America, it is generally four days later for each degree of latitude towards the north. The same plants flower at Zurich 6 days later than at Parma ; at Tubingen, 13 days later ; at Jena, 17 ; at Berlin, 25 ; at Hamburgh, 33 ; at Greifswald, 36 ; and at Christiania, no less than 52 days later than at Parma. In the Berlin

district, an elevation of 1,000 feet renders vegetation 10 to 14 days later: so also in regard to fruiting. The wheat harvest begins

At Naples.....	In June.
In Central Germany	July.
In the South of England.....	August

Ripe Cherries are to be had

At Naples.....	First days of May.
At Paris.....	End of June.
In Central Germany.....	do.
In the South of England.....	22d July.

II.—DISSEMINATION OF PLANTS.

1.—AGENTS EMPLOYED IN THEIR DISSEMINATION.

1142. Some plants are disseminated generally over the globe, while others are confined within narrow limits. Some of the common weeds in Britain, such as Chickweed, Shepherd's-purse, and Groundsel, are found at the southern extremity of South America. *Lemna minor* and *trisluca*, *Convolvulus sepium*, *Phragmites communis*, *Cladium*, *Mariscus*, *Scirpus lacustris*, *Juncus effusus*, and *Solanum nigrum*, are stated by Meyen to be common to Great Britain and New Holland. *Nasturtium officinale*, and *Samolus Valerandi*, are very extensively diffused, and they may be reckoned true cosmopolites. They are both natives of Europe, and they occur, the former near Rio Janeiro, the latter at St. Vincent. The lower the degree of development, the greater seems to be the range. Some Cryptogamic plants, as *Lecanora subfusca*, are found all over the world. Man has been instrumental in diffusing widely culinary vegetables, such as the potato, and the cereal grains, as well as many other plants useful for food and manufacture. Corn plants, such as Barley, Oats, Rye, Wheat, Spelt, Rice, Maize, and Millet, are so generally cultivated over the globe, that almost all trace is lost of their native country. They can arrive at perfection in a great variety of circumstances, and they have thus probably a wider geographical range than any other kind of plant. As regards these plants, the globe may be divided into five grand regions—the region of Rice, which may be said to support the greatest number of the human race; the region of Maize; of Wheat; of Rye; and lastly, of Barley and Oats. The first three are the most extensive, and Maize has the greatest range of temperature. The grains extending farthest north in Europe are Barley and Oats. Rye is the next, and is the prevailing grain in Sweden and Norway, and all the lands bordering on the Baltic, the north of Germany, and part of Siberia. Wheat follows Rye; it is cultivated in the middle and south of France, England, part of Scotland, part of Germany, Hungary, Crimea, and the Caucasus. We next come to a district where wheat still abounds, but no longer exclusively furnishes bread,

rice and maize becoming frequent. To this zone belong Portugal, Spain, part of France, Italy and Greece, Persia, Northern India, Arabia, Egypt, the Canary islands, &c. Wheat can be reared wherever the mean temperature of the whole year is not under 37° or 39° F., and the mean summer heat, for a period of at least three or four months, is above 55° . It succeeds best on the limits of the subtropical region. In the Scandinavian Peninsula, the cultivation of Barley extends to 70° north latitude, Rye to 67° , and Oats to 65° . The cultivation of Rice prevails in Eastern and Southern Asia, and it is a common article of subsistence in various countries bordering on the Mediterranean. Maize succeeds best in the hottest and dampest parts of tropical climates. It may be reared as far as 40° north and south latitude on the American continent on the western side, while in Europe it can grow even to 50° or 52° of latitude. It is now cultivated in all regions in the tropical and temperate zones, which are colonized by Europeans. Millet of different kinds is met with in the hottest parts of Africa, in the south of Europe, in Asia Minor, and in the East Indies. Henslow gives the following table to show the range of Wheat and Barley, and the mean temperature required for them:—

Lat.	Winter Mean.	Summer Mean.	Annual Mean.	
$62\frac{1}{2}$ Ferøe.....	39°	51°	45°	} Barley.
70 Lapland.....	22	46	33	
$67\frac{1}{2}$ Russia.....	9	46	32	
$57\frac{1}{2}$ Siberia.....	0	60	32	
58 Scotland.....	36	57	46	} Wheat.
64 Norway.....	23	59	39	
62 Sweden.....	23	59	39	
$60\frac{1}{2}$ Russia.....	15	60	37	
30 Cairo.....	57	88	72	
22 Macao.....	64	82	73	}
22 Rio Janeiro.....	68	78	74	
23 Havannah.....	71	82	77	
21 Bourbon.....	71	80	77	

Winds, water, and animals, are also instrumental in disseminating plants. Many seeds with winged and feathery appendages are easily wafted about; others are carried by rivers and streams, and some can be transported by the ocean currents to a great distance, with their germinating powers unimpaired.

2.—GENERAL AND ENDEMIC DISTRIBUTION OF PLANTS.

1143. While some plants are generally diffused, it is found that the different quarters of the globe are each characterized by more or less distinct floras. Europe, Asia, Africa, North America, South America, and Australasia, may be regarded as separate provinces of the vege-

table kingdom, possessing species, genera, and families of plants, which give to each division its distinctive features. Humboldt and Bonpland, in their travels in equinoctial America, did not see an exogenous plant which was found equally in the New and the Old World; the only plants which they discovered common to both being some grasses and sedges. Among 4,160 species met with in New Holland by Brown, 166 only were to be found in Europe.

1144. Some plants live in society, occupying exclusively large tracts of ground, from which they banish all other vegetables. These are called by Humboldt *Social* plants. They give a peculiar feature to the countries and districts in which they grow. To this class belong many species of Seaweed in the ocean; Cladonias and Mosses in the waste levels of Northern Asia; Grasses (Bamboos), and some Cactuses, Mangroves, and Avicennias in tropical countries; Ferns in the South Sea Islands; Banksia speciosa in New Holland; Cinchonas in certain parts of South America; Coniferous trees and Birches in the Baltic and Siberian plains.

1145. Some plants are very much restricted in their distribution over the globe; a few are confined to single localities, while others have a limited longitudinal range. The species of the genus Erica, Heath, which extend from northern regions to the Cape of Good Hope, are scattered over a surface very narrow compared to its length; in other words, while their latitudinal range is great, their longitudinal range is very much restricted. Calceolarias occur chiefly on the western side of the Cordilleras of Chili. Lobelia Dortmanna is found principally in the western countries of Europe. Camellias are also limited in longitudinal direction, so also Phalangium bicolor, and Raymondia pyrenaica. Arbutus Unedo, Erica mediterranea, and Menziesia polifolia, whose chief seat is in the Pyrenees and the mountains of Asturias, migrate in a north-westerly direction, and appear in Ireland. It is said that Azaleas, Rhododendrons, Magnolias, Vacciniums, Actæas, and Oaks, which form prevailing genera on the east of the Rocky Mountains, scarcely appear on the western side. Epacridaceæ are confined to New Holland; Cinnamon, Cloves, and Nutmeg, are the produce of the Indian Archipelago; Gentians and Saxifrages form a characteristic feature of the European Alps; Bejarias and Cinchonas of the Peruvian Cordilleras; Schizanthuses of Chili; Polemoniaceæ of California and Oregon; yellow and brown Papilionaceæ of Australasia; Disa grandiflora is a rare orchid peculiar to Table Mountain at the Cape of Good Hope; and Pringlea antiscorbutica is a cruciferous plant peculiar to Kerguelens-land. It is said that Origanum Tournefortii is found only in a small island in the Grecian Archipelago. The vegetation of islands removed from continents presents often peculiar features, the ocean acting as a barrier to the dissemination of plants. The island of St. Helena was originally inhabited by a most peculiar

vegetation, although its productions now are completely changed by the destruction occasioned by cattle, and by the introduction of European and other plants, especially fruit trees. Such may also be said of the plants found in the Sandwich Islands, the Society Islands, and the Canaries. The island of Madeira has 672 Phanerogamous plants, of which 85 are peculiar to it.

8.—CONJECTURES AS TO THE MODE IN WHICH THE EARTH WAS ORIGINALLY CLOTHED WITH PLANTS.

1146. It is an interesting question to determine the mode in which the various species and tribes of plants were originally scattered over the globe. Various hypotheses have been advanced on the subject. Linnæus entertained the opinion, that there was at first only one primitive centre of vegetation, from which plants were distributed over the globe. Some, avoiding all discussions and difficulties, suppose that plants were produced at first in the localities where they are now seen vegetating. Others think that each species of plant originated in, and was diffused from, a single primitive centre, and that there were numerous such centres situated in different parts of the world, each centre being the seat of a particular number of species; they thus admit great vegetable migrations similar to those of the human races. Those who adopt the latter view, recognize in the distribution of plants some of the last revolutions of our planet, and the action of numerous and varied forces which impede or favour the dissemination of vegetables in the present day. They endeavour to ascertain the primitive flora of countries, and to trace the vegetable migrations which have taken place. Daubeny says, that analogy favours the supposition that each species of plant was originally formed in some particular locality, whence it spread itself gradually over a certain area, rather than that the earth was at once, by the fiat of the Almighty, covered with vegetation in the manner we at present behold it. The human race arose from a single pair, and the distribution of plants and animals over a certain definite area, would seem to imply that the same was the general law. Analogy would lead us to believe, that the extension of species over the earth originally took place on the same plan on which it is conducted at present, when a new island starts up in the midst of the ocean, produced either by a coral reef or a volcano. In these cases the whole surface is not at once overspread with plants, but a gradual progress of vegetation is traced from the accidental introduction of a single seed, perhaps of each species, wafted by winds, or floated by the currents. The remarkable limitation of certain species to single spots on the globe, seems to favour the supposition of specific centres. Professor E. Forbes says, the hypothesis of the descent of all the individuals of a species, either from a first pair or from a single individual, and the consequent theory of specific centres

being assumed, the isolation of assemblages of individuals from their centres, and the existence of *endemic* or very local plants, remain to be accounted for. Natural transport, the agency of the sea, rivers, and winds, and carriage by animals, or through the agency of man, are insufficient means in the majority of cases. It is usual to say, that the presence of many plants is determined by soil or climate, as the case may be; but if such plants be found in areas disconnected from their centres by considerable intervals, some other cause than the mere influence of soil or climate must be sought to account for their presence. This cause he proposes to seek in an ancient connection of the outposts or isolated areas with the original centres, and the subsequent isolation of the former through geological changes and events, especially those dependent on the elevation and depression of land. Selecting the flora of the British islands for a first illustration of this view, Professor Forbes calls attention to the fact, well known to botanists, of certain species of flowering plants being found indigenous in portions of that area, at a great distance from the nearest assemblages of individuals of the same species in countries beyond it. Thus, many plants peculiar in the British flora to the west of Ireland, have the nearest portion of their specific centres in the north-west of Spain; others, confined with us to the south-west promontory of England, are, beyond our shores, found in the Channel Isles and the opposite coast of France; the vegetation of the south-east of England is that of the opposite part of the continent; and the Alpine vegetation of Wales and the Scotch Highlands is intimately related to that of the Norwegian Alps. The great mass of the British flora has its most intimate relations with that of Germany. He believes, therefore, that these isolated outposts were formerly connected together by chains of land, and that they have been separated by certain geological convulsions. Islands may be considered as the remains of mountain chains, part of the flora of which they still exhibit, and the further they are from continents, the more likely are the plants to be peculiar.

1147. All the vegetable productions of the globe are distributed according to harmonious laws, which are by no means fully developed. The greater number of families is distributed over the whole globe; individual representatives of the groups appearing in different regions. The regions of the globe, as regards their vegetable productions, are related either in the families, the genera, or the species of plants which they produce. By *families*, Hinds remarks, the most distant or general resemblances are established, constituting *analogy*. One family may occupy the place of another in certain regions. Thus, the Mesembryaceæ of South Africa are represented in America by Cactaceæ; and in the south of Europe, only by a few species of *Sempervivum* and *Sedum*. By *genera*, a closer approximation is established—that of *affinity*. The *Cistuses* of Spain and Portugal are represented by the

Helianthemum of the north of Europe; and the genera of *Abies* and *Pinus*, in arctic and temperate regions, have their representatives in the genera *Araucaria*, *Ephedra*, and *Dammara* of the south. By *species* again, the most perfect accordance of characters or *identity* is established.

1148. Meyen states, that the species of a genus, and genera, and natural orders, proceed from a point, and range themselves round it in concentric circles, or spread out from it like rays in all directions; or are distributed in belts of greater or less breadth, which are parallel to the meridians, or to the parallels of latitude. A genus or family predominates in certain regions, and attains its maximum there, while in others it is at its minimum. Hence, regions are distinguished by the names of plants which attain their maximum there. *Palmae*, *Musaceae*, *Piperaceae*, and *Scitamineae*, attain their maximum in the torrid zone, although representatives of them extend to high latitudes, or to the temperate zone. Thus, the Palm called *Chamærops humilis* is found in 49° north latitude. The *Ericaceae* of the old world have their maximum in the south of Africa. A single form, *Calluna vulgaris*, common Heather, is predominant in the north; and a shrubby species, *Erica arborea*, represents the order in the south of Europe. *Acacias* attain their maximum in New Holland, while *Acacia heterophylla* represents the family in the Sandwich Islands. The *Lauraceae* of the tropics have *Laurus nobilis* as their representative in Europe.

4.—DISTRIBUTION OF PLANTS CONSIDERED PHYSIOGNOMICALLY AND STATISTICALLY.

1149. The distribution of plants over the globe may be considered either *Physiognomically*, as regards the prevalence of certain vegetable forms which give a general character to the landscape of a country; or *Statistically*, as regards the numerical proportion which different groups bear to each other, or to the whole known plants.

1150. **Physiognomy of Vegetation.**—In prosecuting this department of botanical geography, it is necessary to specify those vegetable forms which give a character to the landscape. This has been done more especially by Meyen, who gives the following series:—

1. Gramineous or Grassy Form. This is illustrated in northern countries by meadows and pastures. The cereal grains also have a great influence on the aspect of countries. Under this form are included *Cyperaceae*, *Restiaceae*, and *Juncaceae*. In the torrid zone some arborescent forms occur, as Bamboo; and along with these are associated Sugar-cane and Rice. Barley is an extra-tropical form, while *Carex* extends to cold regions.

2. Scitamineous Form. This includes the Ginger, Arrow-root, and Plantain family, some of which attain a large size. They contribute to give a character to the torrid zone.

3. Pandanus or Screw-pine Form. A tropical form illustrated by Screw-pines and *Dracenas*.

4. Pine-apple Form. Illustrated by the *Bromeliaceae* of warm climes.

5. The Agave or American Aloe Form. Chiefly tropical and subtropical.
6. The Palm Form. Under this are included also the Cycadaceous family. They give a character to the hotter regions of the globe. Some of the Palms are social, as the Date and Coco-nut. *Chamarops humilis* represents this form in Europe.
7. Filical or Fern Form. True Ferns, in an especial manner, affect the landscape in tropical and warm regions.
8. Mimosa Form. This includes Leguminous plants in general. The finely cut foliage of some has a resemblance to Ferns. Modifications of this form occur both in warm and cold regions. Acacias in New Holland give a peculiar feature to the landscape.
9. Coniferous Form. The Abietinæ are characteristic of northern regions, and the Cupressinæ of southern.
10. The Protea, Epacris, and Erica Forms. These forms supply the place of Coniferæ in the southern hemisphere. The Protea and Epacris forms occurring in Australasia, and the Erica form at the Cape of Good Hope.
11. Myrtle Form. Some of these, such as *Melaleuca* and *Eucalyptus*, characterize New Holland scenery; others, as Guavas, are tropical.
12. Forms of Dicotyledonous trees. Some with broad and tender leaves, as Birch, Alder, Poplar, Oak, Lime, Elm, Beech, and Horse-chestnut, giving a character to the physiognomy of the colder half of temperate climates; while others, with thick, leathery, and showy leaves, as Olives and Laurels, are characteristic of warmer climates; and a third division, with large, beautiful leaves, *Cecropia*, *Artocarpus*, and *Astrapæa*, abound in the hottest climates.
13. Cactus Form. This form is developed chiefly in America, especially in Brazil.
14. Form of Succulent plants. Seen in the Mesembryaceæ of South Africa.
15. Lily Form. This includes Liliaceæ, Amaryllidaceæ, and Iridaceæ. Modifications of this form occur in warm and temperate climates.
16. Forms of Lianas or Climbing-plants. These forms are chiefly tropical, and are illustrated by Passion-flowers, *Paullinias*, *Aristolochias*, and *Bauhinias*.
17. Pothos Form. This is a tropical form, and is illustrated by various species of Araceæ.
18. Orchideous Form. This is seen in the splendid Epiphytes of warm climates. Terrestrial species chiefly occur in cold zones.
19. The Moss Form.
20. The Lichen Form. Both these forms characterize cold regions chiefly.

Besides the forms of plants, it is found that the prevalent colours sometimes give a character to the vegetation. White or pale-coloured flowers are said to be more abundant in northern latitudes than in the tropics, and in alpine situations they are of more frequent occurrence than in the plains. The xanthic series of colours, Hinds states, is abundant within the tropics in the autumn, on the plains over the mountains. The flowers of the cyanic series, especially intense blues and violets, delight in the clear skies of subtropical regions. Hinds gives the following tabular view of the relative proportion of colours :

	Cyanic.	Xanthic.	White.
Central America.....	12	30
Sandwich Islands.....	12	31	7
Alashka.....	26	13	11
California.....	25	19	6
New Guinea.....	12	23	15
Hong-Kong.....	13	27	10

Geyer says, that vivid colours mark the basaltic plains of Upper Oregon; blue and purple, eastward; scarlet with golden-yellow, westward; glaucous green reigns in the herbage over the plains; deep saturated green in the valleys.

1151. Statistics of Vegetation.—The number of known vegetable families differs in different latitudes. In examining the distribution of the great classes of the Vegetable Kingdom, it will be found that certain relative numerical proportions have been ascertained. It is not easy to estimate the proportion which Cryptogamous bear to Phanerogamous plants. From data already given, it may be estimated that the proportion for the whole world is as 1 to 7. This proportion varies in different regions; the Cryptogamous plants increasing in their proportion in the northern parts of the temperate zone. Ferns are to known Phanerogamous plants as 1 to 20. This proportion is least in the middle of the temperate zone, and becomes larger towards the equator, and towards the poles. Ferns, however, attain their absolute maximum at the equator, and their absolute minimum in the arctic zone. At North Cape, there are only four species of Ferns found, and yet their proportion to Phanerogamia is 1 to 7 there; and in Greenland, 1 to 10 (Meyen). Humboldt says, that in the torrid zone, Monocotyledons are to Dicotyledons as 1 to 6: in the temperate zone, as 1 to 4; and in the arctic zone, as 1 to 3. Monocotyledons increase in proportion to Dicotyledons as the latitude becomes higher. Some natural orders are very generally diffused, as Leguminosæ, Malvaceæ, Ranunculaceæ, Caryophyllaceæ, Cruciferae, and Umbelliferae. Cellular plants have also a wide range, and so have aquatics. Juncaceæ, Cyperaceæ, and Gramineæ, increase in proportion to all the Phanerogamous plants, as the latitude becomes higher; while Restiaceæ, Leguminosæ, Euphorbiaceæ, and Malvaceæ, decrease. Cruciferae, Umbelliferae, and Compositæ, are highest in their proportion in the temperate zone, diminishing towards the equator and the poles. Hinds gives the following statement as to certain families which are almost exclusively confined to one of the six great divisions of the globe:—

In Europe,—Globulariaceæ a section of Selaginaceæ, Ceratophyllaceæ.

In Asia,—Dipterocarpaceæ, Aquilariaceæ, Camelliaceæ, Moringaceæ, Stilaginaceæ.

In Africa,—Bruniaceæ, Brexiaceæ, Belvisiaceæ, Penæaceæ.

In North America,—Sarraceniaceæ.

In South America,—Rhizobolaceæ, Monimiaceæ, Simarubaceæ, Vochysiaceæ, Calyceraceæ, Escalloniaceæ, Humiriaceæ, Lacisternaceæ, Papayaceæ, Gilliesiaceæ, Gesneraceæ.

In Australasia,—Tremandraceæ, Epacridaceæ, Goodeniaceæ, Stackhousiaceæ, Brunniaceæ.

He also gives the following list of natural orders, as prevailing in the northern hemisphere and southern hemisphere:—

In the northern hemisphere, the following natural orders abound, or are predominant:—Aceraceæ, Aurantiaceæ, Artocarpeæ a section of Urticaceæ, Amentaceæ, Berberaceæ, Boraginaceæ, Caryophyllaceæ, Cistaceæ, Cruciferae, Coniferae, Campanulaceæ, Caprifoliaceæ, Dipsacaceæ, Elæagnaceæ, Fumariaceæ, Grossulariaceæ, Hypericaceæ, Hippocastaneæ a section of Sapindaceæ, Hamamelidaceæ, Magnoliaceæ, Onagraceæ, Orobanchaceæ, Papaveraceæ, Rosaceæ, Ranunculaceæ, Rutaceæ, Resedaceæ, Saxifragaceæ, Umbelliferae, Vacciniaceæ, Alismaceæ.

In the southern hemisphere, the following natural orders are predominant:—Atherospermaceæ, Cactaceæ, Crassulaceæ, Capparidaceæ, Diosmeæ a section of Rutaceæ, Dilleniaceæ, Geraniaceæ, Heliotropæ a section of Ehretiaceæ, Myrtaceæ, Melastomaceæ, Mesembryaceæ, Myoporineæ a section of Verbenaceæ, Malpighiaceæ, Oxalidaceæ, Pittosporaceæ, Polygalaceæ, Proteaceæ, Scævoleæ a section of Goodeniaceæ, Spigelleæ a section of Loganiaceæ, Stylidiaceæ, Amaryllidaceæ, Hæmodoraceæ, Iridaceæ, Restiaceæ.

It is sometimes difficult to tell in what division of the globe a family may be said to be chiefly represented, inasmuch as the species and genera are nearly equal in different countries. When a group of plants occurs only in one of the six great divisions of the globe, it is said to be *monomic* (μόνος, one, and νομός, a region). Thus, Vochysiaceæ, being confined to South America, is a monomic family; and *Cliffortia*, whose shrubby species are all indigenous to South Africa, is a monomic genus. Again, a natural family, common to all the divisions, is *polynomic*; and so also genera, as *Viola* or *Ranunculus*. If restricted to two or more divisions, the groups are *dinomic*, *trinomic*, &c. Aceraceæ, found in Europe, Asia, and North America, are trinomic.

5.—PHYTO-GEOGRAPHICAL DIVISION OF THE GLOBE.

1152. The subject will be considered in two points of view:—1. In respect to the horizontal or latitudinal range of vegetation; and 2. In respect to its vertical or altitudinal range.

1153. **Horizontal Range of Vegetation.**—Various attempts have been made to divide the globe into zones or kingdoms, founded on the characters impressed upon them by the nature of the vegetation. Willdenow, Treviranus, De Candolle, Schouw, and Meyen, have each proposed arrangements. Those of Schouw and Meyen chiefly deserve attention.

1154. Schouw, in his divisions, proceeds on the principle of the predominance of certain characteristic forms or families of plants. His system is founded on the three following requisites:—1. That at least one-half of the known species of plants of that part of the earth, constituting a botanical region, should be peculiar to it. 2. That one-fourth part of the genera of the region should be peculiar to it, or at least should have so decided a maximum as to be only represented in other regions. 3. That individual families should either be peculiar to the region, or at least reach their maximum in it. The regions are divided into provinces according to minor differences in the vegetation; one-fourth

of peculiar species, or some peculiar genera, being sufficient to form a province.

1155.—*Schouw's Phyto-Geographic Regions.*

1. The Region of *Saxifragaceæ* and *Musci*, or the Alpine Arctic Flora.—This region is characterized by the abundance of Mosses and Lichens, the presence of *Saxifragaceæ*, *Gentianaceæ*, *Caryophyllaceæ*, *Cyperaceæ*, *Salices*; the total absence of tropical families; a notable decrease of the forms peculiar to the temperate zone; by forests of Fir and Birch; the small number of annual plants, and the prevalence of perennial species; and finally, a greater liveliness in their simple colours. In this region there is no cultivation. The region is divided into two provinces:—1. The province of the *Carices*, or the Arctic Flora, which comprehends all the countries within the polar circle, with some parts of America, Europe and Asia, which are to the south of it, more especially Lapland, the north of Russia, Siberia, Kamtschatka, New Britain, Canada, Labrador, Greenland, and the mountains of Scotland and Scandinavia, mean temperature 36° to 41° . 2. The province of *Primulaceæ* and *Phyteumæ*, or the Alpine Flora of the south of Europe, which embraces the flora of the Pyrenees, Switzerland, the Tyrol, Savoy, the mountains of Greece, the Appenines, and probably the mountains of Spain. Mean temperature, 47° to 66° .

2. The Region of *Umbelliferae* and *Cruciferae*.—These tribes are here in much greater number than in any other region; *Rosaceæ*, *Ranunculaceæ*, *Fungi*, *Amentaceæ*, and *Coniferae*, likewise very numerous; the abundance of *Carices*, and the fall of the leaves of almost all the trees during winter, form also important features of this division. It may be separated into two distinct provinces:—1. The province of the *Cichoraceæ*, which embraces all the north of Europe, not comprehended in the preceding region, namely, Britain, the north of France, the Netherlands, Germany, Denmark, Poland, Hungary, and the greater part of European Russia. 2. The province of the *Astragali*, *Halophyta*, and *Cynarocephalæ*, which includes a part of Asiatic Russia, and the countries about the Caucasian and Altai mountains. Mean temperature, 36° to 57° . The cultivated plants are—Rye, Barley of different kinds, Oats, Wheat and Spelt, Maize, Millet, the Potato, Buck-wheat, Apple and Pear, Quince, Cherry, Plum, Apricot, Peach, Mulberry, Walnut, Vine, Gooseberry and Currant, Strawberry, Cucumber and Melon, Cabbage, Mustard, Pea, Bean, Beet, Spinach, Carrot, Flax, Hemp, Trefoils and Vetches, Rye-grass, &c.

3. The Region of *Labiatae*, and *Caryophyllaceæ*, or the Mediterranean Flora.—It is distinguished by the abundance of the plants belonging to these two orders. *Compositæ*, *Galiaceæ*, *Boraginaceæ* also occur in considerable quantity. Some tropical families are also met with, such as Palms, Laurels, *Araceæ*, *Anacardiaceæ*, grasses belonging to the

genus *Panicum* (millet), and some *Cyperaceæ*. *Solanaceæ*, *Malvaceæ*, *Leguminosæ*, *Urticaceæ*, and *Euphorbiaceæ* increase. The forests are composed chiefly of *Amentaceæ*, and *Coniferæ*, as birches, oaks, firs, &c., the copses, of *Ericaceæ* (the heath tribe,) and *Anacardiaceæ*, as the mastich. We meet in this region with a great number of ever-green trees. Vegetation never ceases entirely, but verdant meadows are more rare. Schouw divides this region into five provinces:—1. The province of the Cistuses, including Spain and Portugal. 2. The province of the *Salviæ* and *Scabiosæ*, the south of France, Italy, and Sicily. 3. The province of the Shrubby *Labiataæ*, the Levant, Greece, Asia Minor, and the southern part of the Caucasian countries. 4. The Atlantic province, the north of Africa, of which he does not yet know any distinctive character. 5. The province of *Sempervivæ*, the Canary Isles, and probably also the Azores, Madeira, and the north-west coast of Africa; many *Sempervivums*, and some *Euphorbias* with naked and spiny stems particularly characterize this province. *Erica arborea*, and *Pinus canariensis* are found here. Mean temperature, 54° to 72°. Cultivated plants are the same as in the second region, with the addition of Rice, Guinea Corn, Italian Millet, Fig, Almond, Orange and Lemon, Water Melon, Olive, Cotton. Rye and Buckwheat are only cultivated in the mountainous regions.

4. The Region of *Asters* and *Solidagos*.—This is marked by the great number of species belonging to these two genera, by the great variety of Oaks and Firs, the small number of *Cruciferaæ* and *Umbelliferaæ*, *Cichoraceæ*, and *Cynarocephalæ*, the total absence of the genus *Erica*, or heath, and the presence of more numerous species of *Vaccinium*, or whortleberry, than are to be met with in Europe. It comprehends the whole of the eastern part of North America, with the exception of what belongs to the first region. It has been divided into two provinces:—1. That of the south, which embraces the Floridas, Alabama, Mississippi, Louisiana, Georgia, and the Carolinas. 2. That of the north, which includes the other states of North America, such as Virginia, Pennsylvania, New York, &c. Mean temperature, 54° to 72°. In the northern districts, down to the parallels of 55° or 50°, there is no cultivation. South of this line, the cultivation is the same as in the second region. Maize is cultivated to a greater extent in North America than in Europe.

5. The Region of *Magnolias*, comprising the most southern parts of North America, between 36° and 30°. The tropical forms which show themselves more frequently than on a similar parallel of the old continent, are the chief feature in the vegetation. Thus we meet with *Anonaceæ*, *Sapindaceæ*, *Melastomaceæ*, *Cactaceæ*, and *Zingiberaceæ*. This region has fewer *Labiataæ* and *Caryophyllaceæ* than occur in corresponding latitudes in the Old World. It presents more trees with fine blossoms, and shining, sometimes pinnated, leaves, as Mangolia,

Tulip-tree, Horse-chestnut, Robinias or False Acacias. Mean temperature, 59° to 72° . The same plants cultivated as in the third region. Rice is much cultivated. In the southern district, the Sugar-cane is productive; and, in the eastern districts, Cotton is grown to a great extent.

6. The Region of *Camelliaceæ* and *Celastraceæ*, or the Japanese region.—This region is as yet too little known to enable us to determine accurately its characteristic features. It embraces the eastern temperate part of the old continent, namely Japan, the north of China, and Chinese Tartary, between lat. 30° and 40° north. Its vegetation appears to occupy a middle place between that of Europe, and that of North America, approaching more to the tropical than to the European. It has an affinity to the Indian Flora, as shown by the occurrence of Bananas, Palms, *Zingiberaceæ*, *Anonaceæ*, *Sapindaceæ*, and *Cycadaceæ*. We meet with *Eriobotrya japonica* or Loquat, Tea, and various species of *Citrus*. Mean temperature, 54° to 68° . The cultivated plants are—Rice, Wheat, Barley, Oats, Millet, Buckwheat, Apple and Pear, Quince, Plum, Cherry, Apricot, Peach, Loquat, Orange and Shaddock, Melon, Tea, Hemp, Paper-Mulberry, Cotton, and False Sago.

7. The Region of *Zingiberaceæ*, or the Indian Flora.—*Zingiberaceæ* here are much more numerous than in America, as well as *Leguminosæ*, *Cucurbitaceæ*, and *Tiliaceæ*, although in a less degree. In consequence of the imperfect state of our knowledge, we cannot subdivide this region into provinces. It comprehends India, east and west of the Ganges, the island of Ceylon and the south-eastern Peninsula, to the height of 4,500 to 5,500 feet above the level of the sea. Mean temperature, 65° $75'$ to 81° $50'$. The cultivated plants are—Rice, Coco-nut, Tamarind, Mango, Ginger, Cinnamon, Mangosteen, Peppers, Indigo, Cotton, Coffee, Bananas, Guava, Orange and Shaddock, Sugar-cane, Cloves, Turmeric.

8. The Emodic Region, or the Mountains of India.—This comprises the Alpine region south of the ridge of the Himalaya. It includes Sirmore, Gurwal, Kamaon, Nepal, and Bhotan, to a height of from 4,500 to 10,700 feet above the level of the sea. Some tropical plants grow in the lower parts of the region. Deodar, *Pinus excelsa*, *Webbiana*, and other *Conifera*, are met with. Mean temperature, 66° to 37° . Some European grains and fruit are cultivated, along with Mountain Rice.

9. The Region of the Asiatic Islands.—This includes the mountainous districts of the islands between the south-eastern Peninsula and Australia, to the height of 5,500 feet above the level of the sea. Mean temperature, 66° to 84° . The cultivated plants are those of the Indian region (7); also, Bread-fruit, Cassava, Nutmeg, Camphor, Papaw, and Dammar.

10. The Region of Upper Java.—This embraces those districts of the island of Java and the islands of the Indian Archipelago which

have an elevation of 5,500 feet above the level of the sea. The vegetation of this region has not been sufficiently ascertained.

11. The Polynesian Region.—This includes all the islands of the Pacific Ocean within the Tropics. Mean temperature, 72° to 82° . The cultivated plants are—Bread-fruit tree, Coco-nut, Double Coco-nut, Yams, Plantain, Cabbage, Palm, Paper-Mulberry.

12. The Region of *Aymridaceæ*, or of Balsam trees.—This comprehends the Persian or Arabian Flora, especially the south-western part of the highlands of Arabia or Yemen. In this region are many trees yielding gums and balsamic resins, as species of *Mimosa*, *Acacia*, *Balsamodendron*, *Boswellia*. Cultivated plants are—Maize, Millet, Date-palm, Coco-nut, Fig, Apricot and Peach, Plum, Apple, Quince, Vine, Coffee-tree, Tamarind, Papaw, Sugar-cane, Ginger, Cotton, and Indigo.

13. The Desert Region.—This includes Northern Africa, to the south of the mountains of Atlas, between lat. 30° and 15° N., and the northern part of Arabia. *Phoenix dactylifera*, or the Date-palm, and *Cucifera thebaica*, or Doom-palm, are found here, but little is known in regard to the vegetation of the region. Mean temperature, 72° to 86° . Cultivation is confined to the valley of the Nile and the Oases. We meet with Guinea Corn, Wheat and Barley, and the South European and Indian grains.

14. The Region of Tropical Africa.—This includes that part of Africa lying between the parallel of 15° , and the tropic of Capricorn, or between the northern and southern limits of periodical rains, with the exception of Abyssinia and the unknown countries of the interior. The Flora of the western part of this region is characterized in part by the presence of *Adansonia*, or the Boabab, the largest known tree. We also meet with the *Elais guineensis*, a palm which furnishes oil. Other characteristic plants are *Sarcocephalus esculentus* and *Schmiedelia africana*. The vegetation of Guinea and Congo is a mixture of the Floras of Asia and America, though most resembling the former. The eastern part of the region, including Madagascar, has a peculiar Flora, distinguished chiefly by the genera *Danais*, *Ambora*, *Dombeya*, *Dufourea*, *Didymomeles*, and *Senacea*. Mean temperature, 72° to 86° . Cultivated plants: Maize, Rice, Guinea Corn and Millet, Yams, Cassava, Banana, Mango, Papaw, Pine-Apple, Cashew, Tamarind, Coffee, Sugar, Cotton, Ginger, Cardamoms, Earth-nut.

15. The Region of *Cactuceæ* and *Piperaceæ*.—This embraces Mexico, New Grenada, Guiana, and Peru. These natural orders are here predominant, both as regards the number of species and the individual plants. Murichi or Ita Palm, *Phytelephas* or Ivory Palm, and *Victoria regia*, are peculiar to Guiana. Mean temperature, 68° to 84° . Cultivated plants: Maize, Guinea Corn, Cassava, Yams, Batatas, Arracacha, Arrow-root, Plantain, Mango, Custard Apples, Guava, Coco-nut, Papaw, Avocado Pear, Pine Apple, Cashew, Tamarind,

Grenadilla, Vine, Indian Fig, Jambos, Chocolate, Vanille, Coffee, Sugar, Capsicum, Cochineal-Cactus, Cotton, Earth-nut.

16. The Region of the Highlands of Mexico.—This embraces the districts which have an elevation of more than 5,500 feet above the level of the sea. Many European plants are cultivated here, as well as Maize. *Pinus religiosa* and other *Coniferae* are found. Mean temperature, 78° to 64°.

17. The Region of *Cinchonæ*, or Medicinal Barks.—This comprises a part of the elevated regions or Cordilleras of South America, included in the torrid zone, the Andes from 5,500 to 9,000 feet. The *Cinchona* belongs exclusively to this region, and forms its principal feature. In the higher regions the Potato and Quinoa are cultivated, as well as some European grains and fruits. In the lower districts, Maize and Coffee are still cultivated. Mean temperature, 68° to 59°.

18. The Region of *Escalloniæ* and *Calceolariæ*.—It embraces the highest parts of South America, or that portion of the chain of the Andes, which has more than 9,600 feet of elevation. Besides the plants mentioned, we meet with alpine plants, as Saxifrages and Gentians, and species of *Draba*, *Arenaria*, *Carex*, *Lobelia*, and *Salvia*, besides some European genera belonging to the orders *Gramineæ* and *Cichoraceæ*, such as *Bromus*, *Festuca*, *Poa*, *Apargia*, and *Hypochaeris*. Mean temperature, 59° to 34°.

19. The West Indian Region.—This includes the whole district of the Great and Little Antilles. Bananas, Plantains, Mangos, Guava, Avocado Pear, Tamarind, and many other useful plants are met with. The Flora is intermediate between that of Mexico and the northern parts of South America. Mean temperature, 59° to 78°. Cultivated plants the same as those in the fifteenth region.

20. Region of *Palmæ* and *Melastomaceæ*.—It embraces Brazil and that part of South America which lies to the east of the chain of the Andes, between the Equator and the Tropic of Capricorn. *Vellozia* and *Lichnophora* give a decided feature to the peculiar vegetation of some of the mountainous parts. Here, also, numerous large peculiar species of *Eriocaulon* occur. Species of *Croton*, *Dorstenia*, and *Heliconia* are also met with, along with arborescent *Solanums*. In place of the few mosses and lichens which cover the trunks or branches of forest trees in temperate climes, in Brazil they are bearded from the roots to the very extremities of the smallest branches with Ferns, *Araceæ*, *Tillandsias*, Cactuses, Orchids, *Piperomias*, and *Gesneras*. Mean temperature, 59° to 82°. Same plants cultivated as in the fifteenth region.

21. The Region of Arborescent *Compositæ*.—The great number of arborescent *Compositæ*, and of plants belonging to the order *Calyceraceæ*, forms the chief feature of this Flora, which approaches in a remarkable manner to that of Europe, whilst it differs entirely from

the floras of Chili, the Cape, and New Holland. This region comprehends the lower part of the basin of La Plata, and the plains which extend to the west of Buenos Ayres and Chili, between the Tropic of Capricorn and latitude 40° south. (The Flora of Chili approaches that of New Holland, the Cape of Good Hope, and New Zealand, in *Goodenia*, *Araucaria*, *Proteaceæ*, *Gunnera*, *Ancistrum*.) Mean temperature, 59° to 75°. European plants form here objects of culture. Wheat, the Vine, and the Peach, are widely extended.

22. The Antarctic Region.—This includes the countries near the Straits of Magellan, Terra del Fuego, and the Falkland Islands. There is a considerable resemblance between the vegetation here and what is seen in the north temperate zone. Polar forms display themselves in the species of Saxifrage, Gentian, Arbutus, and Primrose. There is also a resemblance between the plants of this region and those of the mountains of South America, of Chili, the Cape, and New Holland. *Fagus antarctica* and Winter's Bark occur here. Many of the common British weeds are found at the southern extremity of South America. In the Falkland Islands is found *Dactylis cæspitosa*, the Tussac grass, which furnishes most valuable fodder. Mean temperature, 41° to 46°. No cultivation.

23. The Region of *Mesembryanthema* and *Stapsiæ*.—These two genera, as well as the *Ericææ*, Heaths, are very abundant. The latter family is found in greater quantity here than anywhere else. The region embraces the southern extremity of Africa. *Iridaceæ*, *Pelargoniums*, *Alioneæ*, *Bruniaceæ*, and *Selaginaceæ*, and various *Gnaphaliums* and *Elichrysums*, occur in this region. Mean temperature, 55° to 73°. Cultivated plants: European kinds of grain, fruit, and vegetables; also Batatas, Plantains, Tamarind, Guava, and Shaddock.

24. The region of *Epacridaceæ* and *Eucalypti*.—It comprehends the temperate parts of New Holland and Australia beyond the tropics, with the Island of Tasmania or Van Diemen's Land. Besides the plants whence it receives its name, it is characterized by the orders *Stackhousiaceæ* and *Tremandraceæ*, and by the presence of a great number of *Proteaceæ*, *Myrtaceæ*, *Stylidiaceæ*, *Restiaceæ*, *Diosmeæ*, *Casuarinææ*, and *Acacias*. *Araucaria*, or *Eutassa excelsa*, the Norfolk Island Pine, forms one of the features of the region. It is one of the most peculiar Floras. Mean temperature, 52° to 72°. In the British Colonies, the European kinds of grain and fruit are cultivated.

25. The Region of New Zealand.—This Flora, besides the plants peculiar to New Zealand, as *Phormium tenax*, New Zealand Flax, comprehends several others which belong to the extremities of America, Africa, and Australia. We find in these islands *Corypha australis*, the Australian or Southern Palm, Tree Ferns, and *Dracænas*, forests of *Dammara australis*, one of the *Conifereæ*, and many *Myrtaceæ*.

Mean temperature between latitude 34° and 36° south, from 61° to 63° . Many of the European plants are cultivated.*

1156. Meyen divides the horizontal range of vegetation into zones, taking for his basis the three ordinary divisions of the torrid, the temperate, and the frigid zone, and subdividing each hemisphere into eight smaller zones.

1157.—*Meyen's Phyto-Geographical Zones.*

A.—TORRID ZONE.

1. Equatorial Zone.—This extends 15° on both sides of the equator, and has a mean annual temperature of $78\frac{1}{2}^{\circ}$ to $82\frac{1}{2}^{\circ}$ F. The forms characteristic of this zone are chiefly Palmæ, Musaceæ, aborescent Gramineæ, Pandanus, Scitamineæ, Orchids, and Lianas; besides plants belonging to the orders Malvaceæ, Anonaceæ, Anacardiaceæ, Artocarpeæ a section of Urticaceæ, Lecythidaceæ, Malpighiaceæ, Sapindaceæ, Cæsalpinieæ a section of Leguminosæ, Cedrelaceæ, and many others.

2. The Tropical Zone.—This reaches from the 15th degree on each side of the equator to the tropics, in 23° latitude. Mean temperature $73\frac{1}{2}^{\circ}$ to $78\frac{3}{4}^{\circ}$. Summer temperature, $80\frac{1}{2}^{\circ}$ to 86° ; winter temperature in the eastern coast countries, 59° . Besides many equatorial forms, as Palms, Musaceæ, Scitamineæ, Meliaceæ, Anonaceæ, Sapindaceæ, Orchidaceæ, Araceæ, and Lianas, there are in this zone Tree-ferns, and plants belonging to Convolvulaceæ, Melastomaceæ, and Piperaceæ.

B.—TEMPERATE ZONE.

3. Subtropical Zone.—This extends from the tropics, 23° to 34° of latitude. Mean temperature, $62\frac{1}{2}^{\circ}$ to $71\frac{1}{2}^{\circ}$; summer temperature, $73\frac{1}{2}^{\circ}$ to $82\frac{1}{2}^{\circ}$. There is a number of tropical fruits in this region. The winters are mild, and vegetation is green throughout the year. In the northern division of the zone, Palms and Bananas grow on the plains. The Date-palm, Doom-palm, Chamærops Palmetto, many succulent Mesembryaceæ and Crassulaceæ, arborescent Euphorbias, Camellia, Thea, Aucuba, and Magnolias, are met with. In the southern division are Proteaceæ, Myrtaceæ, Epacridaceæ, Ericaceæ, many Compositæ, Diosmeæ, Zamias, and Cactaceæ.

4. The warmer Temperate Zone.—It embraces the space between 34° and 45° of latitude, including the southern part of Europe, Asia Minor, north of China, and Japan. Mean temperature, $53\frac{1}{2}^{\circ}$ to $62\frac{1}{2}^{\circ}$. Summer temperature in North America, 77° ; in Europe, $75\frac{1}{4}^{\circ}$ to 68° ; in Eastern Asia, $82\frac{1}{4}^{\circ}$; Winter temperature in the New World, $44\frac{1}{2}^{\circ}$ to $32\frac{1}{2}^{\circ}$; in Europe, 50° to $34\frac{3}{4}^{\circ}$; in Eastern Asia, $26\frac{1}{2}^{\circ}$. Many subtropical forms occur. Evergreen dicotyledonous trees and

* See a Coloured Delineation of these Regions in Johnston's Maps.

shrubs, Cistuses, many species of Ericaceæ, Lauraceæ, and Myrtaceæ, and the Vine, are met with. In some parts of the zone, Solidagos and Asters, Magnolias, and Smilacææ, abound; while in others there are representatives of the Mimosa form, Myrtaceæ, and Proteaceæ.

5. The colder Temperate Zone.—This includes a belt from 45° to 58° latitude. Mean temperature, 43° to $53\frac{1}{2}^{\circ}$. Minimum summer temperature on the West Coast, $56\frac{1}{4}^{\circ}$; in the interior of the Continent, 68° ; Minimum winter temperature in the interior of Europe, 14° . England, the north of France, and Germany, supply the characteristics of the vegetation of this zone. It embraces the region of Umbelliferæ, and Cruciferæ of Schouw. Meyen selects plants having a more marked physiognomic effect, such as ordinary Dicotyledonous trees, along with Abietinææ, and heaths covered with *Calluna vulgaris*.

6. The Subarctic Zone.—This reaches from 58° latitude to the arctic circle, 66° . Mean temperature, $39\frac{1}{4}^{\circ}$ to 43° . Summer temperature in the New World, $66\frac{1}{4}^{\circ}$; in the Old World, $60\frac{3}{4}^{\circ}$ to 68° ; Winter temperature of the former, 14° ; of the latter (Western Europe), $24\frac{1}{4}^{\circ}$; of the interior of Russia, 14° to $10\frac{1}{2}^{\circ}$. It is characterized by Firs and Willows in the northern hemisphere.

C.—FRIGID ZONE.

7. The Arctic Zone.—This extends from the Arctic circle, 66° to 72° . Mean temperature, $28\frac{1}{2}^{\circ}$ to 32° , and towards the eastern and continental portions, far below the freezing point. The Birch, and some Coniferæ, may be said to characterize this zone.

8. The Polar Zone.—This includes all lands from 72° to 82° latitude. The mean temperature of one point in this zone, viz., Melville Island, is $1\frac{1}{2}^{\circ}$. In the Old World, the mean temperature is $16\frac{1}{2}^{\circ}$. Summer temperature of the New World, $37\frac{1}{2}^{\circ}$, and of the Old, $38\frac{1}{4}^{\circ}$; Winter temperature, -28° in the New, and $-2\frac{1}{4}^{\circ}$ in the Old World. No trees nor bushes grow in this zone. Some Saxifrages, and numerous Cryptogamic plants, as Lichens, prevail in it.

1158. **Vertical Range of Vegetation.**—Under this head we consider the changes produced in the physiognomy of vegetation on ascending mountains. It has reference to the distribution of plants in an altitudinal or hypsometrical point of view. This geographical range is best seen in the high mountains of tropical countries, where all gradations are met with, from the heat of the torrid zone to the cold of the frigid zone. Humboldt, in describing South American scenery, remarks:—"In the burning plains, scarce raised above the level of the Southern Ocean, we find Bananas, Cycadaceæ, and Palms in the greatest luxuriance; after them, shaded by the lofty sides of the valleys in the Andes, Tree Ferns; next in succession, bedewed by cool misty clouds, Cinchonas appear. When lofty trees cease, we come to Aralias, Thibau-

dias, and myrtle-leaved *Andromedas*; these are succeeded by *Bejarias* abounding in resin, and forming a purple belt around the mountains. In the stormy regions of the *Paramos*, the more lofty plants and showy flowering herbs disappear, and are succeeded by large meadows covered with grasses, on which the *Llama* feeds. We now reach the bare trachytic rocks, on which the lowest tribes of plants flourish. *Parmelias*, *Lecidias*, and *Leprarias*, with their many-coloured sporules, form the flora of this inhospitable zone. Patches of recently fallen snow now begin to cover the last efforts of vegetable life, and then the line of eternal snow begins."

1159. On the mountains of temperate regions the variety is rather less, but the change is not less striking. "We begin to ascend the Alps, for instance, in the midst of warm vineyards, and pass through a succession of oaks, sweet chestnuts, and beeches, till we gain the elevation of the more hardy pines and stunted birches, and tread on pastures fringed by borders of perpetual snow. At the elevation of 1,950 feet, the vine disappears; and at 1,000 feet higher, the sweet chestnuts cease to grow; 1,000 feet farther, and the oak is unable to maintain itself; the birch ceases to grow at an elevation of 4,680, and the spruce fir at the height of 5,900 feet, beyond which no tree appears. The *Rhododendron ferrugineum* (the Rose of the Alps) then covers immense tracts to the height of 7,480 feet, and *Salix herbacea* creeps 200 or 300 feet higher, accompanied by a few *Saxifrages*, *Gentians*, and *Grasses*, while *Lichens* and *Mosses* struggle up to the imperishable barrier of perpetual snow." In central and southern Europe, the proportion of *Monocotyledons* to *Dicotyledons*, which is as 1 to 4 in the plains, decreases with the elevation on dry mountain slopes, till at the height of 8,526 feet, it is as 1 to 7. Moist mountain slopes favour *Monocotyledons*, the proportion on them being as 1 to 3.

1160. The following table shows the height at which corn and trees grow in different quarters of the globe:—

	Torrid Zone.		Temperate Zone.			Frigid Zone.
	Andes 0° Lat.	Mountains of Mexico, 20° Lat.	Caucasus, 42° 30'	Pyrenees, 43°	Alps, 46°	Lapland, 67°—70°
Inferior limit of per- petual snow.....	15,200 feet	13,478	9,900	8,400	8,220	3,300
Upper limit of trees,	10,800 —	12,000	6,700	7,020	6,000	1,500
Distance between trees and snow....	4,400 —	1,478	3,200	1,380	2,220	1,800
Distance between snow and corn....	5,200 —		3,780		4,200	2,700

In the Himalaya, the upper limit of trees on the south side is marked by *Quercus semecarpifolia*, at 11,500 feet, and on the north side by *Betula alba*, at 14,000 feet. The Birch also forms the limit on the

Caucasian mountains. On the Pyrenees and Alps, the limit of trees is marked by *Coniferæ*; on the Pyrenees, by *Pinus uncinata*; on the south side of the Alps, by *Larix europæa*, the Larch; and on the north side, by *Abies excelsa*. In Lapland, the Birch forms the upper limit of trees. The upper limit of shrubs is determined by the *Rhododendrons* in the Old World, on the Pyrenees at 8,312 feet, and on the Alps, at 7,480; and by *Bejarias* on the Andes, at the height of 13,420 feet. On the south side of the Himalaya, *Juniperus*, *Salix*, and *Ribes*, form the upper limit of shrubs, at 11,500 feet; on the north, *Genista versicolor* ascends to 17,000 feet.

1161. The following have been given by some authors as the zones of Alpine vegetation:—

1. Region of Lowland cultivation. Its extent of elevation is at the spot where the prevailing cultivated plants of the latitude cease to be productive. In Mount *Ætna*, it rises to 3,300 feet; on *Teneriffe*, to nearly 3,000 (zone of vines). It embraces two zones of the *Cactus* and *Euphorbia* in the *Canaries*. In *Madeira*, it embraces two regions of *Spix* and *Martius*; the region of tropical plants reaching to 700 feet, and the region of the vine, fruit, and corn, to 2,300. In *Norway*, *Sweden*, and *Finmark*, it is narrow. In the *Carpathians*, it rises to 1,500 feet. Within the tropics, it is a broad and important region. On the *Andes*, at *Quito*, it only ceases at 5,000 or 6,000 feet.
2. Region of Woods. A magnificent region in all Alpine districts, and well characterized on the *Andes* and *Himalaya*. *Humboldt* notices it in *Teneriffe*. In *Ætna*, it extends to 6,200 feet. In the *Canaries*, to 4,080. In *Madeira* (region of *Chestnut*), to 2,950. In *Lapland*, it extends to 800 feet. In *Finmark*, 70° north latitude, to 730.
3. Region of Shrubs. Region of *Retama* (*Spartium nubigenum*) in *Teneriffe*. On the *Pyrenees* and *Mount Blanc*, it is extensively covered to about 9,000 feet with *Rhododendrons*. In the *Andes*, about *Quito*, it reaches 13,000, and is conspicuous for its shrubby compositæ. In *Madeira*, it embraces *Kuhl's* regions of *Spartium* and *Heath*. On the mountains of *Lapland*, it attains 1,000 feet, and is characterized by *Betula nana*, *Vaccinium*, and *Salix*. In *Finmark*, its limits are 1,100.
4. Region of Grasses. These predominate in certain Alpine situations, and in certain parallels of latitude. In *South Shetland*, none of the islands exhibit any *Phanerogamous* vegetation, with the exception of straggling grass. In *Melville Island*, 75° north latitude, the proportion of grasses to *Phanerogamous* plants is 1 to 5; in *Great Britain*, 1 to 12½. In the *Andes*, the region is traced to *Paramos*, and occupies a space of 13,000 to 14,500 feet; here are large cattle farms. In the *Himalaya*, a fine green sward is often seen at 14,600 feet. In *Teneriffe*, it is distinguished by *Humboldt*. On the *Swiss Alps*, *Poa annua* exists at an elevation of 7,400 feet.
5. Region of Cryptogamous plants. This is well marked in many places. *Colonel Hall*, in *Chimborazo*, under the *Equator*, at nearly 16,000 feet, found *Draba arctoides*, and *Culcitium rufescens*; still higher, a moss, which may be considered as having attained the highest limit on the globe at which vegetable life exists. *Lichens* are the latest plants met with in ascending *Teneriffe*, the *Himalaya* mountains, and the *Alps*.

1162. In the mountains of the torrid zone, the following regions

are described by Meyen, corresponding to the zones given in his horizontal range of vegetation:—

1. The region of Palms and Bananas, extending from the level of the sea to 1,900 feet of altitude. It corresponds to the Equatorial zone.
2. Region of Tree Ferns, and species of Ficus, extending from 1,900 feet to 3,800. (Tropical zone.)
3. Region of Myrtaceæ and Lauraceæ, extending from 3,800 to 5,700 feet. (Subtropical zone.)
4. Region of Evergreen Dicotyledonous trees, extending from 5,700 to 7,600 feet. (Warm Temperate zone.)
5. Region of Deciduous Dicotyledonous trees, extending from 7,900 to 9,500 feet. (Cold Temperate zone.)
6. Region of Abietineæ, the Pine and Fir, extending from 9,500 to 11,400 feet. (Subarctic zone.)
7. Region of Rhododendrons, extending from 11,400 to 13,300 feet. (Arctic zone.)
8. Region of Alpine plants and Lichens, extending from the upper limit of shrubs to the snow line at 15,200 feet. (Polar zone.)*

1163. **Zones of Marine Vegetation.**—An analogy is found to exist between marine and terrestrial plants. The former as well as the latter manifest a diversity of appearance, according to their position, surrounding medium, relative degrees of pressure, and exposure to light. Harvey has observed that some sea weeds seem to be unaffected by circumstances of this nature, and are found equally abundant under opposed latitudes and in extremes of temperature. The lower we descend in the scale of marine vegetation the better is this illustrated. Many of the Diatomaceæ seem to be generally distributed.

1164. Lamouroux has estimated the marine Algæ as ranging from 5,000 to 6,000, which he has shown are distributed in various regions. Their distribution is much influenced by the degree of exposure to light, as well as by the motion of the waves. Great depths of the ocean are observed to exercise an influence on marine vegetation, similar to that which high mountains have on land plants. Some species, as the Laminariæ, are confined to the colder regions of the sea, while others, as the Sargassa, are only found where the mean temperature is considerable. The colour of Algæ may be regarded as being in a measure indicative of their depth of growth, the Algæ of green colour being generally found either in fresh water or in the shallower parts of the sea; the olive-coloured Algæ abound most between tide marks; those of a red colour occur chiefly in the deep and dark parts of the sea.

Marine vegetation is found to vary both in its horizontal and vertical range. This difference is less decided than that which is observable amongst land plants, owing probably to the greater uniformity

* On the subject of Botanical Geography, the student may consult Meyen's work printed by the Ray Society; Hooker's Treatise in Murray's Encyclopædia of Geography; Malte Brun's Physical Geography; Humboldt's Narrative, &c.; and Berghaus's Maps, edited by A. K. Johnston,—a cheap quarto edition of which has been published.

of the ocean's temperature. The ocean has been divided into the following provinces of marine vegetation:—1. The Northern Ocean, from the Pole to the 60th parallel of north latitude. 2. The North Atlantic, between the 60th and 40th parallels, the province of the species of *Fucus* proper. 3. The Mediterranean, which is a sub-region of the warmer temperate zone of the Atlantic, lying between the 40th and 23d northern parallels. 4. The tropical Atlantic, in which *Sargassum* abounds. 5. The Antarctic American regions from Chili to Cape Horn, and the whole circum-polar ocean south of 50° of latitude. 6. The Australian and New Zealand province. 7. The Indian Ocean and Red Sea. 8. The Japan and China seas, besides certain provinces in the Pacific.

1165. Forbes remarks of the vertical range, that one great marine zone lies between high and low water mark, varying in species according to the nature of the coast. This zone is generally uniform throughout the northern hemisphere. A second zone begins at low water mark, and extends to a depth of 7 to 15 fathoms. The first of these, or the littoral zone, has been divided into sub-regions, in which certain marine species are found to prevail. 1. The sub-region of *Fucus carnaliculatus*. 2. The sub-region of *Lichina*. 3. The sub-region of *Fucus articulatus*, *F. nodosus*, and *Corallina officinalis*. 4. The sub-region of *Fucus serratus*. The second or Laminarian zone includes the great Tangle sea-weeds and deep-water Fuci. The lowest form of vegetable life does not, in British seas, extend deeper than 50 fathoms. In the Mediterranean and the *Ægean* Sea, the Coralline zone ranges from 50 to 100 fathoms. The lowest forms of marine vegetation are alone met with in the deepest waters.

1166. Marine vegetation is equally various in its horizontal range. In the North Sea and the British Channel, *Chorda Filum* is found to constitute beds of 15 to 20 miles in length, and about 600 feet in breadth. *Sargassum bacciferum* constitutes the Gulf-weed of the Atlantic. It occupies the eddy caused by the revolution of the Atlantic current, and extends over a space of 260,000 square miles. It appears to derive its sustenance wholly from the ocean, inasmuch as it has never been found attached. The *Macrocystis pyrifera*, and the *Laminaria radiata* are remarkable for the size and extent of their range. Immense green masses of the *Macrocystis* are met with in every latitude. The tribe *Fucoideæ* are met with in abundance towards the poles, in which regions they are observed to attain their greatest bulk, diminishing and ceasing as they approach the Equator. *Cystoseiræ* follow a course in the higher latitudes of the southern hemisphere similar to the *Fucoideæ*. *Laminarias* inhabit the antarctic ocean, and stretch northwards to the Cape of Good Hope. The red, green, and purple Lavers of the British seas are found at the Falkland Islands. In like manner, special characteristics are found

to appertain to marine vegetable growth as met with in other localities.

Species which are found to abundantly inhabit one sea, may be scarcely present in another. This is seen in the difference between the marine vegetation of the Red Sea as compared with that of the Mediterranean, and that of the Mediterranean as contrasted with the Atlantic. Sargassa, which abound in the Red Sea, and Fuci, which are numerous in the Atlantic, are equally rare in the Mediterranean.

1167. Distribution of Plants in Britain.—The climate of Britain is influenced by its geographical position, and the form and elevation of its surface. The eastern coasts partake more of the continental climate, while the western experience the insular or more equable climate. The mean temperature varies from 46° to 52° F. In ascending the mountains, there is a fall of 1° of the thermometer on an average for every 230 or 240 feet of ascent. The number of Phanerogamous plants amounts to nearly 1,600, of Cryptogamous to about 4,800. Mr. H. C. Watson, who is our chief authority on the geography of British plants, in his earlier works gives the following distribution in ascending regions:—

1. The region of the Plains. This includes all the low and open country from the south coast of England to the borders of the Highlands, in lat. 56° 56½°, terminating at the sea level on the shores of the Clyde and Tay. It contains the common weeds and wild flowers of the country, and among trees, the Oak, Ash, &c.
2. Upland Region. This is marked by the occurrence of *Vaccinium* *Vitis*-*Idæa*, *Arctostaphylos* *Uva*-*Ursi*, *Polygonum viviparum*, *Trientalis europæa*, *Linnaea borealis*, &c.
3. Median Region. A narrow belt just above the upper limit of cultivation, and the growth of *Amentaceæ* such as the Oak and Hazel.
4. Subalpine Region. This commences at from 1,500 to 2,400 feet above the level of the sea. *Juncus trifidus*, *Saxifraga nivalis*, and *Gnaphalium supinum* occur here.
5. Alpine Region. This region is only truly seen in Scotland. *Saxifraga cernua* and *rivularis*, *Draba rapestris*, *Luzula arcuata*, *Stellaria cerastoides*, and *Alsine rubella*, may be said to characterize it.

Of late years, Mr. Watson has proposed the following geographical division of British plants:—

1. Agrarian. Plants found only within the limits of cultivation, or within the limits of *Pteris aquilina*, in waste land, the elevation or climate of which would not forbid successful cultivation.
2. Agro-Arctic. Plants found within the region of cultivation, and also above these limits.
3. Arctic. Plants found only above the limits of cultivation.

1168. In considering the distribution of British plants in connection with geographical or local position, Watson suggests the following types:—

1. Atlantic type: embracing species found in the south-west of England and Wales, as *Erica ciliaris*, *Sibthorpia europæa*, and *Euphorbia Peplis*.
2. Germanic type: including species chiefly seen in the south-west of England, the chalk plants, as *Phyteuma*, *Ophrys*, *Hippocrepis*, and *Onobrychis*.
3. The English type: consisting of species chiefly or exclusively found in England, and decreasing in frequency northwards, as *Acer campestre*, *Tamus communis*, *Iris foetidissima*, *Orchis Morio*.
4. British type: comprehending species widely spread over Britain, as *Bellis*, *Calluna*, *Corylus*.
5. Scottish type: embracing species prevalent chiefly in Scotland, or the north of England, as *Trollius*, *Linnaea*, *Andromeda*, *Primula farinosa*.
6. The Highland type: containing species either limited to the Scottish Highlands, or extending to the mountains of the north of England and Wales, as *Salix herbacea*, *Azalea*, *Arctostaphylos*, *Lobelia*, *Subularia*.
7. Hebridean type: composed of a few species peculiar to the extreme north and west of Scotland, or at least chiefly seen there, as *Eriocaulon septangulare*, *Ajuga pyramidalis*, *Primula scotica*.

1169. Professor E. Forbes has adopted Mr. Watson's view. He thinks that the vegetation of the British islands may be said to be composed of five floras:—1. A west Pyrenean, confined to the west of Ireland, and mostly to the mountains of that district. 2. A flora related to that of the north-west of France, extending from the Channel Isles, across Devon and Cornwall, to the south-east and part of the south-west of Ireland (Watson's Atlantic type). 3. A flora common to the north of France and south-east of England, and especially developed in the chalk districts (Watson's Germanic type). 4. An Alpine flora, developed in the mountains of Wales, north of England, and Scotland (Watson's Highland type). 5. A Germanic flora, extending over the greater part of Great Britain and Ireland, mingling with the other floras, and diminishing, though slightly, as we proceed westwards, indicating its easterly origin and relation to the characteristic flora of northern Germany (Watson's British, English, and Scottish types). In attempting to account for the peculiarities of these floras, most of which had been previously noticed by Mr. Watson, Professor Forbes enters into some geological speculations, which, although entirely conjectural, are well worthy of consideration. He numbers in ascending order these floras, according to their magnitude as to species, and also, in his opinion, according to their relative age and period of introduction into the area of the British islands. His conclusions on this point are the following:—

"1. The oldest of the floras now composing the vegetation of the British isles, is that of the mountains of the west of Ireland. Though an Alpine flora, it is southernmost in character, and is quite distinct as a system from the floras of the Scottish and Welsh Alps. Its very southern character, its limitation, and its extreme isolation, are evidences of its antiquity, pointing to a period when a great mountain barrier extended across the Atlantic from Ireland to Spain.

"2. The distribution of the second flora, next in point of probable date, depended on the extension of a barrier, the traces of which still remain, from the west of France to the south-east of Britain, and thence to Ireland.

"3. The distribution of the third flora depended on the connection of the coast of France and England towards the eastern part of the channel. Of the former existence of this union no geologist doubts.

"4. The distribution of the fourth, or Alpine flora of Scotland and Wales, was effected during the glacial period, when the mountain summits of Britain were low islands, or members of chains of islands, extending to the area of Norway through a glacial sea, and clothed with an arctic vegetation, which in the gradual upheaval of those islands and consequent change of climate, became limited to the summits of the new-formed and still existing mountains.

"5. The distribution of the fifth, or Germanic flora, depended on the upheaval of the bed of the glacial sea, and the consequent connection of Ireland with England, and of England with Germany, by great plains, the fragments of which still exist, and upon which lived the great elk and other quadrupeds now extinct.

"The breaking up or submergence of the first barrier led to the destruction of the second; that of the second to that of the third; but the well-marked epoch of migration of the Germanic flora indicates the subsequent formation of the straits of Dover and of the Irish Sea, as now existing.

"To determine the probable geological epoch of the first or west-Irish flora—a fragment, perhaps with that of north-western Spain, of a vegetation of the true Atlantic—Forbes seeks among fossil plants for a starting-point. This he gets in the flora of the London clay, or Eocene, which is tropical in character, and far anterior to the oldest of the existing floras. The geographical relations of the Miocene sea, indicated by the fossils of the crag, give an after-date certainly to the second and third of the above floras, if not to the first. The epoch of the red or middle crag was probably coeval with the second flora; that of the mammaliferous crag with the third. The date of the fourth is too evident to be questioned; and he regards the glacial region in which it flourished as a local climate, of which no true traces—as far as animal life is concerned—exist southwards of his second and third barriers. This was the newer Pliocene epoch. The period of the fifth flora was that of the post-tertiary, when the present aspect of things was organized.

"Adopting such a view of the relations of these floras in time, he thinks that the greatest difficulties in the way of changes of the earth's surface and destruction of barriers—deep sea being found where land (probably high land) was—are removed when we find that those greater changes must have happened during the epoch imme-

diately subsequent to the Miocene period; for we have undoubted evidence that elsewhere, during that epoch, the Miocene sea-bed was raised 6,000 feet in the chain of Taurus, and the barriers forming the westward boundary of the Asiatic Eocene lakes so completely annihilated, that a sea several hundred fathoms deep now takes their probable place. The changes required for the events which he would connect with the peculiar distribution of the British flora are not greater than these. Professor Forbes thinks that the peculiar distribution of endemic animals—especially that of the terrestrial mollusca—bears him out in these views.”

1170. The observations of Watson and Forbes lead to the conclusion that, with the exception of *Eriocaulon septangulare*, the British islands do not contain a single plant which is not found on the continent of Europe. These islands, therefore, cannot be considered as a centre of vegetation, but as having been colonized by successive vegetable migrations. Their opinion as to the origin of British plants, as condensed by Martins, is, that these islands have been peopled by many colonies successively leaving the continent of Europe, from the epoch of the middle tertiary formation up to our own. When a vast continent extended from the Mediterranean regions to the British islands, the plants of the Asturias, and those of Armorica, peopled the south of England and Ireland. To this period succeeded the glacial epoch, during which the lands were immersed to a depth of about 1,300 or 1,400 feet. This is the period of the migration of the arctic plants, which still inhabit the tops of the Scottish mountains. When these lands emerged anew, England was united to France, the temperature being such as it is at present. At this time, the great German floral invasion took place, absorbing, so to speak, all the rest, and leaving very slight remains of them. Thus, while the Asturian plants, those of the south, are reduced to a small number of species confined to the south-west of Ireland, the hardy vegetables of the north completed their conquest. The colonization being completed, England became separated from the continent.

1171. Martins agrees with Watson and Forbes in their general views of the British flora, adopting the following types:—

1. Asturian type—the remains of a Peninsular flora.
2. Armorican type—vegetation like that of Brittany and Normandy.
3. Boreal type—flora like that of the Swiss Alps, Lapland, Iceland, and Greenland.
4. Germanic type—the basis of the vegetation of England.

He thinks, that while Europe has had the principal part in the colonization of the British islands, a great vegetable migration has also taken place from America; and that the arctic plants originating in Greenland, have propagated themselves across Iceland, Feroe, and Shetland, as far as the mountains of Scotland. These mountains have

therefore derived their flora partly from Norway and partly from Greenland, by a sort of double migration. His opinion is founded on the fact, that the relative proportion of plants, exclusively European, which enter into the flora of Shetland, is $\frac{1}{4}$; into that of Feroe, $\frac{1}{7}$; and into that of Iceland, $\frac{1}{10}$; all the rest being common to Europe and America. In proportion, therefore, as we remove from Europe, the number of vegetables peculiar to that continent diminishes; but at the same time the proportion of the Greenland plants increases in nearly the same ratio. Martins, however, does not agree with Forbes in his bold and novel hypothesis, but attributes the colonization to the transport of seeds from America and Europe, by means of the gulf stream. This stream, he thinks, has thrown *Eriocaulon septangulare* on the shores of the Hebrides, and, running along the coast of Scotland, carries seeds to the sandy shores of Shetland, Feroe, and Iceland. He considers it the principal agent in the diffusion of European plants in these islands. Winds, aerial currents, and birds, he thinks, have also contributed to the dissemination of species.*

1172. Acclimatizing of Plants.—It is commonly supposed that by length of time plants may be rendered fit to endure a climate which they could not stand in the first instance. It has been said, that by slow degrees tender plants may become acclimatized to cold climates. Such a view, however, is totally inconsistent with the facts of the case. Each species of plant naturally bears a certain range of temperature, and it is impossible to extend that range. Many plants originally placed in greenhouses, and subsequently planted out, are held up as cases of acclimatization. *Aucuba japonica*, coming from a warm climate, was at first treated in this country as a stove-plant, and was afterwards planted out, and was found to endure the climate, but no change was made in the constitution of the plant. It was capable from the first of enduring the cold of this climate. *Aponogeton distachyum*, an equatic from the Cape, was cultivated long in the stoves of the Edinburgh Botanical Garden. A specimen was accidentally thrown into the open pond, where it has continued to live and flower for many years. The constitution of the plant is unaltered. It was able to bear a certain range of temperature, but cultivators were not aware of this in the first instance. Plants sent from warm countries, and supposed to be delicate, are often quite hardy, inasmuch as their native locality has been high on the mountains. Such is the case with *Araucaria imbricata* from Chili, and with some Nipal and Japan plants. Again, take the Potato, the Dahlia, Heliotrope, and Marvel of Peru, which have been long cultivated in Britain, and it will be seen that they are not in the slightest degree more hardy than when first

* For further details on the subject of the Geography of British plants consult Watson's *Distribution of British Plants*, and *Cybele Britannica*; Forbes's paper in the Reports of the Geological Survey of Great Britain; Martins' papers in the *Edinburgh Philosophical Journal* for 1849.

introduced; they are injured by the frost just as easily as at first.

1173. Something, however, may be done by the art of the gardener, to render half-hardy species of plants less tender. In this climate, the great risk in such cases, is frequently not so much the degree of cold, as the accession of it at a time when the plants cannot resist it, in consequence of being full of sap. Attention, therefore, should be paid to bringing the plants into as dry a state as possible, at the beginning of winter. Lindley remarks that the only means of effecting this consists in thoroughly drained soil, and an elevated situation—the first preventing a plant from filling itself with moisture during winter, or overgrowing itself in summer, so as to enable it to ripen its wood; and the latter securing it from the action of those early frosts in autumn, or those late frosts in spring, which are so pernicious even to our own wild trees. In an elevated situation, a plant also escapes the risk of being stimulated into growth by a few days' warmth, succeeded by nipping colds, which so often occurs in our variable climate.

PART IV.

FOSSIL BOTANY.

1174. THE history of vegetation could not be considered complete, unless we endeavoured to give some account, however brief, of the plants which existed on the earth in its primeval state, during the extended geological epochs which elapsed before the establishment of the present order of things. This subject is alike interesting to the botanist and the geologist. It has sometimes been called Geo-Botany, and is an important section of Oryctology (*ὄρυκτός*, fossil). "Geology," says Philips, "would never, perhaps, have escaped from the domain of empiricism and conjecture, but for the innumerable testimonies of elapsed periods and perished creations, which the stratified rocks of the globe present, in the remains of ancient plants and animals. So many important questions concerning their nature, circumstances of existence, and mode of inhumation in the rocks, have been suggested by these interesting reliquæ; and the natural sciences have received so powerful an impulse, and been directed with such great success to the solution of problems concerning the past history of the earth, that we scarcely feel disposed to dissent from the opinion, that without fossil Zoology and Botany, or what is denominated Palæontology (*παλαιός*, ancient), there would have been no true Geology." The stratified crust of the globe is full of these monuments of vanished forms of life. They are of various kinds, are in different states of preservation, and occur very unequally in rocks of different kinds and ages. The remains of ancient vegetation are very abundant in the coal measures, the important combustible material derived from them, and which is vegetable matter in an altered form.

1175. **Characters and Arrangement of Fossil Plants.**—From the state in which fossil plants are found, it is by no means an easy matter to determine their nature accurately. Brongniart states that the preservation of vegetables, in a fossil state, may be effected—1st, By

the impression or cast of the plant being discoverable, though accompanied by a complete destruction of the vegetable tissue, or the preservation of but a few of its constituent parts; 2d, By the structure of the tissues of the vegetable organs, through petrification or carbonization, being more or less completely preserved, in consequence of the total change or modification of their nature. The impression or cast, without the preservation of any portion of the organs of the vegetable more or less altered, is rather rare; yet it is the habitual state of fossil vegetables in the variegated sandstone and tertiary limestones.* It is rarely that any of the essential organs are found in such a state of preservation as to furnish distinct characters. It is chiefly from the fragments of stems, and the impressions of leaves, and some fruits, that the fossil botanist can draw conclusions. Besides this, fossil vegetables, thus reduced to some of their insulated organs, scarcely ever present them in such a state of preservation as enables them to be studied in all their constituent parts. Sometimes the internal structure of the stem can be traced, and by examination under the microscope, the nature of its woody tissue may be determined. In this way some fossilized woods have been referred to the Coniferous tribe, in consequence of the presence of punctated woody tissue (fig. 47). Fossil woods have been shown by chemical tests to contain portions of vegetable tissue, cemented into a mass by silica. In some cases the vessels and cells are separately silicified without being united into a compact mass. In these instances the wood breaks down easily. At times the internal structure is obliterated, and it is only from the external configuration, the nature of the outer covering, and the scars of the leaves, that any conclusions can be drawn. The leaves often furnish important and valuable characters, and, in the case of fossil ferns, their form, divisions, and venation, supply distinguishing marks. The leaves, however, are generally isolated, and are rarely found in connection with the stems. Thus, the separation of the different parts of the plant, and, in most cases, their imperfect state of preservation, are great obstacles in preventing the determination of fossil plants by a comparison with those which now exist on the earth. Before, then, endeavouring to compare a fossil vegetable with living vegetables, it is necessary to reconstruct, with as much exactness as possible, according to the parts preserved, and the general data of vegetable anatomy and organography, the portions of the plant under examination; to contrast these portions with the other organs of the same plant, searching for their points of attachment, their forms and vascular connections, being guided generally by traces of structure rather than by exterior form; and by endeavouring to recomplete a vegetable by bringing together all fragments from the same fossil beds, which may belong to the same plant. The connection of

* Vide Translation Brongniart's *Tableaux des Vegetaux Fossiles*.—Edin. Phil. Jour., vol. 48.

the different parts of the same plant is of the greatest importance in vegetable palæontology, as from their fragmentary nature many difficulties arise. These difficulties are increased as we go back to the earliest geological epochs, for the farther they are removed from the present state of things, the greater are the differences between the fossil and living plants. Dr. Hooker remarks, that the knowledge of recent botany required to throw light upon the study of fossil plants, and the origin of coal, must be both varied and extended. "Some acquaintance with systematic botany is the first requisite; through this alone can any approximation to the living affinities of the fossil be obtained. It should embrace not only a knowledge of the principal groups, or natural orders under which all plants are arranged, but a familiarity with vegetable anatomy; for when the stem or trunk alone is preserved, which is often the case, a minute examination of its tissues is the sole method of determining its position in the natural series. There must also be some general ideas of the vegetation both of the tropics and cooler latitudes, of mountain-chains, table-lands, valleys and estuaries; more especially of countries characterized by equable, and by excessive or extreme climates, as compared with continents, and of humid and desert districts; in short, of all the complex associations with, or dependence of botanical characters upon, surface, soil, and climate, which the globe presents."

1176. Many of the fossil plants of the tertiary or recent strata may be referred to genera at present existing, and merely present specific differences; such as pines, elms, beeches, maples, &c. Those of the secondary strata may, in general, be referred to known families, but in most instances require the formation of new genera; while those of the older strata, in numerous instances, cannot be classed in existing families, and must constitute new groups. Such are the Calamites, Lepidodendrons, and Sigillarias of the coal formation. From all the investigations of fossil botanists, however, it appears that the same great types existed in a former state as at the present day, viz., Cellular Cryptogamies, Vascular Cryptogamies or Acrogens, Monocotyledons and Dicotyledons, both Angiospermous and Gymnospermous.

1177. Schlotheim, Sternberg, Brongniart, and others, have proposed arrangements of fossil plants. That of Brongniart is founded on the resemblances which the fossils bear to living plants. He divides them into many branches, comprehending—

1st. Amphigenous Cryptogamous Vegetables, or Cellular Cryptogams, which he subdivides into two classes, the Fungi and Algæ.

2d. Acrogenous Cryptogamous Vegetables, comprehending two classes, the Musci, a numerous class of Filices, itself subdivided into five families,—the Ferns, Marsileaceæ, Characeæ, Lycopodiaceæ, and Equisetaceæ.

3d. Dicotyledonous Phanerogamous Vegetables, of which he enumerates the

several families, indicating the characters which approximate them to the same families of living vegetables.

4th. Monocotyledonous Vegetables *

When the analogy between a fossil and a living plant is such that the difference is not greater than occurs among the individuals included in a species of the living genus, then the fossil and living plant may be considered identical, and the epithet of fossil is applied to the name of the plant. If, on the other hand, the fossil presents distinct specific characters, but does not differ more from living species than these species differ among themselves, then it is looked upon as a new species of the genus. If the differences are well marked, but at the same time the organ which represents them is not of sufficient importance to induce the belief that the plant differs from others of the genus in all its essential organs, then the termination *ites* is added to the name of the genus. Thus, *Lycopodites* is a genus of fossil plants allied to *Lycopodium*, apparently not differing, so far as known, in essential and important parts; so also *Zamites* allied to *Zamia*, *Thuyites* to *Thuya*. If a fossil plant, although presenting several essential characters of a family, yet differs in the fossilized organ from all the known genera of the family, as much or more than these genera do among themselves, then it is to be considered as a new genus different from those actually existing. This will be seen in many of the coal fossils.

1178. Before proceeding to notice some of the more important fossil plants, it will be necessary to give a sketch of the geological constitution of the globe. The rocks of which the earth is composed are either stratified or unstratified; and the former are either fossiliferous or non-fossiliferous (azoic). The unstratified rocks, included under the names of Granitic and Trappean, are considered igneous, plutonic, or eruptive rocks, and do not generally exhibit any fossiliferous deposits. Vegetable structures in an altered condition have, however, been found in the substance of Trap rocks. Distinct evidences of Dicotyledonous structure charred have been discovered in various situations, more particularly near the Giant's Causeway, in Antrim, under the Trap rocks in the Island of Mull, and in those rocks of the same nature which are near Edinburgh. Fossil plants have not been found in certain metamorphic rocks such as Gneiss and Mica-slate, which present a stratified appearance, and appear to have been deposited in peculiar circumstances, probably at a high temperature. The igneous and metamorphic rocks are sometimes included under the comprehensive name of Hypogene. The non-discovery of fossil remains in rocks may, as Forbes has shown, indicate the formation of these rocks in positions destitute of organic life, such as are to be met with in the extreme depths of the ocean.

* Edin. Phil. Jour., vol. 48.

1179. **The Fossiliferous Stratified Rocks** have been divided into—1. Palæozoic (*παλαιός*, ancient, and *ζωή*, life), or those which contain the earliest fossil remains, including what have been called Transition, Primary fossiliferous, and Grauwacke rocks, and extending up to the Magnesian Limestone, immediately above the Coal measures. 2. Secondary, extending to the chalk. 3. Tertiary, extending to the recent deposits. The following tabular view is given by Ansted:—

I. Palæozoic.	
Older Palæozoic.....	Lower and Upper Silurian.
Middle Palæozoic.....	Devonian and old Red Sandstone.
Newer Palæozoic.....	{ Carboniferous System, Lower new Red Sandstone, and Magnesian Limestone or Permian System.
II. Secondary.	
Older Secondary.....	Upper new Red Sandstone or Triassic System.
Middle Secondary.....	Liassic, Oolitic, Wealden.
Newer Secondary.....	Cretaceous.
III. Tertiary.	
Lower Tertiary, or Eocene group.	
Middle Tertiary, or Miocene group.	
Newer Tertiary, or Pliocene group.	
Superficial Deposits, or Pleistocene group.	

1180. **Fossil Plants in Different Strata.**—The plants in these stratified rocks are either of a marine, fluviatile, lacustrine, or terrestrial nature, according to the state of the globe at the period of their deposition. The condition of the strata as regards fossils may depend in some measure on the depth at which they were deposited under the waters of the globe; for Forbes finds that in the present ocean there is, at a certain depth, a zero of animal, and probably of vegetable life. The state of preservation depends much on the nature of the plant in regard to its anatomical structure. Cellular plants, which are easily destroyed, have in a great measure disappeared, while plants which resist well the decomposing action of water and other agents, such as ferns, occur in great abundance. In the Silurian system, the fossils consist chiefly of invertebrate animals. Lignite has been detected by Hugh Miller in the old Red Sandstone of the north, and has been referred to some coniferous plants by Nicol. In the Carboniferous system, fossil plants occur in vast quantity. With the Palæozoic series one great epoch in the Rock formations was concluded, and a change took place so as to usher in the Secondary series. In the new Red Sandstone, the fossil remains are few and local, while in the Oolitic and Cretaceous systems they are more numerous. With the Secondary series of strata a general condition of the globe ended, and a new one commenced with the Tertiary strata. In these we meet with fossil remains nearly resembling or identical with the existing races. The names given to the groups indicate this. In the Eocene group (*ἑως*, dawn or morning, and *καινός*, new) we meet with a certain proportion

of living shells. In the Miocene (*μείων*, less,) the number of living species increases, although still less in number than the extinct ones; while in the Pliocene (*πλείων*, more,) the recent shells outnumber the extinct ones. The differences between the organic contents of one system and another, are in proportion to the interval of geological time elapsed between them; and the older the rocks, the more are the fossils distinct from the plants of the present day. The systems of organic life have always been adjusted to the actual condition of land and sea.

1181. The number of fossil plants known to M. Adolphe Brongniart, in 1836, was 527. In 1845, Goeppert and Bronn stated the number to be 1,792; and as in the 100,000 plants now known to exist in different parts of the globe, a large proportion consists of cellular plants, which would disappear in the process of fossilization, it would seem that the total number of known fossil species bears a considerable proportion to those now existing. Their numerical distribution in the different rocks is stated by Goeppert to be as follows:—

Older and Middle Palæozoic.....	52
Carboniferous.....	819
Permian.....	58
Triassic.....	86
Oolitic.....	234
Wealden.....	16
Cretaceous.....	62
Tertiary.....	454
Unknown.....	11
	<hr/>
	1792

From this table, Murchison remarks, it appears that the newer Palæozoic group contains more than half the known species of fossil plants,—a remarkable circumstance when it is considered that the great herbivorous land quadrupeds had no existence before the tertiary period. The small number of plants in the Cretaceous system is probably owing to the depth of the sea in which these formations were deposited.

1182. Among Dicotyledonous fossils there are numerous Amentacæ, Coniferæ, and Cycadacæ, besides many doubtful species. Among Monocotyledons, there appear to be plants belonging to Liliacæ, Smilacæ, Palmæ, Potamæ, Pandanacæ, &c. Among Acotyledons there is a vast number of genera and species belonging to Equisetacæ, Lycopodiaceæ, and Filices, and a few Marsileaceæ, Musci, and Algæ.

1183. It is impossible in a short treatise like this to allude to many of the fossil species of plants. It will be sufficient to indicate some of the more important genera. The vegetable remains met with in the Silurian and Devonian rocks are few and unimportant, compared with those which characterize the Carboniferous period; and their structure

seems to indicate either that they have been longer exposed to the agency of moving water, or that they are cellular marine species. An exception to this, however, occurs in the Lignite noticed by Miller. It is in the argillaceous and sandy beds of the Carboniferous system we first meet with decided proofs of the existence of land plants. These strata and sands alternate with beds of Coal, a mineral which is of vegetable origin, and which is deposited in various quarters of the globe in hollow troughs, the layers varying from the thickness of a few inches to 10 or 20 feet. It is rare to find coal exhibiting vegetable structure under the microscope; but in certain cases this may be evidently seen. I have seen it in the Arniston coal of this neighbourhood. Some have maintained that each stratum of coal is the product of a peculiar vegetation, frequently different from that which precedes, and from that which follows it. Hence each stratum is often characterized by the predominance of certain impressions. In examining a coal seam, there are frequently evidences of three distinct phases. In the underclay there are roots permeating the mass; then comes the coal, which is either formed out of the plants whose roots are in the clay, or of others which have grown with them, or have been drifted; and lastly, above the coal there is the shale, which appears to have supported a vigorous vegetation. Unger has enumerated 683, and Brongniart 500 plants as belonging to the Coal Flora of Britain. The latter is equal to about 1-20th of the number of species now growing on the surface of European soil.

1184. M. Brongniart, from the investigation of the several geological formations, has arrived at the conclusion, that three distinct periods of vegetation can be established. In the most ancient periods, the predominance of acrogenous cryptogamous vegetables (*Ferns* and *Lycopodiaceæ*), later, the predominance of gymnospermous dicotyledons (*Cycadææ* and *Coniferæ*), without any admixture of angiospermous dicotyledons; and, in the last place, during the chalk formation, the appearance and predominance of angiospermous vegetables, both dicotyledons and monocotyledons, may be considered as characteristic. These differences lead M. Brongniart to recognize three long periods of vegetable growth, which he terms the reign of the Acrogens, the reign of the Gymnosperms, and the reign of the Angiosperms, as indicative of the successive predominance of each of these three great divisions of the vegetable kingdom, rather than the complete exclusion of the others. Each of these three kingdoms is commonly subdivided into many periods, during which traces of the same family and genera are discoverable. These periods comprehend many epochs, during which vegetation does not appear to have undergone any notable changes. Materials are often wanting to precisely establish these subdivisions, either from a want of accuracy in determining the exact geological position of beds enclosing vegetable

impressions, or because the division of the various species in the different beds of the same formation has not been carefully established. M. Brongniart proposes the following general division of the fossil kingdom :—

I. REIGN OF THE ACROGENS.

1. CARBONIFEROUS PERIOD.

(Not susceptible of subdivision into epochs in the present state of our knowledge.)

2. PERMIAN PERIOD.

(Forming only one epoch ?)

II. REIGN OF THE GYMNOSPERMS.

3. VOSGIAN PERIOD.

(Constituting a single epoch.)

4. JURASSIC PERIOD.

Keupric epoch, Lias epoch, Oolitic epoch, Wealdian epoch.

III. REIGN OF THE ANGIOSPERMS.

5. CRETACEOUS PERIOD.

Subcretaceous epoch, Cretaceous epoch, Fucoidian epoch.


6. TERTIARY PERIOD.

Eocene epoch, Miocene epoch, Pliocene epoch.


1185. I. Reign of the Acrogens.—In this period Acrogens and Gymnosperms are found to have existed simultaneously, the former predominating over the latter in number and size. The number of the Fern family, the great development of the Lycopodiaceæ, are the most prominent characters of this epoch. Other anomalous families belonging to the Gymnosperms are also with them discoverable, which evidently differ from the actually existing families of that department.

1186. Fossil Plants of the Carboniferous System.—The great mass of fossil plants of this system belong to Acrogenous Cryptogamies and Dicotyledonous Gymnosperms. Some of the plants called Palms, such as *Nöggerathia*, *Flabellaria*, and *Artisia*, are referred by Brongniart to the latter division. Ferns abound in this system, especially in the clays, ironstones, and sandstones. The species are included under the genera *Sphenopteris*, *Pecopteris*, *Neuropteris*, *Odontopteris*, *Cyclopteris*, *Glossopteris*, and *Lonchopteris*. These plants rarely exhibit any traces of fructification, in consequence of only one surface of the fronds being exposed to view, and they are therefore distinguished chiefly by the shape and venation of their fronds. Thus, *Sphenopteris* (*σφήν*, a wedge, and *πτέρις*, a fern), has a bi-tripinnatifid frond, pinnæ narrowed at the base (cuneate), not adherent to the rachis, lobed, veins bipinnate, somewhat radiating from the base (fig. 793). In *Pecopteris* (*πέκω*, I comb), the frond is pinnatifid or bi-tripinnatifid, pinnæ adnate to the rachis, sometimes confluent, a strong primary vein reaching the apex, the secondary veins being nearly straight, simple or forked, rarely

pinnate, sori rounded at the end of the secondary veins (fig. 794). In *Neuropteris* (*νεῦρον*, a nerve), the frond is pinnate or bipinnate, pinnæ



subcordate at the base, distinct from the rachis, strong primary vein vanishing towards the apex, secondary veins oblique, arched, repeat-



edly dichotomous (fig. 795). Tree-ferns appear to have existed in Britain during the deposit of the coal strata, and to have occupied an

Fig. 793.—*Sphenopteris Hæninghausii*, a fern of the Carboniferous system.

Fig. 794.—*Pecopteris aquilina*, another fern.

Fig. 795.—*Neuropteris Loshii*, another fern.

Fig. 796.—*Sphenophyllum dentatum*, one of the dubious forms of the Carboniferous system, perhaps allied to *Salisburia*.

Fig. 797.—*Annularia brevifolia*, a coal plant, the affinities of which are unknown.


important place in the flora. The stems of these ferns are included under the genus *Caulopteris*. The fronds have not been found attached; but it is probable that some of the fronds found in the coal measures have been connected with these stems. Other plants, considered by some authors as resembling Marsileaceæ, have also been found in certain localities; such as *Sphenophyllum* (fig. 796) and *Annularia* (fig. 797).



1187. Some fossil plants, allied to *Lycopodium*, also occur in the coal measures. Brongniart believes they are more abundant in the ancient beds than in the superior beds of the greater part of the coal formation. These have been included under the genera *Lycopodites* (figs. 798, 799) and *Selaginites*. Some fossil species allied to these are grouped under the genus *Lepidodendron* (λεπίς, a scale, and δένδρον, a tree) (figs. 800, 801). They appear to occupy an intermediate place between Coni-



800



801

- Fig. 798.—*Lycopodites* (*Walchia Schlotheimii*), a plant allied to *Lycopodium*.
 Fig. 799.—*Lycopodites* (*Walchia hypnoides*), another *Lycopodiaceous* plant.
 Fig. 800.—*Lepidodendron crenatum*, with the scars of the leaves on its stem. It belongs to a family of plants apparently intermediate between *Coniferæ* and *Lycopodiaceæ*.
 Fig. 801.—*Lepidodendron elegans*, with its dichotomous trunk, and linear acute leaves.

feræ and Lycopodiaceæ. Their leaves are arranged in the same manner as some of the Coniferæ, and their scars are similar. Their branches bifurcate like Lycopodiaceæ. They occur in the form of dichotomous trunks, 20 to 45 feet high, with linear or lanceolate leaves (fig. 801) like those of some species of *Lycopodium* and *Eutassa*. The stem consists of a thin cuticle, a double cellular zone, a hollow vascular cylinder, and a pith. The tissue in the outer part of the double cellular zone is arranged like that of ferns, the vascular cylinder is about $\frac{1}{8}$ of an inch thick, and consists of polygonal tubes marked with lines, while the pith is composed of fusiform cells. The stems are marked with rhomboid and orbicular scars (fig. 800). Their cone-like fruit is said to occur in a fossil form called *Lepidostrobus*; but under this name, as well as that of *Strobilites*, the fruit of many different plants seems to be included. *Lepidophyllum* is said to be the leaf of some *Lepidodendron*, and *Ulodendron* is an allied genus.

1188. *Stigmaria* (στίγμα, a mark or impression) is a fossil genus, the species of which abound in the coal measures. They occur generally in the bed called the Underclay. *Stigmaria ficoides* (fig. 802) is the common species. It consists of a dome-shaped mass sending forth grooved and pitted branches, which divide dichotomously, and extend 20 to 30 feet. Slender processes are given off, which appear to have been hollow (fig. 802). These processes (called fistular leaves,)



802



803

form an entangled mass traversing the argillaceous lower bed in every direction. In *Stigmarias* three tissues are met with, vascular tissue forming the inner part of the cylinder, ligneous forming the wood, and cellular tissue forming a broad cortical zone, as well as the central


Fig. 802.—*Stigmaria ficoides*; a branch giving off fistular leaves, which traverse the Underclay in all directions.

Fig. 803.—*Sigillaria pachyderma*; the stem marked with scars, and fluted longitudinally.

portion or pith. Some think that the stores of fossil fuel in England and America are mainly due to the presence of this plant. There are no plants of the present day which resemble *Stigmarias*. Some have referred them to such genera as *Cactus* and *Euphorbia*, others to *Isoetes*. King says that they are the roots of the fossil plants called *Sigillaria*.

1189. *Sigillaria* (*sigillum*, a seal,) is another plant which appears to have aided in the formation of coal. It occurs in the form of compressed stems, attaining a height of 40 to 50 feet, and a breadth of 5 feet. The stems are fluted longitudinally, and marked at regular intervals by single or double scars, the remains of the leaf insertions (fig. 803). Some suppose *Sigillarias* to be allied to Tree-ferns, others to *Coniferæ*. Brongniart says they resemble *Zamia integrifolia*, and appear to predominate in the middle and superior beds of the coal formations. King thinks that they are intermediate between Ferns and *Cycadaceæ*, that the fronds called *Neuropteris* constitute their foliage, and that *Stigmarias* are their roots. They have a medullary sheath in the shape of apparently isolated bundles, and vessels intermediate between true spiral and scalariform. The bark is said to be composed of two different layers, thus giving rise to different impressions. The furrows or fluted marks are due to the arrangement of the leaves on the stem. King says, that if in imagination we delineate a channelled stem of any height between 12 and 100 feet, crowned with a pendant fern-like foliage, furnished with wide spreading thickly fibrilled roots, and growing in some densely-wooded swamp of an ancient Mississippi, we will then have formed a tolerably close restoration of a *Sigillaria* vegetating in its true habitat.

1190. *Calamites* (καλαμος, a reed), another coal fossil plant, occurs in the form of jointed fragments, originally cylindrical, and perhaps



804




805

hollow, but now crushed and flattened (fig. 810). The stems are branched (fig. 805), and there appears to have been a distinct wood

Fig. 804.—*Calamites Suckovii*, composed of jointed striated fragments having a bark.
Fig. 805.—*Calamites cannaeformis* giving off branches.

and bark. Their internal cavity seems to have been separated by horizontal partitions at the articulations, the intervals between the articulations becoming smaller towards the ends of the branches. Both stems and branches are ribbed and furrowed (fig. 804). Some refer the numerous species of *Calamites* to *Equisetaceæ*, but the presence of wood and bark has led others to place them among *Dicotyledons*.

Some interesting fossil *Coniferæ*, included under the names *Pinites* and *Araucarites*, are found in the carboniferous sandstone, as in Craigleith Quarry, near Edinburgh. The specimens found in Craigleith have been referred to *Pinites Withami* and *medullare* of Lindley and Hutton (*Araucarites* of Goeppert, and *Dadoxylon* of Endlicher). Some of these seem to be allied to the *Araucaria* tribe; for instance to *Eutassa excelsa*, the Norfolk Island Pine. Their wood, under the microscope, exhibits the punctated appearance and other characters of *Coniferæ* (figs. 806-809), and the disks are often in two or more rows (fig. 808).



Some specimens have been found upwards of 70 feet long. These *Coniferæ* seem not to have been associated with the *Sigillarias* and the other plants which abound in coal seams. They probably flourished in the neighbourhood, and were at times transported to these localities. A peculiar kind of fossil fruit called *Trigonocarpum*, resembling that of a Palm, has been found in some of the carboniferous sandstones. *Nöggerathia* and *Flabellaria* have been referred to species of Palms, but Brongniart considers them as *Cycadaceous*. *Lyginodendron* (λύγινοδενδρον, wicker work,) is a peculiar coal fossil discovered by the Rev. Mr. Landsborough in Ayrshire, and described by Mr. Gourlie. Its im-

Figs. 806-809.—The structure of wood in recent *Coniferæ*, to illustrate the appearances presented by some fossil woods.

Fig. 806.—Transverse section of a piece of Coniferous wood, of the natural size.

Fig. 807.—A section of the same wood seen under the microscope. The medullary rays and woody tubes seen without any large porous vessels.

Fig. 808.—Longitudinal section of the same, in the direction *B C*, magnified. A medullary ray seen crossing the woody tubes, which are marked by disks, in one or more rows.

Fig. 809.—Section of the same in the direction *A B*, perpendicular to the medullary rays, which are seen at intervals between the woody fibres.

pression consists of rounded narrow twigs, which cross each other like the parts of an osier basket. *Sternbergia* is a peculiar fossil, consisting of horizontal plates, which are held together by some connection in the axis. Some look upon it as allied to *Dracæna*. It may be remarked, in general, that the Carboniferous flora is uniform, or nearly so, in all parts of the globe where carboniferous fossils have been obtained, viz. the whole of western, northern, and eastern Europe, North America, from Alabama to Melville Island, various districts of Asia, Eastern Australia, and Van Diemen's Land, and probably the Asiatic Islands.

1191. As the great mass of fossils in the coal formation consists principally of ferns, Brongniart has been led to draw conclusions as to the climate of the globe, at the time when the coal fossils grew. Ferns of the present day thrive best in a moist insular climate, and many of them occur in tropical climates. Hence Brongniart conjectures that at the coal epoch the surface of the earth consisted of a series of islands in the midst of a vast ocean, and that the temperature was higher generally than that of the present day. In the forests of these islands lofty *Lepidodendrons* would occur with their delicate and feathery fronds; *Sigillarias*, with their fluted stems and enormous matted roots; *Calamites*, with their singular branches; Tree-ferns and Coniferous plants, resembling the Norfolk Island Pine, and towering 100 feet above the rest of the forest. He also thinks that the immense deposits of carbon at that epoch, warrant the conclusion that the air contained a large amount of carbonic acid. These conclusions are, of course, mere hypotheses. In regard to the temperature, it may be remarked, that there is no evidence from the nature of the flora, of a marked increase of temperature at the coal epoch. In New Zealand, which is in a latitude the same as that of a great part of Europe, a very large proportion of the vegetation consists of Acrogenous plants. Ferns and their allies, in that country, cover immense districts, replacing the grasses of other countries, and giving a marked character to all the open land. Some of the ferns attain a height of 30 or 40 feet, and occur in groups. *Hemitelia capensis* too, a Tree-fern found at the Cape, was also seen by Gardner, at an elevation of 6,000 feet, on the Organ mountains, thus showing a capability of enduring a great range of climate, and warning us against hasty conclusions on the subject of the temperature of the world at the coal epoch.

Dr. Hooker thinks that the prevalence of ferns may be regarded as a probable evidence of the paucity of other plants, and the general poverty of the whole flora which characterized the formation. He is led to these conclusions from observing the mode in which the ferns in Van Diemen's Land and New Zealand monopolize the soil, choking plants of a larger growth on the one hand, and admitting no undergrowth of smaller species on the other. In New Zealand he has col-

lected 36 kinds of ferns on an area not exceeding a few acres; they gave a most luxuriant aspect to the vegetation, which presented scarcely a dozen flowering plants and trees besides.

1192. Some have supposed that the plants of the coal fields have been drifted into basins, others that they grew in the spots where they are now found. Beaumont thinks that all the vegetables which are now converted into coal, grew in swampy islands, covered with a luxuriant vegetation, which accumulated in the manner of peat-bogs; that those islands having sunk beneath the ocean, were there covered with sand, clay, and shells, till they again became dry land, and that this operation was repeated in the formation of each bed of coal. The occurrence of stems of trees in an erect state (fig. 810), appeared to



810

him to confirm the view that the trees were in situ. Ansted says, that although many trees are found in the coal measures in an erect or highly inclined position, there is no reason for believing that they grew on the spot where they are met with. He rather thinks that they have been caught or stopped in their passage down a rapid stream, and, like the snags in some of the great American rivers, have been detained till the lower portion was firmly embedded in the rapidly forming sandstone. The embedding of stems in strata of sandstone, is similar to what Gardner saw near the mouth of the Rio San Francisco, where

Fig. 810.—Vertical stems of fossil trees. Calamites chiefly, found in the Coal measures of Treuil, near Saint Etienne.

coco-nut trees were found with their stems immersed to the depth of 50 feet or more in the embankment of sand which stretches along the shore. Phillips remarks, that the condition of the plants which compose the coal, the general absence of roots, the fragmentary state of the stems and branches, the dispersed condition of the separable organs, all confirm the conclusion that the plants have been swept down from the land on which they grew by watery currents, often repeated, and deposited in basins and large estuaries of the sea, or perhaps rarely in lakes of fresh water.

1193. The nature of the vegetables which appear peculiar to the Permian period, M. Brongniart believes is far from being determined in a positive manner, in consequence of the uncertainty which is attendant on the period of certain geological formations. On account of these doubts, M. Brongniart indicates three separate floras. 1st, The Flora of the bituminous slates of Thuringia, composed of Algæ, Ferns, and Coniferæ. 2d, The Flora of the Permian sandstones of Russia, which comprehends Ferns, Equisetaceæ, Lycopodiaceæ, and Meggerathiæ. 3d, The Flora of the slates of Lodène, which is composed of Ferns, Asterophyllitæ, and Coniferæ. It is observed that many of the fossils met with in the bituminous slates of Thuringia appear to be marine plants, and that hitherto no species common to these localities has been found.

1194. **II. Reign of the Gymnosperms.**—In the Carboniferous period, the Acrogenous Cryptogams were found to predominate, while the Gymnosperms Dicotyledons, less numerous, showed themselves in unusual forms. At a later period, these anomalous forms were observed to disappear, and Acrogenous Cryptogams and Gymnospermous Dicotyledons to evidently enter into existing families, differing from them only in generic forms; the Ferns and Equisetaceæ, which represent the Acrogens, were less numerous, the Coniferæ and Cycadæ almost equalling them in number, and usually exceeding them in frequency.

The reign of the Gymnospermous Dicotyledons is divided into two periods: the first, in which the Coniferæ predominate, and in which the Cycadæ scarcely appear; the second, in which this family becomes preponderating in the number of species, in frequency and variety of generic forms.

1195. Brongniart believes the Vosgian period not to have been of long duration, and to comprehend only the variegated sandstone, properly so called. It is characterized by the existence and presence of ferns of forms often anomalous, appertaining to genera now extinct, while Gynosperrmous plants are represented by several genera of the Coniferæ, of which the species and specimens belonging to the *Voltzia* and *Haidingeria* are very numerous. The Cycadæ, on the contrary, are very rare.

1196. The Jurassic period of Brongniart includes—1. The *Keupric Epoch*, whose vegetables consist of Algæ, of Ferns and Equisetaceæ, of Cycadeæ and Coniferæ, and also of two doubtful Monocotyledons (*Palæoxyris* and *Preisleria*). 2. The *Lias Epoch*, whose essential characters are the great predominance of Cycadeæ, and the existence among the ferns of many genera with reticulated venation. 3. The *Oolitic Epoch*, which furnishes Algæ, Ferns, Marsileaceæ, Lycopodiaceæ, Equisetaceæ, Cycadeæ, and Coniferæ. The distinctive characters of this epoch are—amongst the Ferns, the rarity of reticulated venation so numerous in the Lias; the greater approach of the Cycadeæ in character to those now existing; the diminution of genera much more remote from living species, and the greater frequency of the Coniferæ. 4. The *Wealdean Epoch*, which affords Algæ, Ferns, Cycadeæ, and Coniferæ, and some Carpolithes, as plants of a doubtful class.

1197. **Fossil Plants of the Secondary Strata.**—The plants of the Secondary series of strata are thus shown to be different from those of the Carboniferous system. The *Stigmarias*, *Sigillarias*, and *Lepidodendrons* cease, and are replaced by ferns and their allies. Plants of the *Zamia* tribe occur along with some Coniferæ. The pinnated frond called *Pterophyllum* (fig. 811), appears to belong to a Cycadaceous plant allied to *Zamia*, while *Voltzia* (fig. 812) and *Peuce* seem to be Coniferous genera. In the secondary formations, generally, there are no true coal fields, although carbonaceous matter, in the form of imperfectly bituminized lignite, is found deposited occasionally. Notwithstanding the absence of true coal, the sandstones and shales of the Oolitic system contain numerous fossil plants, which in their form and character seem to be a transition between the Palæozoic flora, and that of more recent formations. In the upper Oolite at Portland, there is a bed of earthy brown matter, about a foot in thickness, commonly known as the Dirt-bed, in which there is an assemblage of silicified stumps of large trees (fig. 813). They are generally in an erect position (fig. 814), and appear to be in situ. They are probably Cycadaceæ. Plants of this natural family, in the form of



811


Fig. 811.—*Pterophyllum Pleiningerii*, apparently the frond of a fossil Cycadaceous plant allied to *Zamia*.

Zamites (fig. 816), *Mantellia* (fig. 814), *Cycadites*, and *Pterophyllum* (fig. 815), may be said to characterize the fossil flora of the older



812

oolite beds. These plants take the place of the *Lepidodendrons* and *Sigillarias* of the carboniferous period. The stems are found without



813



814

any leaves attached, but some suppose that the fronds called *Otopteris* are the leaves of some of the species. The occurrence of these tropical forms seems to indicate a warmer climate than now exists in Britain. There are 6 existing genera of Cycadaceæ (§ 1047). Two of these are represented in figs. 817, 818, showing the forms of the stems and leaves or fronds. There are 5 fossil ones according to authors. The fossil Cycadaceæ known up to 1845, and distinguished by specific names, amount, according to Goeppert, to 78. Amongst these are 9 stems or stipes, 65 fronds, and 4 fructifications. The genera are :—


Fig. 812.—*Voltzia heterophylla*, one of the fossil coniferæ of the Triassic system.

Fig. 813.—The Dirt-bed of Portland, containing stumps of fossil Cycadaceæ in an erect position.

Fig. 814.—*Mantellia nidiformis* (Cycadeoidea), one of the silicified Cycadaceæ of the Portland Dirt-bed.



815



816



817



818

Fig. 815.—*Pterophyllum Williamsoni*, the frond of one of the fossil Cycadaceæ.

Fig. 816.—*Zamites*. One of the fossil Cycadaceæ.

Fig. 817.—*Cycas revoluta*, one of the species of *Cycas* of the present flora of the globe, with its scale-like stem and pinnate fronds.


Fig. 818.—*Zamia pungens*, one of the Cycadaceæ at present existing on the globe.

		Stems.	Fronds.	Fructification.
Cycadites,	11	4	7	—
Zamites,	28	5	23	—
Zamiostrobus,	4	—	—	4
Pterophyllum,	23	—	23	—
Nilsonia,	12	—	12	—
	78	9	65	4

To this some add *Mantellia* and *Palæozamia*. The greater number of these is found in the Oolite and Lias. In the inferior Oolites are also seen peculiar Coniferæ, belonging to the genus *Brachyphyllum* (fig. 819), and a plant said to be a true *Equisetum* (fig. 820). The fossil



819



820

Flora of the Wealden formation resembles, in many respects, that of the Oolitic system. Fronds of ferns, leaves of Coniferæ (*Sphenophyllum*), and of Cycadaceæ, occur along with some Fucoids. There are also silicified coniferous trunks, and two peculiar genera called *Clathraria* and *Endogenites*. The former is an arborescent plant, having its bark formed by the union of the bases of the leaves, and covered by distinct scales. The latter is apparently a Monocotyledon, perhaps allied to Palms. Some fruits, resembling those of Palms, are also found. In the Cretaceous system, there occur fossilized Dicotyledonous leaves and fragments of wood, marked by perforations of marine animals. Some of the Dicotyledons are Coniferous, others Cycadaceous.

1198. **III.—Reign of the Angiosperms.**—The dominating character of this last transformation of the vegetation of the globe is the appearance of Angiospermous dicotyledons, those vegetables which constitute

Fig. 819.—*Brachyphyllum*, a Coniferous genus of the Oolitic system.

Fig. 820.—*Equisetum columnare*, a fossil species of the Oolite.

more than three-fourths of the vegetable creation of our epoch. M. Brongniart was of opinion that these vegetables did not begin to appear till after the chalk, with the earliest beds of the tertiary formations; but recent investigations have shown him that beds belonging to the chalk formation present some very distinct examples. In this reign, Brongniart distinguishes two great periods—1st, The Cretaceous period, a kind of period of transition, in which a few Cycadæ and Angiosperms are found; 2d, The Tertiary period, presenting all the characters arising from the predominance of Angiosperms, Dicotyledons, and Monocotyledons, and divisible into several epochs. The *Fucoidian epoch* seems to form the natural limit between the cretaceous and tertiary periods, is characterized by deposits rich in Algæ, of a very peculiar form. M. Brongniart has not found land plants mingled with these marine species, nor does he believe that fossil woods have been met with.

1199. The Tertiary period is considered to have its characters the best marked. The abundance of Angiospermous dicotyledons, and different monocotyledonous families, but especially the Palms, immediately distinguish it from the most ancient periods, while in the Cretaceous period the Angiosperms appear nearly equal to the Gymnosperms. In the Tertiary period they are found to greatly exceed them. At the Cretaceous epoch, Cycadæ and Coniferæ, allied to the genera inhabiting tropical regions, existed; during the Tertiary period, the Cycadæ appear to have been completely wanting in Europe. M. Brongniart, having divided the Tertiary period, as regards plants, into the Eocene, Miocene, and Pliocene, believes that he shall find the number of species in the great divisions are thus distributed in these three Floras:—


	Eocene Epoch.			Miocene Epoch.			Pliocene Epoch.	
Cryptogams,	33	—	10	—	13	—
Amphigenous,	—	16	—	6	—	6
Acrocycogenous,	—	17	—	4	—	7
Phanerogams,	—	—	—	—	—	—
Monocotyledons,	33	33	26	26	4	4
Dicotyledons,	143	—	97	—	195	—
Gymnosperms,	—	40	—	19	—	31
Angiosperms,	—	103	—	78	—	164
	209 —			209 —			212 —	

1200. With regard to the characters drawn from vegetable forms during these three epochs, the most remarkable appear to be—1st, In the *Eocene* period, the presence, but rarity of the plants, limited to a small number of the species. The predominance of Algæ and marine Monocotyledons, which is ascribable to the great extent of marine formations during this epoch. The existence of a great number of extra


European forms, resulting especially from the presence of the fossil fruits of Sheppey. 2d, In regard to the *Miocene* epoch, the abundance of Palms in the greater number of localities, belonging, without doubt, to this epoch, and the existence of a number of non-European forms. 3d, The *Pliocene* epoch is known by the great predominance and variety of Dicotyledons, the rarity of Monocotyledons, and, above all, the absence of Palms, as well as the general analogy of the forms of these plants with those of the temperate regions of Europe, North America, and Japan.

1201. Fossil Plants of the Tertiary Strata.—From these observations it may be inferred that with the Chalk formation, the ancient condition of organized beings appears to have ended, and a new one to commence in the Tertiary series. The plants of the Palæozoic and Secondary series are all extinct. Those which occur in the tertiary are totally different from all that have previously appeared. With the chalk, Ansted says, we close, as it were, one great volume of the history of animated creation. Everything up to this point belongs to the past; everything on this side of it may be ranked among indications of the present. New forms, new types of organization, corresponding to different habits and altered circumstances, now replace those which have passed away. The conditions under which animals and vegetables lived were changed, and a new epoch commenced upon the earth.


1202. Except a few doubtful fossils of the lowest organization, there are none common to the secondary and tertiary periods. The tertiary series are well seen in the south of Europe, Asia, and America. In Britain the tertiary deposits are met with in the London clay, in Hampshire and the Isle of Wight, the Suffolk and Norfolk Crag, and



821



822



823

Figs. 821-825.—Structure of ordinary Dicotyledonous stems, to illustrate the appearances presented by some tertiary fossil woods.

Fig. 821.—Portion of a Dicotyledonous (Exogenous) stem cut transversely. Natural size.

Fig. 822.—Section of the same magnified, to show the occurrence of large porous vessels. The ordinary Dicotyledons differ in this respect from Coniferæ.

Fig. 823.—Longitudinal section of the same in the line A B, perpendicular to the medullary plates, showing a large porous vessel, and the rays appearing here and there among the woody tissue.

in the Till of the Clyde. The London clay contains numerous fruits belonging to many hundred species of plants. The first tertiary land




824



825


of which we have knowledge, seems to have been richly clothed with plants. The strata are, generally speaking, rich in fossils. The stems



826



827



828

Fig. 824.—Leaf of an unknown fossil Elm of the middle Tertiary epoch.

Fig. 825.—Leaf of *Comptonia acutiloba*, an Amentaceous plant of the same epoch.

Fig. 826.—Section of a recent Palm stem, to show its structure. The dark dots marking vascular bundles in the midst of cellular tissue.

Fig. 827.—A portion of the same magnified, to show the vascular bundles.

Fig. 828.—*Palmitis Lamanonis*. Leaf of a Monocotyledon resembling a Palm.

and leaves appear to be those of Dicotyledons, little differing from the plants of the present day (figs. 821-825). In the brown coal of this series, the structure of the wood is evident, and distinctly exogenous (figs. 821-823), and there are often associated with it leaves of Poplars, Elms (fig. 824), and other forest trees. The fossil plants of the Isle of Sheppey have been examined by Bowerbank, and have led to the determination of several hundred species of plants, all of them extinct, and all resembling those of warmer climates. Fruits of *Nipadites* (*Pandanocarpum*), a fossil plant, allied to Nipa, one of the Pandanaceæ; *Hightea*, a five-seeded fruit, probably Malvaceous; also the fruit of a Proteaceous plant, and of species allied to Canna, Cucum-ber, and the Leguminosæ and Coniferæ of the present day. To some of them the names of *Cupanoides*, *Wetherellia*, *Cucumites*, and *Mimosites*, have been given. In some of the tertiary formations there occur pieces of wood, which present the structure of that of Pepper-plants and of Palms (figs. 826, 827), and there are also leaves which have the flabelliform appearance of Palm leaves, included under the name of *Palmacites* (fig. 828). Specimens allied to Chara are also found, with their fructification denominated *Gyrogonites*. Above the tertiary strata, and forming a continuous series with the formations of the present day, occur the superficial deposits or the Pleistocene (πλειστός, most,) group. They are ascribed to the violent action of water, and are sometimes called Diluvial.

1203. There have thus occurred in this earth successive deposits and creations of plants and animals. These deposits have been variously altered and distributed, but all have proceeded upon one grand and harmonious principle, and all speak the same language. They tell us of changes which have taken place during the preparation of this world for the abode of man, the noblest of the Great Creator's works. Many are afraid to enter upon the investigations of Palæontology, from a mistaken idea that they are at variance with Scripture. We need not fear, however, that in prosecuting our scientific researches, we shall ever arrive at a point where the knowledge of nature will be found at variance with the truths of Scripture. The more each is studied, the more shall we find occasion to admire the harmony that subsists between them, and the beautiful light of illustration which they reciprocally shed on one another.

For further details on the subject of Fossil Botany, consult Brongniart's *Plantes Fossiles*, Hutton and Lindley's *Fossil Flora*, Witham on *Fossil Plants*, Goeppert's *Fossil Ferns*, Goeppert on the *Flora of the Brown Coal Formation*, in *Arbeiten der Schlesischen Gesellschaft* for 1847, and in the *Quarterly Journal of the Geological Society*, Vol. V., King's Papers on the *Coal Fossils* in the *Edinburgh Philosophical Journal*, Hooker's Papers in the *Report of the Geological Survey of Great Britain*; besides Geological works, such as Ansted's *Geology*.

APPENDIX.

I.—ON THE USE OF THE MICROSCOPE IN BOTANICAL RESEARCHES.

1204. THE Microscope (*μικρὸς*, small, and *σκοπέω*, I see,) is an instrument for enabling the eye to see distinctly objects which are placed at a very short distance from it, or to see minute objects that would otherwise be invisible. It has been used with great success in the examination of vegetable structure. To it we are indebted for a knowledge of the various component parts of plants, the motions of the juices and ciliary movements, as well as for the facts connected with the development of the embryo. It is an instrument, however, which requires to be used cautiously; and the conclusions drawn from it ought to be carefully weighed, more especially when the observations have been made with high magnifying powers.

1205. **Lenses.**—Before proceeding to notice the construction of simple and compound microscopes, it will be advantageous to notice the different kinds of lenses used, and the sources of error which require to be guarded against in their preparation. The chief forms of lenses used are, the *double-convex*, with two convex faces; *plano-convex*, with one face flat and the other concave; *double-concave*, with two concave faces; and *plano-concave*, with one flat and one concave face. Sometimes, also, a *meniscus* is used, with a concave and a convex face, and a sharp edge. In the use of ordinary lenses, there are sources of error from the form of the lens, and the nature of the material of which it is made. When parallel rays fall on a double-convex or a plano-convex lens, they are brought into a focus at a certain distance: but it is found that no lens with a spherical surface can bring the rays of light coming from one point exactly into the focus at another point. Hence arises what is called *spherical aberration*. Moreover, the material of which the lens is made acts differently on the different portions of each ray, and separates the white light into different colours, which have various degrees of refrangibility. This gives rise to *chromatic* (*χρῶμα*, colour,) *aberration*. To remedy these defects, certain combinations of glasses have been adopted, so that the light traversing one lens through the centre may pass through near the margin of another. The confusion produced by these aberrations may be greatly lessened by diminishing the pencil of light; for instance, by employing a stop or diaphragm, which lessens the aperture of the lens, and cuts off the peripheral rays. In lenses of low power, such as are used in the simple dissecting microscope, these aberrations will not cause much confusion. It is only when high powers are required that these aberrations must be done away with, the aperture being increased without interfering with definition. The invention of Wollaston's doublet with two lenses, and Holland's triplet with three, was with the view of diminishing, as far as possible,

these aberrations. In this, however, they were not successful, for coloured images were still produced. Their lenses were constructed of the same kind of material; and it was afterwards found that in order that lenses might present the object uncoloured, or be what is called *achromatic* (α , privative, and $\chi\rho\omega\mu\alpha$, colour), it was necessary to use glasses of two different densities. Achromatic lenses, or such as are nearly free from aberration, are constructed by placing together glasses of different dispersive powers, and of different forms. The usual achromatic consists of a double-convex lens, made of plate or crown-glass, and a plano-concave, made of flint-glass, fitted accurately to it, and cemented by Canada balsam. Sometimes three lenses are used, the middle one being double-concave, and in that case they cannot be cemented.


1206. **Microscopes** are of two kinds—Simple and Compound. By the *Simple microscope*, objects are viewed through a single lens, or through two or three lenses placed together, so as to form doublets or triplets. The glass is arranged so that it can be brought over the object, and adjusted, by means of a rack and pinion, or by some other contrivance, to its exact focal distance, the object, when opaque, being seen by light thrown from above, and when transparent, by light transmitted from below. This instrument, when used with single lenses or doublets, is the best for ordinary botanical investigations, more especially for dissections. The combination of three lenses approaches too near the object to be easily used. A very high power may be obtained by means of the lenses termed Coddington's and by doublets formed of plano-convex glasses. The chief objections to the simple microscope are the fatigue attendant on long-continued investigations, and the small field of view. In the simple microscope, glasses of the following focal lengths may be employed—viz., $1\frac{1}{2}$ inch, $\frac{3}{4}$, $\frac{1}{2}$, $\frac{1}{4}$; and, if very minute objects are to be examined, of $\frac{1}{16}$, $\frac{1}{32}$, or $\frac{1}{64}$ of an inch.

1207. In the *Compound microscope* there are two sets of lenses,—the one called the *object-glass* or objective, the other the *eye-piece* or ocular. The first receives the rays from the object, and bringing them to new foci, forms an image, which the second treats as an original object, and magnifies it just as the single microscope magnified the object itself. In the construction of the object-glasses, great care is taken to render them achromatic. Those made by the most eminent London makers consist of two or three compound lenses, which cannot be used separately, but are fixed together in a tube. In the case of high powers, the object-glasses are also provided with an adjustment for the thickness of the glass covering the object to be viewed. The eye-piece, also, must be so formed as to be free from error. That used is call Huyghen's, and consists of two plano-convex lenses with their plane sides towards the eye, and placed at a distance apart equal to half the sum of their focal lengths, with a diaphragm placed midway between the lenses. In this eye-piece, the lens next the eye is called the eye-glass, the other the field-glass. The eye-pieces supplied with the best microscopes are usually three; and they are so constructed, that, with each of the object-glasses, they give a certain amplification of the object, the powers being in the proportion of 1, 2, and 3, or 1, $1\frac{1}{2}$, and $2\frac{1}{2}$. In the best microscopes there is also an *achromatic condenser* or eclairage, through which the light reflected from the mirror passes. The amplification by means of an eye-piece in the compound microscope enables us to use an object-glass of a lower power than would otherwise be necessary. This kind of microscope, when well constructed, gives a flat and colourless picture of the object, with clearness of definition. The observer can use it for a length of time with less fatigue than when employing the simple microscope.

1208. In examining vegetable structures, an instrument magnifying 150 to 200 diameters is usually sufficient; but in some instances higher powers are required. Achromatic object-lenses of $1\frac{1}{4}$, $\frac{2}{3}$, and $\frac{1}{4}$ of an inch, are recommended as the most essential; and two eye-pieces should be provided, one of about $1\frac{1}{2}$ and the other of $2\frac{1}{2}$ inches in length. The instrument should have both a coarse and a fine adjustment; and it is of importance that it should be made to incline at an angle, or to stand horizontally. A moveable stage is also useful, so that the different parts of the object may be viewed without being touched by the fingers, and a spring-holder to fix the objects on the stage.

The following figure (829) represents an achromatic microscope of modern construction. It consists of a massive circular foot; jointed pillar to admit of the instrument being used either horizontally, vertically, or at any intermediate angle; rack-work adjustment, with sliding stage; Huyghenian eye-piece; set of three achromatic object-glasses, single lens and adapter, concave mirror, stage and spring forceps, animalculæ box, glass sliders for receiving objects to be examined, and pieces of microscopic glass for covering objects, with objects mounted in Canada balsam on sliders.

DIRECTIONS.—The spot best adapted for microscopic observations is one free from tremor, where a good light, either from a white cloud, or Argand lamp, may be obtained. The mirror, *h*, and optical parts of the microscope are to be rendered free from dust by the careful use of a piece of chamois leather, or camel's-hair pencil; the foot, *g*, is to be screwed to the pillar, *f*, and the eye-piece, *o*, having been slid into one end of the body, *a*, and the object-glass, *i*, or *j*, being screwed into the other end; the body is then to be firmly screwed into the arm *c*, and the compound body inclined so as to bring the eye-piece, *o*, in such a position as to be opposite the eye of the observer when sitting in an easy posture. The adjustment of the light is the next step, and this is easily effected by turning the mirror, *h*, until the light reflected by it passes through the hole in the centre of the stage, *e*, and the maximum amount will be obtained by now looking through the instrument, and slightly varying the position of the mirror, until the whole field of view is clearly illuminated. The object to be examined must be placed on a slip of glass, either with or without water, and a thin slip of microscopic glass be laid on it, and then placed on the stage, *e*, in such a position as to be exactly in the axis of the object-glass or power employed; the object is retained from slipping by the sliding-piece, *p*, which serves readily to retain the object in the centre of the stage; the focus will now be easily obtained by turning the milled head, *d*, so




as to raise or lower the body, *a*, as occasion may require, until on looking through the instrument the object appears clearly defined. If the object is very opaque, the light reflected by the mirror will be of no benefit; it will, therefore, be necessary to throw a powerful light on the object by means of the condenser, *r*, the pin of which fits into the hole at the corner of the stage, and to view it with as low a power as possible. The achromatic lenses, *i*, are used together for the highest power, but must be separated, and only one employed if a lower power is required. The plano-convex lens, *j*, is for use in dissecting small objects, and for mounting specimens. If the body, *a*, is removed, and the adapter, *k*, having the object-glass attached to it, be screwed into the arm, *c*, the instrument will be converted into a simple microscope, well adapted for dissecting and examining objects to prepare them for mounting to be viewed by the compound body and higher powers.

A more powerful instrument is shown by fig. 830. This has a compound body, sliding stage, revolving diaphragm, rack-work coarse adjustment, tangent screw fine adjustment, concave and plane mirrors, double pillar support on mas-

sive tripod base, two sets of achromatic object-glasses $\frac{1}{4}$ and $\frac{3}{4}$ inch focus, shallow and deep eye-pieces, adapter for use in dissections, mounted condenser, spring and stage forceps, box for live insects, selenite stage, micrometer, polarizing apparatus, and objects mounted in balsam.

DESCRIPTION.—The stand, or base, consists of a strong tripod, *a*, supporting two upright pillars, *b*, *b*, between the upper parts of which an axis works. This carries the whole of the optical parts of the instrument which can be adjusted to any inclination, horizontal, vertical, or intermediate. The stage,



830

d, *e*, is firmly attached to the axis, as is also the double mirror, *f*. The triangular bar, *g*, has a rack on its posterior part, which is worked by a pinion, the milled heads of which are seen at *h*, *h*. The body, *i*, screws firmly into the arm, *j*; the object-glasses contained in the boxes, *k*, *l*, screw into the body at *m*; the eye-piece, *n*, or *o*, slides into the other end of the body. The mirror is plane on one side, and concave on the other, and is fitted with a universal

movement, so as to be inclined in any desired position. The milled heads, *h*, *h*, by being revolved, raise or lower the body, *i*, and constitute the coarse adjustment; the fine adjustment is effected by turning the milled head, *p*. The object to be examined is placed on the stage, *d*, and retained in the required position by the sliding piece, *e*. The quantity of light admitted through the instrument may be modified by the diaphragm, *r*, which consists of a plate of brass with four apertures of different diameters, made to revolve on a central pin or axis fixed to the bottom of the stage. In addition to the four holes mentioned as needed to admit the requisite amount of light, the diaphragm is furnished with a fifth hole, into which the mounted Nicol's prism, *s*, screws, forming the polarizer; the analyzer, *t*, is screwed into the upper part of the adapter of either, *k* or *l*, previous to its being attached to the body, *i*. The polarizer, *s*, is mounted in a double tube, so as to be capable of being revolved by turning the large milled head at the bottom. The condensing lens, *u*, is for illuminating opaque objects, and fits by a pin, *c*, into the hole at the corner of the stage; it is so fitted that it can be used in any required position or angle. The forceps, *v*, are for lifting small objects. The stage forceps, *w*, fit the hole in the corner of the stage, *d*, and serve to hold any opaque or transparent object while being examined. The live, or animalculæ box, *x*, is for retaining insects, &c. in the field of view, and for the examination of a drop of water containing animalculæ. The adapter, *y*, may have the object-glass, *k* or *l*, screwed into it, and be itself screwed into the arm, *j*, in the place of the body, *i*, when it is desired to form a simple microscope.

Among the objects furnished with the microscope will be found a plate of selenite, which, if laid under many animal and vegetable structures while being examined by polarized light, will cause them to assume the most gorgeous colours, although without this addition they may be perfectly colourless.

The achromatic object-glasses, *k*, consist of three plano-convex lenses, and the set *l*, of two. All of each set are used together when the highest magnifying power is required, but where a lower magnifying power is needed, one lens only may be used; care must be taken not to mix the lenses of one set with those of the other.

1209. **Micrometer.**—In measuring the size of objects, a *micrometer* (*μικρός*, small, and *μέτρον*, a measure) is employed. The stage micrometer consists of a piece of glass, ruled with fine lines by means of a diamond point, at some known distance apart, such as the $\frac{1}{100}$ or $\frac{1}{1000}$ of an inch. This is laid on the object-glass plate, and the substance to be measured is placed on it. It is necessary that the object and the lines should be seen at the same time, which cannot be always accomplished. The eye-piece micrometer consists of an eye-piece having a glass divided by lines, varying from $\frac{1}{15}$ to $\frac{1}{100}$ of an inch apart, placed either in the focus of the eye-lens, or below the field-lens. Other kinds of micrometers are also employed, such as the cobweb micrometer, where, by the motion of a screw fine wires or cobwebs are made to separate from each other.

1210. **Microscopic Apparatus.**—In delineating minute structures, it is useful to have the image thrown on paper by means of a *camera-lucida*, or small prism, which can be easily attached to the microscope. A *polarizing apparatus* is a useful appendage. In the apparatus sent along with microscopes will be found a *compressorium*, for the purpose of applying pressure to objects whilst they are under examination, troughs for holding such plants as *Chara*, which are to be seen in water, and various instruments for the dissection and examination both of animal and vegetable structures. In testing the power of the instru-

ments, certain minute objects are used, such as the scales of *Podura plumbea* (common Springtail), of *Lepisma saccharina*, of *Hipparchia Janira* (common Meadow Butterfly), the hair of *Dermestes*, Muscular fibrillæ, and the beautiful marking of the Diatomaceæ. Certain markings occur in these *test-objects* which can only be seen by good microscopes.

1211. In viewing objects under the microscope, they must be placed on slips or slides of glass, which should be of a uniform size, not less than three inches by one; and they should be covered with pieces of very thin glass, $\frac{1}{80}$ to $\frac{1}{100}$ of an inch thick, and about $\frac{3}{4}$ of an inch square. The slides ought to be made of thin plate-glass, and the covers of very thin crown or plate-glass. In examining recent vegetable structures, it is best to moisten them with water. When the parts are dry, thin selections may be made either by means of slicing instruments, or by a sharp knife. Many dry objects are well seen when immersed in Canada balsam. To preserve objects in a moistened state, the substances used are alcohol, solution of salt, alum and corrosive sublimate, water containing a small quantity of creasote (5 grains to the ounce), and glycerine. The objects, in such instances, are placed in shallow glass cells, or they are laid on the slides and covered with thin glass, which is cemented by means of japanner's gold size, or black japan varnish. The methods of proceeding are afterwards described.

The following details are chiefly condensed from Schacht's admirable treatise on the microscope, and from the works of Hanover, Quekett, Jabez Hogg, and Beale:—

1212. In proceeding to use the microscope for the first time, with the view to make it a subject of study, it is necessary to have a variety of tools and apparatus to aid in preparing objects for investigation. These may be arranged beside the observer in such a way that they shall be always within his reach. A small tray or box, with divisions, containing a pair of needles in handles (such as are used for crochet needles), a sharp knife or razor, a section-knife (such as that invented by Valentin, and which bears his name), scissors, and a pair of sharp or fine needle-pointed forceps, about three inches long, are among the most essential instruments required. Glass slides may be arranged also upon the same tray for common use, and the thin glasses for covers should be kept in a small box by themselves. For manipulating with the object to be examined, and reacting upon it, another set of materials is required,—namely,

1213. **Re-agents.** The essential re-agents are, 1. Distilled water; 2. Alcohol in the strong state, and also diluted in the proportion of about 1 part to 10 of distilled water. It is the best preserving agent; it removes colour and also air. 3. Ether, which dissolves resins, fats, and oils. 4. A solution of liquor potassæ diluted to about 1 to 20. It *swells* up, and sometimes separates membranes of cells and tubes when they exist in condensed layers. 5. A solution of iodine in iodide of potassium diluted to the following strength,—namely, 1 grain of iodine to 3 grains of iodide of potassium and an ounce of distilled water. 6. Chromic acid diluted in the proportion of about 1 to 30 or 40 of distilled water. These two last re-agents chiefly act by colouring the cell walls or the contents of the cells. 7. Sulphuric acid. 8. Oil, such as the finest of that obtained from coal, and known as mineral oil, is to be recommended for examining and preserving objects in. It does not become rancid, nor has it any affinity for oxygen. For the examination of pollen and spores there is nothing better. 9. One part of dry muriate of lime and 3 of water makes also an excellent solution for preserving objects


which do not contain starch. 10. Glycerine is the best preserving agent for cells containing starch. 11. Canada balsam; and 12. Turpentine, are most useful re-agents and preservative materials for many dry preparations. 13. A strong solution of Carbonate of soda. 14. A solution of Hydrochloric acid may also be found useful in removing deposits of carbonate of lime. 15. A solution of acetic acid. 16. A solution of carbonate of potass. These sixteen substances should be arranged in stoppered glass bottles, fitting into a stand or box, so as to be of easy access, and little brushes, pipettes, and glass rods, should be arranged beside these bottles, in order to apply the fluid to the object. Lastly, the student should provide himself with a small note-book of good drawing-paper, on which he ought constantly to practise the delineation of the forms or outlines of the objects seen, and he should endeavour to colour them also when required.

1214. Numerous other requisites and appliances will suggest themselves during the course of investigations, and especially such as will secure cleanliness of the object and of everything used in the research. 1. The microscope, when not in use, must be kept under cover, generally a glass shade. 2. Its lenses must be cleansed when necessary by a cloth which is used *only* for that purpose, or by dry elder pith. The cloth best adapted for this purpose is old and frequently washed muslin. 3. A separate cloth of a coarser kind is to be used for drying and wiping the slides and covers. 4. Covers of a middle size, from concave disks, such as watch-glasses up to the size of a wine-glass without the stem, or other bell-shaped jars, are also required to protect the objects, if it is necessary to leave them for any length of time.

1215. **General Rules.**—The microscope is used to the best advantage in a room which receives its light from the north, or west, or both. The light which is reflected from a white and motionless cloud opposite to the sun, is the best that can be obtained. If gas-light is to be used, it ought to be corrected or modified by passing it through blue glass shades before reaching the mirror; but for exact observation daylight is always to be preferred. In conducting microscopic observations great steadiness of the instrument is required, which should accordingly be set upon a very firm and sufficiently large table, so that all the apparatus hitherto mentioned shall be within reach of the observer. It is proper also to begin the examination of objects with the lower magnifying powers, and to pass gradually from them to the use of the higher powers. By such means a far larger portion of the object is seen, and a more correct idea is obtained of the relations of the parts when considered as a whole. Object-glasses, varying from 30 to 50 diameters, are the best to begin with. The eye-glass of lowest power, that is, the longest one of the series, is also the one which ought generally to be used in the first instance, and as long as the power can be increased by object-glasses of greater magnifying power, any more powerful eye-piece should not be used, because by the use of such eye-pieces the image is rendered more obscure, while less light is obtained for its display.

1216.—**Necessity of Studying the Sources of Errors of Observation.**
—**FIRST**—Extraneous or accidental objects may be present, and may be derived from various sources; such as water too long used may bring before the eye both plants and animals of the lowest forms, which otherwise would not have been present. Fresh water is absolutely necessary, and it may be even recently distilled. Particles of dust, or fibres from the cloths used in cleaning the glasses, may also add to the confusion. These consist generally of

fibres of paper, linen, woollen, cotton, or silk fabrics, or minute hairs from the brushes used in manipulation. **SECONDLY**—Air-bubbles are almost invariably a source of wonder, admiration, and annoyance, to the microscopic observer in his first attempts; but once seen and studied, they begin no longer to distract the attention, and the microscopist soon gets into the habit of disregarding their presence. They generally appear in the form of circles of larger or smaller diameter, with a dark black-looking rim, seen by transmitted light; while with incident or reflected light, their rim appears of a white colour. Pressure under a glass cover causes them sometimes to assume very irregular shapes, but possessing the same properties in their margin or outline in their behaviour with the light. It is also necessary to become familiar with the appearances of the lowest forms of animal and vegetable life, such, for instance, as the common forms of infusoria, the yeast, and such like plants; and the different forms of mould. It must not be forgotten also, that the various menstrua used in manipulation also contain elements which properly belong to them, and which must therefore be distinguished. For example, the epithelium of the saliva, the blood corpuscles which may be in serum, and crystals which may be deposited in various fluids which are used as re-agents. A peculiar motion, known as “molecular motion,” is also a phenomenon which must be recognized. It is peculiar to all very small particles or molecules, when they float in a very thin fluid medium. It is best seen in the fine granules of milk when mixed with water, in the milky juices of the plants, and may be observed very distinctly in solutions of India-rubber, or caoutchouc, when dissolved in ammonia. A magnifying power of 400 or 500 diameters is the best for this observation. Another class of deceptions originates in the eye itself. These are the “*muscæ volitantes*,” the nature of which is described as follows by Dr. W. Mackenzie:—



831

“The vision of objects on the surface or in the interior of the eye has attracted attention, chiefly in relation to a symptom, to which the name of *muscæ volitantes* has been given. Any spectrum, or visual appearance, which is apt to impose on the mind, and lead one to think that flies are moving before the eye, is called a *musca volitans* (fig. 831).”

“The condition comprehends those sensations which arise from, 1. The layer of mucus and tears on the surface of the cornea; 2. Corpuscles between the external surface of the cornea and the focal centre of the eye; 3. Corpuscles between the focal centre of the eye and the sensitive layer of the retina.”

“In hanging the head over the microscope, especially if one is affected with catarrh at the time, the globules, by gravitating to the centre of the cornea, not unfrequently appear to the observer so as to impede his view of the object, till

by the act of nictitation he clears them away. In telescopic observations, also, the muco-lacrymal spectrum is apt to prove a source of annoyance. Thus, in looking at the sun through a tinted glass, the observer may be unable to distinguish the spots on that body, being perplexed by what seems the reflection of some part of his own eye interposed between it and the sun. This is caused by the layer of mucus and tears on the surface of the cornea.

"If one looks at the flame of a candle two or three feet distant, or at the sky, through a hole made in a blackened card with the point of a fine needle, or through a convergent lens of short focus, such as the eye-glass of a compound microscope, on steadily regarding the luminous field presented to view, four sets of spectra will be seen (fig. 831), independent of the muco-lacrymal spectrum. The most remarkable appears nearest to the eye, and consists of twisted strings of minute pearly globules, hung across the field of view (fig. 831 *a*). The second in point of remarkableness, and the farthest of the four from the eye, consists of watery-like threads, destitute of any globular appearance, and depending chiefly from the upper part of the field (fig. 831 *b*). I call the former the *pearly spectrum*, and the latter the *watery spectrum*. In two distinct planes, between those occupied by these two spectra, float two sets of globules, not aggregated into threads, but insulated. These constitute what I call the *insulo-globular spectra*. The individual globules of the set farther from the eye, being hazy and ill-defined, may be compared in appearance to small grains of sago (fig. 831 *c*). The globules of the set nearer to the eye are clear in their centre, exteriorly to which they present a sharp black ring, and still more exteriorly a lucid circumference (fig. 831 *d*). These four sets of spectra never mingle with one another, so as to change the order in which they stand before the eye; but the pearly spectrum always appears the nearest; then the sharply defined insulo-globular; then the obscurely defined globules; and farthest away the watery threads.

"Almost every eye, even the most healthy, and which has never attracted the possessor's attention by *muscæ volitantes*, exhibits the pearly spectrum, on being directed towards a luminous field, through a fine pinhole, the eye-glass of a compound microscope, or a convex or concave lens of short focus. I have given it the same name of the *pearly spectrum*, from its resemblance to a string of pearls. Prevost had already called it *appareance perlée*, or simply *perles*.

"The lines of the pearly spectrum are hung across the field of vision as often transversely as vertically. On first directing the eye towards the luminous field, in one or other of the methods just mentioned, perhaps only a very few small pearly globules are perceived; but after steadily regarding it for a short time, numerous strings of them are discovered, generally twisted in different forms, and presenting a variety of knots, loops, and agglomerations. Sometimes they are so numerous as to form an extensive shower or cloud. The pearly threads are of different lengths; some of them very short, others stretching across the whole field. Not unfrequently some of them end abruptly in a sort of bulb. The globules or pearls forming the threads or rosaries, seem joined together merely by apposition, without being contained in any tube. Sometimes, however, the globules are rather indistinct, and then the threads approach to a tubular appearance. The globules are always in single rows. They appear destitute of any nucleus. They are not all of one diameter, but are all smaller than the globules of the insulo-globular spectra. I have not satisfied myself that all the pearly threads occupy the same plane, although it is very evident that they are behind the insulo-globular spectra.

"That portion of the pearly spectrum which appears in the centre of the field of view has but little real motion, less perhaps than the watery spectrum which is seen beyond it. Both partake, of course, in the motion of the eyeball; and this gives to both a wide apparent motion. But if the field be examined towards its circumference, or if the eye be suddenly rotated upwards, other pearly spectra appear, which it is difficult or impossible for the observer to bring directly before him; and which, when he succeeds in some measure in doing so, quickly subside again out of view, partly by a real motion of their own, partly by a wide apparent motion, owing to their obliquity in respect to the axis of vision. It is these last spectra, chiefly, which produce the pearly *muscæ volitantes*."

There are also various optical phenomena caused by refraction, and which are necessary to be attended to. They depend, for the most part, upon a bad adjustment of the focus, or illumination of the object. The appearances are also most frequently associated with an increase of the magnifying power, and especially with the use of powerful eye-glasses. Large grains of potato-starch, pollen-grains, the thickened substance of wood-cells, and the cells of cartilage, are among the most common objects which exhibit such optical phenomena, which consist in a feeble and generally yellowish colouring of the edges of the objects when seen with particular foci.

1217. Focal Adjustment of the Microscope.—The regulation of this adjustment is based on the fact, that the microscope can only afford a view of one surface of an object at any given time, so that nothing is distinctly seen which lies above or below such a focal plane at that time; and the more plane the field of vision, the clearer and better will be the view of objects in that plane if the adjustment is correct. The more perfect the object-glass, and the greater the angle of aperture, the more exact is this focal plane, and the more sensitive is the instrument to any small alteration of focus. The focal adjustment is made and varied by what is called a fine adjustment screw, and which is sometimes graduated; and the accurate adjustment of the object is judged of by the sharpness of the delineation of the image, as well as by the fineness and clearness of the outline. An experienced microscopic observer always uses the instrument with his finger and thumb grasping the fine adjustment screw, and would not be content with his observation, although it was limited to a mere peep of the object, unless he had made the fine focal adjustment for himself.

1218. How and what to Observe.—Rigid and accurate observation must be methodically cultivated from the first, and logical reasoning, carefully guarding against all sources of fallacy, must accompany every step taken, judgment given, or conclusion arrived at by the investigator. The physical characters which distinguish microscopic objects are, 1. Shape; 2. Colour; 3. The edge or border; 4. Size; 5. Transparency or opacity; 6. Surface; 7. Contents; 8. Effects of re-agents. These physical properties all in their turn claim the attention, so that the mind of the observer becomes impressed with positive or negative results. The mind must be constantly awake, and as long as the objects are within the sphere of our vision, they must make a due impression on our mental faculties. By a careful examination of the circumstances to be distinguished, so as to understand them thoroughly, and be familiar with them in all their bearings, we shall seldom require to say of an object, "I must have seen—the object, but did not observe—its properties." Of these properties we shall treat in their order, and

1. As regards shape, it is obviously a property which immediately strikes the

eye, especially if it is a peculiar one, and not well known as a shape familiar to the naked eye. In describing shape, it is to be done by reference to well-known objects, such as mathematical figures, and to the shapes of natural and artificial familiar things, such as by the words "pear," "heart," "kidney," or "bladder-like;" "dome," "lattice," "ladder," "spiral like," and so on.

2. Colour. The observation of this physical property is one of great interest, considering that the cause or source of colour is so variously modified. The quantity of the object, for example, modifies the colour. Blood, which is ordinarily described as red, is seen to be yellow when individual corpuscles are observed. Different colours are also produced by different modes of illumination, and the linear arrangement of markings is sometimes associated with the most beautiful iridescence. Re-agents also modify colour, as when a solution of iodine reacts upon starch, or inuline, or on the albumen of the seeds of many plants; or when alcohol reacts upon chlorophyll, and so on.

3. As regards the edge of the object, its darkness, abruptness, fineness, softness, delicacy, smoothness, or irregularity, must be specially noted.

4. The size of the object must be determined with exactness by actual measurement. For this purpose a micrometer is necessary. There are various forms of this instrument, which, of course, are variously used (see par. 1209). A mode of ascertaining the magnifying power of the compound microscope is founded on the assumption, that the naked eye sees most clearly and distinctly at the distance of ten inches. If a divided scale be placed on the stage, and distinctly seen magnified through the instrument, let a rule be held at ten inches distance from the right eye, while the observer uses, at the same time, his left eye in looking at the other scale through the microscope, and let the rule be gently moved so that it is seen to overlap or lie by the side of the magnified picture of the other scale,—a comparison as to how many of its known divisions correspond with a number of those on the magnified scale, will indicate the magnifying power. Upon a similar principle a pair of compasses may be substituted, whose points being placed on the stage are separated till they cover or mark off so many spaces as magnified by the instrument. If they cover 1 magnified space, and correspond to 2, 3, or more known spaces on the rule, then the instrument is said to magnify 2, 3, or 400 times linear that known space. If $\frac{1}{1000}$ of an inch is found to cover 3 inches on the rule, the instrument magnifies 300 times; if 4 inches, 400 times, and so on; if 2 inches, 200 times, and so on. In this way is determined the magnifying power of any combination of lenses, and the scale which is magnified is called the glass or object-glass micrometer. The size of objects may be measured by placing them directly on this micrometer; but it is obvious that they cannot under high powers be brought into focus at the same time as the lines of the micrometer. An instrument called the eye-piece micrometer, is therefore generally used. It consists of a known scale ruled on glass, and placed in the focus of the upper glass of the eye-piece. It is observed with it how many of its divisions correspond with one division magnified by the microscope. Thus, with an instrument which is known to magnify 250 diameter linear, it is found that 1 space of the magnified micrometer corresponds with 5 of the eye-piece micrometer, the size of the object will be $\frac{1}{5}$ of the known scale. If the scale be $\frac{1}{1000}$ th part of an inch, and the object is seen to correspond to 1, or to 4, or to 5 of those spaces, then its size is $\frac{1}{1000}$ th, or $\frac{4}{1000}$ th, or $\frac{5}{1000}$ th part of an inch.

The exact size of objects can only be determined by actual measurement; but at the same time the size may sometimes be conveniently compared with that of any other familiar microscopic object, such as a human blood disc, or the like.

5. Transparency is a physical property, which varies much from a condition perfectly diaphanous to one of more or less opacity. The rays of light may be completely prevented from passing through the object, it then appears black by transmitted light, when really it is white by reflected light. A luminous appearance is produced by objects such as fat and oil globules, which refract the rays of light.

6. The *surface* of an object ought to be examined in the first place, and by gradually adjusting the focus to different planes below this transparent surface, the structure throughout may be observed.

7. The contents of enclosing structures must be observed. We notice thus, 1st, Whether a cell has the appearance of being empty, as when it contains air, as is the case with perfect vessels and wood-cells. 2d, Whether its contents are fluid, with solid particles floating or contained in the fluid. 3d, The nature of the contents, as ascertained by chemical re-agents, such as of substances held in solution. For example, sugar, gum, dextrine, oil, resin, or saline matters. The solid contents of enclosing structures, such as cells or tubes, consist for the most part of starch, inuline, and chlorophyll, and sometimes crystallized products. In the case of crystals, their forms are sometimes alone sufficient to determine the nature of the substance crystallized. Oxalate of lime, for instance, exists as octahedral crystals, and as stellate-shaped crystals—of a four-sided prismatic form—to which the name of raphides has been given. If form alone is not sufficient to indicate the nature of the crystal, then the use of chemical re-agents must be resorted to. Carbonate of lime, for instance, is known by its effervescence and solution, under the influence of sulphuric or hydrochloric acid. The test must be carefully applied to the edge of the glass cover by means of a fine rod of glass.

Crystals may be always found in the outer coat of the onion, in the garden rhubarb, and in the squill. The acicular crystals are generally phosphate of lime. Others having lime as a basis, are combined with tartaric, malic, or citric acids. Such are easily destroyed by acetic acid. They vary in size from $\frac{1}{100}$ to $\frac{1}{1000}$ of an inch, and are to be found in all parts of the plant in the interior of the cells. The cactus tribe of plants contains large quantities of raphides. The brittleness of many plants is due to them. In the bark of the hickory (*Carya alba*), masses of the flattened crystalline prisms are to be seen.

Silica constitutes another crystalline formation, to be seen in plants such as the Gramineæ. Chlorophyll is a substance of a resinous or waxy nature, and is the source of the greenness of plants, being especially abundant in the cells of the leaves. Alcohol removes the green colour. Inuline (a name derived from the plant *Inula*) is a substance resembling starch, prevalent in the Compositæ. Its difference from starch consists in its not containing so large a proportion of the elements of water, and in being coloured pale yellow or brown with iodine. Starch is characterized by the blue colour given to its granules on the application of free or uncombined iodine. An example is to be found in the potato.

To detect sugar, there are no certain chemical re-agents known applicable to microscopic observation, although it has been presumed that the red colouring of the contents of cells, consequent on the use of concentrated sulphuric acid, may be a reaction on sugar.

Gum and dextrine are coagulated by alcohol. Resins, and oils or fats, are removed by ether.

1219. Elements of the Tissues of Plants.—In order to examine these elementary structures, as well for the purpose of acquiring proficiency in the use of the microscope, as to obtain a scientific knowledge of structural botany or vegetable histology, the student ought to examine systematically; 1st—Cells, as they are the foundation of all the organs of plants. Cells are found to exist in a great variety of forms, and therefore it is better to begin with the most simple as seen in (1.) *Cellular plants*, such as *Conferva*, *Diatomace*, *Closterium*, and the like; (2.) *Cellular tissue* of more complex plants, such as may be seen in sections from the pith of the Elder, the Rice Paper plant, the large nucleated cells of the Onion bulb, the stellate cells of the *Juncus*, and the twisted cells in the hairs of Cotton; (3.) Spiral cells, in which a spiral filament seems as if coiled up in the interior of the cell, or from a thickened portion of the cell-wall. The *Oncidium*, *Jungermannia*, *Collomia*, *Acanthodium*, *Lophospermum*, &c. furnish beautiful illustrations; (4.) the Pollen cells, as in the *Acacia*, the *Lily*, the *Passion flower*; (5.) the spores of *Equisetum*, *Fern*, *Moss*, *Jungermannia*, *Agaricus*, *Zygnema* in conjugation, spermatozoa cells of *Chara*; (6.) the cells of the cuticular apparatus, the glands, scales, and hairs of plants. The *Pelargonium*, *Oncidium*, the *Wheat Straw*, and the *Equisetum*, furnish beautiful material for observing the structure of the cuticle. The leaf of the *Mentha*, and the *Sweet Briar*, will supply examples of the glands and scales, and the cells of the hairs are best studied in *Cynoglossum*, *Sterculia*, *Trichinium*, *Tradescantia*, *Antirrhinum*, *Nettle*, &c.

In the simple forms of the cell it will always be found that, when isolated, each consists of a little sac, closed on all sides, and containing fluid in their interior. Each cell also generally contains an evident nucleus or cytoblast, consisting of a granular round or oval mass, either free or attached to the wall of the cell. In the interior of this nucleus may sometimes be seen a nucleolus, or a variety of nucleoli. Iodine sometimes serves to render the internal contents of cells more obvious, as in the case of a very transparent nucleus which is not seen unless coloured with the solution of iodine. The nucleus is never wanting in young cells, although it may be often concealed by the granular contents of the cells. By means of diluted sulphuric or nitric acid, upon the fresh cell, its inner mucilaginous coating, which is generally granular, is made to coagulate and usually contracts, so as to enclose the solid contents of the cell. This is the primordial utricle of Mohl, which is found in all young cells, in the cells of all succulent tissue, such as fruits, the leaves of the *Aloe* and *Agave*, and in the fresh leaf of the *Liver Wort*. In very thick cells it can scarcely be discovered, nor in those altered by deposits of solid substances in the interior of the original cellulose of the cell-wall, and its presence cannot be determined after fossilization.

1220. Growth of the Cell—Modification of Form and Arrangement with respect to one another.—The elementary form of the cell is a closed sac, of a more or less rounded shape, which increases in size in various ways. Its form is modified by growth, and by pressure of adjoining cells. Cells which increase equally on all sides and at all points of their periphery are seen in the reproductive cells, in spores, and in pollen grains, and in the pulp of ripe succulent fruits. Pressure more frequently changes the form of cells into polygonal figures. The pressure of adjoining cells being mutual, the number of the angles and sides of the polygon depends on the number and arrangement of the contiguous cells. Such cells constitute the parenchyma of vegetable structures, and, for the most part, exhibit the forms of pentagons and hexagons, or more numerous-sided figures.

Sometimes the cell increases in growth to a greater extent in one direction than in another. Wood cells are examples of cells extending lengthways; they are elongated in the direction of the axis of the stem. The cells of the medullary rays extend breadthways. The wood cells are generally much extended, thickened and pointed at both ends. Maceration in water is the best method of obtaining these elongated cells, as from the root of the elder, flax, and hemp. Such cells constitute what is called the "woody fibre" of plants. Cells of a stellate form arise when the growth does not increase equally at every point of the periphery. The pith of the *Juncus conglomeratus*, and leaf-stalk of the *Musa*, and many other spongy tissues, furnish examples. Air passages exist in such tissue, in consequence of the cells only touching each other at certain spots.

Cells increase in growth by deposits of new matter in their interior. This new matter may simply thicken the cell-wall by its deposition upon the original cellulose. Sometimes the deposit is laid on so as to assume the form of a spiral. In young growing cells this is frequently observed, but not in old full-grown cells. Fresh twigs of the *Pinus abies*, in spring and in summer, show this spiral band, and, generally, it may be stated that the deposits are laid down within or upon the walls of cells in the form of a spiral. Pits, dots, or discs are modifications of this deposit. They are to be studied in the *Coniferæ*, *Cycas revoluta*, and in the vessels of tropical twining plants. The construction of the pits and discs must be observed from three sides. In *Pinus sylvestris*, for example, when a transverse section is made, or when a section is made, at right angles to the medullary rays (tangential), the pits are seen sideways; the canaliculus of each wood cell is seen, and the lenticular cavities on the sides and between the cells. If a longitudinal section is made in the direction of the medullary rays the pits are seen looking on their faces. They show the form of two circles, one within the other; the larger circle bounds the lenticular cavity, the smaller circle points out the situation of the canaliculus, and when it is very conical or prolonged a third circle may be seen with some focal adjustments, and in some the innermost circle is in the shape of a slit, as in the *Cycas*.

1221. **Vessels of Plants** are distinguished from the other structures by being placed in rows, one above the other, and are rendered continuous by the obliteration, in whole or in part, of their transverse partition walls. The vessels of plants consist of spiral vessels, porous vessels, annular vessels, scalariform vessels, and laticiferous vessels. The spiral is the true type of the vascular tissue, and may be studied best in the *Canna bicolor*, *Nepenthes*, *Banana*, *Cactus*, *Hyacinth*, *Asparagus*, *Mistletoe*, and *Long leek*; in the two last examples the spiral vessels are branched. A dissection of a recent part must be made, and by macerating the specimen in boiling water its tissues will be more easily separable. The wood of the *Yew* tree also furnishes beautiful illustrations of dotted spiral vessels. In the annular vessels the coils of the spiral are separated from each other by the elongation of the cell, and they thus assume the form of rings. In the stems of the *Balsams* the most beautiful transition from the spiral to the annular vessel is to be seen. The root of the *Opuntia* will also furnish examples, while scalariform vessels may be seen in *Osmunda*, *Asplenium*, *Cheilanthes*. The porous or pitted vessels are best studied in the wood of the *Laurus sasafra*s, the *Sugar Cane*, and the *Nepenthes*, while the pits and the spiral band are to be seen together in the *Filia Europea*, and the *Taxus baccata* (*Yew* tree). The vascular tissue must be examined, after being treated with alcohol, in order to expel the air from the interior of the tubes.

Besides the tissues described in the preceding paragraphs, there are some

combinations of structure so commonly spoken of that they require notice. These are "vascular bundles," "tissue of the liber," "intercellular passages," "and laticiferous vessels." In a "vascular bundle" the cambium cells constitute the most essential part. They are the elongated thin-walled cells between the wood and the bark. Besides these cells, "vascular bundles" contain vessels and wood cells. These bundles in dicotyledonous plants can only be studied at the apex of a young shoot, and may be seen in *Viscum*, *Lilia*, and *Pinus*. "The tissue of the liber" never originates in the proper wood. It consists of elongated flexible cells, generally much thickened, and neither possessed of cambium vessels nor wood cells. The liber cells are to be found both isolated and in masses, in the leaves and in the pith of the stem. In the *Apocynæ* and *Asclepiadæ* liber cells are to be found, which contain a milky juice. "Intercellular passages" are to be found where many cells meet. They are chasms filled with air, and sometimes, though less frequently, with fluid. They are beautifully seen in a transverse section of the leaf-stalk of the *Cycus revoluta*. "The laticiferous vessels" form a continuous, net-like tissue, spread round the cells, and consisting of ramifying canals, in which a milk-like and frequently coloured juice is contained. It is not yet settled whether they are formed by the cells or by the thickening of the walls of the intercellular passages. The leaves of the *Chelidonium* and *Euphorbia* will furnish illustrations.

1222. Methods of Examination.—A few practical lessons, from one skilled in the use of the instrument, form the best introduction to private study and personal application of the microscope. Rules at the best can only be regarded as hints, and, as such, they are here laid down. Three things deserve special attention, 1st, Preservation and arrangement of the microscope; 2d, Illumination and choice of magnifying powers; 3d, Selection, preparation, observation and preservation of objects, together with their description, measurements, and delineation.

1223. Preservation of the Instrument.—One who has any regard for his instrument will never suffer it or its lenses to be handled by those unaccustomed to their use. A microscope in constant use is not to be taken to pieces, but preserved under cover of easy access, so as to prevent the injurious effects of the direct rays of the sun. Dust is to be removed from the lenses by a camel-hair pencil, and by methods described at par. 1214. The situation of dirt may be discovered by turning the eye-piece or object-glass. A microscope should never be exposed in a chemical laboratory, unless, perhaps, in the balance room, and under similar restrictions as to its use.

1224. Choice of Magnifying Power.—Weak eye-pieces and strong object-glasses are to be recommended. The eye-piece does not add either clearness or distinctness to the object, and when it is very powerful the field of view becomes too small to take in the whole image formed by the object-glass; for the magnitude of the field of view and the strength of the illumination diminishes according to the magnifying power of the eye-piece employed. The lower powers are of use in searching for the object to be examined, which may thus be more easily found by a higher power. The magnifying power should also be varied as little as possible, in order that the comparative magnitude of different objects may be more quickly determined. For the lower power a linear amplification of from 20 to 50 diameters, and for a higher power a linear amplification of from 300 to 500 diameters at most, will give a sufficiently wide range of powers. The powers are increased by a more powerful eye-piece or object-glass, or by both, or by lengthening the tube of the microscope.

1225. Preparation and Selection of Objects for Examination.—It must be determined whether the object is fit for microscopic investigation. It is not uncommon to hear persons, ignorant of what a microscope can do, request objects to be examined which, from their size and nature, are unfit for such methods of observation. It is also to be laid down as a rule that bodies which are not perfectly fresh are not suited to be observed microscopically. Consistence, dryness or wetness, determine in general the chief modes by which the object is to be examined; a sufficient degree of thinness being necessary to view objects by transmitted light. Opaque objects require merely to be made smooth or level on one side, and to be fixed on the other. If the object is to be viewed by transmitted light a section or slice sufficiently thin must be procured; a common sharp knife, a razor, or Valentin's section knife are the instruments to use. They must be moistened with water, and sometimes it is advisable to make the section under water. If the object is very delicate it may be laid between two pieces of cork, and the whole cut through. If the object is hard, such as a fossil plant, it must be ground down with a file, a grindstone or a lappe, and then polished on one surface. This surface is to be fixed to a glass slide by Canada balsam, and the opposite surface rubbed down and polished in a similar manner, till it is sufficiently thin to be viewed by transmitted light, and it must be of uniform thinness. Sections should be made in all the opposite directions, so that a correct knowledge may be obtained of the relation of the component parts. Maceration in water, and tearing the parts asunder with fine needles, are the best methods for obtaining the ultimate tissues of plants. Attenuating or diluting media are of importance in regard to their influence on the objects examined. Distilled water, solutions of common salt or sugar, the white of egg, the serum of the blood, the fluids of the eye, acetic and chromic acids, dilute alcohol, creasote, oils, turpentine, are amongst the most useful.

Thin glass plates to cover the object under the microscope must be invariably used. They prevent evaporation, and preserve the moisture about the object, they prevent the object-glass being covered with vapour, and so rendered obscure, and, lastly, they produce a slight pressure, by which the elementary parts of the substance may become separated from each other, so as to lie on one plane. The thin covers are not so absolutely necessary where very low powers are used.

In placing the object on the stage care must be taken not to bring it in contact with the object-glass of the instrument. It is also to be remembered that, in a compound microscope, the image is inverted, and that, consequently, the object is moved in a direction contrary to that of the image.

1226. Preservation of Microscopic Objects.—The following apparatus is required, viz., glass slides ground at the edges, and of the requisite standard size, with circular glass covers. Preserving agents, cement, and a circular table for mounting and making cells. The best preserving media, for vegetable substances, are—

A solution of chloride of calcium,
Glycerine,
Copal varnish,

Mineral oil,
Canada balsam.

The best cements are—

Brunswick black,
Japauner's gold size,
Black japan,
Sealing-wax varnish,

Robinson's liquid glue,
Gum mastic and caoutchouc dissolved in chloroform.

Objects are put up (*i. e.* preserved) either as dry or as wet objects. For dry objects, the oils and the Canada balsam are the preservative materials, and they are not suited for wet objects. Before mounting objects in Canada balsam they should be perfectly clean and free from moisture. They may also be soaked in turpentine, especially opaque objects, as it renders them more transparent. In mounting, the balsam should be heated to expel the air, till a fine delicate film is apparent on its surface. By this process the air is expelled.

The solution of chloride of calcium is adapted for the preservation of wood and leaves, and for most species of isolated tissue, even the youngest. The colouring matter in the cells, however, is always more or less altered by it, while grains of starch, if present, swell up and can scarcely be recognized. The strength of the solution is one part of lime to three of water. Glycerine is used in equal parts mixed with camphor water, which prevents the tendency to mildew. The chlorophyll and the grains of starch remain unchanged, and the laminæ of the starch appear more beautiful after a few hours' immersion in the glycerine solution.

Canada balsam and copal varnish are used for the preservation of dry and fossil woods. These sections are cut thin, and the hard woods which contain gum, resin, and such like matters, should be soaked in essential oil, alcohol, or ether, before mounting. If the entire structure of any exogenous wood is required to be examined, the sections must be made in at least three different directions, which may be termed the transverse, the longitudinal, and the oblique, or as they are sometimes called, the horizontal, the vertical, and the tangential, each of which exhibits different appearances. The vertical section should show the number and size of the medullary rays, that is, the small portions of pith which proceed horizontally from the centre, enclosed in a sheath of woody fibres, and which are known to the cabinet-maker as the silver grain of the wood. In coniferous trees, as the pine, this section shows also the beautiful discs on the walls of the fibres. The tangential-vertical section is a slice across the medullary rays, and exhibits the form and arrangement of the cellular tissue within them. These vertical sections show the form, size, and connections of the woody fibres, the spiral, reticulated, and dotted vessels.


In endogenous trees horizontal and vertical sections only are required.

Peat wood requires to be digested in a strong solution of carbonate of soda, and fossil woods which have been converted into carbonate of lime should be digested in hydrochloric acid.

1227. To make Cells and to fix the thin Glass Cover.—The cells are made either round or square by thin layers of cement, according to the depth required. Perhaps the round ones are neater, but they require circular pieces of glass for covers, and by the aid of the circular table the roundness of the mounting can be made with perfect accuracy. The cover is laid gently down, so as to float on the solution in which the object lies, and by pressing gently down on the cover, the superabundant fluid is made to pass out by the edges, and may be taken up by a sponge or blotting paper. A thin layer of Brunswick black, or liquid glue, or gold size, may be placed round the edge, which will gradually harden and completely seal up the preparation.

II.—ON COLLECTING AND EXAMINING PLANTS, AND ON THE FORMATION OF AN HERBARIUM.

1228. **Instruments and Apparatus.**—In examining the characters of plants with a view to classification, the chief instruments required are a lancet-pointed knife, a small pair of forceps, and a lens. In more minute examinations, the simple or compound microscope must be called into requisition. In selecting specimens, care should be taken to have the plants in a perfect state, or with all the characteristic parts present. The entire plant should be taken when practicable; when that is not the case, then those parts should be taken on which the generic and specific characters are founded. The roots should always be carefully washed at the time the plants are gathered. In most cases, particularly in specimens of Umbelliferae, Leguminosae, Compositae, Rosae, &c., it is of importance that both flowers and fruit should be preserved. In the case of Willows, the young shoot, with its fully developed leaves, as well as the male and female flowers, are requisite. In Rubi, specimens of the young shoots must be taken. When bulbs or tubers exist, they should be preserved, either in an entire or split condition; and when there is much mucilaginous matter in them, they may be enveloped in small pieces of paper, so as to prevent them from adhering to the drying paper. In the case of Ferns, two fronds are necessary to make a perfect specimen, showing both surfaces, along with a portion of the rhizome. Entire specimens of Gramineae and Cyperaceae should be collected; these, when long, may be bent into one or more folds, corresponding to the size of the paper on which they are to be fastened, the folds being temporarily retained by small slips of paper having slits in the centre. No bad specimens ought to be preserved. The size of the paper recommended for forming an herbarium is about 17 by 10½ inches. It ought to be stiff paper fit for writing upon. The size of the paper will determine that of the specimens to be dried. Two small tin flagons with wide openings are useful, the one for containing aquatic plants, which it may be desirable to place in water, such



as Algæ, Chara, Lemna, &c., the other, filled with alcohol, will be found convenient for preserving objects which are not wished to be dried, such as isolated flowers, pulpy fruits, &c. &c.

1229. In taking up the roots of plants, a small *Digger* or trowel is used, 7 or 8 inches long (fig. 832); the spnd 2½ inches long, 2½ inches wide at the top, narrowing gradually to 2 inches at the bottom, the lower angles slightly

rounded. This can be put into a leather sheath, and fastened by a strap round the waist, the spade itself being attached to the strap by a long string. A japanned tin box or *Vasculum* is required for the reception of specimens. This should be of sufficient length to receive a plant of the full size of the herbarium paper; it ought to be convex on both sides (fig. 833); and its capacity may vary according to the wish of the collector. In long excursions, where productive localities are visited, it will be found that a vasculum 20 inches long, by 8 or 9 inches wide, and 5 deep, is not too large; and when it is made of thin tin, it is by no means heavy. At one end a good sized thickish handle should be placed, and it is necessary to have wires fixed at each end so as to receive a strap for fastening the vasculum on the shoulders. The lid of the vasculum should be large, and is best secured by a wire which slips into a tin sheath, and so constructed as not to be liable to slip out when the box is held by the handle. For mosses and some Alpine species of plants, a small box may also be carried in the pocket. Many plants will not bear transport; their flowers fall off easily, and they are so delicate that their foliage becomes shrivelled. In such instances it is best to put them at once into paper. This is managed by having a small *Field-book* (fig. 834), which may be put into the pocket or suspended round the neck, secured by straps so as to give pressure, and with an oil-cloth covering which may be used in wet weather. This field-book may be made with two thin mahogany boards on the outside.

1230. Herbarium, and Process of Drying.—Of all botanical works the most instructive and at the same time most agreeable to consult, is a well-arranged herbarium. A specimen gathered with intelligence and necessary care is preferable to the most faithful engraving or drawing; and a glance at such a specimen conveys to the mind an idea of the general form of the plant, infinitely more clear than that which can result from a lecture of the best description. A description, besides, is not intended to be read and studied in the absence of the plant itself, but more with a view to facilitate its study when immediately under observation. If dried specimens are less easy to study than fresh plants, they have the advantage over the latter of being brought into requisition at all seasons of the year; their possession is ever so much the more useful and agreeable, since it is, in general, during winter, when all trace of vegetation seems to have disappeared from the soil, that there is more time to devote one's self to the study of species in the silence of the botanical cabinet or sanctum. Many years after having collected the specimens contained in an herbarium, they vividly recall the beauty or remarkable features of the places of which they were originally the denizens, and the happiness of so many botanical excursions, which must be experienced to be properly appreciated.

1231. The *Paper for drying* should be moderately absorbent, 18 inches long by 11 broad, and arranged in parcels containing not less than four sheets. The paper which is used extensively in Scotland is made by Mr. Weir, Queen-Street, Glasgow. In many respects the Edinburgh botanists prefer it to Bental's. It is of considerable thickness, absorbs moisture rapidly, but does not become too moist, and dries easily. A very thin kind of paper, called crown-tea-paper, is used for holding very delicate plants, which cannot be easily transferred from one paper to another during drying. After being carefully laid out in the folds of this paper, they are placed between the sheets of drying paper, and when the paper is changed they are transferred at once in their thin cover without being disturbed.

1232. In order that pressure may be given, *Boards* are requisite. These should be exactly the size of the drying paper. Some of them are used for outside boards, and these ought to be thicker and stronger than the rest; others are inside boards, about $\frac{3}{8}$ of an inch thick. For every two reams of drying paper, not less than ten boards should be procured; two of which are for the outside, and eight for the inside. Sheets of stout pasteboard are also useful for packing up the plants as they become dry. The pressure is best applied on a botanical excursion by means of a rope put twice round the boards and paper, and tightened by a rack-pin. This is much better than straps, which are apt to give way, and are with difficulty replaced during an excursion. In other circumstances, pressure is best applied by means of heavy weights. The pressure ought not to be less than 100 lbs. This is preferable to a screw-press, in which the pressure is not kept up while the plants are losing their moisture. Boards may be replaced by frames of hard wood, furnished with cross-bars. They are lighter, and have the advantage of leaving a portion of the surface of the packet in contact with the air, and consequently of hastening dessication, at the same time they are less solid than the boards. Sheets of tin may be usefully employed to separate the different layers of plants in process of drying, as they hinder the humidity of one from reaching the other, or the inequalities of the larger injuring the smaller and more delicate. Such sheets of tin will be found very advantageous during an excursion, in multiplying the number of the presses, the more especially when a great quantity of plants have to be dried; in the latter case it will be requisite to have the sheets of tin pierced with circular holes, with a view of furnishing an outlet to the humidity. In the case of plants with strong stems, they must either be split, or a sand-bag, of the same size as the boards, used, so as to equalize the pressure.

1233. **Process of Drying.**—The plants when collected are to be placed on the drying paper. In doing this, a parcel of not less than four sheets is put on one of the outside boards; then the specimens are laid out carefully, preserving as far as possible their natural habits, and laying out the leaves and other parts. Another parcel of drying paper is then placed above these, and the same process is repeated with other specimens until twelve such parcels have been placed together. Then one of the inner boards is laid down, and other layers of paper and specimens are applied, until the whole parcel is of sufficient size to be subjected to pressure. After twelve hours' pressure, in most instances, the paper is changed, the moist paper being hung up to dry; and in transferring the specimens from the wet to the dry paper, a large pair of surgeon's forceps is used. The interval elapsing between the changing of the paper may be increased or diminished according to the nature of the plants, and the state of the weather. In the course of eight or ten days, ordinary specimens will be so dry as to require only very slight pressure, with a moderate circulation of air. Some very dry plants, as grasses, may require only one changing. Succulent plants, such as *Sedum*, *Sempervivum*, &c., continue to grow, however much submitted to pressure and the ordinary methods of dessication already indicated. In order to dry these plants completely and rapidly, it is necessary to disorganize them, and for this purpose boiling water must be had recourse to. Each succulent plant must be immersed for fifteen or twenty seconds, taking care, however, only to plunge the stem up to the point where the flowers commence. The plants thus dealt with are then placed upon a cloth and left to drain for a few instants, after which they must be carefully placed between the folds of the drying paper, not forgetting to lay out, according to the natural direction, any

of the parts which the water may have disarranged. It then remains to disorganize the flowers kept free from the water, which would continue to grow separately under the pressure; the ovaries of the Orchidaceæ, of which the subterranean stock and stem have been warmed, ripen thus in exhausting the other parts of the flower. When the plants have been enclosed between the double sheets of drying paper, then, by means of a common glass bottle, which is used after the manner of a roller, the part of the plant which is still living is gently crushed, so as to disorganize, without, however, entirely flattening it. Orchideous plants are sometimes put into warm paper, and changed frequently, with the view, if possible, of preserving their colours by the rapidity of drying. Scarification has sometimes been adopted with the view of allowing the juice to flow out rapidly. At the time the specimens are laid out on the drying paper, a label shall be inserted with the date of collecting, the name of the station, its elevation above the sea (if it can be ascertained), and any remarks as to soil or geological structure that may be known.

1234. When the specimens (whether Phanerogamous or Cryptogamous) are fully dried, they are then selected for the herbarium, and are fastened upon fine stiff paper, 17 inches by 10½. In large herbaria, which are constantly consulted, the best way of securing the specimens is by means of fine thin glue; the plants, after the glue is put on them, being made to adhere to the paper, by pressure between folds of drying paper. Some use gummed paper, others use thread or narrow ribbon, by means of which the specimens are sewed to the paper. It would appear that insects have an unfortunate predilection for the most beautiful plants. They perforate the specimens at every point, and rarely fail to attack the Umbelliferae at the centre of the umbel, in such a manner, in fact, as to cause the rays to be detached immediately on touching the specimen. There are few botanists who do not know by experience with what chagrin and discouragement one finds that on the very day it is desired to consult the precious specimens, so carefully placed intact in the herbarium, the whole is completely reduced to powder. The plants of certain families are more particularly exposed to the ravages of insects. Those families most frequently attacked are the Cruciferae, Umbelliferae, Gentianaceae, Euphorbiaceae, Salianiae, Liliaceae, &c. Under these circumstances, all plants, whether acquired by exchange or gathered by one's own hand, must never be neglected to be submitted to an alcoholic solution of corrosive sublimate. The immersion of dried plants in the alcoholic solution mentioned, has the inconvenience of discolouring them more or less completely, and making them assume a light brown tint; but there can be no hesitation between the alteration of their colour and the complete destruction with which they are menaced, if not submitted to the above manipulation. In herbarium presses camphor is employed to prevent the attack of insects. The specimens must be kept dry, and frequently examined, and when insects are present they must be retouched with the solution already indicated.

1235. **Ferns, Mosses, and Lichens.**—Ferns are plants generally very easy to prepare and dry, requiring scarcely any particular care. They ought to be gathered in their various stages of development, above all in their condition of fructification. If the plant presents fertile and sterile fronds, it is, perhaps, unnecessary to add that both must be collected. I have nothing specially to observe with regard to Lycopodiaceae, Equisetaceae, Marsileaceae, Isoetideae, &c. Suffice it that the same *modus operandi* applies to them as to the phanerogamous plants in general. As to plants of the families of the mosses and lichens, they present a peculiarity somewhat curious and useful to know relative

to their gathering and preparation. These plants being completely dried and *dead* to all appearance for months and even years, if moistened and exposed to the free air reassume all the semblance of life, indeed, as if their existence had never been interrupted. When dry, however, they must be protected from humidity, for if damp and they ferment, their destruction is inevitable. In order to gather mosses and lichens, if the species grow upon a tree, a portion of the bark must be carefully detached with it; a lichen growing on the damp ground must be taken up with the digger. Some lichens, on stones and rocks, are so encrusted that it is necessary to use a chisel and hammer and detach the piece of the rock on which the specimen grows. After rain, and during damp weather, is the best time for collecting mosses and lichens, as they are then easily detached, and less liable to be broken. Certain mosses, and the major part of the lichens, can be gathered during all the seasons of the year; but it is particularly in autumn, towards the end of winter and the first portion of spring, that the mosses and Hepaticæ are in the best condition, viz.: in fructification. In gathering mosses and lichens the botanist must provide himself with small bags of strong white paper, each specimen being placed in its particular bag, and the name of the plant, if known, with other information in connection with its gathering and habits, written in pencil on the outside. This mode of proceeding is a *sine qua non* with regard to the plants in question, otherwise if placed in the vasculum together they would be sure to mingle and cause confusion. Mosses and lichens being placed between the folds of the drying paper, are then treated like ordinary plants, but their preparation occurring more frequently in winter, when the air is charged with humidity, and being, besides, of a vitality susceptible of long resistance, it is essential to leave them longer between the folds of the paper than would be necessary in summer. The foliaceous and encrusted expansions of lichens demand no especial care; the same may be said of those mosses which are of an elongated and dendroidal form, but those species which are minute, and which, growing upon the damp soil, present a short and serrated tuft or *gazon*, require particular precautions. They must be divided with a knife, and placed upon the paper in such a manner that the soil left may be immediately underneath the moss; the earth adherent to the roots being, at the same time, detached, leaving sufficient, however, to form a *binding matrix*. When these preparations are complete, they are then placed on white paper with a solution of gum arabic, and transferred to the herbarium.

1236. **Algæ and Characææ.**—Algæ are of two classes, fresh water and marine. Our stagnant fresh waters contain a great number of genera and species; some even are found on the humid surface of the earth; but it is in sea water that the largest and most beautiful in configuration and colour are to be found. They are, for the most part, attached to submerged rocks, exposed on the receding of the tide, and also to rocks always submerged at considerable depths. A great portion of species of algæ, both of fresh and salt water, consist of long flexible filaments, or thin delicate membranes; these plants assume the form of a moist hair pencil when withdrawn from the water, and if placed in that state upon paper, in order to be dried, they form a compact mass, in which it would be impossible to distinguish any character or discriminate the form of the living plant. To prepare, properly, the algæ, and possess them in all their beauty, it is necessary to have a wide vessel or basin in which the plant is placed, when the filaments are lightly agitated with a quill or such like instrument, to make them take the direction that is natural to them. A sheet of paper is now introduced under the algæ, agreeably to the size of the specimen, and when

fairly settled and spread out on it, gradually withdrawn in an inclined direction from the base to the extremity. The paper, upon which the alga has spread itself, being withdrawn from the water, it is placed on an inclined board, to allow as much of the remaining water to drip from it, prior to placing it between the folds of the drying paper; before enclosing it, however, it will be necessary to cover the specimen with a piece of tissue paper, slightly greased with fine lard, the superabundant grease being removed by placing the tissue paper first between unsized paper and submitting it to gentle pressure. The object of this greased tissue paper is to prevent the specimen adhering to the drying paper. These covering slips of paper must be, of course, prepared beforehand. After being properly dried in the usual way the specimen may be immediately transferred to the herbarium, and, as insects attack few algæ, it is not necessary to apply the preservatives which are in use for other plants. For the microscope the algæ are placed upon a lamina of mica, prepared for that purpose. The most delicate species of algæ, thus prepared, do not alter by desiccation; they preserve not only their form and all the elegance of their configurations, but likewise their magnificent colours. To study them with the microscope it is only necessary to moisten a fragment, when it will assume the appearance of the living plant. The large species of fucus are prepared like phanerogamous plants, folding the fronds upon themselves, if they exceed the size of the paper; but previously it is proper to wash out the salt which they contain with fresh water. In the case of sea-weeds, it is necessary to avoid such specimens as are in a state of decay. Those which are taken should be spread out in the shade to dry, without washing them with fresh water, and when quite dry, packed loosely in a box. Many species are found thrown upon the beach, and the pools in the rocks at low water are often filled with excellent specimens. The stems of the larger algæ are often covered with parasitic species, which should be dried without separation.

1237. **Fungi.**—Certain fungi are vivaceous, and of a ligneous consistence, and, consequently, may be gathered at all seasons of the year, but it is particularly during autumn that large species are developed, of which the duration is so transient. To preserve the small species, which grow on bark and leaves, and even the vegetations on substances in a state of putrefaction, they are dried with paper like ordinary plants. But if it is desired to devote time and study to this class, they must be submitted to the microscope, prior to drying them, and magnified drawings made with the greatest care; of those minute species, of which the forms are most curious and varied, the drawings must be placed in the herbarium by the side of the specimens themselves. As to the larger species it is not the less important to reproduce them, by means of the pencil, on account of the extreme difficulty of preparing them in such a manner that they may be recognizable. These drawings must all be tinted, for colour is an important character for the distinction of species. The ligneous fungi can be preserved very well, but the majority can be with difficulty placed in an herbarium on account of their volume and thickness; “a thin slice must then be taken from the centre, extending from the top of the pileus to the base of the stipe. This is dried separately to show the gills or spores. The inner cellular portion of the pileus and stipe is then removed, and those parts are dried so as to give the form. Dried fungi are very much exposed to the ravages of insects; they must, therefore, be impregnated occasionally with a solution of corrosive sublimate in alcohol. Many melancholy accidents have occurred from parties mistaking other members of this division for the *Agaricus campestris*, the species

generally used in the culinary department; it may not be irrelevant, therefore, to advise particular attention to the characters which distinguish the chief edible mushroom, otherwise considerable danger is incurred. Travellers visiting foreign countries (although not botanists) will find it an easy matter to preserve mosses, lichens, and sea-weeds in a state fit for after-examination.

1238. Dissections of Plants.—It is much more advantageous to study and dissect a living than a dried plant, for the study of the organs in a fresh state is more easy, and presents many more chances of rigorous exactitude. Unfortunately it is impossible in a number of cases to procure the living plant required, and one must be satisfied with obtaining a dried specimen. In order to submit these dried specimens to study, ramules of flowers, fruits, or any other organ, must be detached and subjected for some minutes to ebullition in water, or, in event of the tissues being very delicate, simply immersed in hot water and then left to cool. It is surprising to observe with what rapidity, under the influence of hot water, the organs deformed by the compression and dessication of years, reassume their form and almost primitive consistence. Fruits, again, become pulpy, seed soften, corollas regain their original form, and the various parts retake their respective and natural dispositions. Dissection is said to be *organic* when it has for its object the isolation of the parts of the flower, of the fruit and seed one from the other, and the study of their forms, situation, and general relations. Dissection is termed *anatomical*, when the end in view is the exact knowledge of the nature of the organs, and the examination of the tissues which constitute them. A simple microscope suffices in the majority of cases for organic dissections and for analytical researches with a view to the discrimination of phanerogamous species, and even the greater portion of cryptogamous ones. For anatomical dissections, on the other hand, it is necessary to have at command the *best* and most powerful microscopes, and the finest scalpels and pointed *needles*. Although the observation of anatomical details require often considerable magnifying powers, it would be in vain to attempt to dissect with such aid, not only because the compound microscope, in reversing the image, renders the movements of the scalpels and *needles* very difficult to direct, but the distance is so small which separates the *objective* from the *object*, as almost to form an insurmountable obstacle to dissection. It is for this reason that with the simple microscope even anatomical dissections must be prepared, and under a considerably diminished magnifying power, with a view of being transferred to the more powerful lenses of the microscope. It must be borne in mind that in the majority of dissections, whether the organs be derived from a living plant, or one softened in water, they must be made in a *drop of water*. The water prevents the objects passing from under the scalpel, facilitates the separation, and hinders the drying, at the same time adds to the transparency, and consequently aids observation. Certain organs are endowed with a hygrometric property, in virtue of which their appendages roll or fold themselves one upon the other under the influence of moisture, and only spread themselves out under the influence of dessication. Such are the teeth of the peristome of the urn of the mosses, the hairs of the tufts of the Compositæ, &c. When, then, these organs have been submitted to softening in water, their state of contraction renders their study almost impossible. This inconvenience is remedied by placing them upon a slip of glass, or, better, upon a slip of strong paper, heated gently, and as rapidly as possible, over the flame of a spirit lamp. On the instant the divisions of the peristomes spread themselves out, and the fascicles of hairs expand, only then can these organs be with utility submitted to observation.

1239. Specimens in a moist state.—In preserving fresh specimens of fruits, and the other parts of plants, the best mode is to put them into a saturated solution of salt and water. They can thus be sent home from foreign countries in jars or barrels. In making a museum of such specimens, they are put into glass jars, the sizes of which should be regular—4, 8, 12, and 16 inches high, with a diameter varying according to the size of the specimen. The glasses may be filled with the following solution, which is nearly the same as that used by Goadby, and which seems to answer well in most instances:—

Bay salt.....	4 ounces.
Burnt alum.....	2 ounces.
Corrosive sublimate.....	20 grains.
Boiling water.....	2 quarts.

Dissolve and filter the solution. Spirit is often used, but it usually makes all colours alike brown. It is useful for delicate specimens which are required for dissection. Diluted acetic acid may be employed in some cases. The mouth of the glass jars may be conveniently covered with India rubber, or in the case of glasses of small diameter, with a watch glass secured by sealing wax, or by circular glass covers cemented by a lute composed of resin 1 part, wax 2 parts, and vermilion 1 part.

1240. Seeds, when sent from abroad, should be collected perfectly ripe and dry, and if possible kept in their entire seed-vessels. Small seeds may be folded in cartridge paper, and should be kept in a cool and airy place during transport. Large seeds and oily seeds, which lose their germinating power speedily, are best transported in earth. A box about 10 inches square, with the sides $\frac{3}{4}$ of an inch thick, answers well. In this may be put alternate layers of earth and seeds, the whole being pressed firmly together. Living plants are best transported in Wardian cases (§ 297), and seeds or fruits may also be scattered in the earth of the cases. Bulbs and rhizomes not in a state of vegetation, cuttings of succulent plants, as aloes and cactuses, and the pseudo-bulbs of Orchideous plants, may be put into a box or barrel with dry moss, sand, peat, or sawdust.

1241. Botanical Study and Room for Herbarium.—The “local” destined for the herbarium must be well aerated, in fact, exempt from humidity at all seasons. If, during winter, the slightest damp is suspected, the chamber must be carefully warmed, as a few weeks of humidity will suffice to destroy the most considerable and well-arranged collection. If, notwithstanding every precaution, some of the specimens disclose moisture, all the parts must be touched lightly with a hair pencil dipped in alcohol. This *local* should be situated as free as possible from noise and exterior motion. The centre of the room being occupied by a large plain table intended for spreading out the plants, either for classification and intercalation, or, as it may be, for comparing and studying them. The walls of the chamber should be occupied by the *cases* or presses to contain the herbarium, and the *books* for reference or consultation be always at hand. A table, smaller than the preceding one, should be placed near a window, with a large field of light, for the purpose of microscopic studies; and, in order not to fatigue the eye, the operator should place himself immediately at the side and not directly in front of the windows. Another table will be found of service if reserved for the instruments not immediately in use.

1242. Hints for Botanical Excursions.—In countries or districts where the means of communication and transport are numerous and accessible, it is not our province to indicate or limit the baggage of the exploring

botanist: we will content ourselves simply with recalling the adage, "If fine weather, take your cloak; if it rains, take it if you like." Nevertheless, for a botanical excursion the strictly necessary will always be the most convenient. In mountainous districts this may be laid down as the absolute rule, for the greater number of the routes are only accessible to travellers on foot and to horses; and botanic material being, in general, very considerable, it must be a point kept in view never to augment its volume or weight, without actual necessity. The indispensable baggage consists of a ream or two of the paper used by botanists for drying and preserving plants, placed between two boards of the same dimensions as the paper, and fastened by two leather straps. Then follow the smaller *field books* in which are placed the smallest and most delicate specimens, together with a larger and smaller sized vasculum. Anything of a more immediate personal nature must be contained in a small portable carpet bag, which must, at the same time, contain a small tin flagon of alcohol, and other articles necessary for the gathering, preparation, and study of the plants. One must be clothed with a view to precaution against cold and not *heat*, for whatever may be the heat of the day, it is always fresh and often cold during the night; besides frequently, during the same excursion, the heat will be excessive, changing to the glacial temperature of the cold winds, which usually reign amidst mountainous quarters; the danger lying in the sudden change from the profuse perspiration of a long walk to a sudden cooling. In order to avoid which extremes, it is necessary to be comfortably clothed, a change or two of stout worsted stockings, with strong double-soled tacketed shoes, are indispensable: a waterproof cape or cloak will also prove a most useful companion. As there is little time for study or extended observations during an excursion, it is only necessary to carry the usual portable lens and forceps, to be chiefly called into use during the evenings in arranging the plants gathered during the day. The only book required is the "Flora," or annotated Catalogue of the plants of the country or district to which you have directed your attention: a thermometer will also be of service if it be wished to determine the different heights at which the various species grow. When an excursion is prolonged to days, make it a rule to rise with the sun, and return only to head-quarters on the day's decline, preparing the plants gathered prior to either eating or sleeping. Preserve always independence whatever friends may be met in places where a halt is made. These halts must be devoted to finishing any work in the preparation of the plants, placing those which are quite dry in separate packets, with a view to having them forwarded to their destination. With the view of transporting dried plants securely in wet weather, it is useful to have a supply of oil cloth to cover them. Each evening also make it a rule to note down any observations in a journal kept for that purpose.

CUI BONO?—CONCLUSION.

How to reply to this question so often repeated by persons strangers to the studies and pleasures of the botanist? How will you be enabled to estimate the satisfaction which the mind experiences in the research and discovery of a natural law, and in the contemplation of the wonders of nature? Will you speak of the inexhaustible and veritable happiness which to the naturalist replaces ordinary pleasures, and of that real beauty which he knows well to discriminate from conventional beauty? Of the sentiment of admiration with which he views the elegant bearing of the most common weed of our fields; the most humble

convolvulus for example, encircling, with its spiral blossoms, the flexible stem of a grass which he prefers to the comparatively best fabrics or articles decorated, as they generally are, with representations of the vegetable world, so incorrect and often ridiculous? Of the pearls and diamonds of the rose which sparkle on the leaves under the influence of the sun's first rays, and of which the effect appears to him in no wise inferior to the lustre of the most precious gems? The interrogator little affected by the beauty of plants, and who has never thought of them in any other point of view than that of alimentary consumption, would look at you with astonishment, and the more benevolent be scarcely able to repress a smile! Yet the botanist not only picks with care a weed, but seems to attach to its possession the utmost importance; detaches by turns an almost imperceptible fragment of its flower, and devotes hours, nay, entire days, to its study. If you mention the services which the profound study of plants and their properties, and the different modes of culture applicable to them render to medical science, to agriculture, horticulture, and domestic economy, you will be comprehended more clearly; but if you do not derive a *material* advantage yourself from their labours, of what use is it?—will only give place to—of what use is it to you? If, even admitted, that these studies may form a very good pretext for spending leisure hours, it is, on the other hand, difficult to convince that you can also devote to the same pursuit an entire life, and with profit. In fine, let us not reproach the cock for preferring the barley pickle to the pearl, but rest content in rendering every justice to the barley pickle, with a view of appreciating the pearl at its value. “Ergo vale, Lector amice: sylvas ruraque lacte peragra, et scientiam amabilem auge.”



GLOSSARY.

A

A (α privative of the Greeks). This letter placed before a Greek or Latin word indicates the absence of the organ: *Ex.* Aphyllus, without leaves; Acaulis, without stem.

ABNORMAL. Form different from the usual form.

ABORTIVE. Said of an organ which has been suddenly arrested in its development.

ABBREVIATION. Consists in words reduced to their first syllables. Names of authors, for example, are abridged thus—Lmk., Lamarck; L., Linnaeus; D. C., De Candolle. R., rare; ed., edition; fl., flora or flower; fr., fruit; fruct., fructification; fig., figure; herb., herbarium; var., variety; s. v., sub variety; vol., volume. Among the signs more commonly used are the following:— ∞ , indicates an indefinite number, as in the case of stamens;

① ⊙ A, an annual plant; ② ⊙ ♂ B, biennial; 24 Δ P, perennial; ♀, ligneous plant, a tree;)=, plant turning from left to right; (, plant turning from right to left; ∞, turning in an indeterminate direction; (o=), cotyledons accumbent, radicle lateral; (O //), cotyledons incur-

bent, radicle dorsal; (O >>), conduplicate, radicle dorsal: these three signs have been principally used in the description of the Cruciferae. ♂, male flower; ♀, female flower; ♂ ♀, hermaphrodite, or having both stamens and pistils; ♂ ♀, unisexual, or male and female flowers separate; ♂ ♀, monœcious, same plant containing male and female flowers; ♂ ♀, diœcious, male and female flowers on different plants; ♂ ♂ ♀, having hermaphrodite and separate male and female flowers on the same plant;

ACI

!, the point of exclamation is the sign of certitude; ?, the point of interrogation is the sign of doubt.

ABRUPT. In an abrupt manner; abruptly acuminate, where a point rises from the broad part of a leaf; abruptly pinnate, synonymous with paripinnate, or terminating in two pinnae.

ABSORPTION. Function of vegetable tissues endowed with the property of absorbing liquids, as for example, in the case of *spongioles*.

ACALYCATUS. Deprived of the calyx; applied to dicotyledonous plants, of which the floral envelope seems to be single.

ACAULIS. Without an apparent stem.

ACCRESCENT. Applied to parts which continue to grow after flowering. In the case of the *Physalis Alkengi* the vesiculous red coloured envelope which contains the bacciform fruit is constituted by an accrescent calyx.

ACCRETE. Said of an accrescent organ which has reached the apogee of its development.

ACCUMBENT. Relates to the embryo of the Cruciferae, and results from the different curvatures of the cotyledons and radicle, and the different positions of the radicle with regard to the cotyledons.

ACEPHALOUS. Without head. *Ex.* Brassica oleracea, variety *Acephala*.

ACEROSE. Terminated by a very sharp point.

ACETABILIFORM. In the form of a cup.

ACHÆNIUM (α privative, and *χαίρω*, to open). The most simple achæmium consists of an isolated and monospermous carpel, as in the case of the *ranunculus*, *anemone*, *rose*, &c.

ACHLAMYDEOUS. Having no floral envelope

ACICULAR. In the form of a needle.

ACINÆIFORM. In the form of a sabre: *Ex.* *Mesembryanthemum aceniforme* (*seldom used*).

ACO

- ACOTYLEDONOUS.** Wanting cotyledons.
- ACROCARPI.** Mosses having their fructification at the summit of their branches.
- ACROGENS, ACROGENOUS.** Plants which increase only by the summit. They present a distinct axis, and often veritable leaves, and consist of vascular and fibrous and cellular tissue, such are the Ferns, Mar-silaceæ, Lycopodiums, and Mosses.
- ACULEATE.** Having prickles.
- ACUMINATE.** Of which the summit terminates abruptly in a sharp point. Sepals, petals, capsules, can be said to be acuminate.
- ACUTE.** Terminating in a small point.
- ADHERENT.** Parts attached or united, as the calyx to the ovary. Advantageously replaces the expression *ovary inferior*.
- ADNATE.** United in all its length to another organ of which the dimensions are relatively large, and in such a manner that the adnate organ seems to be an appendage. *Ex.* The stipules are adnate in the Rosaceæ. Anthers united to the corolla in all their length are also said to be adnate.
- ADRESSED or APPRESSED.** Closely applied or bedded. *Pili Adpressi.* Appressed hairs.
- ADVENTITIOUS.** Organs which do not follow the normal rule. *Ex.* Roots which grow from stems, particularly if aerial.
- AERIAL.** The expression aerial stem is used in contradistinction to subterranean stem.
- ÆSTIVATION.** The disposition or arrangement of the organs which constitute the flower, particularly in the state of bud.
- AFFINITY.** Relation between genera and species.
- AGAMOUS.** Not produced by fecundation. This expression is replaced by Cryptogamous.
- AGGLOMERATED.** Organs are said to be agglomerated when they lie close together in compact masses.
- AGGLUTINATED.** So adherent as to be detached with difficulty.
- AGGREGATED.** Organs together in a single mass. The fruit of the mulberry is composed of fructiferous flowers aggregated.
- AGYRATUS.** Not rounded.
- ALA.** Prolonged membrane in the form of a wing. The two lateral petals of a Papilionaceous flower receive this appellation; also the extension of the carpels of the elm, &c. Certain seeds surrounded by a thin and membranous border are said to have alæ.
- ALBUMEN.** Perisperm and Endosperm. Envelope of the embryo performing the same functions as the white of egg.
- ALBURNUM.** The layer of new wood formed each year by a Dicotyledonous stem.

ANG

- ALGÆ.** Water weeds.
- ALPESTIS.** Applied to plants whose habitat is in alpine regions.
- ALTERNATE.** Said of objects which are arranged in such a manner that each of them is found at the same level of the interval which separates them from two others situated on another level. Leaves are termed alternate when they are spread one by one on the same stem at different levels.
- ALVEOLATE.** Said of a surface which presents regular cavities separated by thin walls similar to the honeycomb. *Ex.* The receptacle of the sunflower (*Helianthus annuus*).
- AMENTUM.** Unisexual spike. The amentaceous inflorescence characterizes the families Juglandæ, Cupuliferæ, Salicinæ, Betulinæ, &c.
- AMYLUM.** Fecula or starch.
- AMNIOS.** The gelatinous liquor which surrounds the embryo and intended for nourishment. The membrane which contains the amnios is called the embryo sac.
- AMORPHOUS.** Presenting no definite form.
- AMPHIBIOUS.** Applied to plants which are both terrestrial and aquatic. *Ex.* Polygonum Amphibium.
- AMPLEXICAUL.** Embracing the stem in the greater part of its circumference.
- AMPULLA.** Applied to the vesicles borne by the leaves of Utricularia.
- ANA.** Greek prefix, signifying *upon* or *to*; used in compound words. *Analogous*, resembling another.
- ANASTOMOSIS.** The confluence of vessels, or union of the nerves or veins of a leaf.
- ANATROPOUS.** (See *Ovule*.)
- ANCEPS.** Double edged.
- ANDRIA (Male).** Monandria, one stamen; Polyandria, many stamens.
- ANDRÆCIUM.** Applied to the stamens, or male organs of the flowers, &c. Analogous to calyx, corolla, and disc.
- ANDROGYNOUS.** A monoecious plant, the male and female flowers being grouped on the same plant, as in some species of *Carex*.
- ANDROPHOROUS.** The tube formed by the filaments united to the stamens; such is the staminal column of the Malvaceæ.
- ANDROSTYLUM = Gynostemium.** The organ resulting from the stamens being united with the style. *Ex.* The Orchideæ.
- ANFRACTUOUS.** Applied to sinuosities; the result of irregular folds; the cotyledons of the nut present anfractuosities.
- ANGIOCARP.** A fruit covered by the surrounding parts which are developed with

ANG

- it, and to which it adheres; fruit of the mulberry for example.
- ANGIOLUM** = Peridium. The membrane which envelops the parts of the fructification of certain Fungi. *Ex.* The Lycoperdon.
- ANGIOSPERMOUS.** (αγγιον, a vessel.) Applied to plants whose seeds are enclosed in a pericarp, in contradistinction to gymnospermous, in which the seeds appear uncovered.
- ANISOS.** (Latin *inequalis*.) Unequal; anisostemonous signifies that the stamens are different in numbers to the petals; isostemonous meaning that the stamens are equal in number to the petals.
- ANNULUS.** A ring, the remains of the veil which in certain agarics or mushrooms remain around the stalk; the annulus elasticus constitutes the circular rim of the sporangia of the ferns.
- ANNUALS.** Plants which last only a year.
- ANTHELA.** Name given to the inflorescence or false panicle of the Juncus, Luzula, and certain Cyperaceæ.
- ANTHER.** (ανθεος, flowered.) The part of the stamen which contains the pollen.
- ANTHERIDIUM.** The male organ in Cryptogamic plants.
- ANTHODIUM.** The capitulum of flowers in the family of the Compositæ.
- ANTHOS.** A flower; (Latin, *flos*.) *Ex.* Polyanthus, many flowered. In compound words from the Latin, *florus* replaces anthus, as Grandiflorus, synonym of Macranthus.
- ANTICUS.** Anteriorly, or in front, synonym of *introrsus*, or introrse.
- ANTITROPAL.** Is said of an embryo, the radicle of which is diametrically opposite to the hilum.
- APETALOUS.** Having no petals or corolla; in apetalous plants the calyx is susceptible of presenting the green colour, as in the families of the Urticaceæ, Cannabineæ, but frequently the calyx is coloured, as in the genera Daphne, Aristolochia, &c.
- APHYLLUS.** Devoid of leaves. The leaves exist in some cases, but are reduced to discoloured or coloured scales (*not green colour*), as in the Neottia nidus avis; *secondly*, the leaves exist under the form of bractæ and floral organs, as in the Cuscuta; *thirdly*, the aerial stem being represented by a single internode, can present leaves in its length only. In this last case this aerial stem is only in general a peduncle arising from a subterranean stem, as in the Primula officinalis, Scilla nutans, &c.

AST

- APICULATE.** Terminated at the summit by a short and sharp point.
- APOPHYSIS.** The swelling at the base of the urn in certain genera of the mosses.
- APOTHECIUM.** In the lichens the receptacle containing the fructification.
- APPENDAGE.** Applied to divers accessory parts or prolongations. The tails of the anthers in the Compositæ are appendages; they may be terminal, lateral, dorsal, or basilar, according to their position.
- APTEROUS.** Without wings, in contradistinction to alatus.
- AQUATIC.** Applied to plants which grow in fresh water, whether plunged completely, or having their inferior parts only submerged.
- ARACHNOID.** Long fine hairs intercrossed, and resembling the web of a spider, as in the rosettes of the Sempervivum arachnoideum.
- ARBORESCENT.** Like a tree.
- ARCHEGONIUM-PISTILLIDIUM** is to the sporangia of the Cryptogams what the ovary is to the fruit in Phanerogamous plants.
- ARCUATE.** Curved in the form of an arch.
- ARENOSUS.** Preferring sandy places.
- AREOLA.** The circular stain or coloured circle occupying the bottom of a corolla. Areolate, marked with an areola.
- ARILLUS, ARILLOA.** The membranous covering of certain seeds, enveloping some more or less completely, as in the mace of the nutmeg.
- ARHIZUS.** Without root: *Ex.* The Lemna arhiza.
- ARISTA.** A prolongation, or filiform appendage, as seen in the glumes of the Gramineæ.
- ARMATUS.** Furnished with an armature of sharp points or prickles.
- ARTICULATIONS.** Joints easily separating.
- ARVENSIS.** That which grows in the fields.
- ASCENDANT.** Stems united at the base are said to be ascendant. In the case of other plants when the stems are procumbent naturally, and only rise in their upper part. *Ex.* Polygonum persicaria. The filaments of the four long stamens of the Crucifere are ascendant, the filaments of the two short stamens on the same are straight.
- ASCIDIUM.** Pitcher or vessel formed by the folded leaf, as in Nepenthes distillatoria, &c.
- ASCUS** (*Plur.* ASCI). Sac containing many spores; a sort of sporangium. Ascigerous, bearing sporiferous sacs.
- ASPERITY.** Roughness, from the surface being covered with short hairs.
- ASTOMOUS.** Without mouth.

AST

ASTYLUS. Without a style; said of an ovary of which the stigma is sessile.

ATROPOUS = Orthotropous.

AURATUS. Gold colour.

AURICULATED. Having ear-like appendages; the foliaceous expansion at the base of a petiole. *Ex.* The leaf of the *Sonchus oleraceus*.

BUL

AXIL. The angle where the leaf and stem unite.

AXILLARY. Rising from the axil of a leaf.

AXIS. The stem, either aerial or subterranean, and the root constitute the axis, and that axis ramifies itself indefinitely into secondary axes.

B

BACCA. A berry; soft and succulent fruit containing many seeds; it may consist of a single carpel, or many carpels united together. Bacciform, or Baccate, in the form of a berry.

BALAUSTA. This name is given to the fruit of the *Punica granatum*, or Pomegranate. It is composed of two superposed rows of carpels arranged in such a manner that in the inferior row the placentas are axillary, and in the superior they appear parietal. These two rows of carpels constitute an adherent fruit, of which the pericarp is coriaceous, and surmounted by the persistent limb of the calyx, which forms a crown with five teeth. The carpels are filled with numerous seeds pressed one against the other.

BARBA (and in words derived from the Greek, *Pogon*. *Ex.* *Andropogon*). Hairs arranged in regular order; barbed, bordered, or terminated by a regular mass of hairs.

BARK (*Cortex*). The exterior covering of the stem which envelops the wood-like case. The wood forms a cylinder in the dicotyledons, composed of fibro-vascular fascicles arranged in circles or concentric layers around the medulla. In the Monocotyledons the fibro-vascular fascicles appear scattered in a mass of cellular tissue.

BASIS. The base of an organ. Basilar, appertaining to the base.

BASIGYNIUM = Podogynium. Support of the ovary.

BEAK (*rostrum*). Terminal prolongation of an organ. In the *Carex*, the fruit (*achænium*) is enclosed in a bractial accrescent covering, named a utricle; this utricle is frequently prolonged into a beak.

BIS, and in compound words with *Bi*. Twice. Bidentate, with two teeth-like processes; bifid, a leaf divided into two lobes to near the middle of its length; bifoliate, with two leaves; bifurcate, like a fork; bi-jugate, with two pair of compound leaves; bilabiate, Gamopetalous corolla with two lips, *Ex.* *Lamium*; trilamellar, presenting two lamellæ; bilobate, having

two lobes; bilocular, with two loculements; binary, arranged by two; bi-partite, cut down in two beyond the middle; bi-pinnate, a leaf twice pinnate; bi-pinnatifid, divisions of the leaf pinnatifid and themselves pinnatifid; biplicate, folded twice; bi-septate, with two divisions; bisexual, with two sexes; bisuturate, with two sutures; biternate, with divisions themselves ternate, or divided into three parts; bivalvular, with two valves; biporose, with two pores.

BOTHRENCHYMA. Dotted vessels which have depressions in the interior of their walls.

BRACHIATE (*Rami brachiati*). Horizontal opposite branches crossing the branches at a right angle, which spring immediately from a superior or inferior level.

BRACT. A leaf from which the flowers proceed, and intermediate between the leaves of the stem and the sepals; bracteoles, or bracteolæ, are bracts which are at the base of the particular support of each flower, as in a multifloral inflorescence.

BRYOLOGY (*Bryologia*) = Muscology. That division of Botany which treats of the structure and classification of mosses. The former is the more proper term, as the latter, Muscology, is somewhat ambiguous, and might be applied to the study of flies.

BULB. A subterranean bud with scales. When the scales cover the bulb, or almost do so, the bulb is said to be tunicated, which is in general the case; when the scales are narrow, as in the genus *Lilium*, the bulb is said to be squamose; in the *Allium fallax* the bulbs constitute the terminal buds of the divisions of a rhizome or underground stem. Bulbous plants belong in general to the division of the Monocotyledons; a small number of Dicotyledons, however, produce bulbous subterranean buds. *Ex.* *Saxifraga granulata*, certain *Oxalis*, &c. Bulbosus, having the character of a bulb; bulbiform, the appearance of a bulb; bulbiferous, bearing bulbs; bulbil (*bulbilus*), a small bulb; the small bulbs which

BUR

develop themselves in the place of flowers upon the umbels of certain species of *Allium*, the *Allium vineale*, for example, are named bulbils; bulbils develop themselves at the axil of the leaves of the *Lilium bulbiferum*.

CAR

BURSICAL. In the description of the flower of certain Orchids, this name is given to a membranous fold forming a small sac, or purse, containing the adhering extremity of the pollenic masses.

BYSSACEUS. Composed of long silky hairs.

C

CACUMEN = Apex. The summit.

CADUCOUS. Applied to organs which detach themselves spontaneously from the stem, disarticulating at their base.

CAERULEUS. Of a blue colour.

CAESPITOSE. Having the appearance of a tuft of grass.

CALATHIFORM. Having the form of a cup. *Calathium* is syn. with *Anthodium* and *Capitulum*.

CALCAR. A tubular prolongation from the base of an organ almost similar to the finger of a glove. In the *Aquilegia*, each of the five petals is prolonged into a *Calcar*. In the *Orchis* it is the labellum which takes this form; in the *Delphinium* it is one of the superior divisions of the calyx; in the case of the *Impatiens* (*Balsaminæ*) and the *Tropæolum* it is the inferior sepal. *Calcarate*, flowers which are furnished in this way.

CALCEIFORM or CALCEOLATE. In the form of a shoe or slipper. *Ex.* The petals of the Orchid *Cypripedium*, and the *Calceolaria*, or Slipper plant.

CALLOSITY. Said of an organ which has a coriaceous thickening or consistence on a limited portion of its extent.

CALYCIFLORÆ. De Candolle's name for the division of polypetalous Dicotyledons, whose flowers are inserted upon the tube of the calyx.

CALYCIFORM. In the form of a calyx.

CALYCVUS or CALICULUS. A bundle or row of small bracts, which forms a verticill immediately beneath the calyx, and resembling an exterior calyx. *Ex.* The Pink. In the *Potentilla* the *caliculus* is formed by the stipules united two by two, which are a dependence of the sepals.

CALYPTRA. The covering of the sporangium of mosses.

CALYX. The exterior verticill of the flower; it is sometimes developed in the form of fine silk hairs, as on some of the *Compositæ*.

CAMBIVM. Elaborated mucilaginous sap which contributes to the formation of the vegetable tissues.

CAMPANULATE. In the form of a bell. *Ex.* The corolla of the genus *Campanula*, and the calyx of the *Hyoscyamus Niger*.

CAMPESTRIS. Growing in the fields.

CAMPULITROPAL or CAMPYLOTROPAL. See OVULE.

CAMPYLOSPERMATE. Of which the seed is curved.

CANALICULATE. Folded longitudinally like a groove: in some leaves the external surface of the petiole is furrowed or grooved.

CANCELLED. In form of a lattice or trellis.

CANDIDUS. Of a pure white.

CANESCENT. Appearing white by reflection.

CAPILLARY. Having the form of a hair.

The word capillary denotes a finer object than filiform, and less fine than setaceous. The stems of the *Sagina* *apetala* are filiform; the beak which surmounts the achæmium of the Dandelion is capillary, and the silky hairs which the beak bears are setaceous.

CAPITATE. Terminated in a rounded head. *Pili Capitati*, capitate hairs, their extremities being rounded.

CAPITULUM. The inflorescence of the *Compositæ*. This name is applied to every inflorescence which forms an aggregation of sessile flowers.

CAPREOLATE (Cirrhus). Having filiform organs or tendrils.

CAPRIFICATION. Maturing of the fig by means of the wild fig.

CAPSULE. Dry vessel containing seed: it may be either single or have many cells, dehiscent or indehiscent.

CHARACTERS are the differences which distinguish one species from the species of the same group, or a natural group from the groups of the same order. Characters are said to be positive when they are founded upon the presence of organs, and negative when founded upon the absence of organs.

CARINA. A keel. The two united inferior petals of the corolla in the *Papilionaceæ*.

CARINAL. Belonging to the carina.

CARNOSE. Fleshy, pulpy.

CARPEL. *Carpellum*, *Carpellium*. A carpel is to the pistil or pistils what a sepal is to the calyx, a petal to the corolla, a stamen to the androeceum,—in fact, a modified leaf.

CARPOPHORE is a prolongation of the axis of the flower, which elevates the base of

CAR

- the ovary, or of the fruit above the point of insertion of the other verticils: in the Cruciferae, for example, the ovary is placed manifestly above the level of the stamens, and thus constitutes a carpophore.
- CARUNCULA.** Fleshy appendage upon a limited extent of some seeds.
- CARTILAGINOUS.** Consistence of cartilage. *Ex.* The leaves of the *Eryngium Maritimum*.
- CARYOPSIS** or **CARIOPSIS.** Dry fruit containing a single seed, which is united by all parts with a thin pericarp; this fruit has the aspect of a seed: such is the fruit (commonly termed seed) in the family of the Grasses.
- CATKIN** = **Catulus** = **Amentum**.
- CAUDEATE.** Having a tail.
- CAUDEX.** Applied to the stem of palms and tree ferns; more frequently stipes is employed instead.
- CAUDICLE.** The diminutive of *Cauda*, and applied to the filiform process which supports the pollen mass in Orchids.
- CAULESCENT.** Provided with an evident stem, in contradistinction to those which have a subterranean stem, however short it may be.
- CAULIS.** An aerial stem.
- CELLULE** = **Utricle**. Cellular tissue; a tissue composed of cells: the fungi, lichens, algæ, &c., are entirely composed of this tissue: *cellulose* is the chemical matter forming the cell wall.
- CENTIMETRE** (French measure). The hundredth part of the mètre; the tenth part of the decimètre; ten times the length of the millimètre, and equal to 0.39371 English inches.
- CENTRIFUGAL.** Flying from the centre. In the centrifugal inflorescence, the centre flower expands first, and successively to the circumference, in contradistinction to centripetal inflorescence, in which the expansion of the flower commences at the circumference, or from the base to the summit.
- CEREAL.** Appertaining to barley, wheat, oats, and such grains.
- CERNUOUS.** Inclined or curved by the weight of the terminal part.
- CHALAZA.** The point at which the funiculus (or the raphe, which is its continuation) ceases, and where the ovule or seed begins; the chalaza is therefore the real base of the ovule or seed.
- CHLOROS.** In compound words derived from the Greek, signifies green, or greenish-yellow.
- CHLOROPHYLE.** The colouring matter of the leaves.

CON

- CICUTRICULA.** Name applied to the cicatrix of the hilum upon the seed, when it is detached from the funiculus.
- CHROMA.** Colour; chromule, the floral colouring matter; chromism, excess of colour.
- CHRYEOS.** Gr., *golden yellow*; Lat., *auratus*.
- CICATRIX.** Mark left upon an organ by the base of a part articulated with it, after the fall of this part. A branch after the fall of the leaves present cicatrices corresponding to the base of the petioles.
- CILIA** (Cilium). The short hairs upon the surface of a leaf, of a petal, on the angles of a stem, and of a fruit. Ciliated, fringed with cilia.
- CINENCHYMA.** Laticiferous tissue.
- CIRCINATE** (*Folia Circinalia*). Leaves rolled up in the form of a crozier. *Ex.* The young leaves of the *Drosera*, and the fronds of the ferns.
- CIRCUMSCISSILE.** Seemingly cut circularly. *Ex.* The pericarp of fruits under the name of *Pixidium*. Circumscissile dehiscent, opening in the manner of a *pixidium*, or lid. The fruit in the *Anagallis*, and the *Hyoscyamus*, is a *pixidium*.
- CLADENCHYMA.** Branching cellular tissue.
- CLADOS**, in words derived from the Greek signifies branch.
- CLASS.** Comprehends a group founded upon characters of the first order; each class reunites several families. Families are to classes what genera are to families. Classification, methodic arrangement.
- CLATHRATUS** = **Canellatus**. In form of a lattice, or trellis.
- CLAVATE.** Like a club, increasing from the base to the summit. Claviform, in the form of a club.
- CLINANTHIUM.** Enlarged summit of a peduncle bearing sessile flowers, as in the *Compositæ*, *Dipsacæ*, &c.
- COCCINNEUS.** Bright scarlet colour.
- COCCUM** and **Coccus**. Kind of fruit. *Ex.* *Euphorbiacæ*.
- COCHLIARIFORM.** In form of a spoon. The aestivation of the *Aconite*, in which one of the petals larger than the others is curved upon, and covers them, is said to be cochlear.
- CENANTHIUM** (*κενὸς*, common, *ανθός*, a flower), is a term applied to the inflorescence of the fig, which is formed by flowers reunited into a common concave receptacle; in other words, a deeply depressed peduncle.
- COHERENT** = **Connatus**. Parts united among themselves, particularly if of the same nature. The word is sometimes used as synonymous with agglutinated. The authors are coherent in the *Balsamineæ*.

COL

- COLEORHIZA.** The sheath which envelops the first normal root in certain plants, the Gramineæ, for example.
- COLLATERAL.** Said of two organs placed side by side. *Ex.* Two ovules in an ovary.
- COLLUM.** Vital node, or neck; the part where the plumule and radicle of the embryo meet.
- COLPENCHYMA.** Wavy cellular tissue.
- COLUMELLA.** Column formed by the prolongation of the axis of the flower beyond the point on which are inserted the carpels. *Ex.* The carpophoræ of the Umbelliferae. This term is also applied to the central column in the sporangia of mosses.
- COLUMN** = **Gynostem.** The organ resulting from the uniting of the stamens and the style in the Orchids.
- COMA.** This term is applied to different organs: to tufts of radicle fibres, or hairs; and to elongated bractæ which pass beyond the inflorescence of certain plants.
- COMMISSURE.** In the Umbelliferae, the term is applied to the two exterior and opposite lines which correspond to the junction of the borders of the two carpels.
- COMOSE.** Furnished with hairs.
- COMPOSITE, or COMPOUND.** In the case of leaves when the limb is divided into several secondary articulated leaves.
- COMPRESSED.** When the form seems modified by a lateral pressure.
- CONCEPTACLE.** In Cryptogamic plants receptacles of various forms and natures which contain either spores or buds.
- CONDUCTOR (Tissue).** Name given to the tissue of the style, and of the placental cord through which the pollen passes to reach the cavity of the ovary.
- CONDUPPLICATE.** Folded double in a longitudinal sense. The young leaves are conduplicate in the cherry and the peach, &c.; not only leaves, but modified leaves may be conduplicate; the cotyledons are conduplicate in the Geranium, the Brassica, the Diplotaxis; the standard, or superior petal, is conduplicated in the bud of the Papilionaceæ.
- CONE.** A dry fruit covering naked seeds, and formed of bracts. *Ex.* The hop.
- CONJUGATE.** Arranged by two, or in pairs. Conjugate pinnate = bipinnate. *Ex.* *Mimosa pudica*, or Sensitive plant.
- CONNATE.** Said of two leaves which are applied to each other and united at their base. *Ex.* The floral leaves of the *Lonicera Caprifolium*, or honeysuckle.
- CONNECTIVE.** The part of the stamen which continues and terminates the filament, and to which part are fixed the anther lobes.

COR

- CONTORTED.** Regularly twisted. Contorted aestivation is seen in the corolla of the Malvaceæ.
- CONTRACTILITY.** The curious vital phenomenon which manifests itself in certain vegetable organs, and analogous to the muscular contractility of animals. *Ex.* The movements in the Sensitive plant, the sudden movement of the stamens in the *Berberis* when touched at the base; etherization and chloroform suspend, in the case of vegetables as well as animals, the phenomenon of contractility.
- CONVOLUTE.** The rolling up of leaves when in the first state of their development. Convolute aestivation is synonymous with imbricated aestivation.
- CORDATE.** In form of a heart. Said of the blade of a leaf or petal, the widest part being next the stem. *Ex.* *Viola odorata*, &c.
- CORDIFORM.** Analogous to the form of a heart, but applied particularly to organs which have a certain thickness; the embryo of *Trapa nutans*, the capsule of *Polygala vulgaris*.
- CORIACEOUS.** A dry membrane analogous to the consistence of leather; dry, thin, pericarps are in general coriaceous.
- CORM.** Underground stem. *Ex.* *Arum*.
- CORNATE, or CORNEOUS.** Having the hardness and appearance of horn; bicornis, or bicornate, having two horns; tricornis, or tricornate, having three horns. The perisperm of the Date and Asparagus are cornute.
- COROLLA.** In dicotyledonous plants the verticil, or envelope between the calyx and the stamens.
- COROLLÆFLORE.** De Candolle's name for gamopetalous hypogynous Dicotyledons. *Ex.* *Pumulaceæ*, *Solanaceæ*, *Labiatae*, &c., &c.
- CORONA (a crown).** The petaloid appendages arranged in a circle which exist at the throat of the corolla of the *Lychnis*, and of the perianth of the *Narcissus*.
- CORRUGATED.** Folded irregularly.
- CORTEX.** See **BARK**.
- CORTINA.** In the genus *Agaricus* the remains of the veil which continues adherent to the rim of the pileus.
- CORYMB.** A raceme in which the lower branches, or peduncles, are the longest, all the flowers, however, coming nearly to a level. The inflorescence of the *Sambucus niger*, or Elder, is a corymb. Corymbiferous, bearing a corymb; Corymbose, when the ensemble presents at the epoch of flowering the aspect of a large corymb.

COSTA (a rib) **COSTATE**. Presenting prominent ribs, as in the appearance of vessels in leaves. Each of the two carpels which form the fruit of the Umbelliferæ present ordinarily five ribs, called primary—one dorsal, two marginal or commissural, and two intermediate, or lateral. Some secondary ribs sometimes exist in the intervals which separate the primary.

COTYLEDON. The two first leaves of the embryo or rudimentary plant.

CRAMPOUS. A name applied to adventitious roots which grow on all the length of certain creeping stems, (*Ex.* *Hedera*, *Helix*, or *ivy*.) and serve to fix it when it climbs against a wall, or tree.

CRASSUS. Thick, fat. *Ex.* The *Crassulaceæ*.

CREMOCARP. Fruit of the Umbelliferæ formed of two achænia.

CRENATE. Finely indented. The leaves of the *Glechoma hederacea*, or ground ivy, are crenate. Crenatures, the indentations.

CRETACEOUS. Growing on chalky soils.

CRESTS. Fleshy appendages of fruits and seeds in the form of a crest. The middle lobe of the inferior petal of the *Polygala* is in the form of a crest.

CRISP. Undulated margins, as in some leaves.

CRUCIFORM. In the form of a cross. *Ex.* The corolla of the *Cruciferaæ*.

CRUSTACEOUS. A thick, hard, and fragile membrane, is said to be of a crustaceous consistence. *Ex.* Some lichens.

CRYPTOGAMS. Name given by Linnæus to the class in his system comprehending the *Cryptogams*. The cryptogams, or acotyledons, include the Ferns, *Lycopodiaceæ*, *Marsileaceæ*, *Equisitaceæ*, *Fungi*, *Algæ*, *Isoetidæ*, *Hipaticæ*, *Musci*, &c.

CRYPTOGAMOUS. Said of the organs of reproduction not constituted by stamens and ovules.

CRYPTOS. Concealed.

CUCULLATE. Like a cowl or hood. *Ex.* *Aquilegia vulgaris*. Cuculliform, in the form of a cowl.

CULM. Stem of the grasses.

CUNEIFORM, CUNEATE. Shape of a wedge. *Ex.* Leaves of a triangular form, the summit of the triangle constituting the base of the leaf.

CUPULA. Organ in the form of a cup; applied particularly to the involucre of the fruit in the family of the *Juglandææ*, the oak, chestnut, &c.

CUSPIS. Long point, large at the base and gradually attenuated. Cuspidate, prolonged into a sharp point.

CUTICLE. The external membrane of the epidermis.

CYATHIFORM. In the form of a goblet. Form of a reversed cone.

CYLINDRICAL. In the form of a cylinder; the stems of grasses are cylindrical; other stems, those of *Labiataæ*, for example, are prismatic.

CYMBÆFORM. Shaped like a boat.

CYME. A definite inflorescence composed of terminal axes, each tending to a single flower. The flowers include either umbels, corymbs, or racemes. Cymose, in the form of a cyme.

CYSTID, CYSTIDIUM. In the class of *Fungi* the name is applied to cells which are found in the receptacle (*Hymenium*).

CYTOS (Greek, a cell). *Cytoblast*, a cell nucleus. *Cytogenesis*, the formation of a cell.

D

DÆDALEUS. Said of organs which present sinuous furrows. *Dædalenchyma*, tortuous cells.

DECA. In Greek, *ten*; Latin, *decem*. *Decagynus*, having ten styles. *Decandrous*, having ten stamens.

DECIDUOUS. Falling. Organs which detach themselves after fulfilling their functions.

DECIMETRE. See *CENTIMETRE*.

DECLINATE. Curving or bending downwards, whether the natural direction, or in virtue of weakness.

DECOLORATE. Having lost its colour.

DECOMPOUND. An organ deeply divided, the divisions themselves being divided.

DEFCORTICATED. Deprived of the bark.

DECRESCENT. When the form of the organs

decreases gradually from the base to the summit.

DECUMBENT. Lying flat by its own weight.

DECURRENT. Parts of certain leaves adherent to the stem beneath the point of their insertion; such decurrent stems are sometimes called winged. *Ex.* *Genista sagittalis*, &c.

DECUSSATE. Opposite leaves; the pairs crossing at right angles.

DEFINITE = *Terminal* = *Centrifugal*. Terminating in a single flower. When stamens are under twenty they are said to be definite.

DEFLEXED. Curved downwards.

DEFOLIATION. Fall of the leaves.

DEGENERATION. Transition from the normal to another state, as when the leaves

DEH

DIV

become petaloid, or the petals foliaceous. In horticulture it signifies the return of a plant changed by cultivation to its original state.

DEHISCENCE. The manner in which fruits open at maturity in order to liberate the seeds; also in the case of the opening of the lobes of the anther.

DELTOID. In form of the Greek letter Δ. This epithet is sometimes applied to leaves, and other organs.

DENDRON (*Gr. a tree*), **DANDROIDAL.** Like a tree.

DENTATE. Toothed. The short and triangular divisions, the result of incisions existing at the margin of leaves. These incisions, or dentate parts, are caused by a failure of parenchyma. The term is likewise applied to the free triangular extremities of the divisions forming a gamosepalous calyx, and a gamopetalous corolla.

DENTICULATE. Very finely toothed.

DEPRESSED. Flattened from above downwards.

DERMA. In compound words from the Greek, *bark*; Latin, *cortex*.

DESMOS. Greek, *fascicle*.

DETERMINATE. Definite, centrifugal (inflorescence).

DEXTORSE. Directing to the right.

DI. Greek, *two*; Latin, *bi*.

DIACHLENUM = *Cremocarp*. Fruit composed of two achenia, as in the *Umbelliferae* and *Galium*.

DIADELPHOUS. Stamens united by their filaments into two bundles. **DIADELPHIA**, in the Linnæan system, plants characterised by diadelphous stamens.

DIALYCARPOUS. Of which the carpels are not united to themselves.

DIALYPETALOUS = *Polypetalous*. When the petals of a corolla are not united.

DIALYSEPALOUS. When the sepals of the calyx are not united.

DIANDRIA. Having ten stamens.

DIAPHANOUS. Pellucid.

DIAPHRAGMA. Transversal closing divisions of a cavity. The fistulous stems of the grasses, and some of the *Umbelliferae* present diaphragma.

DICEPHALOUS. Having two heads.

DICHLAMYDEOUS. Having two envelopes, viz., calyx and corolla.

DICHTOMOUS. Branching in twos.

DICLINOUS. Of which the flowers are unisexual, as the plant may be either monoëcious or diëcious.

DICOTYLEDONOUS. An embryo with two cotyledons.

DIDYMOUS. The union of two organs. A

didymous fruit is composed of two carpels united laterally by their suture. *Ex.* The fruit in the *Galium*. A didymous anther is the result of two lobes united by a very short connective; such is the anther in the genus *Euphorbia*.

DIDYNAMOUS. When in a flower there are two stamens, two of which are shorter than the two others, as in the *Scrophulariaceæ*. *Didynamia*, the class of plants thus characterized by Linnæus.

DIGITATE. A compound leaf, having several leaflets arranged almost like a fan, as in the *Lupinus*.

DIGYNOUS. Having two styles.

DIMEROUS. Composed of two pieces.

DIMEDIATE. Divided or split into parts, as the stamens of the *Salix rubra*.

DIECIOUS. A unisexual plant, in which the male and female flowers are produced by two distinct individuals. *Ex.* The *Salix*.

DIPETALOUS. Two petals. *Diphyllous*, two leaves.

DIPLOPERISTOMA (*Capsula*). Urn of the mosses, with a double peristome.

DIPLOSTEMONOUS. When the stamens double in number the petals.

DISCOID, DISCIFORM. Which has the form of a disc, or flattened sphere.

DISCS. The dottings on coniferous wood.

DISEPALOUS. Having two sepals. *Ex.* *Papaver*, *Chelidonium*, *Fumaria*.

DISKS. In the *Compositæ*, the part formed by the uniting of the tubular florets, of which the summits constitute a plain circular surface, encircled as if by a corona, by the ligulate flowers which occupy the circumference of the capitulum.

DISPERMOUS. Having two seeds.

DISSECTED. Divided into a number of narrow stripes.

DISSEPIMENT. Division in the ovary. True when formed by the edges of carpels, False dissepiments (*dissepimenta spuria*) are seen in the siliqua of the *Cruciferae*.

DISSILIENT. A fruit which bursts with elasticity, and throws its seeds to a distance.

DISTICHOUS. Leaves on opposite sides of a stem or axis, and in two rows.

DIURNAL. Flowers which expand and shoot in the same day.

DIVARICATE. Branches which issue from the stem at right angles.

DIVERGENT. Said of objects that tend in a different direction from one another. *Ex.* Divergent styles.

DIVERSIFORM. Organs of the same nature but of different forms.

DIVISION. A segment, as in the divisions of a leaf; for a leaf is a simple organ, however often cut or divided. The expression—

DOD

the divisions of the calyx, corolla, &c., is different, for the calyx is not a simple organ, it is a reunion of small leaves (SEPALs).

DODECA. In Greek, *twelve*; Latin, *duodecim*. Dodecandria, dodecagynia, twelve stamens and twelve styles.

DORSAL. Belonging to the back. The dorsal part of the carpel corresponds to the external face of the main vein of the carpellary leaf.

DOUBLE. Flowers are said to be so when the stamens become more or less petaloid, as in the Rosaceæ and Malvaceæ; sometimes it results from the transformation of stamens and carpels, as in the Ranunculaceæ, &c. The term double is wrongly applied to certain of the Compositæ, as the Dahlia for example, because the change

ENT

caused by culture is not from the addition of new petals, or from the transformation of different organs into petals, but simply from the amplification of the tubulous corollas or florets, which increase themselves, and often assume new colours.

DRUPE. Fruit composed of a single monospermous carpel, and of which the carpellary leaf becomes fleshy at its external division, and ligneous in its internal division. *Ex.* Peach, cherry. *Drupeols*, small drupes. The fruit of the raspberry is formed by the aggregation of drupeols.

DURAMEN. The heart wood of Dicotyledonous trees.

DYNAMIS. In Greek, *power*. Linnæus characterizes the length of two stamens relatively to others in the same flower thus. Didynamous, having two longer stamens.

E

E, or **Ex.** Privative preposition, corresponds to the Alpha of the Greeks. *Ex.* Ebracteate, deprived of bracts; ecarinate, without keel; eglandulate, without glands; exaristate, without an awn.

EDENTATE. Without teeth.

EFFLORESCENCE. Sort of a glaucous powder. The result of the secretion of a resinous or gummy matter, dried on the surface of the organ which produces it. The fruit of the Vine, and the leaves of the Atriplex are examples.

ELASTICITY. The property of certain organs to distend and contract suddenly. *Ex.* The stamens in the genus *Urtica*.

ELLIPTICAL. Having the form of an ellipse.

EMARGINATE. Superficial and terminal portion taken out.

EMBRYO. The rudimentary plant contained in the seed. Embryo sac, the cellular membrane in which the embryo is enveloped.

EMBRYOTEGIUM. That part of the covering which detaches itself regularly from certain seeds at the commencement of germination, on yielding to the radicle, to which it gives free passage.

EMERGED. A plant is said to be emerged which grows under water, but has its summit only exposed. *Ex.* Potomageton, *Myriophyllum*, &c.

EMPHYSEMATOUS. Said of a compressible tissue, of which the areolæ are distended and ruptured by the air or gases.

ENDOCARP. The inner layer of the pericarp; the membrane enveloping immediately the seed. In Drupes, the Peach, and Cherry, for example, the endocarp is no other thing than the kernel, which is

not, as is commonly supposed, a dependent of the sarcocarp.

ENDOGEN = Monocotyledon. A plant of which the stem is composed of ligneous fascicles isolated in a cylinder of cellular tissue, and not arranged by concentric layers.

ENDOS. Greek, *within*; Latin, *intra*.

ENDOSMOSE. The physical phenomenon which consists in the tendency which two liquids of unequal density have to mingle and become of an equal density. This action operates not only between two liquids in contact, but between two liquids separated by a thin organic membrane, whatever may be the density of the liquids.

ENDOSPERM = Albumen = Perisperm.

ENDORHIZAL. Rootlets from a radicle in a sheath through which its makes way during the period of germination.

ENDOSTOME. The opening of the membrane of the ovule placed under the testa or internal membrane.

ENDOPHLEUM. The inner bark.

ENDOPLEURA. Internal membrane of the episperm or covering of the seed.

ENDOTHECUM. The inner layer of the anther.

ENERVIS. Wanting apparent veins.

ENNEA. Greek, *nine*; Latin, *novem*, *enneandria*, *ennegynia*. Plants with nine stamens and nine styles.

ENODIS. Having no nodes.

ENSIFORM. In the form of a sword. *Ex.* The leaves of the Iris and the Gladiolus.

ENTIRE (Integer). Said of a simple organ, a leaf, for instance, of which the margin or border "is not divided" or cut. The superlative integerrimus is used sometimes

ENV

FIB

- to indicate that there exists not the slightest denticulation.
- ENVELOPES** (Floral). The verticelles surrounding the stamens and ovules (calyx and corolla).
- EPI.** Greek, *upon*; Latin, *supra*.
- EPICARP.** The external layer of the pericarp.
- EPIDERMIS.** The external layer of the bark which covers the surface of a plant.
- EPIGEAL.** Above the ground (Cotyledons).
- EPIGONE** (Epigonium). The membranous (cellular) layer which covers the sporangia of the Mosses and Hepaticæ.
- EPIGYNOUS.** Said of organs of the flower inserted into the summit of a tube adherent to the ovary in such a manner that they appear to spring from the summit of the ovary itself.
- EPIPETALOUS.** Seemingly instated upon a petal. The insertion of the stamens upon the petals is so termed.
- EPIPHRAGM.** In the genus *Polytrichum* (Mosses) this name is given to the membrane which closes the urn after the fall of its operculum.
- EPIHYLLOUS.** Organs which seem inserted upon a leaf or upon a bract.
- EPIPHYTE.** False parasites; plants, such as the Orchids, which in seeking support from trees, derive no nourishment from them.
- EPISPERM.** Envelope composed of two or more membranes which surround the embryo.
- EPISPORE.** The covering of spores in the Uredinæ.
- EQUIANT.** Leaves folded longitudinally. The inferior or exterior embracing the superior or interior. The Iris, for example.
- ERECT.** Rising from the base (ovary).
- EROSTRATE.** Wanting the beak.
- EROSE.** Irregularly toothed leaf, as if bitten.
- ERYTHROS.** Greek, *red*; Latin, *rubrum*.
- ERRUMPENT.** Prominently bursting.
- ETÆRIO.** The group of drupes on the fruit of the Raspberry.
- ETIOLATION.** Blanching, by screening from the light.

- EXO.** Greek, *without*; Latin, *extra*.
- EXALATE.** Deprived of wings = apterous.
- EXALBUMINOUS.** Seed deprived of the perisperm, or albumen.
- EXCENTRIC.** The embryo is said to be excentric when it is situated without the axis of the perisperm.
- EXCIPULUM.** In the Lichens the receptacle which contains the fructification, or thalamium. The excipulum and thalamium constitute the apothecium; excipulate, excipular.
- EXCRETION.** A physiological function, by virtue of which certain liquids, ordinarily of a resinous or gummy nature, and susceptible of concreting, are excreted by organs of a glandular nature, or by accidental fissures.
- EXCURRENT.** Said of the trunk of a tree continued from the base to the extremity of the stem, as in the Pines.
- EXEMBRYONATE.** Containing no embryo.
- EXOGEN** = Dicotyledon. The stem composed of fascicles arranged in concentric layers.
- EXOSMOSE.** The reverse of endosmose, the passing out of a liquid through a membrane.
- EXOSTOME.** The free opening of the external membrane of the ovule.
- EXOTHECIUM.** The external layer of the anther.
- EXsertED.** Stamens are so which extend beyond a gamopetalous corolla.
- EXTIPULATE.** Deprived of stipules.
- EXTINE.** The outer covering of the pollen grain.
- EXTRA-AXILLARY.** Said of a bud which appears situated without or above the axil of the leaf. Extra-foliateous, placed without the leaves; extraneous = exotic, not belonging to Europe.
- EXTRORSE.** Said of anthers which dehisce exteriorly to the flower. *Ex.* Anemone, Clematis.
- EXUNGUICULATE.** A petal which is not attenuated into a claw.

F

- FALCATE.** Falciform, curved like a sickle.
- FAMILY** = Ordo. Natural group of plants formed by the agglomeration of a certain number of genera, and even under circumstances of a single genus.
- FARINACEOUS** = Amylaceous. Containing or having the appearance of starch.
- FASCIATION** (*Fascia*, a band). The flattened ribboned-like form substituted for the cylindrical or prismatic of normal stems.
- FASCICULATED.** Many or several organs

- united into a bundle. Fascicule, a kind of indefinite inflorescence.
- FAVEOLATE.** Presenting little hollows.
- FENESTRA.** The cicatrix which the seed presents after its fall, on the level of the hilum, at maturity. Fenestrate, presenting an opening which has been compared to a window.
- FECULA.** One of the names given to farinaceous or starchy substances.
- FIBROUS.** Composed of fibrous tissue.

FID

Fibro-cellular, spiral cellular tissue. Fibro-vascular, tissue formed by the union of fibres, or fibres and vessels.

FID—**FIDUS**. Terminations indicating that the organ qualified is cut down or cleft. Applied chiefly to adjectives intended to characterize leaves divided down to the middle nerve or vein.

FILAMENT. That part of the stamen which supports the anther. Filamentous, having filaments.

FILICES. Ferns.

FILIFORM. Having the form of a thread. Filipendulus, seemingly suspended to a thread.

FIMBRICE. Fringe. Fimbriated, fringed, or irregularly lacinated at the margin.

FISSURE. A straight cleft or slit.

FISTULOUS. Hollow in the interior, as the internodes of the Grasses, and in some of the Umbelliferæ.

FLABELLIFORM. In the form of a fan. *Ex.* Some of the Palms.

FLAGELLUM. Creeping stem, producing rooting buds, as in the Strawberry.

FLAVEDO. Orange-colour; flavescens, flavidus, of a pale orange or yellow.

FLEXUOUS or **FLEXUOSE**. Bent under the weight of the terminal part, but capable of assuming the natural direction. Said also of an organ which presents alternate curvatures in opposite directions, or zig-zag.

FLOCCUS. A woolly filament. Floccose, covered with woolly filaments or tufts.

FLORA. The whole of the vegetable species of a country. A work in which the vegetation of a country is described and classed.

FLORAL. Relating to the flower. Floriferus, bearing flowers.

FLORUS replaces *flos* in compound words. *Ex.* Multiflorus, bearing a number of flowers; pauciflorus, bearing few flowers.

FLOS. Flower. (Greek, *anthos*.)

FLOSCULOUS. A capitulum of which the florets are tubulous, and with five equal lobes.

FLUITANS. Said of plants which grow in running waters. *Ex.* Ranunculus fluitans. Also of those which float freely, as,

GEM

for instance, the species of the genus Lemna. These latter grow in stagnant waters.

FLUVIAL—**Fluviatile**. Plants which grow on the banks of rivers or streams.

FOLIACEOUS. Having the colour and consistence of leaves.

FOLIATION. The development of leaves. Foliate, furnished with leaves.

FOLIOLUM. A small leaf—applied to the articulated divisions of a compound leaf. Foliated, furnished with small leaves.

FOLLICLE. Fruits formed by a single free carpel, polyspermous, and of membranous consistence. The follicle differs from the legume by its dehiscence, which takes place only by the ventral suture.

FOOT. French, *pied* = 13.110 English inches.

FORAMEN. Small opening or pore, as in the ovule.

FORM—**Formis**. Termination signifying in form of, as squamiform, in the form of a scale.

FORNICATE. Arched in the form of a vault. *Ex.* The superior petal of the Aconitum.

FOVEA. A small hollow. Foveate—foveolate, having small hollows or depressions.

FOVILLA. The granular liquid contained in the pollen grain.

FRONDS. The leaves of the Ferns. These leaves bear the fructification on their under or inferior side. Also applied to the foliaceous thallus of certain Lichens and Hepaticæ. Frondose, said of some Cryptogamic plants (Lichens e.g.) which present foliaceous expansions.

FRUCTIFICATION. The parts composing the fruit.

FRUIT. In botany this term is applied to the fecundated ovary, when it has reached maturity.

FUGACIOUS. Disappearing rapidly and leaving few traces.

FUNICULUS = *Podosperm*. Umbilical cord or support of the ovule. Funiculate, furnished with a long funiculus.

FUNNEL-SHAPED. See *Indibuliform*.

FURCATE. Forked. Divided into two branches; bi-furcate.

FURFURACEOUS. Scurfy or scaly.

FUSIFORM. In the form of a spindle.

G

GALA. (Greek, *milk*; Latin, *lacteus*.) (Compound words).

GAMOS. (Greek, *united with*; Latin, *adherent*.) Gamopetalous = Monopetalous, a flower of which the petals are united. Gamosepalous = Monosepalous, the sepals of the calyx united.

GELATINOUS. Having the consistence of gelatine with water. Some cellular Cryptogames present this consistence.

GERMINATE. Two by two, or in pairs. Gemminiflorate, bearing twin flowers.

GEMMA. Leaf bud. Gemmation, development of leaf buds; gemmiferous, bear-

GEM

ing leaf buds; gemmiparous, reproduction of leaf buds.

GEMMULE = Plumule. (Lindl.) The first bud of the young plant or embryo.

GENERA PLANTARUM. Title of works in which is found the methodic order and characters of genera. (Endlicher.)

GENERIC. Appertaining to a genus.

GENICULATE. Bent like the knee. *Ex.* Alopecurus Geniculatus.

GIBBOSITY. A swelling.

GLABROUS. Smooth, without hairs.

GLANDS. Organs of a cellular nature secreting particular liquids. Certain glands occur on the epidermis. In the nettle the glandular part is situated at the base of the hairs, and the produce of the secretion fills the entire cavity of the hair.

GLARS. The fruit of the oak (acorn).

GLAUCIOUS. Covered with a bluish green pulverulent matter. *Ex.* The grape, chenopodium, &c.

GLOBATE = Globosis. Globulete, of a globular form.

GLOBUS or **GLOBULE.** The fructiferous capsule of plants of the family of the Hepaticæ, the capsuliform involucre of pilularia globulifera—the antheridia of the characæ.

GLOMERULE (Glomerulus). Sort of inflorescence of a definite type. It is a cyme on which all the axes being short, the flowers approach each other like a capitulum.

GLOSSOLOGY = Terminology. That part of the science which has for its object the study of the signification of terms employed to describe the organs and their conditions.

GLUMACEOUS. Of the nature of glumes.

GLUME—**GLUMA.** In the grasses the sterile bracts which are situated at the base of spikelet. *Glumelle*—the palea or fertile glume. *Glumellule*—for the follicles which represent the corolla.

GONIDIUM (zoöis, dust). The reproductive bodies or cells which develop themselves in the thallus of the Lichens and the frond of the Hepaticæ.

HER

GONUS. Greek, Angle. Tetragonous = 4 angled.

GRACILIS. Slender.

GRAIN. A fecundated ovule (containing an embryo) having reached maturity. *The Cereals.* Granum pillinis; the pollen grain.

GRAMINEÆ. The Grasses.

GRAMME. French weight = 15.4440 English inches. Decigramme = tenth part of the gramme. Centigramme, the 100th part. Millegramme, the 1000th part. Décagramme = 10 grammes. Hectogramme, 100 grammes. Centgrammes, Kilogramme = 1000 grammes, (a pound equivalent to 500 grammes or a demi-kilogramme).

GRANULATED. Formed of grains.

GRAVEOLENS. Exhaling a strong disagreeable odour. *Ex.* Ruta graveolens.

GRUMOUS. A substance reduced to small unequal and irregularly rounded masses.

GYMNOS. Greek—naked. Latin—nudus. Deprived of envelopes. Angios is the opposite of Gymnos.

GYMNOSPERMS. Dicotyledons of which the seed is naked, i.e. not contained in a pericarp, but situated at the base of a carpellary leaf. *Ex.* Cycadacæ and Coniferæ.

GYMNOSTOMOUS. In Mosses the capsules deprived of the teeth. *Ex.* Sphagnum.

GYNANDROUS. When the stamens and the style are united into a single body, as in the Orchidacæ.

GYMNOBASE. The fleshy organ to which the ovary appears attached in the Labratæ and Boraginæ.

GYNOCYDIUM. Swelling at the base of the seta of the capsule in the Mosses.

GYMNOPHORE. An organ which results from the prolongation of one or several internodes which constitute the axis of the flower and raises the ovary above the insertion of the calyx.

GYMNOSTEMIUM. In the Orchids the column arising from the union of the stamens with the style.

GYROMES = Gyrate. The elastic ring of the sporangia of Ferns.

H

HABIT. Bearing of a plant i.e. general aspect which it presents. *Habitat* = station.

HASTATE. In the form of a halbert. The leaves are hastate in the Rumex Acetosa.

HAULM. Dead stems of herbs, as of the potato.

HAUSTORIUM. A sucker. In the Cuscuta it serves to absorb the sap from the plant to which it is attached. The extremities

of the parasitic roots of the Orobanchæ and Lathræa are true suckers.

HEMI. Greek, half. Latin, semi.

HEPATICÆ. The Liverworts.

HEPTA. Greek, seven. Latin, septem. Heptendria, having seven stamens.

HERB. A plant with an annual stem, opposed to a ligneous plant. *Herbaceous.* With the consistence of soft leaves.

HER

- HERBARIUM.** Collection of dried plants prepared and classified for study.
- HERMAPHRODITE.** A flower provided with stamens and pistil.
- HESPERIDIUM.** Name given to the fruit of the *Aurantiaceæ* (*Orange*).
- HETEROGAMOUS.** Sign of the Androgynous and polygamous.
- HETEROPHYLLUS.** Presenting two different forms of leaves. *Ex.* *Ranunculus Aquatilis*, of which the submerged leaves are divided into filiform segments, and the floating leaves are suborbicular crenulated.
- HETEROS.** Greek, different. Latin, dissimilis.
- HETEROTROPAL.** (See *Ovule*.)
- HEXA.** Greek, six. Latin, sex. Hexapetalous, having six petals. Hexandria, six stamens.
- HIERNAL.** Growing during winter. The flowering of the *Helleborus Niger* is hibernal.
- HILUM = UMBILICUS.** In the ovule the hilum is the point where the funicle becomes adherent to the testa. After maturation the name is given to the scar which remains at the point of the seed whence the funicle has been detached. In grains of starch the small surface by which the successive deposits which constitute it have commenced.
- HIRSUTE.** Covered with long and round hairs.
- HISPID.** Having strong rough hairs.
- HISTOLOGY.** The study of tissues with the microscope.
- HOMOS**—Similar; Latin, similis; opposed to *Heteros*; Homogamous. In the *Compositæ* a capitulum having all the florets analogous relatively to the sex, as they may be all hermaphrodite or of a single sex.
- HOMOTROPAL = Orthotropal.**
- HOODED.** See *Cucullate*.
- HOROLOGIIUM FLORÆ.** Horological flowers.

INC

- Certain flowers opening and closing at different hours of the day. *Linnaeus* thus formed his floral clock.
- HORTENSIS.** Growing spontaneously in gardens.
- HORTUS.** Garden. *Hortus siccus*, another name for herbarium.
- HUMIFUSE.** Spreading upon the soil.
- HYALINE.** Membranous organs transparent like glass, the glumellules (internal verticelle of the perianth of grapes) are hyaline.
- HYBRID.** Plant produced by the fecundation of one species by another species of the same genus. **HYBRIDATION.** The formation of hybrids.
- HYEMALIS.** Which vegetates and flourishes during winter.
- HYGROMETRIC.** The action of dry or humid air upon the tissues. Certain organs have the property of absorbing easily the humid air, and unroll or expand under that influence.
- HYMEN.** In compound Greek words signifies membrane (*membrana*).
- HYMENIUM** (Fries, &c.) The fructifying membrane of the *Fungi*, particularly the *Agaricus*, the *Boletus*, and allied genera. *M. Leveillé* defines the Hymenium as the membranous part upon which the organs of fructification rest in Mushrooms.
- HYPANTHODIUM.** The kind of inflorescence which the Fig presents (*the receptacle*).
- HYPERTROPHIAL.** Excess of development, an abnormal volume acquired by an organ. The Cauliflower is the result of an hypertrophy of the axis of the common Cabbage.
- HYPO.** Greek, under; Latin, sub.
- HYPOGÆUS.** Under the surface of the soil. The Truffle is an hypogean Mushroom.
- HYPOGYNOUS.** Inserted upon the receptacle, on a level with the insertion of the ovary itself.
- HYPOPHYLLUS.** Inserted under the leaf.
- HYSTERANTHÆÆ.** Plants of which the leaves develop themselves after the flowers.

I

- ICONOGRAPHY** (Vegetable). The art of representing plants and the details of their organization through the means of drawing and engraving. *Icones*, figures drawn, plates engraved.
- ICOS.** In compound Greek words signifies twenty; Latin, viginti. *Icosandrous*, with twenty stamens.
- IMBRICATE** or **IMBRICATED.** Overlying each other in a manner similar to the arrangement of tiles on a roof.
- IMPAR.** Unequal. *Imparipinnate*, compound

- pinnate*, presenting a terminal single leaflet.
- IMPELLUCID.** Not pellucid; imperforate, not perforated.
- INÆQUALIS.** In compound Greek words, derived from the Greek *anisos*, objects unequal in size.
- INCISED.** Cut down, so as not to reach beyond the middle.
- INCLUDED.** Contained within. When the cotyledons are folded longitudinally (cotyl. conduplicate), in such a manner as

INC

to embrace the radicle, this radicle is said to be included. In the case of gamopetalous corollas, when the stamens and style do not project beyond the tube, they are said to be included. These organs projecting are termed *exserted*.

INCRUSTATIONS. Said of the earthy deposits by the water upon certain submerged plants, *Chara* for example. These earthy deposits form around the stems a crustaceous and fragile envelope.

INCUMBENT. The radicle of certain dicotyledonous embryos, folded on the back of one of the cotyledons; if bent upon the commissure of the two cotyledons, is said to be accumbent.

INCURVED. Bent in the superior or upper part.

INDEFINITE = Indeterminate = Centripetal. An inflorescence that commences at the circumference and proceeds towards the centre; stamens exceeding twenty are also said to be indefinite.

INDEHISCENT = not opening. Said of fruits of which the pericarp does not open spontaneously.

INDIGENOUS. Opposite to exotic, and so applied to the spontaneous vegetation of a country.

INDUPLICATE. In æstivation, when the margins of the petals or sepals enter within, instead of being simply contiguous.

INDUSIUM. In the Ferns the covering formed by a layer of epidermis, which in the majority of the genera envelop the sori or groups of sporangia. The indusium is wanting in the genus *Polypodium*. It is reniform suborbicular in the *Nephrodium*; continued with the margin of the divisions of the leaf in the *Pteris*.

INEQUILATERAL. Said of a leaf of which the two longitudinal halves are larger or of different form.

INERMIS. Having no spines, prickles, or any sharp points.

INFERIOR. When the ovary appears to be below the calyx.

INFLATUS. Swelled, dilated by the air, as in the case of the calyx of *Silene inflata*.

INFLORESCENCE. The general disposition and arrangement of the flowers.

INFRA. Beneath or within. Greek, *hypo*.

INFUNDIBULIFORM. In form of a funnel, as in a gamopetalous corolla of which the

JUX

tube is long and straight, and gradually takes the form of a cone reversed. *Ex.* Marvel of Peru.

INNATE. Inserted upon or seeming to spring from.

INNOVATIONS. The buds which are produced each year by a great number of species of Mosses.

INTER. Between. Interfoliar, between two opposite leaves; internodes, the spaces on the stem between the nodes.

INTERPETIOLAR. Stipules are said to be so which come between opposite leaves.

INTERRUPTEDLY PINNATE. When small pinnæ are found placed between larger ones.

INTINE. The inner layer of the pollen grain.

INTRA. Greek, *endos*, within. Intrafoliaceous, between the leaves.

INTRORSE. When the lobes of the anther open next the pistil, and thus the reverse of extrorse.

INVOLUCRE. The bracts situated at the base of the inflorescence in terminal compact heads, called capitula (*Compositæ*).

INVOLUTE. Leaves of which the two longitudinal halves are rolled within in æstivation.

IRREGULAR. Wanting regularity in form, as in the case of leaves or flowers in which there is no symmetry.

ISOCHEINAL = **ISOCHIEIMINAL.** Having the same mean winter temperature. Isocheiminal lines, lines passing through places having the same mean winter temperature. (Humboldt.)

ISOGYNUS. A flower having the carpels and the petals equal in number.

ISOMERIC. Having the same elements in the same proportions, but with different properties.

ISOMEROUS. The parts of the flower equal.

ISOS. Greek, *equal*; Latin, *æqualis*.

ISOSTEMONOUS. The stamens and pistil of a flower equal.

ISOTHERAL (Greek). Having the same mean summer temperature. Isothermal lines, lines drawn through places having the same mean summer temperature. (Humboldt.)

ISOTHERMAL. Having an equal degree of heat. Isothermal lines, lines passing through places of equal mean temperature. (Humboldt.)

J

JUGUM (plur. *Juga*). Longitudinal ribs of the fruit of Umbelliferae—the intervals are named valliculi. Jugum means also a pair of leaflets, or compound pinnate leaves.

JUXTA. Near or against. Juxtaposition—applied near or against, but not adherent.

KEEL. Carina in papilionaceous flowers.

LAB

LABEL = **LOBELIUM**. In orchids, the lower of the three petals of the internal verticille of the printh. It presents frequently very curious forms. This division is in reality superior, but becomes inferior by reason of the twisting of the pedicel which changes the direction of the flower.

LABIATIFLORÆ. A capitulum, of which all the florets are bilabiate.

LABIATE. Gamopetalous corollas divided into two lips; the superior lip is formed by two petals united, and in general bilobate or emarginate; the inferior has three petals united and in general trilobate.

LACINATED. Said of a leaf or petal irregularly cut into segments.

LACTATE—**Lactescens**. Having the aspect and colour of milk or yielding milky juice.

LACUNA. Space between groups of cells. Lacunes are met with frequently in the tissue of aquatic, submerged, or floating plants.

LACUSTRINE. Whose habitat is in lakes or on the border of lakes or ponds.

LEVIGATUS—**Lævis**. Smooth—opposite to asper, rough.

LAMELLÆ. Petaloid appendages at the throat of some corollas, as the *Lychnis* or Gamopetalous corollas, as in a great number of the Boraginæ. Fries gives this name to the gills of the *Agarics*.

LAMINA. The limb or blade of a leaf, petals, &c.

LANA—**Wool**. Lanuginous, woolly.

LANCEOLATE. Oblong straight leaf gradually terminating in a point. *Ex.* *Salix alba*.

LATERAL. Placed upon the side. The lateral nerves of leaves are those which are situated to the right and left of the middle nerve, or they may arise from the base of the leaf.

LATEX. Elaborated liquid contained in the laticiferous vessels.

LATICIFEROUS. The vessels or reservoirs of the latex. They are met with in the *Liber*, during the formation of each of its layers.

LATUS. Wide. *Latiseptæ*, wide folds or divisions in the silicula of the *Cruciferae*.

LEGUME. Fruit of the plants of the *Leguminosæ*. It is a single carpel ordinarily polyspermous, and dehiscing by a ventral and dorsal suture. *Ex.* The *Pea*.

LENTICEL. Small organ or brown rugosity of an oval or elliptical form, presented on the epidermis of some plants.

L

LENTICULAR. Having the form of a double convex lens.

LEPIS. Greek, *a scale*; Latin, *squama*. Lepidote, covered with scales.

LIBER. The part of the bark of dicotyledons composed of fibrous tissue. It is the most internal, and receives likewise the name of endophlæum.

LICHENS. A class of cryptogamous plants.

LIGNINE. The woody matter. Ligneous, of the nature of wood.

LIGULATE. In the form of a small tongue. The exterior florets of the capitata of a number of the *compositæ* are ligulate. *Ex.* *Cicharium Intybus*, the *Dandelion*, *Taraxacum*, *Densleonis*.

LIGULE. The thin transparent membrane at the extremity of the internal face of the petiole of Grasses.

LIB. Flat part of a leaf surmounting the petiole; or of a petal and sepal.

LINE. Measure of length = 12th part of an inch. *Ligne* (Fr.), = 0.091 English inches. Linear, long and very narrow leaf. The limb of the leaf of the greater part of the Grasses, and of the *Cyperaceæ*, is more or less linear.

LITTORAL. Growing on the borders of streams, &c.

LOBE. Deep division of a leaf; applied also to the divisions of an anther separated by a connective. Lobule, small lobe or subdivision of a leaf in a lobate or pinnatifid leaf.

LOCULAR. Appertaining to cavities. Unilocular, bi-triocular, having two or three cavities.

LOCULLICIDAL. Dehiscence of a fruit which takes place by the longitudinal rupture of the dorsal nerve of each carpel.

LOCULUS or **LOCULAMENT**. A cavity, as in the ovary.

LOCUSTA. Synon. of *Spikelet*.

LODICULES. Same as *glumellule*. A scale at the base of the ovary of Grasses.

LOMENTUM and **LOMENTACEOUS**. Said of a fruit composed of a single carpel which divides itself naturally into several false cavities, as, for example, the pod of the *Papilionaceæ*, of the tribe of the *Hedysaræ*.

LUCID. Shining, smooth and wanting hairs.

LUNATE or **LUNULATE**. In form of a crescent or half-moon.

LUTEUS. Yellow.

LYMPH. Aqueous sap.

LYM

MAC

MACRANTHUS. Of which the flowers are large.

MACROS. Greek, *large*; Latin, *magnus-crassus*.

MACULATE. Spotted.

MANUBRIATE. Furnished with a handle. Said of the pollen masses of the Orchidaceæ.

MARESCENT. Said of an organ which after it is withered remains attached to the part from which it has sprung. The leaves of the oak are marescent, the limb of the calyx of the pear and hawthorn. The term is used in contradistinction to caducent.

MARGINATE. Having brims or margins.

MASSES POLLEN. In the Orchidaceæ and Asclepiadaceæ all the pollen grains contained in each of the two lobes of the anther are reunited into a single mass more or less compact.

MATH. A term sometimes used for crop.

MATURATION. The period during which the ovary passes to the state of ripe fruit.

MEDIAN. Occupying the middle part.

MEDULLA. The cellular tissue or pith which occupies the centre of the stem of Dicotyledones. *Medullary Rays*, the expansions of cellular tissue which issue directly or indirectly from the medulla, and are arranged in rays from the centre towards the circumference; the medullary rays separate the fibro-ligneous fascicles longitudinally.

MEIOS. Greek, *less*. Meistemonous, flowers on which the stamens are fewer in number than the petals. (Seldom used.)

MALAS or MALANOS. Greek, *black*; Latin, *niger et ater*.

MEMBRANOUS. Membranaceous.

MEMBRANE. Having the consistence, aspect, and structure of a membrane.

MENISCUS. A lens—convex on the one side and concave on the other, the concavity being less than the convexity.

MERICARP. In the Umbelliferae each of the two actænia formerly united but become free on maturity.

MERITHAL. Sign of internode.

MESOCARP. Sign of sarcocarp; intermediate to the epicarp and endocarp of the fruit.

MESOCHILIUM. The middle part of the labellum of Orchids.

MESOPHLEUM. The cellular middle layer of the bark.

MESOPHYLLUM. That part of the leaf situ-

M

MUC

ated between the lower face and the epidermis of the upper face.

MESOS. Greek, *the middle*. Mesosperm, the middle covering of a seed.

METAMORPHOSIS. Transformation of one organ into another.

METRE = 39.37100 English inches (3 feet 11 lines French).

MICROMETER. Instrument applied to a microscope for measuring objects.

MICROPYLE. The mouth or foramen of the seed. It represents the organic summit of the seed whatever may be its position.

MILLEMETRE. Fr. The 1000th part of the metre = 0.03937 English inches.

MONADELPHOUS. Organs of the same nature united into a single bundle. The stamens are monadelphous in the Malvaceæ.

MONANDROUS. Having one stamen.

MONILIFORM. In the form of a necklace. Composed of globular cells on the same line.

MONOCARPOUS. Bearing fruit but once and drying after fructification.

MONOCLINOUS. Stamens and pistils situated on the same receptacle, or in the same flower. Syn. with hermaphrodite.

MONOCOTYLEDONOUS. Having only one cotyledon.

MONOGYNOUS. Having only one style.

MONÆCIOUS. Stamens and pistils on distinct flowers but growing on the same individual.

MONOPETALOUS = Gamopetalous.

MONOPHYLLOUS. Single leaf.

MONOS. } Gr. *one*; Latin, *unus*; compound words, *uni*.

MONOSOPALOUS = Gamosepalous.

MONOSPERMOUS. With a single seed.

MONOSTACHYA. Having a single spike.

MONSTROSITY. Abnormal development of a flower.

MORPHE. Shape or form of. *Morphology*. The science which has for its object the study of the forms that each organ or parts of an organ assumes.

MUCILAGINOUS. Said of a viscons liquid. The root, the stem, the leaves, and the flowers of the Malvaceæ are mucilaginous, and the majority of bulbs contain a mucilaginous liquid.

MUCRO. A round stiff point terminating abruptly; an extremity more or less large.

Mucronate. Terminating with a mucro. A leaf, a capsule, can be mucronate.

MUCUS. A viscid fluid in certain marine Algae.

MUL

MULTI—Many; Greek, *poly*—Multicostate. said of a fruit presenting a great number of ribs; the Umbelliferae, for example. **Multifid**, a leaf of which the divisions are numerous and extend down to the middle of its blade. **Multipartite**, when the divisions are deeper. **Multilobate**, presenting a great number of lobes. **Multilocular**, with numerous loculements.

MULTIPLE. A compound organ is said to be multiple when it results from the reunion of many or several free organs; thus a multiple ovary or fruit is composed of many free carpels.

OPP

MURICATE. Covered with strong and short points. *Ex.* The ripe carpels of *Ranunculus muricatus*.

MURIFORM. In shape of a wall.

MUSCINEAE. The class of Mosses.

MUSCOLOGY. The study of Mosses. (*Bryology*.)

MYCELIUM. Filaments, simple or more or less complicated, resulting from the vegetation of the spores and serving as supports and roots to the Fungi.

MYCOLOGY. That part of botany which treats of the situation and classification of the family of the Fungi.

N

NAKED. Deprived of envelopes. A naked seed is that which is not contained in a pericarp. *Ex.* The Gymnosperms.

NATANS. Those leaves of plants which grow in stagnant waters, their lower or inferior face being in contact with the water.

NAVICULAR. Hollowed in form of a boat. *Ex.* The fruit of *Valerianella Carinata*.

NECTAR. The sugary liquid secreted by nectariferous surfaces. *Nectariferous*, secreting nectar. Certain petaloid surfaces secrete liquids of this nature, as in the *Fritillaria imperialis*; the Crown Imperial.

NEMNOSAS. Growing in the forests and woods.

NERVATION. Arrangement of the nerves in a leaf.

NITIDUS. Having a smooth and shining surface.

NIVALIS. Said of plants which flourish towards the end of winter, when the snow is scarcely dissipated. *Ex.* *Galanthus Nivalis*, the Snow Drop.

NODE. Swelled articulation corresponding to the point of insertion of a leaf. *Nodose*, presenting nodes or swelled articulations.

NOTORHIZEÆ (*Cruciferae*). Of which the seed presents an embryo to a dorsal radicle.

NUDICAULIS. Naked stemmed.

NUCULANIUM. Fruits composed of a whorl of drupes united amongst themselves, and resulting from an adherent ovary, such as the fruit of the Medlar.

NUCAMENTACEOUS. Dry monospermal fruits, such as those of certain *Cruciferae*.

NUTRITION. The functions which support vegetable life, and particularly absorption and assimilation.

O

OB. Indicates that the base or larger extremity of the object designed is directed in front or above; or, in other words, that this form is reversed. Thus, *cordate* signifies shaped like a heart, the point being above and the broad extremity below; *obcordate*, the point below and the broad part above. *Ex.* The leaflets of *Oxalis Acetosella*; *obconic*, in the form of a cone reversed; *oblong*, from three to four times as long as broad.

OBSELETE. Scarcely apparent.

OBTUSE. Having no point, but roundly terminated.

OBVOLUTE. When parts enrol themselves one upon the other, as in leaves of certain species in the bud.

OCTUEA. A spatterdash. Applied to the complete sheaths which exist at the base of the petiole of certain alternate leaves, and particularly in plants of the family of *Polygonaceae*.

OCTO = eight. *Octandrous* = eight stamens. *Octogynous* = eight styles.

OLKERACOUS. Esculent.

OLIGO = few. *Oligandrous*, when stamens are under twenty.

OMPHALOD. The central point of the hilum; it is the point which corresponds to the passage of the nourishing vessels of the seed.

OOPHORIUM. The capsules with three or four valves which contain three or four subglobular bodies larger than the spores contained in the sporangias, as in *Ly-copodiaceae*.

OPERCULUM. The lid which detaches itself horizontally and circularly from the summit of certain capsules. The capsule or urn of the Mosses opens thus. *Operculate*, furnished with an operculum.

OPPOSITE. Leaves situated in face of each other on the same stem and level. In a flower, the organs of a different verticille

ORB

situated the one in front of the other, are said to be opposite; as is said, the stamens are opposite to the sepals.

ORBICULAR. A leaf which is almost circular.

ORBITICULUS. In the group of *Nidulariæ* (Fungi), a lenticular body at the base of the pedium (which is like a cup) like an egg in a nest.

ORGANOGENY = *Phytogenesis*. The development of an organ during all its successive stages.

ORGANOGRAPHY. All the studies of the organs of plants in the different points of view of their mode of development.

ORTHOS. Greek, *right*; Latin, *rectus*. Orthospermous, of which the seed is right or straight in opposition to campylospermous.

ORTHOTROPAL. The radicle of the embryo directed towards the hilum.

OVAL. A leaf is said to be oval when it assumes the form of an ellipse, of which the

PAR

greatest transversal diameter is beneath the middle.

OVARY. The covering of the ovules.

OVATE, or OVOID. In shape of an egg. Certain seeds and fruits are ovoid.

OVULE. In *Phanerogamous* plants the seed before and during the period of the expansion of the flower. When an ovule is bent on itself, the hilum being in the centre = *amphitropal*. When the ovule is inverted, the micropyle and hilum being close to each other, and the chalazas opposite = *anatropous* or *anatropal*. When the ovule is curved, the micropyle, chalazas, and hilum being near each other = *campulitropal* or *campylitropal*. When the ovule has the foramen opposite to the hilum, also when the embryo has the radicle next to the hilum = *orthotropal* or *orthotropous*.

OVULATE. An ovary containing one or more ovules. *Ex.* *Uniovulate*, *biovulate*, &c.

P

PAGINA. Flat surface. *Pagina superior*, the upper surface of a leaf.

PALÆOPHYTOLOGY. The study of fossil plants.

PALATE. The part which projects of the inferior lip of the *Linaria*, *Antirrhinum*, and closes the entry of the tube of the corolla.

PALEA. The small scarios laminæ which in certain genera of the *Compositæ* occur on the receptacle and separates the florets; also applied to the parts of the perianth of Grasses within the glume.

PALEACEOUS. Covered with scarios laminæ; chaffy.

PALMATIFID. A leaf divided into lobes, like the fingers of the hand or the rays of a fan, the segments reaching the middle of the blade.

PALMATIPARTITE. The leaves are the same as when *palmatifid*, with this difference that the segments reach nearly to the base of the blade.

PALMATISECT. The leaf differs from that which is *palmatifid* in so much as in the former case the divisions divide the blade in all its extent. *Palmate* is the generic term.

PALUDOSE = *Palustris*. Growing in marshy places.

PANICLE. An indefinite inflorescence, distinguished in the Grasses by the long peduncular racemose spikelets. *Paniculate*, constituting a panicle.

PAPILIONACEOUS. The different parts of the corolla in the *Papilionaceæ*. The superior petal is named the standard or vexillum,

the two lateral the alæ, and the two approximated inferior the carina or keel.

PAPILLA. A little conical rugosity, covering certain surfaces. The stigmatic portion of the style in the *Compositæ* is covered with papillæ.

PAPILLOSE. Covered with papillæ.

PAPPUS. A bundle of silky or parallel hairs, more or less divergent, which terminate certain fruits and seeds. In the *Valerianaceæ* and *Compositæ* the fruits are generally terminated by pappus. In certain species of the *Valerianaceæ* the accrescent calyx constitutes a pappus of which the hairs are isolated.

PAPPOSE. Furnished with pappus.

PAPYRACEOUS. Having the consistence of paper.

PARAPHYSES. Articulated filaments which in the Mosses are intermingled with the antheridæ and with the female flowers. Their number is irregular. In some species they are wanting. In the Lichens there exist also paraphyses, each constituted like those of the Mosses. Likewise the greater portion of the organs of *Cryptogamous* plants. The paraphyses must be studied by means of instruments of considerable magnifying power. In order to discern well those of Lichens, a thin vertical slice of an apothecium must be cut and placed between two pieces of glass in the field of the microscope.

PARASITES. Plants which vegetate upon other living plants, and from which they assimilate nourishment. *Ex.* *Viscum album*.

PAR

PER

False parasites (Epiphytes) differ in so much as they do not absorb the sap of the plants which support them.

PARENCHYMA. Pulp part of plants, essentially composed of cellular tissue, which in the leaves fill the meshes between the network of nerves.

PARIETAL. Of or belonging to the walls. Many carpellary leaves, united by their margins, furnished with placental lines, constituting a "parietal placenta," is an application of the term.

PARI-PINNATE. Said of a compound pinnate leaf terminating in a pair of leaflets.

PARTITE. A leaf divided into deep lobes, and reaching near to the base.

PATELLA or PATELLULA. In Lichens a round sessile apothecium, surrounded by a margin belonging to the receptacle and not to the thallus. *Ez.* The Lecidea.

PATENT. Spreading widely. Meaning "open," however, when applied to the corolla.

PATHOLOGY, Vegetable. The study of the diseases of plants.

PATULOUS. Similar to patent, but spreading in a less degree.

PECTINATE. Divided into narrow equal segments, arranged in a single row like the teeth of a comb.

PEDATE. A leaf divided at the top with a leaflet in the fork, and several leaflets on each division.

PEDICEL. The particular support or stalk of each flower. Pedicellate, furnished with a pedicel.

PEDUNCLE. The floral axis, bearing one or a greater number of flowers. Each of its divisions, terminated by a single flower, is the pedicel.

PELLICLE. A thin membrane.

PELLUCID = PUNCTATE. Marked with transparent points, as the leaves of *Hypericum perforatum*; these transparent points are glandular vesicles.

PELORIA. A name given to a teratological phenomenon, which consists in the abnormal or accidental regularity of a flower normally irregular. Such is the deformity seen in the flower of the species of the genus *Linaria*, where five spurs are arranged in a circle instead of a unilateral spur.

PELTATE. In the form of a shield. A peltate leaf has the aspect of a parasol. The leaves are peltate in *Tropaeolum magas*, the *Cotyledon umbilicus*, *Hydrocotyle vulgaris*. Peltiform, said of the frond of an Alga in the form of a peltate leaf; pelta, a sort of apothecium.

PENICILLATE. In the form of a pencil.

PENNIFORM. Frond of an Alga in form of a pinnate leaf.

PENNINERVED. Leaves of which the petiole is prolonged into a middle nerve or vein, emitting, from right to left in all its length, secondary nerves or veins, consequently like a feather.

PENTA. Greek, *five*; Latin, *quinque*. Pentagonal, having five angles; pentagynous, having five styles; pentandrous, having five stamens; pentaphyllus, with five leaves; pentamerous, having equal parts.

PEPO, PEPONIUM, PEPONIDA. The fruit of the Cucurbitaceæ.

PER. The Latin preposition *per* placed before an adjective gives it the value of a superlative. *Pasillus*, slender or delicate; *perpasillus*, very slender or delicate. *Integer*, entire; *perinteger*, very entire.

PERAPHYLLUM. A collective name given to some organs which have no analogy among themselves. For example, the boss or knot of the superior lip of the calyx in the genus *Scutellaria*, the scarious appendage of the calyx in the genus *Salsola*.

PERENNIAL. Living for several years, the opposite of annual.

PERFOLIATE. A leaf is said to be perfoliate when, being alternate, it embraces completely by its insertion the branch upon which it is inserted in such manner as to be traversed by it.

PERI (Greek). Signifies *around*; in Latin, *circa*.

PERIANTH = PERIGONE. The whole of the floral envelopes.

PERICARP. The carpellary leaves at the epoch of the maturity of the fruit.

PERICHÆTIUM. The involucre which surrounds the female flowers in Mosses. The involucral leaves surrounding the Archegonium are called perichætal.

PERICLENIUM = Peripheranthium. The involucre of which the bracts tend to a vertical direction. In this case it is necessary that the peduncle be dilated into a receptacle, and the flowers arranged in the form of a capitulum, as in the *Compositæ*.

PERIDERM (Mohl). The ligneous exterior layer of the bark.

PERIDIUM. In the Fungi, a general envelope, which is ruptured by the progress of the vegetation of the organs which it contains. In the genera *Amanita*, and *Phallus*, the peridium is generally termed the volva.

PERIGONE. Synonymous with perianth.

PERIGYNIUM. Applied by M. Link to the cuticle which envelops the ovary or the fruit in the genus *Carex*. *Perigynous*, when the corolla and stamens are inserted on the tube of the calyx.

PER

PERIPHERICAL. Belonging to the circumference. An embryo, curved, and surrounding the perisperm, is said to be peripheral. *Ex.* The embryo in the Chenopodeaceæ, and in a number of the Caryophyllaceæ.

PERIPERATE. Surrounded by a wing. *Ex.* The fruit of the *Ulmus Campestris*—the seed of the *Drosera Rotundifolia*.

PERISPERM = Albumen. It exists in the Grasses, Liliaceæ, Caryophyllaceæ, Umbelliferae, the Vine, &c., &c.

PERISPORANGIUM. Synonymous with Calyptra.

PERISPORE. Same as Sporangium.

PERISTOME. The margin of the opening of the urn in Mosses after the fall of the operculum, with or without teeth.

PERITHECIUM. In Fungi, certain coriaceous receptacles, containing naked spores, or contained in theca.

PERSISTENT. Said of an organ which does not detach itself spontaneously from the stem—opposed to caducous and deciduous.

PERSONATE. In the form of an ancient mask. It is applied to Gamopetalous corollas, as in *Antirrhinum* and *Linaria*.

PERTUS. Having slits or holes.

PERULAE (*Mirbel*). The envelope formed by the scales of the leaf-bud.

PETALS. The modified leaves which constitute the floral envelope of the corolla. Petaloid, in the form of a petal. Petaloid flowers are those which are doubled by the transformation into petals of the other organs of the flowers.

PETIOLE. The stalk or support of the blade of a leaf. Petiolule, the particular stalk of the leaflet in compound leaves. Peticulate, having a petiole.

PHANEROGAMOUS. Having visible flowers containing stamens and pistils.

PHORANTHIUM = Clinanthium = Common receptacle.

PHRAGMA. Transverse dissepiments in certain fruits; they are of the nature of those which are called false dissepiments. *Ex.* *Cassia Fistula*.

PHYCOLOGY. That part of botany which treats of the structure of the marine Algae.

PHYLLODIUM. Applied to petals deprived of the stalk or limb upon which they are enlarged, and presenting a foliaceous appearance.

PHYLOTAXIS. The study of the laws which govern the arrangement of the leaves on the stem.

PHYLOTOMY. Vegetable anatomy.

PHYSIOLOGY, Vegetable. Study of the vital functions of plants, such as respira-

PLU

tion, circulation, nutrition, fecundation, &c.

PHYTOLOGY = Botany, the study of plants.

PHYTOGENESIS = Organogeny. The study of the development of plants.

PHYTOGRAPHY. The description of plants.

PHYTOS. In compound words from the Greek, *plant*; Latin *planta*.

PILEATE. Resembling a bonnet.

PILEUS. Bonnet. The superior and dilated part of certain Fungi bearing the organs of fructification, *Agaricus Boletus*, for example.

PILIDIUM. Certain receptacles in the Lichens, of which the exterior layer is pulverulent, as in the genus *Calycium*.

PILOSE. Having long unequal hairs on a surface, very little serrated.

PINNATE. Compound leaf, the common petiole, giving insertion right and left to leaflets (petiolated).

PINNATIFID. A leaf cut down to the middle in lateral divisions.

PINNATIPARTITE. A simple leaf of which the segments or divisions extend almost to the middle vein.

PINNULE. The divisions of leaves in the Ferns. Also formerly used in place of leaflets in the description of the leaves of compound pinnate leaves.

PISTIL. The female organ in plants.

PISTILLIDIUM. The female organ in Cryptogamous plants.

PLACENTA. The part of the ovary on which the ovals are inserted. It is considered by many botanists as a prolongation of the axis of the flower.

PLACENTATION. The arrangement of the placenta. It may be either parietal, axillary, or central.

PLATYS. In words derived from the Greek, = *broad*; Latin = *latus*. *Platycarpous*, having large or broad fruit.

PLEIO. Greek, *many*; Latin, *pluri*. *Pleiophyllus*, with many leaves.

PLERIDIUM = Samara.

PLEURENCHYMA. Ligneous tissue.

PLEUROCARPI. That division of the Mosses which have their fruit-stalks always rising laterally from the branch.

PLEURORHIZÆ. Cruciferae, of which the embryo is accumbent.

PLICATE. Folded longitudinally.

PLUMOSE. Resembling a feather; said of a hair bearing two longitudinal rows, composed of short hairs arranged like the divisions of a feather.

PLUMULE. The superior part of the embryo above the insertion of the cotyledon or cotyledons.

PLURIOCCULAR. Having many loculi.

POD

PODETIUM. The pedicel bearing the organs of fructification on several genera of the Hepaticæ.

PODOGYNIUM. Stalk supporting the ovary.

PODOSPERM. = Funiculus. Umbilical cord.

POLLEN. The pulverulent matter contained in the anther. *Pollen-tube*—emitted by the pollen grain when applied to the stigma.

POLYADELPHOUS. Stamens united into several bundles.

POLYANDROUS. Stamens indefinite in number, being above twenty.

POLYCARPOUS. Plants which flourish during many indeterminate years.

POLYCOTYLEDONS. Having many cotyledons.

POLYGAMOUS. Plants bearing at the same time hermaphrodites and male and female flowers. *Ex.* Atriplex Hortensis.

POLYGYNOUS. Having many styles.

POLYPETALOUS. Corolla composed of many petals.

POLYPHYLLUS. An involucre with numerous leaves.

POLYSEPALOUS. A calyx with several sepals.

POLYSPERMOUS. With numerous seeds. Said of a fruit or the loculament of a fruit.

POME. Fruit of plants of the family Pomaceæ (Apple, &c.)

PORE. Foramen, very small opening. Stomata are called sometimes cortical pores.

POROUS. Drilled with pores.

PORRECT. Bearing forwards. Said of a narrow organ erect, relatively to the surface upon which it is inserted; opposed to objects habitually pendant from their own weight. Thus, stamina porrecta, opposed to stamina pendula.

POSTICUS = Extrorse. Directed from the posterior side, in opposition to anticus, directed within. Extrorse anthers are termed antheræ posticæ; introrse anthers are termed antheræ anticæ.

PRÆFLORATION. Arrangement of the parts of the flower during the time it is still in the bud (æstivation).

PRÆFOLIATION. Arrangement of the young leaves in the bud (vernation).

PRÆMORSE. Said of a rhizome which destroys itself successively and rapidly by its posterior extremity, so as to appear truncated and bitten.

PRATENSIS. Growing in the meadows.

PRIMINE = Testa. Name given to the envelope or extreme tunic of the ovule.

PRIMORDIAL. The small leaves which are visible before germination

PRISMATIC = In the form of a prism. The fruit of a great number of Liliacæ (the

PYX

Tulip) and of Iridiæ (Iris, pseudo-Acorns) is prismatic. The stem of a great number of plants. The Liliacæ for example.

PROCESS. Appendage prolonged beyond the organic axis. Under this name are included the appendages which the filaments or the anthers of certain stamens present.

PROCUMBENT = Humifusous. Lying on the ground by weakness or in virtue of a natural direction.

PROEMBRYO. Name given to the false Cotyledons, or articulated and racemose filaments which emanate from the spores in Mosses and Ferns during germination, and preceding the appearance of the true gemmule.

PROLES. Syn. of species. Proliferous or proliferation, teratological phenomenon which consists in the abnormal development of buds.

PROMINENT. Said of an elevated line which rises above the surface.

PRONE. Lying flat on the earth. Pagina prona or infera; said of the dorsal or inferior side of a leaf.

PROPAGULUM. See Gonidium.

PROPHYSES (Willd.). In Mosses the sterile flowers which, mingled with the paraphysis, are inserted at the base of the pedicel of the fertile flower

PROSENCHYMA. Fibrous tissue. This tissue is made up of the ligneous fibres in juxtaposition.

PROTERANTHOUS. Plants of which the flowers develop themselves before the leaves. *Ex.* The Colchicum Autumnale.

PRUINA. Glaucons powder. Pruinose, covered with a glaucous efflorescence; also frosted, or having the appearance of congealed drops of dew.

PRURIENT. Causing itching.

PSEUDO. ψευδής, false; Latin, *spurius*. Pseudo-spermous, fruits having the aspect of a seed; the Achænia and Caryopses, for example.

PUBESCENT. Surface covered with short and soft hairs.

PULVERULENT. Covered with a fine powder like dust.

PULVINATE. In the form of a cushion.

PULVISULUS. Fine powder; the spores in the Lycopodiaceæ; Lycopodium powder.

PUNCTATE. Marked with little spots like points.

PUTAMEN. The hard endocarp of certain fruits.

PYRIFORM. In the form of a pear, pyrum.

PYXIDIA. Dry, capsular, or membranous fruit, opening by a circular lid. *Ex.* Hyoscyamus niger.

QUA

QUADRANGULAR. Prismatic body presenting four longitudinal angles.

QUADRI. In compound words = four times. Greek, *tetra*. Quadricostate, with four sides; quadrifid, with four divisions; quadrijugate, compound pinnatifid leaf with four pairs of leaflets; quadrilateral, with four sides; quadrilocular, with four loculements; quadripartite, with four very deep divisions; quadriphyllus, with four leaves or bractææ.

QUARTINE. Name given by Mirbel to a tegument of the ovule, which he has observed in very rare cases between the tercine and the embryo sac (quintine).

QUATERNATE. An organ of which the parts are arranged by fours. In the case of the

Q

REP

expression "quaternate leaves" it is clearer and better to say "quadrifoliate."

QUINARY. The number five and its multiples.

QUINATE. The parts of an organ arranged by fives.

QUINCUNX. The arrangement of leaves in the bud in which the fifth leaf covers one of the two interior, and has its other margin overlapped by one of the two exterior. Quincuncial, belonging to a quincunx.

QUINQUANGULAR. With five angles.

QUINQUE. In compound words = five. *Ex.* Quinquecostate, quinqueidentate, quinquefid, quinquilobite, quinquipartite, quinquelocular, quinquephyllus, quinquenervis.

QUINTINE (Mirbel) = Embryo sac.

R

RACE, PROLIS. Certain varieties which reproduce themselves indefinitely under the influence of culture, and sometimes even spontaneously.

RACEME. A cluster. Racemose, a plant the inflorescence of which is in the form of a cluster or raceme.

RACHIS. The petiole continued in compound pinnate leaves. The same in the Ferns. Also the axis in Grapes and the axis of a panicle.

RADIANT. Said of the flowers of the circumference in an umbel, these flowers being greater than those of the centre. Said equally of the florets of a capitulum.

RADIATE. Of which the parts are arranged like the spokes of a wheel. The capitulum is radiate in the *Bellis perennis*, *Helianthus annuus*.

RADICAL. Belonging to the root.

RADICLE. The part of the axis which in the embryo is situated beneath the insertion of the leaf or of the cotyledon leaves.

RADII UMBELLÆ (Rays of the umbel). The secondary peduncles of the umbel.

RADIX. The root, the part of the axis increasing by the inferior extremity only.

RAMAL. Appertaining to branches. Ramus, a branch.

RAMENTUM. The brown scale of the Ferns.

RAMOSE. Stem or branch divided into many branches. Ramulus, a little branch.

RAPHE. The jutting or prominent line, which in a reflected ovule commences at the hilum and terminates at the chalazæ.

RAPHIDES. Crystalline substances frequently met with in the cells of many plants.

RECEPTACLE. Enlarged extremity of the

peduncle bearing the bractææ which constitute the involucre and the flowers which form the essential part of the capitulum. The receptacle may be linear, cylindrical, conical, hemispherical, convex, plain, concave, and even tubular. Receptacle of the flower = lower extremity of the pedicel, giving insertion to the different verticils forming the flower.

RECLINATE. Reversed, the extremity curved or inclined towards the base.

RECTANGULAR. Organs which in meeting form right angles.

RECTINERVIS, RECTIVENIUS. Leaves of which the nerves are parallel until near the summit, where they become convergent, as in the Grasses.

RECTISERIAL. Arranged in a rectilinear series, as in the case of leaves.

RECTUS. Straight, right.

RECURVED. Curved, bent in its superior part.

REFLECTED = Anatropal (ovules.)

REFRACTED. Reflected abruptly from the base as if by reason of a fracture. The fructiferous pedicels of the false umbel of the *Holostium Umbellatum*.

REGULAR. Said of a verticil of which all the parts are similar both in form and dimensions.

RELIQUÆ. Remains of withered leaves attached to the plant.

RENIFORM. In the shape of a kidney. The limb of the leaf of the *Asarum Europæum* is reniform.

REPARD. Having an undulated or sinuous margin.

REFENS, = Reptans, creeping—the stems

embedded upon the soil, and emitting adventitious roots.

REPRODUCTION = Propagation. It is oviparous when it takes place by seeds, and gemmiparous when by buds.

RESINOSE. Of the nature of resin.

RESPIRATION. Physiological phenomenon the object of which is the decomposition of the air absorbed by vegetable tissues, the fixation of carbon, and the exhalation of oxygen. The organs of respiration are chiefly on the surface of leaves, and are called stomata.

RESUPINATE. Bent on the back, the result of a twisting.

RETICULATED. In the form of a net, a leaf, of which the nerves anastomose.

RETINACULUM. The glandulous and viscid extremity which terminates the caudicle in certain pollen Mosses.

RETINERVIS. Retivenius (leaves), of which the nerves anastomose like a net.

RETROSE. Reflected, imbricated, leaves from above to below, turned backwards.

RETROSE. Truncated, the extremity cut abruptly.

REVOLUTE. Leaves the margins of which are rolled under during pefoliation.

RHIZOMORPHOUS. Having the appearance of a root.

RHIZOME. An underground stem which creeps horizontally or obliquely under the surface of the soil and terminates in an aerial stem.

RHIZOS. Greek, *root*, or subterranean stem having the aspect of a root.

RHOMBROID. Oval, but slightly angular about the middle.

RIMA. Irregular slit, crevice.

RINGENT. Applied to labiate corollas, and to personate corollas of which the entry of the tube is not closed by a palate.

ROSACEOUS. Applied to corollas the petals of which are shortly unguiculated like those of the Rose.

ROSETTE. The arrangement of numerous leaves in close circles.

ROSTELLUM. The prolongation into a crooked beak of the stigmatiferous and staminiferous column of certain Orchids. Rostellate, furnished with a crooked beak.

ROSTRUM. Prolongation in form of a beak; rostrate, an appendage in form of a beak.

ROTATE. (Rota, a wheel). In form of a wheel; applied to certain Gamopetalons with an almost flat limb. *Ex.* The corolla of the Galium.

ROTUNDUS. Round. Rotundate, in a globular form.

RUBIGINOSE. Colour of rust.

RUDIMENTARY. An organ which exists only in traces, on account of an early arrestment of its development.

RUGA. A wrinkle. Rugose, wrinkled.

RUMINATE. Said of a surface covered with mottles. *Ex.* The Nutmeg.

RUNCINATE. An oblong pinnatifid leaf of which the lobes are pointed and directed towards the base, as in the Taraxacum Densleonis or Dandelion.

RUPESTRIS. Growing on rocks. (Rupes, rocks.)

S

SABULOSE. Growing on sandy soils.

SAC (Embryo) = Sac of the Amnios = Quintine.

SACCATE. In the form of a sac or bag.

SAGITTATE. In the shape of an arrow. That is to say a leaf pointed lanceolate, terminated at its base by two sharp prolongations.

SALICETUM. Place planted with willows.

SAMARA (Gaert). A dry membranous indehiscent fruit with monospermous cells prolonged into membranous wings. *Ex.* The fruit of the Ulmus Campestris or Elm.

SARCOCARP. The fleshy or succulent part of certain pericarps. It is generally the mesocarp which becomes fleshy.

SARCODERM. This is generally applied to the testa or external tissue of the seed, which is fleshy or succulent. In other cases the sarcoderm is a fleshy arillus.

SARMENTOSE. Applied to ligneous stems which attain a greater length relatively to

their diameter, and form runners which find support in neighbouring trees. These stems sustain themselves either in twisting round other stems (*Ex.* the Lonicera Periclymenum or Honeysuckle,) or by means of tendrils of different natures. Vitis Vinifera or Vine, Clematis Vitalba, &c.

SAXATILIS. Growing on stony places, (Saxum, a stone).

SCABISULUS. Scarcely scabrous.

SCABROUS. Covered with short rough hairs. **SCALARIFORM.** In the form of a ladder, as in some vessels.

SCANDENT. Climbing by means of support.

SCAPE. A radical peduncle.

SCARIOUS. Having the consistence of a dry scale. Said of the bractæ of certain involucre, &c.

SCLEROGEN. The hard matter of ligneous cells.

SCORIFORM. In the form of filings. Applied to certain very fine angular seeds.

SCO

SCOBINA. The flexuose rachis of Grasses.

SCORPOIDAL. In the form of the tail of a scorpion. Such is the form of the inflorescence of the *Myosotis*, *Heliotropium*, *Europeum*, &c.

SCROBICULATE. Hollowed, with small depressions

SCUTELLA. Kind of apothecium. Scutum, a shield.

SCYPHUS = *Cyathus*. Scyphiform, in the shape of a drinking cup.

SECRETION. Physiological function, by means of which certain liquids are secreted. They are termed *Excretions* when concreted into a solid substance.

SECUND. Applied to organs turned to one side.

SECUNDINE. The second covering of the ovule.

SEGETALIS. Which grows in harvest.

SEGMENT. A part comprised between two articulations. Also, the divisions of a leaf.

SEGREGATE. Arranged in groups separated one from the other.

SEMI. Half = Greek, *Hemi*. Semi-adherent, applied to an ovary united in its inferior part only to a perigynous tube. Amplexens, half-embracing. Semi-amplexicant, half-amplexicant; semi-flosculus, half floret or ligulate.

SEMI-LOCULAR. Applied to a plurilocular fruit of which the dissepiments are incomplete.

SEMINAL = Cotyledon leaves.

SEMI-PELLUCID. Half-transparent.

SEMI-REFLECTED. Semi-anatropal (*ovule*).

SEMI-VASCULAR. Vascular fascicles surrounded by cellular tissue.

SEMPER VIRENS. Always green. Said of leaves persistent during winter. (Evergreens.)

SEPALS. The modified leaves of which the re-union into a verticil constitutes the calyx, or external floral envelope.

SEPICOLA. Growing on hedges. (*Sepeo*, a hedge.)

SEPTATE. Divided by septa or partitions.

SEPTEM. Seven. Greek, *Hepta*. Septenate, organs approaching in sevens.

SERRALATE. Very finely toothed.

SERRATED. Covered with sharp teeth.

SESSILE. A leaf deprived of a petiole, and consequently having the blade inserted directly on the stem. A flower which has the peduncle very short or absent.

SETA. The pedicel of the capsule, or urn of a Moss.

SIGMOID. A form of the Greek letter *sigma* (σ) in crossing.

SILICULA or SILICULE. A short pod or siliqua, almost as broad as long in the Cruciferae.

SPE

SILIQUA. A long pod in certain genera of the Cruciferae. Siliquose and Siliquae, formed in the shape of a siliqua.

SIMPLE. Not branching or divided. A flower is said to be simple, when it does not present abnormally many rows of verticils; a fruit, when it is the result of the ovary of a single flower; a leaf, as it may be entire or divided, when it is not composed of articulated leaflets.

SINISTORSE. Directed towards the left, in opposition to dextrorse, or towards the right.

SINNATE. With flexuous margins.

SLEEP OF PLANTS. Physiological state under the influence of which certain corollas of plants open and close.

SOCIAL PLANTS. Plants which develop themselves under the same conditions, and grow generally in the same stations, so that the presence of one indicates the probability that others may be near.

SOLID. Full, having no cavity. Solid bulbs are bulbiform rhizomes, short and fleshy, presenting no fleshy scales.

SOLITARY. An organ which is not accompanied by organs of the same nature. The fruit of the Leguminosae is composed of a solitary carpel.

SOREDIIUM. Cells met with on the thallus of certain Lichens.

SOROSIS (Mirbel). The aggregated compound fruits of small succulent drupes half united among themselves, as in a Catkin. *Ex.* Mulberry.

SORUS. Groups of sporangia in the Ferns, inserted at the inferior part of the fructiferous leaves. They may be orbicular, oblong, linear, &c. They are in general covered by an indusium, an organ which is a prolongation of the epidermis of the leaf.

SPADIX. An inflorescence which consists in a radical peduncle terminated by a spike of sessile flowers, furnished at the base with a spathe. *Ex.* Arum and Calla.

SPATHA or SPATHE. An involucre composed of numerous membranous bracts which envelop the inflorescence before the expansion of the flowers. *Ex.* Palms, Arum, Allium, &c.

SPATHACEOUS. Having the aspect and membranous consistence of a spathe.

SPATHELLÆ. Synonym of Glumellæ, (Grapes.)

SPATULATE. Shaped like a spatula; a leaf of a linear form, enlarging at its extremity into an aboval or elliptical blade.

SPECIFIC CHARACTERS. The differences which distinguish species of the same genus.

SPE

- SPECIOSUS.** Beautiful, of an elegant form and brilliant colour.
- SPERMATOZOID.** Vegetable filaments contained in the organs called antheridea, and which exist in the plants of many Cryptogamous families,—Lichens and Hepaticæ for example, and even the Ferns, during the first period of their germination. Each spermatozoid is contained in a membranous cell, from which it issues making spontaneous movements.
- SPERMODERM.** Same as EpispERM.
- SPIKE.** Consists of numerous sessile flowers on an axis. Spikelet, small groups of flowers in Grasses.
- SPINDLE-SHAPED.** Fusiform.
- SPINE.** An abortive branch terminating in a sharp point, such as the spines in the *Prunus Spinosa*.
- SPINESCENT.** In the form of a spine.
- SPINOSE.** Covered with spines.
- SPIROLOBLE.** In the Cruciferae of which the embryo is incumbent, and of which the cotyledons are prolonged and rolled spirally.
- SPONGIOLE.** The cellular extremity of each of the divisions of the root by which absorption takes place.
- SPORANGIUM.** A receptacle which in a great number of Cryptogams contain the spores. They are either membranous, crustaceous, or cartilaginous.
- SPORES.** The reproductive bodies in Cryptogamous plants, analogous to seeds and embryos in Phanerogamous plants, differing, however, in having no appearance of being the result of fecundation; homogeneous, and not composed of distinct parts analogous to those of the Cotyledons. In the Ferns the spores are veritable rudimentary inflorescences, which develop themselves in growing upon the soil.
- SPOROIDIUM.** Spore. Used also as a synonym of Sporozoidium.
- SPOROZOID = Zoospore.** The spores of certain Algæ which seem endued with spontaneous movements after being detached from the mother plant, and which keep in motion for a certain time before fixing upon a solid body in order to germinate.
- SQUAMA.** A scale. Squamiform, in the form of a scale; squamate, provided with scales. Certain bracts are squamiform. Squamulae, little scales = Glumellæ.
- SQUAMOSE.** Covered with spread-out hairs at right angles, as in the case of certain involucres of plants of the Compositæ, of which the bracts are scarious and the extremity spread out or reflected.
- STAMEN.** The male flower of plants, con-

STR

- sisting of the filament and anther. Stamenate, applied to male flowers.
- STAMINODIUM.** In the Orchids the two lateral or sterile stamens, which are reduced to a small fleshy appendage, sometimes indistinct. The genus *Cypripedium* has the middle stamen abortive and the two lateral developed into normal stamens.
- STATION.** The particular place where a species is spontaneously developed. Habitat is synonymous with station in a geological point of view. Locality, a restricted part of a country.
- STICHUS.** In compound words, signifies a row, as distichus, arranged in two rows; tristichus, in three rows; hexastichus, in six rows.
- STIMULUS.** Applied to stinging or urticant hairs. Stimulose, furnished with stinging hairs.
- STIPATE.** Pressed by approached organs.
- STIPEL, or STIPELLA.** Small accessory stipules which accompany the leaflets or segments of certain leaves.
- STIPITATE.** Elevated on a stalk.
- STIPULARY.** Tendrils (*Cirrhii Stipulares*) supposed to be produced by the transformation of the stipules.
- STIPULE.** Accessory leaves, which accompany leaves in a great number of plants.
- STOLO.** A sucker, at first aerial, and then turning downwards and rooting.
- STOLON.** Creeping rooting stems on the level of the base of the leaves, and of rosettes which spring at their axils, more particularly when the internodes of these creeping stems persist more than one year. *Ex.* The Strawberry, the *Ranunculus repens*, and *Potentilla reptans*. *Stoloniferous*, bearing stolons.
- STOMATA (*Stomatis*).** Cortical or epidermal pores, requiring magnifying power to render them visible. They are met with generally on the leaves (inferior side). It is by these openings or sort of mouths that the function of respiration in plants is effected. Under the influence of the light, the external air coming in contact with the cellular tissue of the parenchyma of the leaves, and of the herbaceous envelope of the bark, is exhaled, after having yielded to the vegetable liquids the carbon of its carbonic acid, and the oxygen being set free is thrown into the surrounding air. The stomata are beautifully seen in the leaf of the *Begonia*.
- STRANGULATED.** Contracted and expanded irregularly.
- STRIA.** A very fine line upon a surface. *Striated*, having *Striæ*.
- STRIGOSE.** Furnished with rough strong

STR

hairs, as in the stem and leaves of a great number of the Boraginææ.

STROBILUS—Cone. The aggregated fruit of a great number of the Coniferæ and Cycadaceæ (*Gymnosperous*.) These fruits are composed of imbricated scales, springing at the axil of a membranous bract, sometimes not manifest, and presenting at their base naked seeds, but concealed before the disaggregation of the fruit, by the imbrication of the scales, one upon the other. Such are the fruits on the Pines, Firs, Cedar, &c. &c. *Strobiliform*—having the form of a Strobilus or cone. Said of certain aggregated fruits, which do not belong to the Gymnosperms, *e.g.*, the fruit of the Hop, Alder, &c.

STROPHIOLE. Sort of fleshy arillas, in the form of a crest, constituting an appendage on the surface of the seed, and not enveloping it completely. **STROPHIOLATE**. Said of a seed furnished with a *Strophiole*.

STRUMA. A lateral swelling or rugosity.

STUPOSE, or *Stupaceous*. Having a tuft, (Stupa).

STYLE. The filiform, cylindrical, demi-cylindrical, or prismatic prolongation which surmounts the ovary. The length of the styles is very variable. When wanting, the stigma is said to be sessile.

STYLOPODIUM. An epignous disk or thickening of the base of the style in the Umbelliferæ.

STYLOSTIGIUM (Link.) In the Asclepiadaceæ the enlarged base of the style which covers the ovary like a hood, in the genus *Stapelia*, for example.

STYPTIC. Bitter and astringent.

SUAVEOLENS. Exhaling a sweet odour.

SUB. Almost, scarcely—diminutive associated with qualifying terms, in order to lessen the value. Subinspid, almost inspid; subrotund, almost round. Sub in certain compound words also signifies *under*.

SUBEROSE. Of the nature of cork; suber, presenting the consistence of cork.

SUBMERGED. A plant which vegetates entirely covered with water.

SUBTERRANEAN = Hypogæal. Developing itself under the surface of the soil.

SUBULATE. In the form of an awl; said of a prismatic body terminated gradually by a sharp point, as in the leaves of Firs.

SUCCEDANEOUS. Said of a vegetable product which can replace another in point of its medical properties or economic uses.

SUCCULENT. An organ gorged with an aqueous or mucilaginous juice, whatever may be its taste or properties.

SUCCUS. Juice. The sap more or less elaborated. They may be either aqueous,

SYM

mucilaginous, milky, resinous, opaque, limpid, viscid, &c.

SUCKER. Haustorium. A short adventitious root which implants itself on a living stem and absorbs its sap. The *Cuscuta*, for example.

SUCTION. Phenomenon by which a root absorbs the water into which it is plunged. The extremities of the roots (*Spongioles*) are the points where the suction or absorption takes place with the greatest extremity.

SUFFRUTEX. An under shrub. **SUFFRUTICOSE**, having the characters of an under shrub.

SULCATE. Furrowed—marked with deep grooves. *Sulcus*, a furrow.

SUPERANS. Exceeding adjoining organs in length.

SUPERIOR. Situated above—applicable to the ovary when free and not adherent.

SUPERVOLUTE. Leaves whose margins are rolled beneath during veneration.

SUPINATE. Stems spread upon the soil but not rooting.

SUPRA. In compound words means above, upon. Greek, *epi*. *Supra*-axillary, placed above the axil; said of certain buds.

SUPREME. Situated at the summit.

SUSPENDED. An ovule is said to be suspended, when being inserted towards the superior part of the loculament, its summit is directed towards the base; if the loculament contains many ovules inserted at different heights, and of which the summit is directed downwards, then ovules are said to be pendent.

SUTURAL. Belonging to the suture. *Suturate*, presenting a suture.

SUTURE. The line by which the margins of two leaves are untied. Gamopetalous corollas, gamocarpellous ovaries, &c. present sutures between the margins of different leaves approached in circles. Carpellary isolated leaves present a ventral suture, which results from the union of their two margins. Where the capillary leaves are united in circle they form a plurilocular ovary.

SYCONIUM or **SYCONUS**. (Mirb.) Name given to the inflorescence in the Fig. This inflorescence is constituted in the floriferous state by flowers, and the fructiferous state by free isolated fruits, but contained in close depression of the peduncle, which increases itself in thickness, and becomes fleshy and succulent at maturity.

SYLVESTRIS. Growing in the forests.

SYMMETRICAL. An organ when it is capable of being divided into equal halves; a simple organ—a leaf for example, which is symmetrical is necessarily of a regular

SYN

form; a compound organ—a corolla for example, can be symmetrical without regular simple organs—leaves, sepals, petals, carpels, &c., may or may not be symmetrical; compound organs—calyx, corolla, &c., are always symmetrical, but may or may not be of a regular form.

SYNANTHEROUS. Stamens united by their anthers into a sheath surrounding the style, as in the *Compositæ*. *Synanthos*, flowers united. *Syncarpous*, carpels united. *Syngenesious*, same as synantherous. (*Syn*, together, united.)

TOR

SYNORHIZAL. When the radicle of the embryo is more or less adherent to the perisperm, as in the *Coniferæ*. *Endorhizal*, the radicle of an embryo furnished with a *Coleorrhiza*. *Exorhizal*, when the radicle is developed without a *Coleorrhiza*.

SYSTEM. Classification of plants according to certain principles. The term *Taxonomy* is applied to the study and elucidation of the different classifications which have been successively proposed.

T

TENIÆFORM. Shaped like *tænia* or in the form of a flat band.

TAXONOMY. See *System*.

TEGMEN = *Secundine*. The covering of the ovule situated under the testa.

TEGMENTS or TEGMENTA. Scales (scarious leaves) which in certain trees cover the young leaves during the winter, and are developed in spring.

TEGMENTS or TEGUMENTA. Organs which protect others by covering them. The floral teguments are the bracts. The pericarp should be termed the tegument of the seed, and the epispem, tegument of the embryo; but, by abuse of words, the tunics of the seed (epispem) are called teguments, which form essentially a part of the seed.

TENUIS. Thin, slender, delicate.

TERATOLOGY. The science which has for its object the study of anomalies, or accidental deviations from the normal type.

TERCINE. The third covering of the ovule, in which is developed the embryo sac.

TERNARY. The number three and its multiples. The constituent parts of each of the verticils of the flowers of the monocotyledons are generally ternary. **TERNATE**, arranged by threes. Compound leaves with their leaflets are termed trifoliate, not ternate.

TESTA—*Primini*. The external tunic of the ovule or seed.

TESTICULATE ROOT. Having two oblong tubercles.

TETRA. In compound words derived from the Greek, signifies *four*; Latin, *quadri*. *Tetradynamous*—having four stamens longer than the others, (*two*) as in the *Cruciferae*.

TETRAGONAL. A surface limited by four straight lines.

TETRAGYNOUS. Having four free carpels or styles.

TETRANDROUS. Having four stamens.

TETRAPTEROUS. Four-winged.

TETRASEPALOUS. Having *four* sepals.

THALAMIFLORÆ. De Candolle applies this to flowers with free petals, inserted upon the receptacle on the same level as the ovary.

THALAMUS. Floral receptacle, or extremity of the peduncle, into which are inserted the organs of the flower.

THALLUS—*Fron*d. The foliaceous or fructiferous expansion constituting the stem in the *Lichens*. **THALLOGENS**, plants bearing a thallus.

THECA. A sort of sporangium formed by a prolonged or globular utricle containing the spores. The spores are contained in theca in the *Lichens*, and in a great number of the genera of the *Fungi*.

THECAPHORE. Said of a surface bearing theca, and of a receptacle containing theca, also, as the attenuation of a carpel at its base, into a kind of stalk or pedicel. (This latter term is seldom used.)

THYRSUS. A sort of panicle belonging to mixed inflorescences, with a principal indefinite axis, and lateral definite axes. *Thyrsoïdal*, in the form of a *Thyrsus*.

TOMENTOSE. An organ covered with a cottony pubescence, or soft long flexuose hairs. **TOMENTUM**, cottony pubescence.

TOROSE, TORULOSE. Applied to the silicle of certain *Cruciferae*, which present successive swellings, determined by the presence of superimposed seeds.

TORSIONS, AENORMAL. The accidental twisting of an organ is determined by an inequality of development on its two faces or opposite sides. The torsion may take place in a vertical direction, or produce a rolling up from above downwards.

TORUS. The receptacle of the flower or extremity of the peduncle. It may be flat, concave, convex, conical, and even filiform. Its length is in general proportionate to the number of organs it bears.

TRA

- TRACHEÆ.** Vessels containing one or several fibres, rolled or spiral, and capable of being unrolled by traction. *Trachenchyma*, the tissue composed of these vessels.
- TRANSPIRATION.** Exhalations of liquids through vegetable tissues.
- TRANSVERSAL.** Arranged across. The transversal partitions which divide a cavity are called diaphragms.
- TRAPEZÖIDAL.** Approaching the form of a trapezium or square with unequal sides.
- TRI.** Greek or Latin, *three*. **TRIADELPHOUS**, with three bundles of stamens. *Triandrous*, with three stamens; triangular, having three angles; *tribracteate*, with three bracts. **TRIBRACTEOLATE**, with three bractiolæ.
- TRICHOTOMOUS.** Divided by threes, of which the divisions themselves are divided by three, and so on.
- TRICOCCOUS.** A fruit composed of three carpels.
- TRICOSTATE.** Three-ribbed.
- TROCHLEARIS.** In the form of a pulley; discoid, and presenting a groove upon the margin.
- TUBIFORM.** Like a trumpet, or straight tube widened at its extremity.
- TUBE.** Cylindrical hollow organ, open at

VAG

- its superior extremity. In gamosepalous calyces and certain gamopetalous corollas, the tube is the inferior part when it is cylindrical in form, and the term *Limb* to the spread out superior part.
- TUBER.** An underground fleshy expansion of the stem of plants, usually containing starchy matter, as the Potato.
- TUBERCULES.** Swellings of different natures observed on the axes of subterranean stems.
- TUBEROSE.** Having the shape or appearance of a tubercule.
- TUBEROSITY.** Thickening or nodosity in the form of a tubercule.
- TUNIC.** In the ovule, the testa is sometimes termed the external tunic, and the tegmen the internal tunic. It is also applied to the fleshy leaves constituting the mass of a bulb, as in the case of the Onion. It is then said to be *tunicated*.
- TURBinate.** In the form of a top, or inverted cone, as in the *Brassica rassa*.
- TURIO.** The young shoots which are emitted from a stem during the period it is subterranean. *Ex.* Asparagus
- TYPE.** Said of an individual which reunites all the characters of the species in a high degree.

U

- ULEGINOSE.** Growing in humid, slightly-turfy meadows.
- UMBEL.** An inflorescence belonging to the indefinite type, in which all the branches arise from one point. *Ex.* The *Umbelliferae*. Umbelliform, in the shape of an umbel.
- UMBILICUS.** The hilum. *Umbilicate*, furnished with a hilum.
- UMBO.** Conical protuberance elevating itself upon a surface. *Umbonate*, presenting an umbo or protuberance like the boss of an ancient shield.
- UMBRACULIFORM.** In the form of an umbrella or parasol. *Umbaculum*, organ in the form of an umbrella; for example, the bonnet of the *Agarius* and *Boletus* with central stalk.
- UMBROSUS.** Growing in shady places.
- UNCIA** = Pollex. The thumb. Twelve lines. *Uncialis*, of the length of a thumb.
- UNCINATE.** (*Uncus* = a hook.) Furnished with a crooked point. *Uncinate-aculeate*, having hooked needle points. *Uncinatesetose*, provided with crooked hairs.
- UNCTUOUS.** Having the consistence of oil. Applied to a surface covered with mucilage, as in the case of the surface of the leaves of the *Pinguicula*.
- UNDULATE.** Wavy or sinuous.
- UNGUIS.** Claw; the narrow base of a petal. *Unguiculate*, furnished with a claw.
- URCEOLATE.** In the form of a vase or goblet. Said of a gamopetalous globular corolla, with a narrow opening, as in the *Convullaria Maialis*.
- URENS.** Burning. A term applied to plants of which the sting produces the sensation of burning.
- URN.** The fructiferous capsule of Mosses.
- USTULATE.** Blackened.
- UTRICLE.** An organ resembling a membranous gourd.

V

- VACUUS.** Empty, hollow, sterile.
- VAGINA.** Inferior sheathing part of certain leaves. *Vaginate*, embraced by a sheath. *Vaginula* or *Vaginule*, a small sheath which embraces the base of the stalk of the urns in the Mosses, and

VAL

which is the lower part of the broken epigonium.

VALLECULA. The interval which separates two ribs in the fruit of the Umbelliferæ. *Valleculate*, having vallecules.

VALVATE = *Æstivation*. When the sepals or petals are in contact by their margins without covering themselves mutually.

VALVES. The parts or divisions resulting from the dehiscence of ligneous or membranous fruits. According to the mode of dehiscence the valves represent different parts of the pericarp, scarious bracts forming the involucre of the spikelet or grasses. (See *Glume*.) *Valvule*, a small valve. The lobes of certain anthers open by valvules.

VAXILLARY. The *æstivation* in the Papilionaceæ.

VEGETATION, Organs of. All the organs, with the exception of the flowers, fruits, and seeds, which are termed the organs of reproduction.

VEINS. The secondary nerves in leaves. *Veinules*, very small veins.

VELUM. In the genus *Agaricus* the membrane which arises from the margin of the bonnet, of which it is the continuation, and covers the inferior face furnished with gills, advancing to the stalk, which it embraces in a ring-like manner. In the progress of vegetation the velum is irregularly torn, the debris forming a ring around the stalk.

VELUTINOUS. Having a velvety aspect.

VENOSE. Veined; having fine nerves.

VENTRAL. In carpels, the side next the axis.

VENTRICOSE. Inflated in the manner of a balloon; swelling unequally on one side.

VENUCOSE. Covered with wart-like protuberances.

VERMICULAR. Worm-shaped.

VERNAL. Flourishing in spring.

VERNATION = *Prefoliation* or arrangement of the leaves in the bud.

VERSATILE. Easily turned about on its

XYL

support, as in the case of certain anthers upon their filaments.

VERTEX. The summit.

VERTICAL. Of which the direction is perpendicular to the horizon.

VERTICIL. The whole organs arranged in a circle upon the same level around an axis. **VERTICILLATE**, arranged in verticils or whorls.

VESICLE. Membranous organ in the form of a little bladder. *Vesiculose*, in the form of a vesicle.

VESICOLOR. Applied to an organ which changes its colour many times during the successive phases of its development. For example, the corolla of certain plants of the Borraginææ.

VESPERTINE. Flowering after sunset.

VESSELS. The tissue of plants composed of tubes often cylindrical, more rarely prismatic, long and narrow, and terminated in points more or less attenuated, or obtuse, at the two extremities. They are in general associated with fibres, forming fascicles or bundles, and constituting fibro-vascular tissue. Tracheæ, scalariform, dotted, often moniliform, and such like.

VEXILLUM. Standard. Superior petal of the Papilionaceous corollas.

VIRGATE. In the form of a switch.

VIRIDIS. Green.

VISCOUS, VISCID. Having a syrupy consistence.

VITELLINE. Resembling the yolk of egg.

VITTA (Plur. Vittæ.) Resiniferous tubes situated in the pericarp in the fruit of a great number of plants of the Umbelliferæ. In order to perceive the position of the vallecules it is necessary to cut the fruit across. *Vittate*, having vittæ.

VOLUBLE. A stem or tendril turning itself around other stems which furnish a support to them.

VOLVA. In the *Agaricus*, an envelope in the form of a sac without an opening, which contains the young mushroom during the first period of its existence.

W

WHORL. Synonymous with verticil.

WINGED. Having a broad membranous margin or border as in a Samara. The

wings (Alæ), the two lateral petals of a papilionaceous corolla.

X

XANTHOS. In words derived from the Greek, signifies *golden yellow*; Latin, *aureus*.

XYLON. In words derived from the Greek, signifies *wood*; in Latin, *tignum*.

INDEX.

Paragraph
 Abaxial or abaxile..... 602
 Abbreviations..... 713
 Aberration, spherical.....1192
 chromatic.....1192
 Abies.....1046
 Abietinæ.....1045
 Region of.....1162
 Abnormal.....121
 Abortion or non-develop-
 ment..... 339
 Absorption of fluids by
 plants..... 253
 Inorganic
 matters..... 231
 Acacia.....157, 852
 Acanthaceæ..... 970
 Acanthodium..... 971
 Acanthus..... 971
 Acaules..... 65
 Acclimatizing of plants.....1172
 Accrescent..... 367
 Accumbent..... 601
 Aceraceæ..... 805
 Achenium..... 541
 Achlamydeous.....351, 649
 Achlya.....1129
 Achras..... 937
 Achromatic.....1205
 Acicular..... 150
 Acids, Organic..... 320
 Acinaciform..... 150
 Acinus, one of the divisions
 of the fruit of the Rasp-
 berry..... 545
 Aconita or Aconitina..... 740
 Aconitum..... 740
 Acoreæ.....1092
 Acorus Calamus.....1093
 Acotyledones.....590, 1104
 Acotyledonous stem, 71, 102, 107
 Acramphibrya..... 736
 Acrobrya.....1104
 Acrocarpi.....1115
 Acrogenæ.....1105
 Acrogenous stem, 71, 102, 107
 Acrogens, reign of.....1185
 Actææ..... 739
 Actinenchyma..... 5
 Aculei..... 57
 Acuminate..... 160
 Adam's Needle.....1078
 Adansonia..... 784
 Adherent.....160, 408
 Adhesion of Organs..... 652
 Adnate.....160, 408
 Adpressed hairs..... 60

Paragraph
 Adventitious leaf-buds..... 196
 roots..... 121
 Aerial stem..... 70
 leaves..... 134
 Æsculus..... 807
 Æstivation..... 354
 Æthiopian Lily.....1093
 Affinity.....704, 1147
 Agamous..... 389
 Agaricus.....1124, 1125
 Agathotes Chirayta..... 949
 Agaveæ.....1072
 Ages of trees.....79, 635
 Agglutination of cells... 13
 Aggregate..... 541
 Aggregate fruits..... 538
 Air-cells..... 21
 Air-plants.....121, 251
 Akee fruit..... 807
 Alabastrus..... 354
 Alæ..... 379
 Alangiaceæ..... 864
 Albumen.....557, 588
 Albuminous..... 587
 Alburnum..... 78
 Alder.....1036
 Algæ.....1127
 —, how to prepare.....1236
 Algaroba-bean..... 851
 Alismaceæ.....1088
 Alkaloids..... 321
 Alkanet..... 959
 Allieæ.....1077
 Alligator pear..... 994
 Allium.....1078
 Allspice..... 867
 —, Carolina..... 856
 Almond..... 855
 Aloe, American.....1073
 Aloes.....1078
 Aloes-wood.....1008
 Aloineæ.....1077
 Aloysia..... 969
 Alsinaceous corolla..... 378
 Alsineæ..... 778
 Alsodeleæ..... 768
 Alströmérieæ.....1072
 Althæa..... 782
 Alternæ..... 170
 Altingiaceæ.....1033
 Alumina in the soil..... 236
 Amadou.....1125
 Amanita.....1124
 Amaranthaceæ..... 984
 Amaryllidaceæ.....1072
 Amentaceæ.....1033

Paragraph
 Amentum..... 334
 American cowslip is Dode-
 catheon..... 975
 Anomum.....1062
 Ammonia, source of nitro-
 gen..... 215
 Ammoniacal liquor, as a
 manure..... 246
 Ammoniacal manures..... 243
 Ammoniacum..... 891
 Amnios.....463, 586
 Amniotic sac..... 504
 Ampelidææ..... 813
 Amphibious plants.....1139
 Amphibrya.....1049
 Amphigamæ or Amphi-
 genæ.....1118
 Amphitropal.....468, 603
 Amplexicaul..... 139
 Ampullæ.....120, 164
 Amygdalææ..... 854
 Amygdaline.....310, 855
 Amyridaceæ..... 844
 —, Region of.....1155
 Anacardiaceæ..... 842
 Anacyclus..... 917
 Anagallis..... 975
 Analogy.....1147
 Analysis of soils..... 238
 Anamirta..... 747
 Ananas.....1076
 Anatomy, vegetable..... 1
 Anatropal.....467, 584
 Anatropous.....467
 Andira..... 850
 Androecium..... 390
 Androphore.....399
 Anemoneæ..... 739
 Anethum..... 892
 Anfractuose..... 406
 Angelica..... 890
 Anglenchyma..... 28
 Angiospermæ..... 981
 Angiospermous..... 576
 Angiosperms, reign of.....1198
 Angustiseptæ..... 760
 Angustura or Angostura
 bark..... 828
 —, False..... 947
 Anise..... 892
 Anisostemonous..... 392
 Annual.....633, 713
 Annular cells..... 11
 — vessels..... 35
 Annularia.....1186
 Annulate ferns.....1108

	Paragraph		Paragraph		Paragraph
Annulated root	12	Artichoke, Jerusalem	917	Bane-berry is Actæa	7 9
Anonacæe	744	Artificial System	715	Banksia	998
Anterior, applied to parts		Artocarpeæ	1019	Banyan	1021
of the flower	357	Asarabacca	1010	Baobab	108, 784
Anthemis	917	Asarum	1010	Barbadoes Cherry	804
Author	395, 402	Ascending ovule	471	Barbadoes Gooseberry	883
—, colour of	413	Ascherson on Cytogenesis	22	Barbed	58
—, coverings of	403	Ascidia	164	Bark	72, 82
—, dehiscence of	410	Ascioplacæe	912	—, false in Endogens	92
—, development of	404	Ascomycetes	1123	—, functions of	109
—, form of lobes of	406	Ascus	459, 1119	—, mode of its increase	86
—, union of	417	Ash	941	—, rupture of	87
Anthericæe	1077	Ashes of plants	220	—, separation of outer	
Antheridium	427, 492	— Bean, Pea, and		layers of	88
Antthesis, opening of the		Rye-Grass	223	Bark-bread	1046
flowers	354	Oats	223	Bark, Peruvian	933
Anthocarpous	538, 556	Wheat	223	—, false	913
Anthodium	336	Asparagæe	1077	—, fever	903
Anthotaxis	324	Asparagus	1078	Barley	1142
Antiaris	1022	Aspen	1034	Barosma	828
Antice	412	Asphodelæe	1077	Barringtoniæe	806
Antirrhinidæe	904	Assafetida	891	Barry on Cell-formation	22
Antitropal	603	Asteracæe	910	Basil, see Basilar.	
Aperispermic	605	Asters, Region of	1155	Bases, organic	321
Apetalous	647	Astragalus	850	Basidia	1122
Aphyllanthæe	1077	Atherospermæe	1027	Basil	967
Apicæe	888	Atractenychma	5	Basilar	451
Apicilar	451	Atriplex	987	Bassila	927
Apicular	530	Atropa	961	Bassorin	303
Apiculate	528	Atropa	961	Bast, or Bass	29, 788
Apocarpous	434, 538	Atropous or atropal	465	Bast layer	83
Apocynacæe	944	Attar of Roses	835	Batatas	956
Apophysis	1114	Aucuba	896	Bay-Laurel	855
Apothecia	1119	Aurantiacæe	795	Beaded-vessels	87
Appendicular organs	57	Auricula	975	Bean	830
Appendiculate	383, 898	Auriculæ	150	Bean-caper	826
Apple	855	Autumn Crocus	1089	Bearded	60
Appressed hairs	60	Avellano	998	Bebeeru	994
Aquatic plants	1139	Avocado Pear	994	Beech	1037
Aquifoliacæe	934	Axe-shaped	150	Beech-drops	963
Aquillariacæe	1003	Axial	602	Beet	987
Arabine	803	Axil	140	Beet-sugar	306
Aracæe	1092	Axile	440, 602	Begoniacæe	992
Araliacæe	894	Axillary	140, 169	Bell-shaped	380
Araucaria	1046	Axis, ascending	70	Belladonna	961
Araucarites	1190	—, descending	119	Belvisiacæe	874
Arborescent	68	—, floral	325	Bengal Hemp	850
Arbor-vitæ	1046	Azote, source of	215	Ben-nuts	833
Arbutus	927	Azotized matter in plants	311	Bent or Marram	1103
Archegonium	492	—, Products	310	Benzoïn	932
Archil	1121	Azalea	927	Berberidacæe	748
Aretium	916	Babul-wood	852	Berberry or Barberry fa-	
Arctostaphylos	927	Bacca	550	mily	748
Ardisia	938	Baccate	550	Bergamot	793
Areca	1086	Balanophoracæe	1011	Berry	530
Arecinæe	1085	Balausta	550	Betel-nut palm	1086
Argel	943	Balm	967	Betel or Betle-pepper	1032
Arilode	581, 996	Balm of Gilead	845	Betulineæ	1033
Arillus	581	Balsam, bursting of seed-		Bhang	1020
Arlineæe	1092	vessel	27	Biennial	653, 713
Arista, an awn, as in		— family	819	Bifid	372
grasses	1099	— of Copalva	851	Bifurcate	406
Aristate	1099	— of Peru	850	Bignoniacæe	950
Aristolochiacæe	1009	— of Tolu	850	Bijugate	153
Armeria	977	— of Umiri	810	Bikh	740
Arnica	917	Balsamacæe	1033	Bilablate	382
Arnotto family	764	Balsaminacæe	819	Bilamelar	457
Aracacha	890	Balsamodendron	845	Bilateral	456
Arrow-root	1064	Balsams	318	Bilimbi fruit	821
—, Brazilian	1018	Bamboo	1102	Bilocular	405, 410
—, spurious	1048, 1062	Banana	1066	Bindweed family	955
Artanthe	1092	Bananas, region of	1160	Bipartite	372, 454
Artemisia	917			Bipinnate	152
Artichoke	916			Bipinnatifid	147

Paragraph	Paragraph	Paragraph
Bipinnatifid 147	Budding 181, 573	Camphor of Borneo 790
Birch 1036	Buds, latent 189	Campotropical 466
Bird-lime 899	— on leaves 199	Campylospermæ 8-8
Bird-pepper 961	Bulb, scaly or naked 195	Campylotroual 466, 584
Birthwort family 1009	—, solid 195	Campylotrouous 466
Biscuit-root 1073	—, tunicated 194	Camwood 851
Bistort 991	Bulbils 193	Canada-Balsam 1046
Bitternate 154	Bulblets 193	Canada-Rice 198
Bitter-Almond 855	Bulrush tribe 1092	Canary-seed 198
Bitter-Apple 872	Burdock 916	Candleberry 1035
Bitter-Sweet 961	Burgundy-pitch 1046	Cane-Sugar 304
Bivalvular 530	Bariti-palm 1086	Canella 810
Bixææ 766	Burmanniaceæ 1069	Canker 692
Black-Alder 841	Bursereæ 844	Cannabls 1020
Black-Bryony 1052	Bursicula 1058	Cannabineæ 1019
Bladder-nut 839	Bush, defined 68	Cannaceæ 1063
Bladder-Senna 850	Butea 850	Cance-birch 1036
Bladder-wort 973	Butomaceæ 1089	Caoutchouc 319, 945, 1018, 1021
Blade of leaf 139	Butter of Cacao 786	Caper family 763
Blanching 300, 690	— of Canara 790	Caper-spurge 1017
Bletting 567	Butterfly-weed 943	Capillaire 1110
Blight 693, 1126	Butter-tree of Park 937	Capillary 297
Blood-root 1071	Butter-wort 973	— attraction 267
Boards for pressing plants . 1205	Bytteriaceæ 785	Capitate 58
Bohun-Upas 1022		Capitulum 336, 348
Bombacææ 783	Cabbage 762	Capparidaceæ 763
Bombax 784	Cabbage-Palm 1086	Caprifrication 489
Bones, as a manure 246	Cabbage-tree 850	Caprifoliaceæ 900
Bonplandia 828	Cable-cane 1086	Capsicin 961
Borage 959	Caobomaceæ 749	Capsicum 961
Boraginaceæ 953	Cacao-beans 786	Capsule 554
Borassineæ 1085	Cactaceæ 832	Caraway 892
Borneo-Camphor 790	—, Region of 1155	Carboy in plants 212
Botanical excursions 1242	Cactus family 882	Carbon, fixed by light 236
Botany, object defined 1	Caducous 366, 385	Carbonic acid given off in
—, departments of 1	Cæsalpinieæ 848	— darkness 287
Bothrenchyma 10, 36	Caffeine 905	— decomposed by
Bottle-Gourd 872	Cajeput 867	— aquatics 289
Boucherie's method of pre-	Calabash-tree 951	— given off by
serving timber 270, 699	Calamites 1190	— flowers 475
Box-tree 1018	Calamus 1190	Carboniferous fossils 1186
Brachyphyllum 1197	Calathium 336	Cardamoms 1062
Bract 324, 344	Calcar 363	Cardoon 916
Bracteole 333	Calcarate 363, 373	Carex 1098
Bractlet 333	Calceolaria 965	Caricææ 873
Branches 188	Calceolarieæ, Region of . 1155	Carina 379
Branching of trees 67	Calceolate 381	Carinal 355
Branchlets or Twigs 188	Calculate 347, 364	Cariopsis 542
Brassicææ 758	Calluna 927	Carnahuba palm 1086
Brazil-nuts 867	Calophyllum 799	Carnation 779
Brazil-wood 851	Calotropis 943	Carob-tree 851
Bread-fruit 1022	Calumba 747	Carolina Allspice 856
Bread-tree 1048	Calumbin 747	— Pink-root 947
Brexiacea 824	Calycanthaceæ 856	Carpathian Balsam 1046
Bristles 57	Calyceraceæ 909	Carpel 430
Britain, Flora of 1169	Calyceifloræ 390, 835	Carpology 537
British plants, distribution 1167	Calyptra 1144	Carpophore 437, 533, 543
Brocoll 762	Calyptrate 366	Carrageen 1130
Bromeliaceæ 1075	Calyptrimorphous 164	Carrion flowers 943
Brongniart, Fossil Flora . . 1193	Calyx 350	Carrot 890, 892
Broom 850	—, forms of 358	Cartilaginous 583
Broom-rape 252, 963	—, tube of 362	Caruncles 582
Brucea 947	—, limb of 362	Carunculate 582
Bruniaceæ 886	—, venation of 359	Caryophyllaceæ 778
Brunoniaceæ 919	Cambium 81	— Region of 1155
Bryony 872	— cells 109	Caryophyllaceous corolla . 378
Bryophyllum 199, 880	Cambogia Gutta 799	Caryopsis 542
Buck-bean 949	Camellia 793	Cascarilla 1018
Buck-eye 807	Camelliaceæ, Region of . 1155	Caseine, Vegetable 310
Buckthorn 841	Camera-lucida 1210	Cashew-nut 326, 843
Bucku or Buchu 828	Campanulaceæ 922	Cassava 302, 1018
Buck-wheat 991	Campanulate 380	Cassia 851
Bud, leaf 180	Campeachy-wood 851	Cassia-bark 994
—, floral 324	Camphor 994	Cassia Fistula 851

	Paragraph		Paragraph		Paragraph
Cassowary-tree	1035	Characeæ	1123	Clinandrium	421
Cassythæ	993	Characters of plants	711	Clinanthium	328
Castor-oil	1017	—, essential	711	Clove-pink	779
Casuarinæ	1033	—, natural	711	Clove-tree	867
Catechu	852	—, in the natural system	720	Clover	850
Catingas in Brazil	205	—, subordination in value		Cloves, young bulbs	194
Catkin	334	of	721	Club-moss family	1112
Caudate	541	Charcoal as a manure	246	Club-shaped	58
Caudex	65	Chelidonium	261, 756	Clusiaceæ	793
Caudicle	421	Chemical composition of		Cluster	333
Caulcescent	65	plants	208	Clustered	650
Caulicle	590, 593	Chenopodiaceæ	936	Coal	1183
Cauliculus	590	Cherry	855	Coal Epoch, climate of	1191
Cauliflower	762	Cherry-Laurel	855	Coal of Humus	234
Caulinary	160	Chervil	890	Coca	802
Cauline	168	Chestnut	1037	Cocci	533
Cauls	65	Chian Turpentine	843	Coculus	747
Caulopteris	1186	Chick-pea	63	Cochineal-Cactus	883
Cayenne-pepper	961	Chickweed family	778	Cochleariform	373
Cedar	1046	Chicory	918	Cockscomb	985
Cedrelaceæ	811	Chill-nettle	870	Coco	1093
Celandine	756	Chilies	961	Coco-nut Palm	1086
Celastraceæ	837	Chinese-grass	1020	Cocoa	786
—, Region of	1155	China-bark	903	Cocoinæ	1085
Celery	890	Chiretta or Chiraita	949	Coddington lens	1206
Cell-germ	22	Chive	1078	Codela	756
Cell-nucleus	16	Chienaceæ	791	Cælospermæ	888
Cells	4	Chloranthaceæ	1029	Coffee	905
—, absorption of fluids by	27	Chlorophylle	19, 316	Coir-rope	1086
—, agglutination of	13	Chocolate	786	Colchicæ	1079
—, annular	11	Chorization	383	Colchicia	1080
—, containing air	21	Christ's-thorn	841	Colchicum	1080
—, cortical	84	Chromatometer	671	Colcorhiza	127
—, deposits in	15	Chromogen	676	Collateral	472
—, development of	22	Chrysobalanæ	854	Collecting hairs	69, 453, 499
—, dotted	11	Chyrrus	1020	Collecting of plants	1202, 1229
—, epidermal	48	Cicatrices	104	Collenchyma	13
—, fibrous	11	Cicatricula	140	Collochia	953
—, forms of	5	Cichoraceæ	912, 914	Collum	119
—, functions of	27	Cicuta	893	Colocynth	872
—, nucleated and non-nucleated	22	Cilia	7, 657	Coloquintida	872
—, porous	10	Ciliated	7, 58	Colours of plants	671
—, spiral	11	Cinchona	903	— of epidermis	49
Cellulares	8, 1104, 1118	—, Region of	1155	— of flowers in different orders	684, 685
Cellular envelop	84	—, stipules of	160	—, changes in	683
— tissue	4	Cinchonæ	902	— in autumn	675
Cellules	4	Cinchonia	903	— in a garden	686
Cellulose	2, 14	Cinchyma	38	— prevailing in certain countries	1150
Centaur	949	Cinenchymatous	261	Colouring matters	322
Centrifugal	332	Cinnamon	994	Colpenchyma	5
Centripetal	331	Circinate	185	Coltsfoot	917
Cephælis	904	Circulation in Exogens	264	Columbine	373
Ceramiceæ	1128	— Endogens	265	Columella	532
Ceraxylon	1086	Circumscissile	411, 535	Columellifacæ	930
Cetraria	1121	Circumscription of leaf	139	Columna, column	400
Cetrarin	1121	Cirrhus	157, 201	Columnar	5
Cevadilla	1080	Cistaceæ	767	Combrétacæ	862
Chaillotiaceæ	1004	Cistoma	52	Commelynacæ	1087
Chalaza	465, 583	Citron	796	Commisüre	543
Chalk-fossils	1197	Citrus	796	Composite	910
Chalk-plants	1139	Cladenchyma	5	—, Region of	1155
Chamaeluciæ	866	Classes, defined	710	Compound Organs	43
Chamærops	1086	Classification, principles of	704	Compressorium	1210
Chamomile	917	Clathraria	1197	Conceptacles	1127
Chandeller-tree	1091	Clavate	58, 396	Concostate	144
Chara	1130	Claw	370	Condenser	1207
—, carbonate of lime in	227	Clearing-nut	947	Conducting Tissue	452
—, rotation in	276	Cleft	146	Conduplicate	185, 600, 601
		Clematidæ	739	Cone	334, 559
		Cleomeæ	763	Conenchyma	5
		Climate, continental	1136	Confervacæ	1128
		—, insular	1136	Conia	893
		Climate and season, their effects	1141		

Paragraph	Paragraph	Paragraph
Conical cells 5	Crape-paper 1021	Cysts 20
— root 124	Crassulaceæ 880	Cytinaceæ 1041
Conifera 1044	Cremocarp 533, 543	Cytoblast 22
Coniomyetes 1123	Crenata 146	Cytoblastema 14
Coniothalamææ 1120	Crescenticæ 950	Cytogenesis 22, 25
Conium 893	Cress 762	
Conjugation 492	Cress-wort family 758	
Connaraceæ 846	Crested 372	Dadoxylon 1184, 1190
Connate 163	Crisp 150, 374	Dædaleuchyma 5
Connective 405, 408	Crocus 1068	Daffodil 1071, 1073
Connivent 359	Croton-oil 1017	Dahlia 655
Continuous 140	Crowberry 1013	Dammara 1046
Contorted 124, 354	Crowfoot family 738	Danææ 1109
Contrayerva 1021	Crown-Imperial 1078	Dandelion 918
Convallaria 1077	Crown of the root 70	Daphne 1002
Convolute 185, 355	Crucifera 758	Daphnææ 1001
Convolvulaceæ 935	—, Region of 1155	Darnel-grass 1101
Copaifera 851	Cruciform or Cruciate cor- olla 378	Date-palm 1086
Copaiva 851	Cryptogamic Reproduc- tion 492	Date-plum 933
Copper in plants 229	Cryptogamous plants 323, 1104	Datisacææ 1012
Coquilla-nut 1086	—, Region of 1161	Datura 961
Coral-flower 850	Crystals in cells 18	Day-lily 1078
Corrallina 1129	Cubebin 1032	Deadly-nightshade 961
Coralline-root 125	Cubebæ 1032	De Candolle's System 728
Corculum 592	Cuckoo-pint 1093	Decandrous 394
Cordate 150, 374	Cucumber 872	Deciduous 366, 385
Cordiaceæ 957	Cucumites 1202	— Trees 205
Coriander 892	Cucurbitaceæ 871	Declinate 420
Coriariaceæ 834	Cudbear 1121	Decomound 152
Cork 6	Culm 65	Decurrent 163
—, cells of 6	Cultivation, its effects on flowers, &c 655	Decussate 169
Cork-oak 1037	Cumin 892	Deduplication 383, 654
Corky layer of bark 85	Cuneate 150	Definite Inflorescence 338
Corn 70, 195	Canoniææ 885	— Ovules 472
Cornogenæ 1105	Cupanoides 1202	— vascular bundles 91
Cornaceæ 896	Cupressinææ 1045	Degeneration of Organs 651
Cornel family 896	Cupula 347	Dehiscence 410, 530, 532
Corolla 530	Cupuliferææ 1033	Dehiscent fruits 530, 532, 546
—, forms of 368	Cureuma 1062	Delphinium 740
—, irregularity in 387	Curled 150	Dentate 146
—, irregular monopeta- lous or gamopetalous 381	Currant 884	Deodar 1046
—, irregular polypeta- lous 279	Currant of Anstralia 929	Deposits in cells 15
—, regular monopeta- lous or gamopetalous 380	Curvembryæ 960	Desmidiææ 1128
—, regular polypetalous 378	Cuscutææ 955	Desmodium, Movements of 663
Corollifloræ 390, 697	Cuspidate 372	Dextrin 302
Corolline hairs 61	Cuspis 372	Dhak-tree 850
Corona, crown 384, 942	Custard-apple 745	Diachænum 543
Corpuscles of Brown 510	Cutch, same as Catechu 45	Diachyma 137
Corsican Moss 1130	Cuticle 679	Diadelphous 399
Cortical Integument 92	Cuticle of Petals 369	Dialycarpous 538
Cortical System 82	Cyanic series of colours 679	Dianthus 394
Corylaceæ 1033	Cycadaceæ 1047	Diapensiææ 954
Corymb 333	— fossil 1197	Diastase 307, 310, 621
Corymbifera 912, 914	Cycadites 1197	Diataxis 704
Coryphineæ 1085	Cyclamen 975	Diatomaceæ 1128
Costa 144	Cyclanthææ 1090	Dichlamydeous 351
Costate, ribbed 8	Cycle 171	Dichotomous 99, 338
Cotton 8, 29, 782	Cyclogens 77	Diclinous 388, 648
Cotton-grass 1098	Cyclosis 261, 262	Dicotyledones 736
Cotyledons 168, 586, 590, 599	Cylindrenchyma 5	Dicotyledonous 71, 590
—, folding of 600	Cylindrical 5	Dictyogena 101, 1048
Cowbane 893	Cymbiform 373	Didymocarpeæ 950
Cowberry 928	Cyme 339	Didynamous 419
Cowitch or Cowhage 850	Cymose 339	Digestion of plants 233
Cow-plant of Ceylon 943	Cynarocephalææ 912, 914	Digger 1203
Cowslip 975	Cynarrhodum 541	Digitalform 380
Cow-tree 1022	Cyperaceæ 1097	Digitalin 965
Cow-tree of Demerara 945	Cypress 1046	Digitalis 965
Cow-wheat 965	Cypella 542	Digitate 148
Cranberry 928	Cyrtandrea 950	Digiti-partite 148
Cranes-bill 816	Cystidia 1122	Dilamination 383

- | | <i>Paragraph</i> | | <i>Paragraph</i> | | <i>Paragraph</i> |
|--|------------------|---|------------------|---|------------------|
| Dimidiate | 1114 | Duramen | 78 | Endosmometer | 254 |
| Dinomic | 1151 | Durian | 784 | Endosmose | 27, 254, 255 |
| Dioecious | 389, 648, 713 | Durmast | 1037 | Endosperm | 587 |
| Dionæa | 664, 771 | Durra | 198 | Endospermic | 587 |
| Dioscoreaceæ | 1051 | Dust-brand | 694 | Endostome | 464 |
| Diosmeæ | 827 | Dutch Reed | 226 | Endothecium | 402 |
| Dipetalous | 375 | Dutch Rushes | 1167 | Enceandrous | 394 |
| Dipleciolebas | 759 | Dwale | 961 | Ennobling fruit-trees | 574 |
| Diplod | 137 | Eagle-wood | 1003 | Ensiform | 150 |
| Diplostemonous | 392 | Earcocks | 700 | Entangled | 5 |
| Dipsacaceæ | 908 | Earth-nut, or Pig-nut | 890 | Entire | 146 |
| Dipterocarpaceæ | 789 | Ebenaceæ | 933 | Entophytic | 696 |
| Disa | 1058 | Ebony | 78, 850, 933 | Envelopes of the flower | 351 |
| Dischidia | 943 | Ebracteated | 845 | Eocene | 1180 |
| Disciform | 75 | Egg-apple | 961 | Epacridaceæ | 929 |
| Discoid | 75, 911 | Ehretieæ | 998 | —, Region of | 1155 |
| Discostate | 144 | Elæagnaceæ | 999 | Epiblema | 47, 120 |
| Discs | 80 | Elæocarpæ | 787 | Epicalyx | 364 |
| Diseases of plants | 689 | Elæodendreæ | 837 | Epicarp | 524 |
| — prevention of | 699 | Elat-rium | 872 | Epichilium | 1058 |
| — caused by fungi | 693 | —, bursting of cap-
sule of | 27 | Epidermis | 45 |
| — caused by insects | 700 | Elaters | 1113 | —, appendages of | 57 |
| Disk | 428 | Elatinaceæ | 777 | —, cells of | 48 |
| Dissected | 148 | Elder | 901 | —, colour of | 49 |
| Dissection of plants | 1238 | Elecampane | 917 | —, functions of | 64 |
| Dissemination of plants | 1142 | Electricity in germination | 614 | —, papillæ of | 57 |
| — agents employed | 1142 | Elemi | 845 | —, silica in | 49 |
| Dissepiment | 439 | Elephant's-foot | 1052 | —, texture of | 49 |
| — spurious | 447 | Ellipsoidal | 5 | —, wax on | 49 |
| Dissilent | 533 | Elliptical | 150 | Epigeal | 629 |
| Distichous | 170 | Elm | 1020 | Epigynous | 390, 449 |
| Distractile | 409 | Emerginate | 150, 372 | Epipetalous | 390 |
| Distribution of plants over
the globe | 1146 | Embryo, acotyledonous | 591 | Epiphleum | 82, 85 |
| — endemic | 1143 | —, development of | 502 | Epiphragm | 1114 |
| Dithecal | 405 | —, dicotyledonous | 597 | Epiphytes | 121, 251, 1139 |
| Dittany | 828 | —, direction of its
parts during germination | 624 | Epipollæ | 281 |
| Divergent | 359 | —, formation of | 586 | Epirrhæology | 1132, 1134 |
| Divi-divi | 851 | —, growth of | 622 | Episperm | 578, 579 |
| Divisions of Vegetable
Kingdom | 723 | —, monocotyledon-
ous | 595 | Epithelial | 463 |
| Dodder | 252, 956 | —, parts of 590, 592, 593 | 593 | Epithelium | 431 |
| Dodecahedral | 5 | —, polycotyledonous | 598 | Equal | 146 |
| Dodecandrous | 394 | —, position in the
seed | 603 | Equisetaceæ | 1106 |
| Dogbane family | 944 | —, pseudo-monoco-
tyledonous | 598 | Equisetum, fossil | 1197 |
| Dog-tooth Violet | 1078 | —, fixed | 181 | —, silica in | 226 |
| Dolabriform | 150 | Embryo-buds | 196 | Equitant | 186 |
| Doom-palm | 99, 1084 | Embryo-sac | 463 | Erect | 408 |
| Dorsal-suture | 433 | Embryology | 501 | —, ovule | 471 |
| Dorsiferous ferns | 1103 | Embryonic vesicle | 507 | Ergot | 696, 1103, 1126 |
| Dorstenia | 1021 | Embryotega | 583 | Ericaceæ | 926 |
| Dotted cells | 11 | Emetin | 769, 904 | Eriocaulon | 1095 |
| — vessels | 35 | Empetraceæ | 1014 | Eriogonæ | 990 |
| Double coco-nut | 1086 | Emulsine | 310, 835 | Erythrophylle | 675 |
| Doublet | 1192, 1205 | Endemic distribution of
plants | 1145 | Erythroxyllaceæ | 802 |
| Dracæna | 1078 | Endive | 918 | Escallonææ | 685 |
| Dragon's-blood | 1086 | Endlicher's System | 730 | —, Region of | 1155 |
| Draining | 616 | Endocarp | 524 | Eschscholtzia | 366, 756 |
| Drifted plants | 1192 | Endochrome | 493 | Essences | 317 |
| Drims | 743 | Endogenæ | 1049 | Essential organs of the
flower | 351 |
| Droseraceæ | 770 | Endogenites | 1197 | Eterio | 545 |
| Drupaceæ | 854 | Endogenous, or monocoty-
ledonous stem 71, 91, 94, 107 | 107 | Etalology | 300 |
| Drupe | 545 | —, anomalous | 101 | Eucalypti, Region of | 1155 |
| Drupels | 545 | —, branching | 99 | Eucalyptus | 867 |
| Drying plants | 1206, 1233 | —, subterranean | 100 | Eunymæ | 837 |
| Dry-rot | 697, 1126 | Endophleum | 82, 83 | Eunymus | 838 |
| Duckweed | 1101 | Endopleura | 578, 580 | Euphorbiaceæ | 1014 |
| Ducts, closed | 34 | Endorhizal | 127, 628 | Euphorbium | 1016 |
| Dulcamara | 961 | | | Evening Primrose | 88 |
| Dulse | 1139 | | | Evergreen Oak | 1037 |
| Dumb-cane | 1101 | | | Exalbuminous | 587 |
| Dumose | 68 | | | Exannulate Ferns | 1108 |
| Dumus | 68 | | | Excentric | 603 |
| Durability of heartwood | 78 | | | Exhalation | 203 |
| | | | | Exintine | 422 |

<i>Paragraph</i>		<i>Paragraph</i>		<i>Paragraph</i>	
Exogenæ.....	736	Fimbriated.....	372	Fruits, arrangement of.....	560
Exogenous stem.....	71	Firs.....	1046	—, preserved.....	1239
Exogonium.....	956	Fissiparous.....	493	Frutex, shrub.....	63
Exorhizal.....	126, 629	Fistular.....	164	Fruticose.....	63
Exosmose.....	27	Flabellarla.....	1190	Fucaceæ.....	1128
Exostemma.....	903	Flacourtiaceæ.....	766	Fuchsia.....	868
Exostome.....	464	Flagellum.....	191	Fuller's Teazel.....	908
Exothecium.....	403	Flax.....	29, 818	Fumariaceæ.....	757
Expansion of flowers.....	331	—, New Zealand.....	29	Fumitory family.....	757
—, centrifugal.....	332	—, Pita.....	29	Fungi.....	1122
—, centripetal.....	331	Fleshy.....	583	—, causing diseases.....	1126
Exserted.....	417	Floral axes.....	325, 327	—, edible.....	1124
Exstipulate.....	158	Floral calendar.....	481	—, how to prepare.....	1237
Extline.....	422	— clock or watch.....	483	—, in cutaneous disease.....	1126
Extra-axillary.....	196	Floral Envelopes.....	351	—, on caterpillars.....	1126
Extrarius.....	602	—, development of.....	386	—, phosphorescent.....	1125
Extrorse.....	412	—, functions of.....	473	—, poisonous.....	1124
Eye-bright.....	965	Floral leaf.....	324, 344	Fungus melitensis.....	1103
Eye-piece.....	1207	Florets.....	382	Funiculus.....	462
Eyes of potato.....	70	Florideæ.....	1128	Fusiform.....	5, 29, 124
		Flosculus.....	911	Fustic, morus tinctoria.....	1021
Fabaceæ.....	847	Flower.....	3 0		
Fairy Rings.....	1125	—, position of its parts.....	357	Galaciæ.....	880
Fall of the leaf.....	205	Flower-buds.....	324, 354	Galactodendron.....	1022
False-bark.....	92	Flowering, period of.....	478	Galangale.....	1062
— Trachææ.....	35	—, mode of accelerat- ing.....	479	Galanthus, Snow-drop.....	1072
Falsely-discoid.....	911	Flowering-ash.....	941	Galbanum.....	891
— radiate.....	911	Flowering plants.....	823	Galbulus.....	559
Fan Palm is Corypha.....		Flowering-rush.....	1089	Gale.....	1035
Farnaceous.....	588	Flowerless plants.....	343	Galeate.....	361
Fasciated.....	197, 327	Fluids of plants, special movements of.....	275	Galiææ.....	902
Fascicle.....	339	Foliaceous.....	359	Galipea.....	823
Fascicled.....	650	Folliola.....	145, 151, 358	Galls.....	701, 1037
Fasciculate.....	175	Follicle.....	547	Gama-grass.....	1100
Fasciculated.....	114, 125	Fool's Parsley.....	893	Gamass or Squamash.....	1078
Fasculation of branches.....	190	Foramen.....	464	Gambeer.....	905
Faux.....	377	Forked.....	406	Gamboge.....	799
Feather-grass is Stipa pen- nata.....		Fossil Botany.....	1174	Gamogastrous.....	435
Feather-veined.....	144	—, plants.....	1174	Gamopetalous.....	376, 380
Feathery.....	365	—, their structure.....	1175	Gamophyllous.....	358
Fecundation.....	496	—, state of preservation.....	1175	Gamosepalous.....	358
Fenestrate.....	138, 555	—, arrangement of.....	1176, 1177	Gangrene.....	692
Fennel.....	890, 892	—, their number.....	1181	Garlic.....	1073
Fenugreek, a species of Trigonella.....		— of Carboniferous system.....	1186	Garryacææ.....	1040
Ferns.....	1108	— of Secondary strata.....	1197	Gases, their effects.....	292
—, fossil.....	1186	— of Tertiary strata.....	1201	—, irritant.....	293
—, during coal epoch.....	1191	Fossiliferous strata.....	1179	—, narcotic.....	293
—, how to prepare.....	1235	Forbes Marine algæ.....	1165	Gasteromycetes.....	1123
Fertile flowers.....	649	Forbidden-fruit.....	796	Gasterothalamææ.....	1120
Fertilization.....	496	Forget-me-not.....	959	Geissolomææ.....	10 0
Fescue-grass.....	198	Fovilla.....	423	Gemination.....	184
Feverfew.....	917	—, its use.....	504	Gemmule.....	590, 593
Fibre in cells.....	11	Foxglove.....	965	Genera, defined.....	708
—, in spiral vessels.....	34	Francoeææ.....	880	Geniculate.....	397
—, woody.....	29	Frankeniaceæ.....	776	Gentianacææ.....	948
Fibrine, vegetable.....	310	Frankincense.....	1046	Gentian.....	949
Fibrous cells.....	11	Fraxineææ.....	940	Geo-Botany.....	1174
— root.....	125	French Berries.....	841	Geoffroya.....	850
Fibro-cellular Tissue.....	11	Friar's-balsam.....	932	Geography of plants.....	1131
Fibro-vascular Tissue.....	31	Fringed.....	372	— Influence of external agents on.....	1122
Fleoideæ.....	881	Fritillary.....	1078	— of temperature.....	1135
Ficus.....	1021	Frogbit Family.....	1057	— of moisture.....	1138
—, Region of.....	1162	Frond.....	1108	Geraniaceæ.....	815
Field-book.....	1229	Fruit.....	520	Germen.....	429
Fig.....	1021	—, changes during its formation.....	522	Germinal vesicle.....	510
Fig-wort.....	965	—, parts which form it.....	520, 527	Germination.....	608
Filament.....	395	—, maturation of.....	561	Germination, acotyledon- ous.....	629
—, adhesion of.....	399	—, how accelerated.....	563	—, chemical changes during.....	621
—, form and structure.....	396	Fruits, analysis of.....	566	—, dicotyledonous.....	629
Filbert, see Hazel.....				—, length of time re- quired for.....	632
Filices.....	1108				
Filiform, thread-like.....	453				

	Paragraph		Paragraph		Paragraph
Germination, monocotyledonous	627	Guano, British	245	Helleboreæ	739
—, progress of	622	Guarana	807	Helleborus	740
—, requisites for	608	Guava	867	Helwingiaceæ	1040
Gesneraceæ	925	Guelder Rose	901	Hemerocallidæ	1077
Gibbous	373	Guernsey Lily, <i>is Nerine</i>		Hemicarp	543
Gilliesiaceæ	1081	<i>sarniensis</i>	1037	Hemidesmus	943
Gills	1122	Gunfweed	1129	Hemlock	893
Ginger	1062	Gum	303	Hemp	1020
Ginko	1046	Gum-Arable	852	—, fibres of	29
Ginseng	895	Gum-Dragon	850	—, Indian	1020
Glabrous	60	Gum-lac	850	Henbane	961
Glands	62	Gum-tree of New Holland	867	Henfrey on Cytogenesis	23
—, lenticular	63	Gunjah	1020	Henna	858
—, nectariferous	63	Gutta percha	319, 917	Henslow's comparative	
—, sessile	62	Guttiferæ	798	view of different natural	
—, stalked	62	Gymnogens	732	systems	734
—, vesicular	63	Gymnospermæ	1043	Hepaticæ	1116
Glandular hairs	61, 62	Gymnospermous	576	Heterandrous	394
Glandular woody tissue	30	Gymnosperms, Reign of	1194	Herbaceous	359
Glans	514	Gynandrous	400	Herbarium	1208, 1230
Globe-amaranth	985	Gynizus	433, 458	— room for	1241
Globule, in Chara	427	Gynobase	452	Herbs, defined	68
Glochidiate	58	Gynecium	429	Hermaphrodite	388, 713
Glomerulus	336	Gynophore	437	Hernandiæ	1001
Glossology	704	Gynostegium	942	Hesperidium	552
Glucose	807	Gynostemium	400	Heterocephalons	911, 914
Glumacæ	1096	Gyrate	340	Heterochromous	910
Glumellæ	1099	Gyration	275	Heterodromous	173
Glumellulæ	1099	Gyrocarpæ	862	Heterogamous	911, 914
Glumes	349	Gyrogonites	1202	Heterorhizal	128, 629
Gluten	310	Habitat	713	Heterotropical	464
Glycyrrhiza	850	<i>Hæmatoxylon</i>	851	Hexagonal	5
Gnetaceæ	1045	<i>Hæmodoracæ</i>	1070	Hexagonienchyma	5
Goldfussia	971	Hairs	57	Hexandrous	394
Gongyli	1119	—, circulation in	64	Hiatus	381
Gonidia	1119	—, collecting	59, 64	Hibernacula	182
Goodeniaceæ	920	—, corolline	61	Hibiscæ	781
Gooseberry	884	— of cotton	59	Hickory-nut	1069
Gossypium	782	— of cowitch	58	Hiigtea	1202
Gourd	872	—, functions of	64	Hilum	462, 583
Graft	569	—, glandular	61	Hippocrateacæ	801
Grafting	569	—, irritable	58	Hirsute	60
views of Knight		—, irritable	58	Hirtus	60
—, different kinds of	571	—, lymphatic	61	Hispid	60
—, its effects	572, 573	—, petaline	370	Histology	1199, 1212
Grains, cereal	1106	—, radical	61	Hog-plum	843
Grains of Paradise	1062	—, ramentaceous	57	Holly	935
Gramineæ	1099	—, retractile	59	Hollyhock	782
Grape-sugar	307, 814	Hairy	60	Homaliacæ	1006
Grasses	1099	Halbert-shaped	150	Homochromous	918
—, Region of	1161	Half-equitant	186	Homodromous	173
—, for Pasture	198	Halophytes	228, 987	Homogamous	911, 914
—, fistular stem of	99	Haloragacæ	869	Homotropical	603
Grass of Parnassus	771	Hamamelidacæ	887	Honeysuckle	901
Grass-tree	1078	Hand-plant	784	Hooked	58
Green colour, effects of		Hare-bell family	922	Hop	1020
light on	672	Hare's foot fern	1110	Horehound	967
Green colouring matter of		Haschisch	1020	Hornbeam	1037
plants	19	Hastate	150, 374	Hornwort family	1023
Greenhearted-tree	994	Hazel	1087	Horse-chestnut	807
Green snow	1129	Heart-shaped	150, 374	Horse-radish	762
Greffé des Charlatans	569	Heart's-ease	769	Horse-radish tree	853
Grenadilla	876	Heart-wood	78	Horse-tail family	1106
Grevillea	998	Heat during flowering	475	Hottentot's fig	881
Grossulariaceæ	884	Heath	927	Houseleek	980
Ground-nut	850	Heather	927	Hoya	843
Guaco	917	Hebradendron	799	Huanuco-bark	903
Guaiacum	826	Hedge-hyssop	965	Humulus	1020
Guano	243	Helicoidal	340	Humus	212, 234
—, analysis of different		Helictæcæ	783	Hungarian balsam	1046
kinds	244	Heliotrope	959	Hura	1016
		Heliotropiæ	958	Hyacinth	1073
		Hellebore, white	1080	Hyaline of Barry	16, 22

[illegible]

	Paragraph		Paragraph		Paragraph
Leaves, anomalous forms of	162	Ligulate	382, 911	Madder	905
—, arrangement on the		Ligule	161	Madhuca tree	937
—, stem	169	Ligulifloræ	913	Madia	917
—, cauline	168	Lilac	941	Magnoliaceæ	742
—, compound	151	Liliaceæ	1077	Magnolias, ilegion of	1155
—, deciduous	140	Lilium	1078	Mahogany	812
—, divergence of	171	Lily of the Valley tribe	1077	Maiden-hair fern, is Adiantum	1110
—, effects on the air	284	Limb of leaf	139	Maize	1142
—, evergreen	140	— petal	370	Malesherbiæ	875
—, fall of	205	Lime-fruit	796	Male-shield fern	1110
—, fistular	164	Lime in plants	227	Malicorium	550
—, floral	168, 824	—, in soils	237	Mallow family	781
—, forms of	145	—, phosphate of, as a manure	216	Malombo bark	832
—, functions of	203	Lime-tree family	787	Malpighiaceæ	803
—, in acotyledons	167	Linacææ	817	Malvaceæ	781
—, in monocotyledons	166	Linaria	965	Mammee Apple	799
—, in dicotyledons	165	Linden-tree	788	Manchineel	1016
—, influence on the circulation	271	Linley's system	731	Mandragora	961
—, primordial	168	Linear	150, 374	Mandrake	961
—, radical	168	Linen	29	Manganese in plants	229
—, ramal	168	Ling	927	Mangifera	843
—, seminal	168	Linnea	901	Mango	843
—, simple	145	Linnæan system	715	Mangold-wurzel	987
—, skeleton	135	—, its Imperfections	718	Mangosteen	799
—, spiral arrangement	170	Linseed	818	Mangrove	860
—, submerged	138	Lip	379	Manihot	1018
—, useful in forming wood	117	Liquidambar	1035	Manioc	1018
Lecotropical	466	Liquorice	850	Manna	867, 941, 1044
Lecythidææ	866	Lirellæ	1119	— of Mount Sinai	775
Leechee, see Litchi.		Litchi fruit	807	Manna-Ash	941
Leek	1078	Litmus	1121	Manna-Sugar	306
Legume	548	Lizard's-tail family	1030	Mannite	306, 941
Legumine	310	Loasacææ	870	Mantellia	1187
Leguminosæ	847	Lobed	148	Manures, active Ingredients of	243
—, geographical distribution of	848	Lobeliacææ	923	—, ammoniacal	243
Lemon	796	Localities or stations of plants	1139	—, application of	240
Lemon-grass	1102	Loculaments	440	—, comparative value of	242
Leises	1295	Loculicidal	532	—, experiments with	249
Lentibulariaceæ	972	Locust-tree	850, 851	—, liquid	248
Lenticels	63	Locusta	334	—, natural	241
Lenticular glands	63	Lodiceulæ	1099	—, simple	241, 246
Leopard's bane	917	Loganiacææ	946	Manuring with green crops	247
Lepidocaryinæ	1085	Logwood	851	— sea weeds	247
Lepidodendron	1187	Lomentacææ	760	Maple	805
Lepidophyllum	1187	Lomentaceous	548	Maple-sugar	306
Leptospermææ	866	Lomentum	548	Marantacææ	1063
Lepidostrobis	1187	Longan fruit	807	Maraschino	855
Lepidote, scaly	57	Lonicerææ	900	Marcescent	367
Lepia, a scale	57	Loosestrife	858	Marcegraviacææ	800
Letter-wood	1022	Loquat	855	Marchantiææ	1117
Lettuce	918	Loranthacææ	898	Mare's-tail family	869
Lianas, or Lianes	66	Loranthæ, same as Lesta.		Marginal placenta	438
Liber	83	Lotus	841	Marginate calyx	364
Lichenes	1119	Lotus of Egypt	753	Marine plants	1139
Lichenin	892, 1121	Love-apple	961	Marine vegetation, zones of	1163
Lichens	1119	Love-lies-bleeding	985	Maritime plants	1139
—, how to prepare	1125	Lucern	830	Marjoram	967
—, Region of	1162	Luminosity of plants	669	Marking nut	843
Lid, see Operculum.		Lupulin	1020	Marmalade fruit	937
Life of plants, its duration	633	Lychnophora	917	Marsh-Mallow	782
Light, as affecting plants	1140	Lycopode	1113	Marsh-Trefoil	949
—, effects on flowers	496	Lycopodiaceæ	1112	Marsileacææ	1111
—, effects on respiration	284	Lycopodites	1187	Marvel of Peru	983
—, effects of different rays	290	Lysinodendron	1190	Masked, see Personate.	
—, effects of different rays on the colour of plants	672, 673	Lymphatic hairs	61	Mastic	843
Lign aloes	1003	Lyrate	147	Maté	935
Ligneous tissue	29	Lythracææ	837	Matico or Matica	1032
Lignine	15, 308	Macahuba palm	1086	Maturation of fruit	561
Lignum-vitæ, same as Gualacum.		Mace	581, 596	—, changes during its progress	563
		Macropodus	595		

- | | Paragraph | | Paragraph | | Paragraph |
|-----------------------------|-----------|-----------------------------|--------------------|------------------------------|---------------|
| Mattulla..... | 57, 36 | Mimosites | 1202 | Mummy-wheat | 617 |
| May-Apple | 740 | Mimulus | 965 | Munjeet | 905 |
| Meadow plants | 1139 | Mint | 967 | Muriatic acid gas, effect on | |
| Meadow-saffron | 1089 | Miocene | 1180 | plants | 295 |
| Mealy | 588 | Miostemonous | 392 | Muriform cells | 89 |
| Meconic Acid..... | 756 | Misletoe | 252, 899 | Muriform cellular tissue.. | 5 |
| Medick | 850 | Mitriform | 1114 | Musacææ | 1065 |
| Medlar | 855 | Mordeceæ | 873 | Muscadine | 1126 |
| — of Surinam | 937 | Mohl on Cytogenesis | 22 | Musci | 1114 |
| Medullary Rays or Plates | 89 | Moisture, as affecting the | | Mushroom | 1124 |
| —, position of | 72 | distribution of plants .. | 1133 | Mustard | 762 |
| —, functions of | 109 | Molluginæ | 778 | Mustard-tree | 987 |
| Medullary Sheath | 76 | Momordica | 872 | Muticous, without an awn | 1099 |
| —, its functions | 109 | Monadelphous | 389 | Mycellum | 1122 |
| Melostemonous, see Mio- | | Monandrous | 394 | Myoporineæ | 968 |
| stemonous. | | Monembryony | 585 | Myosotis | 959 |
| Melaleuca | 897 | Moniliform | 393 | Myrica | 1033 |
| Melanthiaceæ | 1079 | — hairs | 57 | Myristicacææ | 995 |
| Melastomacææ | 863 | — vessels | 37 | Myrobalans | 872 |
| —, Region of | 1155 | — roots | 125 | Myronic acid | 762 |
| Melogneta pepper | 1062 | Monimiacææ | 1026 | Myrosine | 762 |
| Meliacææ | 869 | Monkey-bread | 784 | Myrrh | 845 |
| Melilot | 830 | Monkey-pot | 867 | Myrsinacææ | 938 |
| Mellocææ | 878 | Monkshood | 740 | Myrtacææ | 866 |
| Melon | 872 | Monk's Rhubarb | 991 | —, Region of | 1162 |
| Memecylcææ | 863 | Mnocarpicææ | 634 | Myrtle | 867 |
| Meniscus | 1205 | Monochlamydeæ | 980 | Nabæe | 740 |
| Menispermacææ | 746 | Monochlamydeous | 351 | Naegeli on Cytogenesis .. | 24 |
| Mentha | 967 | Monoclinous | 648 | Naladacææ | 1094 |
| Menyanthes | 948 | Monocotyledonous | 590, 1047 | Napiform | 124 |
| Merencilyma | 5 | Monocotyledonous stem | 71, 91 | Narcissææ | 1072 |
| Mericarp | 543 | Mono-Cryptogamææ | 1104 | Narcotine | 756 |
| Merismatic | 493 | Monocneous | 389, 618, 713, 911 | Narthex | 891 |
| Merithal | 169 | Monugamia | 717 | Naseberry | 937 |
| —, laminar | 139 | Monomic | 1151 | Natural System | 719 |
| —, petiolar | 139 | Monopetalous | 376, 380, 381 | Necklace-like | 398 |
| —, tigellary | 139 | Monophyllous | 358 | — vessels | 37 |
| —, vaginal | 139 | Monosepalous | 358 | Nectandra | 994 |
| Merithall | 639 | Monothecal | 405 | Nectarics | 383, 416, 428 |
| Mesembryacææ | 881 | Monotropææ | 926 | Nectariferous glands | 63 |
| Mesembryanthema, Region | | Monstrosities, causes of .. | 646 | Nectarine | 855 |
| of | 1155 | Moon-seed family | 746 | Nelsoniææ | 970 |
| Mesembryanthemum | 881 | Mora wood | 851 | Nelumbicææ | 752 |
| Mesocarp | 524 | Moreæ | 1019 | Nepenthecææ | 1011 |
| Mesochilum | 1058 | Morel | 1124 | Nerium | 945 |
| Mesophlæum | 82, 84 | Morinda | 905 | Neroli oil | 795 |
| Mesophyllum | 137 | Morindine | 905 | Nervation | 141 |
| Mesosperm | 578 | Moringacææ | 853 | Netted | 113 |
| Metamorphosis | 641 | Morphia | 756 | Nettle | 1020 |
| Meteor flowers | 484 | Morphology | 641 | Nettle-tree | 1020 |
| Mezereum | 1002 | Morphology of Mosses .. | 1114 | Neur-pteris | 1186 |
| Micrometer | 1200 | Mosses | 1114 | New Zealand Flax | 29, 1078 |
| Micropyle | 464, 583 | —, how to prepare | 1235 | — spinach | 681 |
| Microscope | 1206 | —, Region of | 1155 | Nicker-tree | 851 |
| —, compound | 1207 | Movements in sea weeds .. | 657 | Nicotiana | 961 |
| —, simple | 1206 | — in sensitive plants | 659 | Nicotina | 961 |
| —, focal adjust- | | Moving-plant, see Desmo- | | Night-flowering Cereus .. | 883 |
| ment of | 1217 | dium. | | — plants | 484 |
| Microscopes, how to use .. | | Mosa | 917 | Night-scented flowers | 637 |
| 1212, 1214, 1215, 1216 | | Mucilage | 303 | Nightshade | 961 |
| Microscopic apparatus .. | | Mucronate | 150 | Nilsonia | 1197 |
| 1208, 1210 | | Mucus, definite | 46 | Nipadites | 1202 |
| Microscopic objects, pre- | | Mudar | 943 | Nitrates as manures | 246 |
| sentation of | 1226 | Mulberry | 1021 | Nitrogen in plants | 215 |
| Microscopic preparations | 1225 | Mullein | 965 | —, sources of | 215 |
| Microscopic re-agents | 1213 | Multicostate | 144 | Nodes | 67, 129 |
| Midrib | 141 | Multifid | 148 | Nodulose or Nodose | 125 |
| Mignonette family | 765 | Multifingate | 153 | Nöggerathia | 1190 |
| Mildew | 695 | Multilocular | 440, 522 | Nomenclature | 704, 712 |
| Milk-tree | 945 | Multiple fruits | 538, 556 | Non-fossiliferous rocks .. | 1178 |
| Milk-vessels | 38 | Multiplication of parts of | | Notfolk Island Pine | 1046 |
| Milk-wort family | 772 | flowers | 653 | Notorhizææ | 601, 759 |
| Millet | 1240 | Multivalvular | 580 | Noyau | 855 |
| Mimoseæ | 848 | Mummy-cloth | 29 | | |

Paragraph	Paragraph	Paragraph
Nuc'coll..... 16	Orange family..... 795	Palæozoic plants..... 1186
Nucleus of a cell..... 16, 22	Orbicular..... 149	Palate..... 381
Nucleus or kernel..... 577	Orchidaceæ..... 1038	Palææ of Compositæ..... 348
Nucleus of ovule..... 463	Orchidaceous corolla..... 379	Palææ of grasses..... 1099
Nuculanum..... 553	Orchil..... 1121	Palm-stein, formation of..... 94
Nucule..... 459	Orders, defined..... 709	——, dichotomous..... 99
Nucumetacææ..... 760	Organic constituents..... 218	——, direction of vascular bundles of..... 95
Number of known plants..... 1133	——, tabular view of..... 208	——, mode of increase of..... 97
Nut..... 544	——, sources of..... 212	——, rings of..... 98
Nutmeg..... 993	Organic elements..... 208	Palm of the Bible..... 1086
Nutrition of plants..... 206	Organogeny..... 442	Palma Christi..... 1017
—— of air plants..... 217	Organography or Structural Botany..... 1	Palmacites..... 1202
Nutritive organs..... 43	Organs, appendicular..... 57	Palmææ..... 1085
—— products..... 312	——, compound..... 2, 43	——, Region of..... 1155, 1162
Nux-Vomica..... 947	——, elementary..... 1	Palmate..... 148
Nyctaginacææ..... 982	——, essential of the flower..... 351	Palmatifid..... 148
Nymphæacææ..... 750	——, essential, their functions..... 488	Palmitt..... 1017
Oak..... 1037	——, nutritive..... 43	Palm-oil..... 1086
Oak-apples..... 701	——, reproductive..... 43, 323, 383	Palo De Vaca..... 1022
Oak-spangles..... 701	Original vegetation of the globe..... 1146	Panax..... 895
Oats..... 1142	Ornus..... 941	Pandanacææ..... 1090
Obcordate..... 150, 372	Ornubanchacææ..... 962	Pandanocarpum..... 1203
Object-glass..... 1207	Orris-root..... 1068	Panduriform..... 147
Oblique..... 146, 372	Orthoplocææ..... 601, 759	Panglææ..... 873
Oblong..... 150	Orthospermææ..... 888	Paniclææ..... 333
Oblong symmetry..... 644	Orthotropææ..... 465, 584	Pansy..... 769
Obovate..... 150	Orthotropus..... 465	Papaveracææ..... 755
Obsolete..... 264	Oryctology..... 1174	Papaw..... 873
Obvolute..... 186	Oryza, see Rice.....	Papayacææ..... 873
Ochnacææ..... 833	Oscillatoria..... 1129	Paper for drying..... 1231
Ochrea..... 158	Osier is Salix viminalis.....	——, for Herbarium..... 1234
Ochro..... 782	Osundææ..... 1109	Paper-mulberry..... 1021
Ochthous..... 294	Otopteris..... 1197	Papilionacææ..... 848
Octangular..... 69	Ouaris..... 947	Papilionaceous corolla..... 379
Odours of flowers..... 687	Oval..... 5, 150	Papillæ of epidermis..... 57
—— in different natural orders..... 688	Ovary or germen..... 429, 431	Pappus..... 365
—— in connection with colour..... 688	——, divisions in it..... 439	Papyrus..... 75, 1096
Oenanthe..... 693	——, half-inferior..... 449	Paracorolla..... 384
Offset..... 191	——, inferior or adherent..... 449	Paraguay Tea..... 935
Oil, cells containing..... 20	——, superior or free..... 449	Paraphyses..... 1114
Oils, fat..... 314	Ovate, egg-shaped..... 150	Parasites..... 252
——, drying..... 314	Ovenchyma..... 5	——, nutrition of..... 252
——, fixed..... 314	Ovular tubes..... 505	——, on roots..... 252
——, in different seeds..... 315	Ovule..... 488, 460	Parasitic Fungi..... 252, 1126
——, solid..... 314	——, ascending..... 471	——, plants..... 1139
——, volatile or essential..... 317	——, coverings of..... 464	Parastemonææ..... 384
Oleacææ..... 794	——, development of..... 463	Parcira-brava..... 747
Oleæ family..... 794	——, erect..... 471	Parenchyma..... 4, 5
Oleacææ..... 940	——, forms of..... 465	——, imperfect..... 12
Oleander..... 945	——, horizontal..... 471	——, perfect..... 12
Oleaster family..... 999	——, inverted..... 471	Parietal..... 440
Olibanum..... 845	——, naked..... 461	Parietalin..... 1121
Oligandrous..... 394	——, pendulous..... 471	Pariglin..... 1054
Oligospermous..... 546	——, position in the ovary..... 471	Parl-plinnate..... 153
Olive..... 941	——, seminude..... 461	Paris..... 1055
Omphalode..... 583	——, suspended..... 471	Parnassiacææ..... 770
Onagracææ..... 868	Oxalate of lime in raphides..... 18	Paronychiacææ..... 879
Onion..... 1078	Oxalidacææ..... 820	Paropsacææ..... 875
Oolitic Fossils..... 1197	Oxidation in flowers..... 473	Parsley..... 690
Oophordium..... 1112	Oxlip..... 975	Parsnip..... 890
Operculate..... 366, 425, 595	Oxycoecus..... 928	Partito or Parted..... 146
Operculum..... 1114	Oxygen in plants..... 213	Passifloracææ..... 875
Ophelia Chirata..... 949	——, sources of..... 214	Passion-flower..... 876
Ophlocaryon..... 807	——, consumed by flowers..... 474	Pastilles..... 932
Ophloglossææ..... 1109	Oyster-plant..... 959	Pastinaca..... 890
Opium..... 756	Oyster-plant of America..... 918	Patchouli..... 967
Opoponax..... 891	Pæoniææ..... 739	Patellææ..... 1119
Opposite..... 169	Pagina of leaf..... 139	Patulous..... 359
Opuntia..... 883	Palæontology..... 1174	Pea..... 850
Orach..... 987	Palæozamia..... 1197	Peach..... 855
Orange..... 796		Pear..... 855
		Pecopteris..... 1186

	Paragraph		Paragraph		Paragraph
Pectine	304, 564	Petiole, winged.....	157	Pistachio nut	843
Pectinate	51, 147	Petionles	151	Pistacia	843
Pedalicæ	950	Petiveriæ	988	Pisticæ	1032
Pedate	148	Pence	1197	Pistil	350, 429
Pedatifid	148	Phænogamous or Phanero-		—, formation of.....	430
Pedicel	325	gamous	323	—, various forms of.....	434
Pedicellate	325	Phanerogamous plants, ar-		Pistillary cord.....	438, 460
Peduncles	325	rangement of their organs	638	Pistillate	649
Pedunculate	325	Philadelphaceæ	865	Pistillidia	459, 492
Pelargonium	816	Phleboid tissue	37	Pistilliferous	388, 649
Pellicle, superficial ..	45	Phloridzine	309	Pita-flax	29, 1073
Pellitory	1020	Phlox family	952	Pitaya-bark	903
Pellitory of Spain.....	917	Phoenix	1086	Pitch	1046
Peloria	965	Phoranthium	326	Pitcher-plant family ..	1009
Pelorzation	654	Phormium	1078	Pitchers	164
Peltate	53, 149, 453	Phragmata	447	Pith	8, 72, 75
Pennisæceæ	1000	Phylla	358	—, discoid or disciform..	75
Pencil-cedar	1046	Phyllocladum	157	—, excentric	80
Pendulous ovule	471	Phyllogen	96	—, functions of	109
Penninerved, see Feather-		Phyllod	327	Piton-bark	903
veined	144	Phyllophor	96	Pitted tissue	26
Penny-royal	967	Phyllostic	177	Pittosporaceæ	823
Pentagonal symmetry ..	643	Phyllotaxis	169	Placenta	48
Pentamerous	643	—, in Acrogens	176	—, how formed	438
Pentandrous	394	—, in Eudogens	176	Placentaries	433
Pentapetalous	375	—, in Exogens	176	Placentation	443
Pentaphyllous	361	Physalis	961	Plaited	185
Pentasepalous	361	Physic-nut	1017	Plane, oriental	1066
Pepo	551	Physiognomy of Vegeta-		Plantaginaceæ	978
Peponida	551	tion	1150	Plantain	1066
Pepper	1032	Physiology or Physiologi-		Planting trees	118
Pepper-brand	693	cal Botany	1	Plants, dissection of....	1238
Pepper-corn	700	Phycomycetes	1123	Platanaceæ	1033
Peppermint	967	Phytclephas	1086	Pleiotracheæ	32
Pepperwort family	1111	Phyto-geographical divi-		Pleistocene	1202
Perenchyma of Morren ..	17	sion of the globe	1152	Pleurenchyma	29
Perennial	633, 713	— zones		—, functions of	41
Perfoliate	163	of Meyen	1157	Pleurisy-root	913
Perforations in cells....	9	Phytolaccaceæ	988	Pleurocarpl	1115
Perianth	352	Phytos	116, 181, 639	Pleurorhizæ	601, 759
Pericarp	520	Phytozoa	427, 492	Plicate	185
Perichætal	1114	Piassaba	1086	Pliocene	1180
Pericladium	158	Picotee	779	Plum	853
Periderm	85, 87	Picræna	832	Plumbaginaceæ	976
Perigone	352	Picrotoxin	747	Plumose	365
Perigynium	1097	Pileorrhiza	120	Plumule	590, 593
Perigynous	390, 449	Pig-nut, see Earth-nut.		Pod	548
Periodicity in flowering..	483	Pileus or cap	1122	Podetium	1119
Peripheral	602	Pill	57	Podocarp	533
Perisperm	577	Pilose	365	Podogonium, same as Gy-	
Perispermie	587	Pilosus	60	nophore	437
Perispore	591	Pimento or Pimenta ..	867	Podosperm	462
Peristomatic	51	Pimpernel, see Anagallis.		Podostemaceæ	1024
Peristome	1114	Pimpinella	892	Poison-elder	843
Peritropous	471	Pinaceæ	1044	Poison-ivy	843
Periwinkle	945	Pine-apple	1076	Poison-nut	947
Pernambuco-wood	851	Pinel-tallow	790	Poison-oak	843
Persimmon	933	Pinenchyma	5	Poison-sumach	843
Personate	381	Pines	1046	Poison-vine	843
Perulæ	142	Pinguicula	973	Poisons, their effects on	
Peruvian Bark	903	Pinites	1190	irritability	666
—, Cherry	961	Pink	779	Polarizing apparatus ..	1210
Petaloid	359, 396	Pink-root	947	Polemoniaceæ	952
Petaloidæ	1056	Pinna	152	Pollard-trees	190
Petals	368	Pinnate	152	Pollen	395, 402, 421
—, adhesion of	376	—, alternately	153	—, application to the	
—, metamorphosed	383	—, equally or abruptly ..	153	stigma	494
Petiolar	160, 201	—, interruptedly	153	—, development of....	421
—, merithal	139	—, unequally	154	—, duration of vitality	
Petiolate	62, 139	Pinnatifid	147	of	500
Petiole	159, 155	Pinnatifid	147	—, scattering of....	496
—, anomalous forms of ..	163	Pilus	1046	Pollen-grains, coverings of	422
—, inflated	157	Lipercaceæ	1031	—, number of....	494
—, sheathing	158	—, Region of	1155	—, union of....	421

	Paragraph		Paragraph		Paragraph
Pollen-grains, size and form of	424	Propagulum	141	Ramal	148
Pollen-masses	421	Proper-vessels	38	Ramenta	57
Pollen-tubes	426	Prosenchyma	5, 29	Ramentaceous	57
—, extent to which they penetrate	504, 512	Proteaceæ	997	Ramplon	922
—, functions of	503	Proteins	310	Range of Vegetation, horizontal	1153
Pollen-utricles	421	Protococcus	1129	—, vertical	1153
Pollinia	421	Protoplasm	14	Ranunculaceæ	738
Pollinaria	477	Pruning trees	113	Rape	702
Polyadelphous	399	Prunus	855	Raphæ	467, 583
Polyandrous	394	Pseudo-bulb	70	Raphidean cells	18
Polycarpic	634	Pseudo-monocotyledonous	598	Raphides	18
Polychroite	1068	Pseudo-parasitic	1139	Raspberry	855
Polycotyledonous	598	Pseudo-spermous	531	Ratafia	855
Polyembryony	509, 556	Pteris aquilina	105, 1167	Rattans	1086
Polygalaceæ	772	Pterocarpus	850	Rattoons	304
Polygamous	359, 648	Pterophyllum	1197	Ravenala	1066
Polygonaceæ	990	Pubescent	60	Rays, medullary	89
Polyhedral	4	Puccoon	756	Receptacle of flower	320, 323
Polynomic	1151	Pucha-pat, see Patchouli.		Receptacles of secretion ..	20
Polypetalous	376, 378, 379	Pulp in fruits	523	Reclinato	185
Polypylous	358	Pulvulus	156	Rectembryæ	960
Polypodiæ	1109	Pumpkin	872	Red-snow plant	1129
Polysepalous	853	Punctated woody tissue ..	30	Reduplicate	354
Polyspermous	546	Punica	867	Regma	554
Polystemonous	392	Purging-nut	1017	Reign of Acrogens	1185
Pome	553	Purples	700	— Gymnosperms	1194
Pomeæ	854	Purslane	878	— Angiosperms	1198
Pomegranate	867	Putamen	524	Reindeer moss	1121
Pumpkin	872	Putty-wort	1060	Reniform	150
Pumpkinseed	796	Pyreneæ	553	Replam	447, 534
Pondweed family	1094	Pyrethrum	917	Reproduction in Cryptogamous plants	492
Fontederiaceæ	1082	Pyridium, same as Pome.		— in Phanerogamous plants	494
Poplar	1034	Pyroloæ	926	Reproductive organs	43
Poppy family	755	Pyrus	855	Resedaceæ	765
Porwort family	774	Pyxidium	554	Reservoirs of oil	20
Porous cells	10	Pyxis	554	Resinous matters	20
— dehiscence	536	Quadrangular	69	Resins	318
— vessels	86	Quadrifurcate	406	Respiration of plants	282
Portland-clirt-bed	1197	Quadrilocular	405, 440	Restiaceæ	1095
Portland sago	1093	Quadrilpartite	362	Reve of Nourse	678
Portugal Laurel	855	Quartine	464, 469	Reticulated cells	11
Portulacaceæ	878	Quassia	832	— vessels	35
Posterior, applied to parts of the flower	357	Quaternate	154	Reticulum	57, 153
Posticæ	412	Quercitron	1037	Refinacula	421
Potameæ	1094	Quercus	1037	Refractile hairs	59
Potash in plants	228	Quick-grass	1001	Retuse	150
Potash and soda as manures	246	Quillaire	854	Revolute	185
Potato	961	Quillata-bark	855	Rhabarberin	991
Potato-disease	691	Quina	903	Rhamnaceæ	840
Potentillæ	854	Quinate	154	Rhatany	773
Pounce	1046	Quince	855	Rheum	991
Præfloration	354	Quineuncial	355	Rhinanthideæ	964
Præfoliation	184	Quineunx	176	Rhizanthææ	1041
Præmorse	124	Quinoa	987	Rhizanthæ	252
Prangos	890	Quinquangular	69	Rhizobolaceæ	808
Pressing plants	1205, 1232	Quinquefid	143, 362	Rhizocarpeæ	1111
Prickles	57	Quinqueocular	440	Rhizogens	64
Prickly Pear	883	Quinquepartite	362	Rhizome	70
Primine	464	Quintine	464	Rhizophoraceæ	839
Primordial	168, 593	Raceme	353	Rhododendron	927
Primroses	975	Races	705	Rhododendrons, Region of ..	1162
Primulaceæ	974	Rachis	102, 325, 331	Rhubarb	991
Prince's feather	985	Radiate	911	Rhus	843
Prismatic	5	Radial	911	Rib, costa	144
Prismenchyma	5	Radial	5	Rib-wort family	978
Privet	941	Radical excretions	132	Rice	1117
Procumbent	66	Radicle	590, 592	Rice	1200
Products of plants	300	Radii	335	Rice paper	8, 552
Progression of sap, cause of ..	266	Radish	762	Rice paper plant	75
Proliferous leaves	159	Rafflesia	1042	Ricinus	1017
— plants	631	Rafflesiaceæ	1041	Rictus	381


	Paragraph		Paragraph		Paragraph
Riga-balsam	932	St. John's bread	851	Schouw's Phytogeographi-	
Ringent.....362, 381		St. John's wort family....	797	cal Regions.....1155	
Ripening of fruit	568	Salep.....1060		Scilla or Squilla	1078
Rocamboles.....1078		Salicine.....309, 1034		Scilleæ.....1077	
Roccella	1121	Salicineæ.....1033		Scillitina	1078
Rock plants	1139	Saline Plants	1139	Scimitar-shaped	150
Rock-Rose family.....	767	Salpiglossideæ	964	Selon	569, 573
Root	119	Salsafy	918	Scitamineæ.....1061	
—, abnormal.....121		Salvadora	937	Scleranthææ.....879	
—, of Acotyledons or		Salver-shaped	380	Sclerogen	15
Acrogens.....123		Salvia	967	Scorpioidæ.....340	
—, adventitious.....121		Samara	542	Scorzonera	918
—, aerial.....121		Samaroid	542	Scotch Fir	1046
—, of Dicotyledons or		Sambucææ.....900		Scotch Thistle	916
Exogens.....126		Samphire.....890		Screw-pine	121, 1001
—, excretions from....	132	Samydacææ.....1005		Scrophulariææ.....	964
—, fleshy.....131		Sand-plants.....1139		Scurf.....57	
—, forms of.....124		Sandal-wood	1007	Scurvy-grass	762
—, functions of.....129		Sandarach.....1046		Scythian Lamb.....1110	
—, mode of increase of	120	Sandbox-tree.....1016		Sea-buckthorn	999
—, subterranean.....120		Sanguisorbææ.....854		Sea-kale	762
—, structure of.....120		Sanalacææ.....1005		Sea-side grape	991
—, of Monocotyledons or		Sap, ascending.....257		Sea-pink	977
Endogens.....127		—, changes in its com-		Sea-weeds	1127
—, young.....119		position.....260		Sea-wrack	1123
Root-like processes of		—, course by which it		Sebesten plums.....957	
Ivy.....122		ascends.....257		Secretions of plants.....	300
Root-parasites.....252, 963		—, crude.....257		—, receptacles of....	20
Root-stock.....70		—, descending.....261		Secundine	464
Rosacææ.....854		—, elaborated.....261		Sedge Family	1097
Rosaceous corolla.....	378	—, height to which it		Sedum	880
Rose.....855		ascends.....274		Seeds.....575	
Rose-apple.....867		Sap-wood	78	—, forms of.....585	
Rosemary.....967		Sapindacææ.....806		—, functions of.....605	
Rose of Jericho.....762		Saponine.....779		—, integuments of....	577
Rosewood.....850		Sapotacææ.....936		—, lobes of.....586	
Rostellum.....421		Sapan-wood.....851		—, position in the ovary	585
Rotate.....380		Sappodilla.....937		—, sowing of.....611	
Rotation of crops.....	232	Sapucaya nut is the pro-		—, dissemination of....	606
Rotation in cells.....	275	duce of a species of Le-		—, maturation of.....	605
—, velocity of the		cythis.....1129		—, mode of preserving..	618
currents.....280		Sarcinula.....524, 545		—, mode of transmission	1240
—, cause of.....281		Sarcocarp.....1000		—, vitality of.....617	
Rough, see Scabrous.		Sarcoderm.....578		Selaginææ.....968	
Rounded, rotundatus....	150	Sarcosperm.....578		Selaginites.....1187	
Rubia.....905		Sargassum.....1129		Semi-anatropal.....	467
Rubiaceæ.....902		Sarmentum.....66		Semifloclous.....911	
Rue.....828		Sarraceniacææ.....754		Seminal leaves.....	599
Ruellia.....971		Sarsaparilla.....1054		—, lobes.....599	
Rumex.....991		Sarza.....1034		Seminude.....576	
Ruminated.....588		Sassafras.....994		Sempervivææ.....880	
Runcinate.....147		Satin-wood.....812		Sempervivum.....880	
Runner.....191		Sauracææ.....1030		Senega.....773	
Rupturing.....536		Savin.....1046		Senna.....851	
Rush Family.....1084		Savoury.....967		Sensitive plants.....	659, 852
Rust.....695		Savoy.....762		Sepals.....358	
Rutacææ.....827		Saxifragacææ.....835		—, adhesion of.....	362
Rye.....1142		Saxifrages, Region of...1155		—, forms of.....359	
Sabadilla.....1080		Scabrous.....58		Septate.....427	
Sac of the embryo.....464, 577		Scævoleæ.....920		Septemfid.....148	
Saccate.....373		Scalariform cells.....11		Septenate.....154	
Sack-tree.....1022		—, vessels.....35, 104		Septicidal.....532	
Safflower.....916		Scales.....57, 346		Septifragal.....532	
Saffron.....1068		Scammony.....936		Septulææ.....760	
Sagapenum.....891		—, adulterations of....	946	Septum.....408, 439	
Sage.....967		Scandent.....66		Sericous.....60	
Sagittate.....150, 374		Scape.....342		Serpentaria.....1010	
Sago.....302		Scar.....140		Serrate.....146	
—, false.....1048		Scarlet-runner.....850		Fessile.....62, 139, 325, 370	
—, Portland.....302, 1093		Scepacææ.....1014		Sesuvææ.....881	
Sago-palm.....1086		Schinus.....843		Setaceous.....57	
Saintfoin.....850		Schleiden on Cytogenesis	22	Setæ.....57, 459	
St. Ignatius's bean.....	947	Schultz, view of Latex...	41	Setose.....57	
				Sexes of plants.....	489

	Paragraph		Paragraph		Paragraph
Sexual system	715	Sour-gourd, same as Bao-		Staminidia, same as An-	
Shaddock	796	bab.		theridia.	
Shallon	927	Southern-wood	917	Stamiferous.....	398
Shallot	1078	Row-bread	975	Staminodlum	415, 651
Shamrock.....	821, 850	Spadix	334	Standard	375
Sheath of leaf.....	139	Spanish-broom, is Spartium		Stapella	943
She-oak	1035	Juncum.		Stapelia. Region of.....	1155
Shield-like	58	Spatha, spathe	349	Staphyliceæ.....	839
Shrubs	68, 713	Spathella	349	Star-anise	743
Sidæ	781	Spathulate	150	Star-apple	937
Sidesaddle-flower.....	754	Spawn	1122	Starch-grains.....	17
Sigillaria	1189	Spear-mint	967	Starch in plants	301
Sileneæ	778	Species, meaning of the		— changed to sugar 309, 477	
Silica in plants.....	226	term	705	Starch-sugar	307
— in epidermis.....	49	—, doubts as to their		Staticæ	976
— in the soil.....	235	limits.....	706	Statistics of vegetation	1151
Silicula	555	—, transmutation of.....	707	Stavesacre	740
Silqua	534, 555	Specimens for Herbarium.....	1223	Steeping seeds	248
Silk-cotton	784	Spermoderm	577, 578	Stellatæ	502
Siliqueosæ	760	Sphærenchyma.....	5	Stellate	5, 57, 380
Silver fir	1046	Sphæria	1126	Stem, forms of	69
Silver-grain	89	Sphenophyllum	1186, 1197	—, acrogenous or acoty-	
Simarubacæ.....	831	Sphenopteris	1186	—, ledonous	71
Sinapis	762	Spheroidal	5	—, aerial	70
Sinapisin	762	Spider-wort	1087	—, endogenous or mono-	
Sinuuous	5	Spigellææ	948	cotyledonous	71
Siritch	951	Spike	334	—, exogenous of dicoty-	
Sissoo	850	Spikelet	334	—, ledonous	71, 107
Skirret	890	Spikenard	907	—, functions of different	
Skunk-Cabbage	1093	Spike-oil	967	parts of.....	109
Sleep of plants	658	Spinage or Sphnach	987	—, internal structure of.....	71
Slipper-like	381	Spindle-tree	832	—, size of.....	108
Sloe	855	Spines	200	—, subterranean	70
Smilacææ	1053	Spiræææ	854	Steniless	65
Smilacin	1054	Spiral cells	11, 403	Sterculiaceæ	733
Smut	694	— vessels	32, 41	Sterigmata	1122
Smut-balls	693	Spiral cycles described by		Sterile flowers	649
Snake-nut-tree	807	leaves.....	171	Sternbergia	1190
Snake-root	773	Spiral veneration	186	Stigma	429, 433
—, Virginian.....	1010	Sprouidea	32	Stigmæria	1188
Snake-wood	947, 1022	Sporobolæ	759	Stilaginacææ	1025
Snowball-tree is Viburnum		Sporoglobo	120	Stings	62
Opulus.		Spongloles	120	Stipe	65, 102
Snowberry	901	—, absorption by	253	Stipels	161
Snow-drop, see Galanthus.		Sporangia	459	Stipitate.....	62, 335, 437
Snow-drop-tree.....	932	Spore	459, 591	Stipulate	154
Snowflake is Leucocum.		Sporidium	591	Stipules	139, 158
Snow-line	1137	Sporocarp	1127	Stock	569
Soap-berries	807	Sporophores	1122	Stomata	50
Soap-wort family	806	Sporule, see Spore.		—, development of.....	55
Stoboles	70	Spruces	1046	—, forms of.....	54
Social plants	1144	Spur, calcar	373	—, functions of	64
Soda in plants	223	Spurge family	1014	—, in Equisetum.....	1107
Soils	233	Spurge-laurel	1002	—, in leaves	136
—, as affecting the distri-		Spurious tracheæ	35	—, number in square	
—, bution of plants.....	1139	Spurred	363	—, inch of surface	56
—, required for different		Squamæ	346	Stoncrop	880
plants	239	Squamash	1078	Stone-pine is Pinus Pineæ.	
Solanacææ	960	Squamulæ	1099	Storax	932
Solanin	961	Square cells	5	—, liquid	1095
Solidagos. Region of	1155	Squash	872	Stramonium	961
Solomon's-seal is Conval-		Squill	1078	Strapshaped	382
laria.		Squitting cucumber	872	S-tratiotæ	1037
Solubility	536	Stachouseacææ.....	836	Strawberry.....	855
Sooranjee	905	Stalk of leaf	139	Strawberry-tree	927
Soot as a manure	246	Stamens	350, 390	Strobilites	1187
Soredia, gonidia breaking		—, definite	394	Strobilus	334, 559
through the thallus of		—, indefinite	34	Strophilote	582
Lichens	1119	—, number of.....	399	Strophioles	582
Sori	1108	—, parts of	395	Stroma	156
Sorosis	557	—, relation to the petals 393		Strumose	398
Sorci	991	—, structure and form.....	391	Stryclineæ	919
Souari-nuts.....	808	—, in Cryptogamic plants 427		Stryclinia	947
Sour-sop	745	Staminal	649	Stryclinos	947

	Paragraph		Paragraph		Paragraph
Stupose	60, 398	Synantherous	417	Thistle	916
Style	429, 432	Synaptase	310	Thorn	200
—, forms of	453	Syncarpous	493, 538, 549	Thorn-apple	961
—, situation of	451	Syngenesia	911	Thrift	977
Style and stigma, irrita-		Syngenesions	417	Throat of corolla	377
bility of	665	Synochreate	160	Thunbergiæ	970
Style-wort family	921	Synonymes	713	Thus	1046
Stylidaceæ	921	Syringa	865	Thuya	1046
Stylopodium	883	Systems of classification ..	703	Thuyites	1177
Styracaceæ	929	—, artificial	715	Thyme	967
Subclasses defined	931	—, natural	719	Thymelæaceæ	1003
Suberic acid	710			Thyrus	333
Suberous layer	816	Tabasheer	226	Tienté, see Upas.	
Subgenus, defined	85	Tabular cells	5	Tigellary merithal	139
Suborders, defined	708	Tacamahac	799, 845, 1034	Tigelle	590, 593
Subterranean plants	709	Tagna, a name for the Ivory		Tiliaceæ	787
— stem	1139	Palm		Tiliæ	737
Subulate	70	Tallow-tree	799, 1017	Tillandsia	1076
Suckers in Dodder	150, 396	Tamaricaceæ	775	Tissue, cellular	4
Succory	122	Tamarind	851	—, fibro-cellular	11
Suffrutex	918	Tamarisk family	775	—, fibro-vascular	21
Suffruticose	68	Tanghin	945	—, ligneous	29
Sugar	68	Tangle	1130	—, phleboïdal	37
—, from Beet	304	Tansy	917	—, pitted	36
—, import of	306	Taphrenchyma	36	—, vascular	28
—, from the maple	305, 1201	Tapioca	302, 1018	—, vasiform	36
Sugar-bush	306	Tap-root	124	—, woody	30
Sugar-cane	998	Taraxacine	918	Tissues, vegetable	1
Sugar-maple	1101	Taraxacum	918	—, arrangement of	42
Sulphur as a manure	805	Tarragon	917	Tobacco	961
Sulphuretted hydrogen,		Tartarian Lamb	1110	Tomato	961
effect on plants	246	Taxineæ	1045	Tomentose	60
Sulphurous acid gas, effect		Taxonomy	703	Tomentum	60
on plants	294	Tea plants	793	Tonka-bean	850
Sumach	843	— of Paraguay	935	Toothache-tree	895
Sumatra-Camphor family ..	789	Teak, African	1018	Toothwort	977
Sun-dew family	770	—, Indian	969	Tormentil	855
Sunflower	770	Teazel	908	Tortoise plant	1052
Sun- Hemp	917	Teel seeds	951	Torns	329
Superior, applied to the		Tegmenta	182	Tous-les-mois	302
parts of the flower	782	Temperature of plants	667	Toxicodendron	843
Supervolute	257	—, effects of	1135	Trachææ	31
Suppression of organs	185	—, as regards altitude	1137	—, closed	34
Supradecomound	647	—, — latitude	1136	—, false	35
Suspended ovule	152	Tendrils	157, 201	—, spurious	25
Suspensor	471	Teratology	646	Trachenchyma	31
Sutural	463, 504, 514	Tercine	464, 669	Tradescantia	1087
Sutres	532	Terminaliæ	862	—, rotation in	278
Swallow's-nest, edible	529	Terminology or Termono-		Tragacanth	850
Sweet-almond	1130	logy	704	Transpiration	203
Sweet-bay	154	Ternate	154	Transportation of plants in	
Sweet-briar is Rosa rubi-		Ternstroemiaceæ	792	close cases	298
ginosa.	994	Tertiary-fossils	1202	Traveller's joy is Clematis	
Sweet-cane	1102	Test-objects	1210	Vitalba.	
Sweet-cassava	1018	Testa	578	Tree-beard	1076
Sweet-fern	1035	Tetradymanous	419	Tree-ferns, Region of	1162
Sweet-flag	1033	Tetragonal, symmetry	643	—, structure of stems	102
Sweet-pea is Lathyrus		Tetragoniceæ	881	—, stem, dichotomous	105
odoratus.		Tetramerous	643	Tree-lily	1071
Sweet-potato	956	Tetrandrous	394	Tree-nettle	1020
Sweet-sop	745	Tetrapetalous	375	Trees, defined	68
Sweet-william is Dianthus		Tetraspore	591	—, ages of	637
barbatus.	363	Tetrathecal	405	Trefoil	850
Sword-shaped, see Ensiform.		Thalamus	329, 350	Tremandraceæ	774
Sycamine of Scripture	1021	Thalamifloræ	390, 737	Triadelphous	399
Sycamore	805	Thallogeniæ	1118	Triandrous	394
— of Scripture	1021	Thallophens	65, 106	Triangular	69
Syconus	558	Thallophyta	1104	Tribes, defined	709
Symbols	713	Thallophytes	65	Tribuleæ	825
Symmetrical flower	713	Thallus	65	Trichotomous	338, 339
Symmetry of organs	375	Thecæ	459	Tricoccus	533
Symplocææ	643	Thecaphore	437	Trientalis	975
Synantheræ	931	Theine	793, 935	Trifid	148, 362
	910	Theobromine	786	Trigonal symmetry	643

- | | Paragraph | | Paragraph | | Paragraph |
|------------------------|---------------|---------------------------|-----------|------------------------------|-----------|
| Trigonocarpum..... | 1190 | Univalvular..... | 530 | Veratres..... | 1079 |
| Trigonous..... | 69 | Unilining..... | 383 | Veratria..... | 1080 |
| Trilocular..... | 440 | Unsymmetrical..... | 375 | Verbascum..... | 965 |
| Trilliacæ..... | 1055 | Upas Antlar..... | 1022 | Verbenacæ..... | 968 |
| Trimerous..... | 643 | Upas Tieuté..... | 947 | Verjuice..... | 814 |
| Trinomic..... | 1151 | Urceolate..... | 380 | Vermiform vessels..... | 37 |
| Tripartite..... | 362 | Urginea..... | 1078 | Vernal grass..... | 1100 |
| Tripe de Roche..... | 1121 | Urn-shaped..... | 380 | Vernation..... | 184 |
| Tripetalous..... | 375 | Urticacæ..... | 1019 | Vernica..... | 965 |
| Triplinnae..... | 152 | Utricle..... | 4, 542 | Verruca, warts..... | 63 |
| Triplinnafid..... | 147 | ——, internal..... | 14 | Versatile..... | 408 |
| Triplet..... | 1205 | ——, primary..... | 504 | Verticil..... | 169 |
| Triquetrous..... | 69 | ——, primordial..... | 14 | Verticillaster..... | 339 |
| Trisepalous..... | 361 | Utricularia..... | 973 | Verticillate..... | 169 |
| Tristichous..... | 170 | Uva..... | 550 | Verticils or whorls of the | |
| Triternate..... | 154 | Uvularicæ..... | 1079 | flower..... | 250 |
| Trivial name..... | 708 | | | Vervain..... | 969 |
| Tropeolacæ..... | 822 | Vacciniacæ..... | 923 | Vesicle, germinal..... | 510 |
| Trophosperm..... | 462 | Vagina Merithal..... | 139, 158 | Vesicles or bladders..... | 3 |
| Truffe..... | 1124 | Vaginula..... | 1114 | Vesicular calyx..... | 367 |
| Trumpet-wood..... | 1022 | Valerian..... | 907 | ——, glands..... | 63 |
| Truncate..... | 150, 362 | Valerian, Greek..... | 953 | Vessels, annular..... | 35 |
| Truncus..... | 65 | Valerianacæ..... | 906 | ——, beaded..... | 37 |
| Tryma..... | 545 | Valleculæ..... | 888 | ——, development of..... | 39 |
| Tuber..... | 70, 193 | Vallisneria..... | 1057 | ——, dotted..... | 35 |
| Tubercular..... | 125 | ——, rotation in its | | ——, laticiferous..... | 38 |
| Tubercules..... | 131 | cells..... | 277 | ——, mode of com- | |
| Tuberosæ..... | 1078 | Vallisneriæ..... | 1057 | munication..... | 40 |
| Tubes..... | 3 | Valonia..... | 1037 | ——, moniliform..... | 37 |
| Tubular..... | 380 | Valvate..... | 186, 354 | ——, necklace-like..... | 37 |
| Tubulifloræ..... | 913, 914 | Vanille..... | 1060 | ——, porous..... | 36 |
| Tulipeæ..... | 1077 | Variation..... | 676 | ——, reticulated..... | 35 |
| Tulip tree..... | 743 | Varieties, defined..... | 705 | ——, scalariform..... | 35 |
| Tunicated bulb..... | 194 | Varnishes..... | 843 | ——, vermiform..... | 37 |
| Turbinate..... | 362 | Vascular bundles in Acro- | | Vexillary..... | 355 |
| Turio..... | 193 | gens..... | 103 | Vexillum..... | 379 |
| Turmeric..... | 1062 | ——, definite..... | 91 | Victoria..... | 751 |
| Turneracæ..... | 877 | ——, in Endogens..... | 91 | Villi..... | 57 |
| Turnip..... | 762 | ——, in Exogens..... | 77 | Villous..... | 60 |
| Turnsole..... | 1018 | ——, functions of..... | 109 | Vine..... | 814 |
| Turpentine..... | 1046 | ——, indefinite..... | 77 | Violacæ..... | 763 |
| , Chilan..... | 843 | ——, simultane- | | Violet family..... | 763 |
| Tussac-grass..... | 1100 | ous..... | 103 | Virginian Snake-root..... | 1010 |
| Tussilago..... | 917 | Vascular tissue..... | 28 | Vis a fronte in the circula- | |
| Tutsan family..... | 797 | Vasculum..... | 1229 | tion..... | 269 |
| Twisted vernation..... | 186, 354 | Vasiform tissue..... | 36 | Vis a tergo in the circula- | |
| Typhineæ..... | 1092 | Vegetable Ivory..... | 1086 | tion..... | 269 |
| | | Vegetable marrow..... | 872 | Viscum..... | 899 |
| Ulmacæ..... | 1019 | Vegetation, alpine..... | 1161 | Vitacæ..... | 813 |
| Ulmine..... | 234 | ——, altitudinal zones | | Vitellus..... | 580 |
| Ulodendron..... | 1187 | of Meyen..... | 1162 | Vittæ..... | 20 |
| Ulva..... | 1130 | ——, horizontal range | | Vivianlacæ..... | 780 |
| Umari..... | 850 | of..... | 1153 | Viviparous plants..... | 631 |
| Umbel..... | 335 | ——, vertical range..... | 1158 | Vochysiaceæ..... | 861 |
| Umbelliferæ..... | 838 | Vell..... | 1122 | Volzia..... | 1197 |
| , Region of..... | 1155 | Veinless..... | 141 | Voluble..... | 66 |
| Umbellule..... | 335 | Veinlets..... | 141 | Volva..... | 122 |
| Umbilical cord..... | 462 | Veins, costal..... | 141 | | |
| Umbilicus..... | 583 | ——, primary..... | 141 | Wake-Robin..... | 1093 |
| Uncaria..... | 905 | Vellozia..... | 1071 | Wall-flower..... | 762 |
| Uncinate..... | 58 | ——, stem of..... | 112 | Walnut..... | 1039 |
| Undershrub..... | 68 | Velum..... | 1122 | Ward's portable conserva- | |
| Undulated..... | 150 | Velutinus..... | 60 | tories..... | 297 |
| Unequal..... | 146 | Velvety..... | 60 | Warty glands..... | 63 |
| Unequally pinnate..... | 153 | Venation..... | 141 | Water-bean family..... | 752 |
| Unguiculate..... | 370 | ——, hidden..... | 141 | Water-chestnut..... | 868 |
| Unguis..... | 370 | ——, furcate..... | 144 | Water-cress is Nasturtium | |
| Unicorn plant..... | 951 | ——, netted..... | 141 | officinale..... | |
| Unicostate..... | 144 | ——, parallel..... | 142 | Water-fennel..... | 1129 |
| Unifugate..... | 153 | ——, radiating..... | 148 | Water-hemlock..... | 893 |
| Unilateral..... | 466 | ——, reticulated..... | 141 | Water-lily..... | 750 |
| Unilocular..... | 405, 440, 522 | ——, of petals..... | 371 | Water-melon..... | 872 |
| Unipetalous..... | 375 | Ventral suture..... | 438 | Water-net..... | 1129 |
| Unisexual..... | 388, 618 | Venus' fly-trap..... | 771 | Water-pepper family..... | 777 |

	Paragraph		Paragraph		Paragraph
Water-plantain	1088	Witch-hazel family	887	Yam	1052
Water-shield family	749	Wood, formation of	111	Yew	1046
Waved	5	Woodbine, same as Honey-		Yucca	1078
Wavy	150, 374	suckle.			
Wax, vegetable	316	Woodruff	905	Zabucajo nuts are the pro-	
Wax-flower	943	Wood-sorrel	821	duce of <i>Lecythis</i> Zabu-	
Wax-myrtle	1035	Woody-fibre	29	cajo.	
Wax-palm	1086	— in the bark ..	83	Zamia	1048
Waxy matter in epider-		Woody layers	77	Zamiostrobus	1197
mis.	49	Woody masses, anomalous	113	Zamites	1197
Way-bred	979	Woody stems, peculiar Bra-		Zanthoxylaceæ	829
Wealden fossils	1197	zilian	90	Zebra-plant	1064
Wedge-shaped	150	Woody tissue, glandular ..	30	Zebra-wood	846
Weeping Willow	1034	—, punctated ..	30	Zingiberaceæ	1061
Wetherellia	1202	Woody wedges in pepper ..	90	—, Region of ..	1155
Wheat	1142	Woolly	60	Zizyphus	841
White wood	78	Woorali, or Wourali	947	Zones of wood	77
Whorl	169	Wormseed	917	— in Exogens ..	79
Whorled	350	Wormwood	917	Zones of vegetation	1156
Whortleberry	928	Wych-Elm is <i>Ulmus</i> mon-		Zoocarps, same as Zoo-	
Willow	1034	tana.		spores.	
Winged fruit	992	Xanthic series of colours ..	679	Zoosperms, same as Zoo-	
Wings	379	Xanthophylle	675	spores.	
Wintereæ	742	Xanthorrhæa	1078	Zoospores	224
Winter-cherry	961	Xanthoxylaceæ	829	Zygophyllaceæ	825
Wintergreen	926	Xyridaceæ	1083		
Winter's bark	743				



Cabinet Edition of the Encyclopædia Metropolitana—continued.

Vol. XXV., price 10s. 6d., cloth lettered,
HISTORY OF THE ROMAN EMPIRE. By the Rev. THOS. ARNOLD, D
 late Head Master, Rugby School, and Others. With Numerous Illustrations.

Vol. XXVI., price 10s. 6d., cloth lettered,
HISTORY OF THE DECLINE AND FALL OF THE ROMAN POW
 By the Right Rev. Bishop RUSSEL, &c. With Numerous Illustrations.

Vol. XXVII., price 6s., cloth lettered,
HISTORY OF GREEK AND ROMAN PHILOSOPHY AND SCIENCE.
 the Right Rev. CHARLES BLOMFELD, D.D., Bishop of London; Rev. Dr. WHEWELL,
 other Contributors.

* A more interesting and instructive volume than this has not been offered to the public for a long time
 exists.

Vol. XXVIII., price 3s. 6d., cloth lettered,
MORAL AND METAPHYSICAL PHILOSOPHY—THE PHILOSOPHY
THE FIRST SIX CENTURIES. By the Rev. F. D. MAURICE, M.A.

Vol. XXIX., price 7s. 6d., cloth lettered,
HISTORY OF THE OTTOMAN EMPIRE; including a Survey of the GR
EMPIRE and the CRUSADES. By Colonel PROCTER, Rev. J. E. RIDDLE, and JAMES M^C
 ECHT, Esq.

—The best work on Turkey with which we are acquainted."—*Courier*.

Vol. XXX., price 2s. 6d., cloth lettered,
TRIGONOMETRY. By GEORGE B. AIRY, F.R.S., Astronomer Royal. Ed
 by HUGH BLACKBURN, M.A., Professor of Mathematics in the University of Glasgow.

Vol. XXXI., price 6s., cloth lettered,
COMPENDIUM OF THE OCCULT SCIENCES: Containing the Traditi
 and Superstitions of past times, and the Marvels of the present day, in a series of Histor
 Sketches, by the Rev. EDWARD SMEDLEY, M.A., E. RICH, Esq., and other Contributors.

"A repository of curious information. . . most instructive and interesting reading."—*Literary Ga*
 —Full of real learning."—*Press*.

Vol. XXXII., price 12s. 6d., cloth lettered,
GEOLOGY—PRACTICAL AND THEORETICAL. By JOHN PHILLIPS, M.A., F.R
 F.G.S., Deputy Reader in Geology in the University of Oxford. With Numerous Illustrati
 "A most admirable text-book."—*Westminster Review*.

Vol. XXXIII., price 5s., cloth lettered,
HISTORY OF THE CHRISTIAN CHURCH, from the Fourth to the Twe
Century. By Rev. J. M. GULDING, Rev. J. B. CARWITHEX, and Others.

Vol. XXXIV., price 5s., cloth lettered,
CHRONOLOGICAL TABLES; FIRST DIVISION. From the Earliest Recd
 to the End of the Fifteenth Century. Edited by ISAIAH M'BURNET, B.A., F.S.A.

Vol. XXXV., price 5s., cloth lettered,
CHRONOLOGICAL TABLES; SECOND DIVISION. From the Beginning
 the Sixteenth Century to the present day. Edited by SAMUEL NEIL, Esq.

Vol. XXXVI., price 5s., cloth lettered,
MORAL AND METAPHYSICAL PHILOSOPHY. MEDLEVAL PHILOS
PHY. By the Rev. F. D. MAURICE, M.A.

Vol. XXXVII., price 7s. 6d., cloth lettered,
HISTORY OF THE REFORMATION, AND ITS CONSEQUENCES.
 the BISHOP OF HEREFORD; the Rev. J. E. RIDDLE, and Others.

Volumes in Active Preparation.

SCULPTURE. By RICHARD WESTMACOTT, F.R.S., A.R.A., with Numer
 Illustrations, revised.

GLOSSOLOGY; or, The Historical Relations of Languages. By SIR JO
 STODDART, LL.D.

MORAL AND METAPHYSICAL PHILOSOPHY—MODERN PHILOSOPH
 By Rev. F. D. MAURICE, M.A., Chaplain of Lincoln's Inn.

ARITHMETIC. By Rev. DR. PEACOCK, Dean of Ely. Revised.

APPLIED MECHANICS. By PROFESSOR J. MACQUORN RANKINE.

PRACTICE OF MEDICINE. By DRs. WILLIAMS AND AITKEN.

Valuable Works of Reference.

BOOK OF ANCIENT HISTORY, from the CREATION of the World to the
of the WESTERN EMPIRE, by SIR JOHN ECCLES, Bart. 8vo. 18s. cloth.
Plans of Cities, &c.

BOOK OF MODERN HISTORY, from the END of the
Dissertations on the
s. cloth.

BOOK OF MODERN HISTORY, from the END of the
Dissertations on the
s. cloth.

BRITISH EMPIRE, by
CAL, by name
y. Map and plan

PÆDIA OF UNIVERSAL HISTORY, by
taucous Events
ed Chronologie
AMUEL NEIL. 8vo.

PÆDIA OF MODERN HISTORY, by
Remarkable Incidents
d by numerous

PÆDIA OF MODERN HISTORY, by
TEER OF THE
Post 8vo, 18s.

ALCYCLOPÆDIA OF MODERN HISTORY, by
AL History,
ATURE, illustr
LL.D. With
s. Sixth Edition

TICAL CO-OPERATION, by
presented by
D.D., LL.D.

PÆDIA OF MODERN HISTORY, by
ations of the
s THOMSON
l. cloth.

PÆDIA OF MODERN HISTORY, by
Dynamics, Electricity, &c.

Dynamics, Electricity, &c.
matics, Meteorology, Optics, &c.
With numerous Illustrations. One large volume, 8vo, 18s. cloth.

XII.
PÆDIA OF THE NATURAL SCIENCES, comprising Botany, Conchology,
ology, Geology, Mineralogy, Palæontology, and Zoology. By WM. BAIRD,
F.L.S., British Museum. With numerous Illustrations. 8vo,—in preparation.

XIII.
ULARY OF PHILOSOPHY, MORAL, MENTAL, and METAPHYSICAL, with
ous References and Quotations. By WILLIAM FLEMING, D.D., Professor of
Philosophy in the University of Glasgow. Foolscap 8vo, 7s. 6d. cloth.

University of Toronto Library

DO NOT
REMOVE
THE
CARD
FROM
THIS
POCKET

Acme Library Card Pocket
LOWE-MARTIN CO. LIMITED

70061

Balfour, John Hutton
Manual of botany

DATE

Bot
B

THE UNIVERSITY OF
CHICAGO PRESS



THE UNIVERSITY OF
CHICAGO PRESS
Applied Sciences
116 (56)